

MAT1830

Lecture 13: Functions

Functions - why should you care?

Functions - why should you care?


The concept of a function is extremely important in both computer science and maths.


Functions - why should you care?




The concept of a function is extremely important in both computer science and maths.




- ▶ Functions (subroutines) in programming are closely related to functions in the mathematical sense.
- ▶ In the case of functional programming languages (eg. Lisp, Haskell, Rust) they are exactly functions in the mathematical sense.
- ▶ Functions are used to define a lot of important concepts in maths and theoretical computer science.

A function can be thought of as a “black box” which accepts inputs and, for each input, produces a single output.













Inputs:




.....
.....




Allowed
outputs:

.....
.....


















- Each input produces exactly one output.
(Always the same output for a given input.)






- Each input produces exactly one output.
(Always the same output for a given input.)






- Each input produces exactly one output.
(Always the same output for a given input.)






- Each input produces exactly one output.
(Always the same output for a given input.)






- Each input produces exactly one output.
(Always the same output for a given input.)






- Each input produces exactly one output.
(Always the same output for a given input.)




- Each input produces exactly one output.
(Always the same output for a given input.)






- Each input produces exactly one output.
(Always the same output for a given input.)




- Each input produces exactly one output.
(Always the same output for a given input.)






- Each input produces exactly one output.
(Always the same output for a given input.)



- Each input produces exactly one output.
(Always the same output for a given input.)



- Each input produces exactly one output.
(Always the same output for a given input.)




- Each input produces exactly one output.
(Always the same output for a given input.)

Domain:

$$X = \{\dots\dots\}$$

Codomain:

$$Y = \{\dots\dots\}$$



- Each input produces exactly one output.
(Always the same output for a given input.)

Domain:

$$X = \{\dots\dots\}$$

Codomain:

$$Y = \{\dots\dots\}$$

input	output
x_1	y_1
x_2	y_2
x_3	y_3
\vdots	\vdots

- Each input produces exactly one output.
(Always the same output for a given input.)

Domain:

$$X = \{\dots\dots\}$$

Codomain:

$$Y = \{\dots\dots\}$$

$$\{(x_1, y_1), (x_2, y_2), \\ (x_3, y_3), \dots\dots\dots\}$$

- Each input produces exactly one output.
(Always the same output for a given input.)

Domain:

$$X = \{\dots\dots\}$$


Codomain:

$$Y = \{\dots\dots\}$$

$$\{(x_1, y_1), (x_2, y_2), \\ (x_3, y_3), \dots\dots\dots\}$$

A set of ordered pairs from $X \times Y$ that contains exactly one ordered pair (x, y) for each $x \in X$.

Remember: The domain and codomain are part of the function and must always be defined.



13.1 Defining functions via sets

Formally we represent a function f as a set X of possible inputs, a set Y so that every output of f is guaranteed to be in Y , and a set of (input,output) pairs from $X \times Y$. The vital property of a function is that each input gives exactly one output.

A function f consists of a *domain* X , a *codomain* Y , and a set of ordered pairs from $X \times Y$ which has exactly one ordered pair (x, y) for each $x \in X$.

When (a, b) is in this set we write $f(a) = b$.

The set of y values occurring in these pairs is the *range* of f .

Note that the range of a function is always a subset of its codomain but they may or may not be equal.

If the range of a function is equal to its codomain, we say the function is *onto*.

Formally, a function consists of a domain X , a codomain Y , and a set of ordered pairs from $X \times Y$ which has exactly one ordered pair (x, y) for each $x \in X$.

Formally, a function consists of a domain X , a codomain Y , and a set of ordered pairs from $X \times Y$ which has exactly one ordered pair (x, y) for each $x \in X$.

The set of y values occurring in these ordered pairs is called the *range* of the function.

Formally, a function consists of a domain X , a codomain Y , and a set of ordered pairs from $X \times Y$ which has exactly one ordered pair (x, y) for each $x \in X$.

The set of y values occurring in these ordered pairs is called the *range* of the function.

The range is always a subset of the codomain but they may not be equal. If they are equal we say the function is *onto*.

Formally, a function consists of a domain X , a codomain Y , and a set of ordered pairs from $X \times Y$ which has exactly one ordered pair (x, y) for each $x \in X$.

The set of y values occurring in these ordered pairs is called the *range* of the function.

The range is always a subset of the codomain but they may not be equal. If they are equal we say the function is *onto*.

“ f is a function with domain X and codomain Y ” is shortened to

$$f : X \rightarrow Y.$$

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.

x	$f(x)$
0	0
1	2
2	4
3	6

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.

x	$f(x)$
0	0
1	2
2	4
3	6

$\{(0, 0), (1, 2), (2, 4), (3, 6)\}$

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.

x	$f(x)$
0	0
1	2
2	4
3	6

$\{(0, 0), (1, 2), (2, 4), (3, 6)\}$

Example Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = 2x$.

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.

x	$f(x)$
0	0
1	2
2	4
3	6

$\{(0, 0), (1, 2), (2, 4), (3, 6)\}$

Example Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = 2x$.


$\{(x, 2x) : x \in \mathbb{R}\}$

Arrow diagrams

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.


Arrow diagrams

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.




Arrow diagrams

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.



Arrow diagrams


Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.



The range of f is $\{0, 2, 4, 6\}$.

Arrow diagrams

Example Let $f : \{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 4, 5, 6\}$ be defined by $f(x) = 2x$.



The range of f is $\{0, 2, 4, 6\}$. (So f is not onto.)

Why don't we always set the codomain equal to the range?

Why don't we always set the codomain equal to the range?

Think about $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = x^8 + 102x^7 - 7x^5 + 20x^4 - 100x + 7.$$

Why don't we always set the codomain equal to the range?

Think about $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = x^8 + 102x^7 - 7x^5 + 20x^4 - 100x + 7.$$

We've set the codomain to \mathbb{R} and that's fine - certainly $f(x) \in \mathbb{R}$.

Why don't we always set the codomain equal to the range?

Think about $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = x^8 + 102x^7 - 7x^5 + 20x^4 - 100x + 7.$$

We've set the codomain to \mathbb{R} and that's fine - certainly $f(x) \in \mathbb{R}$.

What is range of f ?

Why don't we always set the codomain equal to the range?

Think about $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = x^8 + 102x^7 - 7x^5 + 20x^4 - 100x + 7.$$

We've set the codomain to \mathbb{R} and that's fine - certainly $f(x) \in \mathbb{R}$.

What is range of f ? Hard to find and probably ugly.

Why don't we always set the codomain equal to the range?

Think about $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = x^8 + 102x^7 - 7x^5 + 20x^4 - 100x + 7.$$

We've set the codomain to \mathbb{R} and that's fine - certainly $f(x) \in \mathbb{R}$.

What is range of f ? Hard to find and probably ugly.

Another reason is that " $\mathbb{R} \rightarrow \mathbb{R}$ functions", for example, make a nice class to consider.

Question What set of ordered pairs does $f : \{0, 1, 2, 3\} \rightarrow \mathbb{N}$ defined by $f(x) = x^2$ correspond to?

Question What set of ordered pairs does $f : \{0, 1, 2, 3\} \rightarrow \mathbb{N}$ defined by $f(x) = x^2$ correspond to?

$\{(0, 0), (1, 1), (2, 4), (3, 9)\}$.

Question What set of ordered pairs does $f : \{0, 1, 2, 3\} \rightarrow \mathbb{N}$ defined by $f(x) = x^2$ correspond to?

$\{(0, 0), (1, 1), (2, 4), (3, 9)\}$.

Question Does $\{(0, 7), (2, \pi)\}$ correspond to a function $f : \{0, 1, 2\} \rightarrow \mathbb{R}$?

Question What set of ordered pairs does $f : \{0, 1, 2, 3\} \rightarrow \mathbb{N}$ defined by $f(x) = x^2$ correspond to?

$\{(0, 0), (1, 1), (2, 4), (3, 9)\}$.

Question Does $\{(0, 7), (2, \pi)\}$ correspond to a function $f : \{0, 1, 2\} \rightarrow \mathbb{R}$?

No - it doesn't have an ordered pair with first coordinate 1.

Question What set of ordered pairs does $f : \{0, 1, 2, 3\} \rightarrow \mathbb{N}$ defined by $f(x) = x^2$ correspond to?

$\{(0, 0), (1, 1), (2, 4), (3, 9)\}$.

Question Does $\{(0, 7), (2, \pi)\}$ correspond to a function $f : \{0, 1, 2\} \rightarrow \mathbb{R}$?

No - it doesn't have an ordered pair with first coordinate 1.

Question Does $\{(0, 7), (1, 4), (2, \pi), (2, 3)\}$ correspond to a function $f : \{0, 1, 2\} \rightarrow \mathbb{R}$?

Question What set of ordered pairs does $f : \{0, 1, 2, 3\} \rightarrow \mathbb{N}$ defined by $f(x) = x^2$ correspond to?

$\{(0, 0), (1, 1), (2, 4), (3, 9)\}$.

Question Does $\{(0, 7), (2, \pi)\}$ correspond to a function $f : \{0, 1, 2\} \rightarrow \mathbb{R}$?

No - it doesn't have an ordered pair with first coordinate 1.

Question Does $\{(0, 7), (1, 4), (2, \pi), (2, 3)\}$ correspond to a function $f : \{0, 1, 2\} \rightarrow \mathbb{R}$?

No - it has two ordered pairs with first coordinate 2.


square : $\mathbb{R} \rightarrow \mathbb{R}$

Examples.

1. The squaring function $\text{square}(x) = x^2$ with domain \mathbb{R} , codomain \mathbb{R} , and pairs

$$\{(x, x^2) : x \in \mathbb{R}\},$$

which form what we usually call the *plot* of the squaring function.




The range of this function (the set of y values) is the set $\mathbb{R}^{\geq 0}$ of real numbers ≥ 0 .


$$\text{sqrt} : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}$$

2. The square root function $\text{sqrt}(x) = \sqrt{x}$ with domain $\mathbb{R}^{\geq 0}$, codomain \mathbb{R} , and pairs

$$\{(x, \sqrt{x}) : x \in \mathbb{R} \text{ and } x \geq 0\}.$$




The range of this function (the set of y values) is the set $\mathbb{R}^{\geq 0}$.



$\text{cube} : \mathbb{R} \rightarrow \mathbb{R}$

3. The cubing function $\text{cube}(x) = x^3$ with domain \mathbb{R} , codomain \mathbb{R} , and pairs

$$\{(x, x^3) : x \in \mathbb{R}\},$$



The range of this function is the whole of the codomain \mathbb{R} , so it is onto.

Question 13.1 Which of the following rules define functions?

Question 13.1 Which of the following rules define functions?

- ▶ For each non-empty set S of natural numbers, let $f(S)$ be the least member of S .

Question 13.1 Which of the following rules define functions?

- For each non-empty set S of natural numbers, let $f(S)$ be the least member of S .

Yes.

Question 13.1 Which of the following rules define functions?

- ▶ For each non-empty set S of natural numbers, let $f(S)$ be the least member of S .

Yes.

- ▶ For each set X of real numbers between 0 and 1, let $g(X)$ be the least member of X .

Question 13.1 Which of the following rules define functions?

- ▶ For each non-empty set S of natural numbers, let $f(S)$ be the least member of S .

Yes.

- ▶ For each set X of real numbers between 0 and 1, let $g(X)$ be the least member of X .

No - $g(\{x : x \in \mathbb{R} \text{ and } \frac{1}{2} < x < 1\})$ is not defined.

Question 13.1 Which of the following rules define functions?

- ▶ For each non-empty set S of natural numbers, let $f(S)$ be the least member of S .

Yes.

- ▶ For each set X of real numbers between 0 and 1, let $g(X)$ be the least member of X .

No - $g(\{x : x \in \mathbb{R} \text{ and } \frac{1}{2} < x < 1\})$ is not defined.

- ▶ For each circle C in the (x, y) plane, let $h(C)$ be the minimum distance from C to the x axis.

Question 13.1 Which of the following rules define functions?

- ▶ For each non-empty set S of natural numbers, let $f(S)$ be the least member of S .

Yes.

- ▶ For each set X of real numbers between 0 and 1, let $g(X)$ be the least member of X .

No - $g(\{x : x \in \mathbb{R} \text{ and } \frac{1}{2} < x < 1\})$ is not defined.

- ▶ For each circle C in the (x, y) plane, let $h(C)$ be the minimum distance from C to the x axis.

Yes.

Question 13.1 (cont.) Which of the following rules define functions?

Question 13.1 (cont.) Which of the following rules define functions?

- ▶ For a pair A, B of sets of real numbers let $s(A, B)$ be the smallest set which has both A and B as subsets.

Question 13.1 (cont.) Which of the following rules define functions?

- ▶ For a pair A, B of sets of real numbers let $s(A, B)$ be the smallest set which has both A and B as subsets.

Yes (depending on your interpretation of “smallest”).

Question 13.1 (cont.) Which of the following rules define functions?

- For a pair A, B of sets of real numbers let $s(A, B)$ be the smallest set which has both A and B as subsets.

Yes (depending on your interpretation of “smallest”).

$$s(A, B) = A \cup B.$$

Question 13.1 (cont.) Which of the following rules define functions?

- ▶ For a pair A, B of sets of real numbers let $s(A, B)$ be the smallest set which has both A and B as subsets.

Yes (depending on your interpretation of “smallest”).

$$s(A, B) = A \cup B.$$

- ▶ For a pair A, B of sets of real numbers let $t(A, B)$ be the largest set which is a subset of both A and B .

Question 13.1 (cont.) Which of the following rules define functions?

- ▶ For a pair A, B of sets of real numbers let $s(A, B)$ be the smallest set which has both A and B as subsets.

Yes (depending on your interpretation of “smallest”).

$$s(A, B) = A \cup B.$$

- ▶ For a pair A, B of sets of real numbers let $t(A, B)$ be the largest set which is a subset of both A and B .

Yes (depending on your interpretation of “largest”).

Question 13.1 (cont.) Which of the following rules define functions?

- ▶ For a pair A, B of sets of real numbers let $s(A, B)$ be the smallest set which has both A and B as subsets.

Yes (depending on your interpretation of “smallest”).

$$s(A, B) = A \cup B.$$

- ▶ For a pair A, B of sets of real numbers let $t(A, B)$ be the largest set which is a subset of both A and B .

Yes (depending on your interpretation of “largest”).

$$t(A, B) = A \cap B.$$

13.2 Arrow notation

If f is a function with domain A and codomain B we write

$$f : A \rightarrow B,$$

and we say that f is from A to B .

For example, we could define

$$\text{square} : \mathbb{R} \rightarrow \mathbb{R}.$$

We could also define

$$\text{square} : \mathbb{R} \rightarrow \mathbb{R}^{\geq 0}.$$

Likewise, we could define

$$\text{cube} : \mathbb{R} \rightarrow \mathbb{R}.$$

However we could not define

$$\text{cube} : \mathbb{R} \rightarrow \mathbb{R}^{\geq 0},$$

because for some $x \in \mathbb{R}$, $\text{cube}(x)$ is negative.

For example, $\text{cube}(-1) = -1$.

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

$$x^2$$

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

x^2 Yes.

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

x^2 Yes.

$\frac{1}{x}$

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

x^2 Yes.

$\frac{1}{x}$ No - undefined for $x = 0$.

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

x^2 Yes.

$\frac{1}{x}$ No - undefined for $x = 0$.

$\log(x)$

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

x^2 Yes.

$\frac{1}{x}$ No - undefined for $x = 0$.

$\log(x)$ No - undefined for $x \leq 0$ (because $e^x > 0$ for all $x \in \mathbb{R}$).

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

x^2 Yes.

$\frac{1}{x}$ No - undefined for $x = 0$.

$\log(x)$ No - undefined for $x \leq 0$ (because $e^x > 0$ for all $x \in \mathbb{R}$).

\sqrt{x}

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

x^2 Yes.

$\frac{1}{x}$ No - undefined for $x = 0$.

$\log(x)$ No - undefined for $x \leq 0$ (because $e^x > 0$ for all $x \in \mathbb{R}$).

\sqrt{x} No - undefined for $x < 0$.

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)

x^2 Yes.

$\frac{1}{x}$ No - undefined for $x = 0$.

$\log(x)$ No - undefined for $x \leq 0$ (because $e^x > 0$ for all $x \in \mathbb{R}$).

\sqrt{x} No - undefined for $x < 0$.

$\sqrt[3]{x}$

Question 13.2 Which of the following functions can be defined on the whole of \mathbb{R} , so that the function values also lie in \mathbb{R} ?

(In other words, which can be $\mathbb{R} \rightarrow \mathbb{R}$ functions?)


x^2 Yes.

$\frac{1}{x}$ No - undefined for $x = 0$.

$\log(x)$ No - undefined for $x \leq 0$ (because $e^x > 0$ for all $x \in \mathbb{R}$).

\sqrt{x} No - undefined for $x < 0$.

$\sqrt[3]{x}$ Yes.



13.3 One-to-one functions

A function $f : X \rightarrow Y$ is *one-to-one* if for each y in the range of f there is only one $x \in X$ such that $f(x) = y$.

For example, the function $\text{cube}(x)$ is one-to-one because each real number y is the cube of exactly one real number x .


The function $\text{square} : \mathbb{R} \rightarrow \mathbb{R}$ is *not* one-to-one because the real number 1 is the square of two different real numbers, 1 and -1 . (In fact each real $y > 0$ is the square of two different real numbers, \sqrt{y} and $-\sqrt{y}$.)

On the other hand, $\text{square} : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}$ is one-to-one because each real number y in $\mathbb{R}^{\geq 0}$ is the square of only one real number in $\mathbb{R}^{\geq 0}$, namely \sqrt{y} .


The last example shows that the domain of a function is an important part of its description, because changing the domain can change the properties of the function.

Question Is the function pictured below one-to-one?

Question Is the function pictured below one-to-one?




Question Is the function pictured below one-to-one?




Yes.

Question Is the function pictured below one-to-one?

Question Is the function pictured below one-to-one?




Question Is the function pictured below one-to-one?



No.

Question Is the function pictured below one-to-one?



No. $f(x_2) = f(x_3)$.

13.4 Proving a function is one-to-one

There is an equivalent way of phrasing the definition of one-to-one: a function $f : X \rightarrow Y$ is one-to-one when, for all $x_1, x_2 \in X$,

$$f(x_1) = f(x_2) \Rightarrow x_1 = x_2.$$

This can be useful for proving that some functions are or are not one-to-one.

Example. The function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = 6x + 2$ is one-to-one because

$$\begin{aligned} & f(x_1) = f(x_2) \\ \Rightarrow & 6x_1 + 2 = 6x_2 + 2 \\ \Rightarrow & 6x_1 = 6x_2 \\ \Rightarrow & x_1 = x_2. \end{aligned}$$

Example. The function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = x^2 + 1$ is not one-to-one because $f(-1) = 2$ and $f(1) = 2$ and so

$$f(-1) = f(1).$$

To show that a function $f : X \rightarrow Y$ is one-to-one we must show that, for all $x_1, x_2 \in X$,

if $f(x_1) = f(x_2)$ then $x_1 = x_2$.

To show that a function $f : X \rightarrow Y$ is one-to-one we must show that, for all $x_1, x_2 \in X$,

$$\text{if } f(x_1) = f(x_2) \text{ then } x_1 = x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = 3x - 2$ one-to-one?

To show that a function $f : X \rightarrow Y$ is one-to-one we must show that, for all $x_1, x_2 \in X$,

$$\text{if } f(x_1) = f(x_2) \text{ then } x_1 = x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = 3x - 2$ one-to-one?

Suppose that

$$f(x_1) = f(x_2) \quad \text{for some } x_1, x_2 \in \mathbb{Z}.$$

To show that a function $f : X \rightarrow Y$ is one-to-one we must show that, for all $x_1, x_2 \in X$,

$$\text{if } f(x_1) = f(x_2) \text{ then } x_1 = x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = 3x - 2$ one-to-one?

Suppose that

$$\begin{array}{lll} f(x_1) & = & f(x_2) \\ \text{Then } 3x_1 - 2 & = & 3x_2 - 2. \end{array} \quad \text{for some } x_1, x_2 \in \mathbb{Z}.$$

To show that a function $f : X \rightarrow Y$ is one-to-one we must show that, for all $x_1, x_2 \in X$,

$$\text{if } f(x_1) = f(x_2) \text{ then } x_1 = x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = 3x - 2$ one-to-one?

Suppose that

$$f(x_1) = f(x_2) \quad \text{for some } x_1, x_2 \in \mathbb{Z}.$$

Then

$$3x_1 - 2 = 3x_2 - 2.$$

So

$$3x_1 = 3x_2.$$

To show that a function $f : X \rightarrow Y$ is one-to-one we must show that, for all $x_1, x_2 \in X$,

$$\text{if } f(x_1) = f(x_2) \text{ then } x_1 = x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = 3x - 2$ one-to-one?

Suppose that

$$f(x_1) = f(x_2) \quad \text{for some } x_1, x_2 \in \mathbb{Z}.$$

Then

$$3x_1 - 2 = 3x_2 - 2.$$

So

$$3x_1 = 3x_2.$$

So

$$x_1 = x_2.$$

To show that a function $f : X \rightarrow Y$ is one-to-one we must show that, for all $x_1, x_2 \in X$,

$$\text{if } f(x_1) = f(x_2) \text{ then } x_1 = x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = 3x - 2$ one-to-one?

Suppose that

$$f(x_1) = f(x_2) \quad \text{for some } x_1, x_2 \in \mathbb{Z}.$$

Then
$$3x_1 - 2 = 3x_2 - 2.$$

So
$$3x_1 = 3x_2.$$

So
$$x_1 = x_2.$$

This shows that f is one-to-one.

To show that a function $f : X \rightarrow Y$ is *not* one-to-one we must show that there exist $x_1, x_2 \in X$ such that

$$f(x_1) = f(x_2) \text{ and } x_1 \neq x_2.$$

To show that a function $f : X \rightarrow Y$ is *not* one-to-one we must show that there exist $x_1, x_2 \in X$ such that

$$f(x_1) = f(x_2) \text{ and } x_1 \neq x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = x^2 + 1$ one-to-one?

To show that a function $f : X \rightarrow Y$ is *not* one-to-one we must show that there exist $x_1, x_2 \in X$ such that

$$f(x_1) = f(x_2) \text{ and } x_1 \neq x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = x^2 + 1$ one-to-one?

No.

To show that a function $f : X \rightarrow Y$ is *not* one-to-one we must show that there exist $x_1, x_2 \in X$ such that

$$f(x_1) = f(x_2) \text{ and } x_1 \neq x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = x^2 + 1$ one-to-one?

No. $f(1) = 2$ and $f(-1) = 2$

To show that a function $f : X \rightarrow Y$ is *not* one-to-one we must show that there exist $x_1, x_2 \in X$ such that

$$f(x_1) = f(x_2) \text{ and } x_1 \neq x_2.$$

Question Is $f : \mathbb{Z} \rightarrow \mathbb{Z}$ defined by $f(x) = x^2 + 1$ one-to-one?

No. $f(1) = 2$ and $f(-1) = 2$ (and obviously $1 \neq -1$)


Question 13.3 Find the range of the following functions, given that the domain of each is \mathbb{R} .

Question 13.3 Find the range of the following functions, given that the domain of each is \mathbb{R} .

$$e^x$$


Question 13.3 Find the range of the following functions, given that the domain of each is \mathbb{R} .

$$e^x$$



Question 13.3 Find the range of the following functions, given that the domain of each is \mathbb{R} .


$$e^x$$



The range is $\{y : y \in \mathbb{R} \text{ and } y > 0\}$

Question 13.3 Find the range of the following functions, given that the domain of each is \mathbb{R} .

$$e^x$$



The range is $\{y : y \in \mathbb{R} \text{ and } y > 0\}$

The rest of these are left as an exercise.