# Numerical Analysis - Winter 2019

## Assignment #3

Issued: Nov.27, 2019 Due: Dec.14, 2019

Please hand in the C or Matlab code (.m files), graphics, and a brief description of your reasoning as well as comments if any. You should pack all of your files into a .rar or .zip file, titled as "xxxxxxx(your student ID)\_Homework\_3", and then submit it by uploading to ftp server or sending to TA before 11:59pm of the due day.

Please upload to the 'hw3' directory if you submit your homework in time.

#### **Problem 1:**

The following linear systems  $A\mathbf{x} = \mathbf{b}$  have  $\mathbf{x}$  as the actual solution and  $\tilde{\mathbf{x}}$  as an approximate solution. Compute  $\|\mathbf{x} - \tilde{\mathbf{x}}\|_{\infty}$  and  $\|A\tilde{\mathbf{x}} - \mathbf{b}\|_{\infty}$ .

$$\mathbf{a.} \quad x_1 + 2x_2 + 3x_3 = 1,$$

$$2x_1 + 3x_2 + 4x_3 = -1,$$

$$3x_1 + 4x_2 + 6x_3 = 2,$$

$$\mathbf{x} = (0, -7, 5)^t$$

$$\tilde{\mathbf{x}} = (-0.2, -7.5, 5.4)^t$$
.

**b.** 
$$x_1 + 2x_2 + 3x_3 = 1$$
,

$$2x_1 + 3x_2 + 4x_3 = -1,$$

$$3x_1 + 4x_2 + 6x_3 = 2$$
,

$$\mathbf{x} = (0, -7, 5)^t$$

$$\tilde{\mathbf{x}} = (-0.33, -7.9, 5.8)^t$$
.

### **Problem 2:**

Show that if *A* is symmetric, then  $||A||_2 = \rho(A)$ .

#### **Problem 3:**

Implement the algorithm of Gaussian elimination with scaled partial pivoting, and solve the following linear systems.

$$\mathbf{a.} \quad 0.03x_1 + 58.9x_2 = 59.2,$$

$$5.31x_1 - 6.10x_2 = 47.0.$$

Actual solution [10, 1].

**b.** 
$$3.03x_1 - 12.1x_2 + 14x_3 = -119,$$

$$-3.03x_1 + 12.1x_2 - 7x_3 = 120,$$

$$6.11x_1 - 14.2x_2 + 21x_3 = -139.$$

Actual solution  $[0, 10, \frac{1}{7}]$ .

### **Problem 4:**

Implement the Jacobi iterative method and list the first three iteration results when solving the following linear systems, using  $\mathbf{x}^{(0)} = \mathbf{0}$ .

$$\mathbf{a.} \quad 4x_1 + x_2 - x_3 = 5,$$

$$-x_1 + 3x_2 + x_3 = -4,$$

$$2x_1 + 2x_2 + 5x_3 = 1.$$

**b.** 
$$-2x_1 + x_2 + \frac{1}{2}x_3 = 4$$
,

$$x_1 - 2x_2 - \frac{1}{2}x_3 = -4,$$

$$x_2 + 2x_3 = 0.$$

# **Problem 5:**

Use the Jacobi method and Gauss-Seidel method to solve the following linear systems, with TOL = 0.001 in the  $L\infty$  norm.

**a.** 
$$3x_1 - x_2 + x_3 = 1$$
,

$$3x_1 + 6x_2 + 2x_3 = 0,$$

$$3x_1 + 3x_2 + 7x_3 = 4.$$

**b.** 
$$10x_1 - x_2 = 9$$
,

$$-x_1 + 10x_2 - 2x_3 = 7,$$

$$- 2x_2 + 10x_3 = 6.$$