# Boron Ion Beam Production for the Future CERN BIO-LEIR Facility

Joshua Stafford-Haworth jstaffor@cern.ch







# Boron Ion Beam Production for the Future CERN BIO-LEIR Facility

Joshua Stafford-Haworth jstaffor@cern.ch














- Hadron therapy mostly conducted with protons
- lons offer different radiobiological properties
- Comprehensive study needed to evaluate different species
- Protons -> Neon
- Re-use LEIR synchrotron at CERN for light ion medical research facility.





- Hadron therapy mostly conducted with protons
- Ions offer differenteadio biedogical properties








#### Hamblow Emerapyon esieg (CLEIR)

- Hadron therapy mostly conducted with protons
- Ions offer different madio bie dogical properties







#### Hamblow Emerapyon esieg (CLEIR)

Hadron therapy mostly teoreticted with protons

· lons offer different properties

- Comprehensi different sevaluate
- Protons
- Re-use CERNefor light ion med.

  Others

AD Antiproton Decelerator





© CERN 2013

Protons

Carbon

proton/antiproton conversion

PS Proton Synchrotron

ed WAKefield Experiment ISOLDE Isotope Separator OnLine DEvice


#### Hamblow Emerapyon esieg (CLEIR)

Hadron therapy mostly teoreticted with protons

· lons offer different properties

- Comprehensi different servaluate
- Protons
- Re-use ion med

AD Antiproton Decelerator




ed WAKefield Experiment ISOLDE Isotope Separator OnLine DEvice

Protons






#### Hamblow Emerapyon resign (CIEIR)






#### Hate Early Emply Emply Emply Emply (the IR)



© CERN 2013



#### Hatel Early Emply Emply Emply Emply (the IR)







## Light Ion Injectors



#### Light Ion Injectors

- New Linac?
- Cyclotron?
- Re-use heavy ion injector, Linac 3?
- A new source is needed that can deliver p->Ne
- Pantechnik Supernanogan currently used at CNAO, HIT, MedAustron.
- Permanent Magnet 14.5 GHz ECR capable of oven technique with gas mixing.



#### **Boron Production**




#### **Boron Production**


- JYFL (Jyväskylä) and FLNR (Dubna) have previously delivered Boron
- Both ECR sources, both much larger than Supernanogan, and not permanent magnet designs
- Verification required
- MIVOC technique with Supernanogan at Helmholtz-Zentrum, Berlin.












- MIVOC chamber
- Needle Valve
- Standard Valve
- RF waveguide
- Source plasma chamber






- MIVOC chamber
- Needle Valve
- Standard Valve
- RF waveguide
- Source plasma chamber






- MIVOC chamber
- Needle Valve
- Standard Valve
- RF waveguide
- Source plasma chamber






- MIVOC chamber
- Needle Valve
- Standard Valve
- RF waveguide
- Source plasma chamber





- MIVOC chamber
- Needle Valve
- Standard Valve
- RF waveguide
- Source plasma chamber







- MIVOC chamber
- Needle Valve
- Standard Valve
- RF waveguide
- Source plasma chamber












Atomic Formula	B <sub>10</sub> H <sub>14</sub>
Melting Point	373 K
<b>Boiling Point</b>	486 K
Vapor Pressure	0.269 mbar
Phase at 300k	Crystalline






Atomic Formula	B <sub>10</sub> H <sub>14</sub>
Melting Point	373 K
<b>Boiling Point</b>	486 K
Vapor Pressure	0.269 mbar
Phase at 300k	Crystalline


**Boron** 






Atomic Formula	B <sub>10</sub> H <sub>14</sub>
Melting Point	373 K
<b>Boiling Point</b>	486 K
Vapor Pressure	0.269 mbar
Phase at 300k	Crystalline






















































































































































































Distribution Favours <sup>11</sup>B<sup>3+</sup>










































## Success!



#### Success!

- Supernanogan can deliver Boron for medical research!
- M-carborane without support gas can deliver up to 50 μA of <sup>11</sup>B<sup>3+</sup>
- Decaborane, less successful. Pyrolysis makes a dirty source!



# Boron Ion Beam Production for the Future CERN BIO-LEIR Facility

Joshua Stafford-Haworth jstaffor@cern.ch







# Boron Ion Beam Production for the Future CERN BIO-LEIR Facility

Joshua Stafford-Haworth jstaffor@cern.ch





