

PERFORMANCE OF A LOW FREQUENCY QWR-BASED SRF GUN


Wisconsin Free Electron Laser (WiFEL) Electron Gun Cavity

Gongxiahui Chen, Troy Petersen, Michael P. Kelly, Mike V. Fisher, Mark Kedzie, Tom Reid, ANL, Lemont, Illinois, USA
 Xueying Lu, Philippe Piot, Northern Illinois University, DeKalb, Illinois, USA


ABSTRACT

Superconducting radio-frequency (SRF) electron guns are generally considered to be an effective way of producing beams with high brightness and high repetition rates (or continuous wave). In this work, the 199.6 MHz quarter wave resonator (QWR)-based Wisconsin Free Electron Laser (WiFEL) superconducting electron gun was recently refurbished and tested at Argonne (ANL). The field performance of the e-gun was fully characterized. During this time, multipacting (MP) conditioning was performed for over 20 hours to overcome the hard MP barrier observed in the accelerating voltage range of 8 to 40 kV; the presence of multipacting is expected to operationally important for future e-guns. Here we simulated and studied the effect using CST [1] Microwave Studio and Particle Studio and compare with the measured data.

WiFEL CAVITY GEOMETRY AND THE FIELD DISTRIBUTION




Q-CURVE




- $Q_0 = 2.3 \times 10^8$ @ highest stable CW E_{peak} of 15 MV/m.
- Low field Q -drop was found NOT caused by Q-disease, more likely related to physical imperfection.

MULITPACTING (MP) CONDITIONING



- Measured MP turn-on @ ~8 kV.
- Strong MP observed in between 8 kV and 40 kV.
- Total conditioning time: >20 hours.

CST Simulation



Fit to the exponential function:

$$N_e(t) = N_0 e^{\alpha t}$$



where,

N_e - total number of particles



N_0 - primary number of particles

α - growth rate

• Cavity may have a high SEY.



PARTICLE EVOLUTION



- A 'delayed' MP behavior shows up after extending the running time.
- Lower MP turn-on voltage is found (<10 kV). In good agreement with the experimental results!

ACKNOWLEDGEMENT

This work was supported by the U.S. Department of Energy, Office of Nuclear Physics, under Contract No. DE-AC02-06CH11357. This research used resources of ANL's ATLAS facility, which is a DOE Office of Science User Facility.

REFERENCES

- [1] CST Studio Suite, version 2020, <https://www.cst.com>.