

New upper bounds for the chromatic numbers of Euclidean spaces:

$$\chi(\mathbb{R}^4) \leq 43, \quad \chi(\mathbb{R}^5) \leq 132, \quad \chi(\mathbb{R}^7) \leq 1029, \quad \chi(\mathbb{R}^9) \leq 7203, \quad \chi(\mathbb{R}^{10}) \leq 45619$$

Leonid L. Ivanov*

Nadezhda V. Glushkova†

September 2026

Abstract

A coloring of $\mathbb{R}^n$ is *proper for the forbidden distance segment* $[1, \ell]$ if no two points of the same color are at a distance from $[1, \ell]$; the minimum number of colors is $\chi(\mathbb{R}^n, [1, \ell])$, and $\ell = 1$ gives the classical chromatic number $\chi(\mathbb{R}^n)$ of the Nelson–Hadwiger problem. We prove the new upper bounds $\chi(\mathbb{R}^4) \leq 43$, $\chi(\mathbb{R}^5) \leq 132$, $\chi(\mathbb{R}^7) \leq 1029$, $\chi(\mathbb{R}^9) \leq 7203$, $\chi(\mathbb{R}^{10}) \leq 45619$, improving the previously known 49, 140, 1372, 17253 and 3^{10} ; in particular, this refutes the conjecture of Arman, Bondarenko, Prymak and Radchenko that 49 and 140 are optimal among all lattice colorings of $\mathbb{R}^4$ and $\mathbb{R}^5$. The first four bounds come from explicit rational lattices — an Eisenstein lattice in $\mathbb{R}^4$, a lattice in general position in $\mathbb{R}^5$, and laminations of the Eisenstein colorings $E_6^*/343$ and $E_8/2401$ in $\mathbb{R}^7$ and $\mathbb{R}^9$ — and each is reduced, by one verification protocol, to a finite list of inequalities between explicitly written rational numbers checked in exact arithmetic. The fifth bound is analytic: we prove that for every Eisenstein lattice Λ the distance between same-colored cells of $(3 + \omega)\Lambda$ equals $\sqrt{7/3} \lambda_1(\Lambda)$, which gives the exact widths of all known colorings with $7^{n/2}$ colors, and a product rule $\sum_i 1/d_i^2 \leq 1$ for the widths of orthogonal products; together they yield $45619 = 2401 \cdot 19$, the first bound in $\mathbb{R}^{10}$ below 3^n , as well as $\chi(\mathbb{R}^{25}) \leq 4 \cdot 7^{12}$ and $\chi(\mathbb{R}^{26}) \leq 19 \cdot 7^{12}$. We also show that no sublattice of E_8 of index below 2401 defines a proper coloring. All code, exact certificates and data are open.

MSC 2020: 52C10 (primary); 05C15, 52C07, 11H31.

Keywords: chromatic number of Euclidean space, Nelson–Hadwiger problem, forbidden distance interval, lattice colorings, Voronoi cell, covering radius, Eisenstein lattices, E_8 .

1 Introduction

The *chromatic number* $\chi(\mathbb{R}^n)$ is the minimum number of colors needed to color the points of $\mathbb{R}^n$ so that no two points at distance exactly 1 receive the same color (the Nelson–Hadwiger problem; for the plane only $5 \leq \chi(\mathbb{R}^2) \leq 7$ is known [7, 15]). For $\ell \geq 1$ let $\chi(\mathbb{R}^n, [1, \ell])$ be the minimum number of colors of a coloring in which no two points of the same color realize any distance from the segment $[1, \ell]$ [10]; forbidding more distances cannot require fewer colors, so

$$\chi(\mathbb{R}^n) = \chi(\mathbb{R}^n, [1, 1]) \leq \chi(\mathbb{R}^n, [1, \ell]) \quad (\ell \geq 1), \quad (1)$$

and every upper bound for the interval version bounds $\chi(\mathbb{R}^n)$. For the history of the problem see the surveys [3, 12, 13, 14].

All known upper bounds in small dimensions come from *lattice colorings*: the space is cut into the Voronoi cells of a lattice Λ , and the cell of $v \in \Lambda$ receives the color of the coset $v + \Gamma$ of a sublattice $\Gamma \subset \Lambda$ of index k . The coloring is proper for $[1, \ell]$ whenever its *width* — the ratio of the smallest distance between same-colored cells to the diameter of a cell — exceeds ℓ (Section 2). The best previous values in dimensions 4–10 are due to Arman, Bondarenko, Prymak and Radchenko [1] (ABPR below): 49, 140, 343, 1372, 2401, 17253 and 3^{10} . In dimension 4 the bound 49 (Coulson; proved, with the minimality of the index 49 on the lattice D_4 , in [1]) improved 54 of [11], while the lower bound is 9 [8] (after 7 [9]). ABPR conjecture (Open question 1) that 49 and 140 are optimal among all lattice colorings of $\mathbb{R}^4$ and $\mathbb{R}^5$. This paper refutes that conjecture.

Theorem 1.1 (main theorem).

$$\chi(\mathbb{R}^4) \leq 43, \quad \chi(\mathbb{R}^5) \leq 132, \quad \chi(\mathbb{R}^7) \leq 1029, \quad \chi(\mathbb{R}^9) \leq 7203, \quad \chi(\mathbb{R}^{10}) \leq 45619.$$

More precisely, $\chi(\mathbb{R}^n, [1, \ell]) \leq k$ for all $\ell \leq \ell_0$ for the triples (n, k, ℓ_0) of Table 1.

Table 1: The five proven bounds; ℓ_0 is the proven width of the forbidden segment ($< x$: proven for every $\ell < x$). Previous bounds are from [1].

n	previous	new	ℓ_0	proof	where
4	49	43	1.00411	exact rational certificate	Theorem 3.1
5	140	132	101/100	exact rational certificate	Theorem 3.2
7	1372	1029	103/100	exact rational certificate	Theorem 3.3
9	17253	7203	< 1.0166127	exact rational certificate	Theorem 3.4
10	3^{10}	45619	$< \sqrt{217/214}$	analytic (product rule)	Theorem 4.7

For $n = 4, 5, 7, 9$ we exhibit a rational Gram matrix and an integer sublattice and reduce the inequality $d > \ell_0$, by one protocol (Proposition 2.3), to finitely many inequalities between rational numbers verified in exact arithmetic; floating point only guided the search. For $n = 10$ the proof is analytic: the *Eisenstein identity* of Section 4 gives the exact width $\sqrt{7/6}$ of the block $E_8/2401$, and the *product rule* $\sum_i 1/d_i^2 \leq 1$ leaves room for a planar block with 19 colors ($6/7 + 4/31 < 1$). The same rule gives $4 \cdot 7^{12}$ and $19 \cdot 7^{12}$ colors in $\mathbb{R}^{25}$ and $\mathbb{R}^{26}$. Section 5 shows that no sublattice of E_8 of index below 2401 is admissible. All code, exact certificates and data are open at <https://github.com/ivaleo/Chromatic> (MIT license), together with the detailed Russian manuscript (search algorithms, ladders of widths, negative screens, and numerical candidates such as an index-28812 coloring of $\mathbb{R}^{10}$, which we do *not* claim as a bound because its diameter certificate is numerical).

2 Width and the certificate protocol

Let $\Lambda \subset \mathbb{R}^n$ be a lattice, $\Gamma \subseteq \Lambda$ a sublattice of index $k = [\Lambda : \Gamma]$, and $V_0 = \{x : \|x\| \leq \|x - b\| \ \forall b \in \Lambda\}$ the Voronoi cell of the origin, a centrally symmetric convex polytope with $\text{diam } V_0 = 2R$, R the covering radius. Color the cell $v + V_0$ by the coset $v + \Gamma$. Two points of the same color lie either in one cell or in cells whose centers differ by $v \in \Gamma \setminus \{0\}$. Put

$$D(v) = \text{dist}(V_0, v + V_0), \quad D(\Gamma) = \min_{v \in \Gamma \setminus \{0\}} D(v), \quad d(\Lambda, \Gamma) = \frac{D(\Gamma)}{\text{diam } V_0}; \quad (2)$$

the open interval $(\text{diam } V_0, D(\Gamma))$ is then free of same-colored distances.

Proposition 2.1. *If $d(\Lambda, \Gamma) > 1$, then $\chi(\mathbb{R}^n, [1, \ell]) \leq [\Lambda : \Gamma]$ for all $\ell < d(\Lambda, \Gamma)$; in particular $\chi(\mathbb{R}^n) \leq [\Lambda : \Gamma]$.*

Proof. A scale $s > 0$ with $s \text{diam } V_0 < 1$ and $sD(\Gamma) > \ell$ exists iff $\ell < d$; for $s\Lambda$ the free interval contains $[1, \ell]$. Use (1) for $\ell = 1$. $\square$

Lemma 2.2 (midpoint lemma; [10, Lemma 2]). $D(v) = 2 \text{dist}(v/2, V_0)$.

Proof. If $x \in V_0$, $y \in v + V_0$, then $x' = v - y \in V_0$ and $\|x - y\| = 2\|(x + x')/2 - v/2\| \geq 2 \text{dist}(v/2, V_0)$ by convexity; for the point $x \in V_0$ closest to $v/2$ and $y = v - x$ equality holds. $\square$

Since V_0 lies in the ball of radius R , $D(v) \geq \|v\| - \text{diam } V_0$, so $D(v) \geq \ell \text{diam } V_0$ can only fail inside the *window*

$$\|v\| < (1 + \ell) \text{diam } V_0 = 2(1 + \ell)R, \quad (3)$$

a provably complete search bound.

Proposition 2.3 (certificate protocol). *Let $Q \in \text{Mat}_n(\mathbb{Q})$ be symmetric with positive leading principal minors, $\Lambda = \mathbb{Z}^n$ with $\langle x, y \rangle = x^\top Q y$, and $\Gamma = C\mathbb{Z}^n$ for an integer matrix C . Suppose a finite computation in exact fraction arithmetic has established: (i) $|\det C| = k$; (ii) for each nonzero class of $\Lambda/2\Lambda$ the shortest representatives are found, and V_0 is cut out by $\langle x, v \rangle \leq \frac{1}{2}\langle v, v \rangle$ over the classes with a unique pair $\pm v$ (Voronoi's relevant vectors [5, Ch. 21]); (iii) all vertices of V_0 are enumerated, completeness being proved by solving all n -subsets of facets or by exact closure along the 1-skeleton, and $R^2 = \max \langle x, x \rangle$ over the vertices; (iv) all $v \in \Gamma \setminus \{0\}$ with $\langle v, v \rangle < W$ are listed, where $W \geq (2(1 + \ell_0)R)^2$ is rational; (v) for each such v a point $x^* \in V_0$ and multipliers $\mu \geq 0$ of the active facets with $v/2 - x^* = \sum_j \mu_j v_j$ are exhibited (the KKT conditions of the projection), whence $D(v)^2 = 4\|v/2 - x^*\|^2$ exactly; (vi) $D_{\min}^2 - \ell_0^2 \cdot 4R^2 > 0$, $D_{\min}$ the minimum over the window. Then $d(\Lambda, \Gamma) > \ell_0$ and $\chi(\mathbb{R}^n, [1, \ell]) \leq k$ for all $\ell \leq \ell_0$.*

*E-mail: leo.ivanov@gmail.com.

†E-mail: glushkova.nv@phystech.edu.

Proof. (ii) is Voronoi's theorem; (iii): the maximum of $\langle x, x \rangle$ over a polytope is attained at a vertex and $\text{diam } V_0 = 2R$; (iv) is (3); (v): KKT conditions are sufficient for projection onto a polytope, and $D(v) = 2 \text{dist}(v/2, V_0)$ by Lemma 2.2; (vi) gives $D(\Gamma) > \ell_0 \text{diam } V_0$, and Proposition 2.1 applies. $\square$

Floating point serves only as an oracle proposing vertices and active sets; every accepted quantity is recomputed over $\mathbb{Q}$, so an oracle error can stop a verification but cannot make it wrong.

3 Four explicit constructions

Each theorem below is proved by the protocol of Proposition 2.3; we give the data and the quantities the protocol produces. The full lists of vertices, window vectors and KKT certificates are in the files named in Section 6.

3.1 Dimension 4: $\chi(\mathbb{R}^4) \leq 43$

Let $S \in \text{GL}_4(\mathbb{Z})$ have order 3 and characteristic polynomial Φ_3^2 . A lattice with $S\Lambda = \Lambda$ is a $\mathbb{Z}[\omega]$ -module of rank 2 ($\omega = e^{2\pi i/3}$), an *Eisenstein* lattice, and the S -invariant forms are the real parts of Hermitian forms $\begin{pmatrix} a & c \\ \bar{c} & b \end{pmatrix}$; in the $\mathbb{Z}$ -basis $(e_1, \omega e_1, e_2, \omega e_2)$, with $u = \text{Re } c$ and $s = \frac{\sqrt{3}}{2} \text{Im } c$, the Gram matrix is

$$Q(a, b, u, s) = \begin{pmatrix} a & -\frac{a}{2} & u & -\frac{u}{2} - s \\ -\frac{a}{2} & a & -\frac{u}{2} + s & u \\ u & -\frac{u}{2} + s & b & -\frac{b}{2} \\ -\frac{u}{2} - s & u & -\frac{b}{2} & b \end{pmatrix}. \quad (4)$$

An S -invariant sublattice is a $\mathbb{Z}[\omega]$ -submodule, so its index is the norm of an ideal; $43 = N(7 + \omega)$, and there are exactly 88 submodules of index 43 (against 81 400 sublattices). Restricting the search to this three-parameter family (instead of nine parameters) produced the construction; in the full space of quaternary forms the search had stalled below the threshold.

Let $\Lambda_\star = \mathbb{Z}^4$ with the form (4),

$$a = 1, \quad b = \frac{39261}{25000}, \quad u = \frac{37381}{100000}, \quad s = \frac{3141}{12500} \quad (200\,000\,Q \in \text{Mat}_4(\mathbb{Z})), \quad (5)$$

and let $\Gamma_\star$ be generated by the rows of

$$T = \begin{pmatrix} 1 & 0 & 0 & 41 \\ 0 & 1 & 0 & 14 \\ 0 & 0 & 1 & 37 \\ 0 & 0 & 0 & 43 \end{pmatrix} \quad (\text{coordinates in the basis of } \Lambda_\star), \quad \det T = 43. \quad (6)$$

Then $\lambda_1(\Lambda_\star)^2 = 1$ and

$$\begin{aligned} \text{diam}^2 V_0 &= \frac{5374343884337}{2019775849050} = 2.660861544\dots, \\ D(\Gamma_\star)^2 &= \frac{7396075258066147}{2756867811550000} = 2.682781969\dots, \\ d(\Lambda_\star, \Gamma_\star)^2 &= \frac{298768283679964998359742207}{296327113258585430253847000} = 1.008238093\dots, \quad d = 1.004110598\dots \end{aligned}$$

Theorem 3.1. $\chi(\mathbb{R}^4, [1, \ell]) \leq 43$ for all $1 \leq \ell \leq \ell_0 = 1.00411$; in particular, $\chi(\mathbb{R}^4) \leq 43$.

Proof. The 15 nonzero classes of $\Lambda_\star/2\Lambda_\star$ give 30 facets (the maximum in $\mathbb{R}^4$); all $\binom{30}{4} = 27\,405$ rational 4×4 systems yield exactly 120 vertices (f -vector $(120, 240, 150, 30)$, $\text{vol } V_0 = \det \Lambda_\star$) and $\text{diam}^2 V_0$ above. The window (3) for ℓ_0 is enumerated, each $\text{dist}(v/2, V_0)$ is certified by KKT multipliers in fractions, and the minimum of D^2 is the value above, attained on three pairs $\pm v$ forming one orbit of $\text{Aut}(\Lambda_\star) \cong \mathbb{Z}/6$. Margin:

$$D(\Gamma_\star)^2 - \ell_0^2 \text{diam}^2 V_0 = \frac{3559631641205182621936510913}{111365100495840332930550000000000} = 3.196 \cdot 10^{-6} > 0. \quad \square$$

A second, independent certificate, which uses a supporting direction for each window vector instead of the projection, bounds

$$D(\Gamma_\star)^2 \geq \frac{869307639027355969}{324032161065600000} = 2.682781969\dots$$

from below, and an exhaustive scan of all 81 400 sublattices of index 43 confirms that (6) maximizes d on $\Lambda_\star$. The point (5) is a rationalization (denominator 10^5) of the numerical optimum $d = 1.00411205721\dots$ of the family.

3.2 Dimension 5: $\chi(\mathbb{R}^5) \leq 132$

The previous bound 140 was proved by ABPR to be minimal among sublattices of A_5^* ; the construction below changes the parent form. Let $\Lambda = \mathbb{Z}^5$ with $\langle x, y \rangle = x^\top Q y$,

$$Q = \frac{1}{100000} \begin{pmatrix} 79327 & 86284 & 69597 & 36140 & 26501 \\ 86284 & 254538 & 216808 & 101257 & 102135 \\ 69597 & 216808 & 296568 & 163667 & 140527 \\ 36140 & 101257 & 163667 & 181733 & 71653 \\ 26501 & 102135 & 140527 & 71653 & 148076 \end{pmatrix}$$

with the leading principal minors

$$79327, \quad 12746807270, \quad 1422467480626358, \quad 124437401434750889345, \quad 9999935596575703407615413,$$

and $\Gamma = \ker \varphi$ for the surjective homomorphism

$$\varphi(x) = 88x_1 + 89x_2 + 127x_3 + 57x_4 + 3x_5 \pmod{132},$$

so $[\Lambda : \Gamma] = 132$ (Smith form $(1, 1, 1, 1, 132)$; φ combines characters mod 11, 3 and 4).

Theorem 3.2. $\chi(\mathbb{R}^5, [1, \ell]) \leq 132$ for all $1 \leq \ell \leq 101/100$; in particular, $\chi(\mathbb{R}^5) \leq 132$.

Proof. The cell has 62 facets and 720 vertices, all solved over $\mathbb{Q}$, with

$$R^2 = \frac{3093570095915736710467707313641}{3999974238630281363046165200000}.$$

For $\ell = 101/100$ the window (3) is $\|v\|^2 < 12.4984\dots$; a Fincke–Pohst enumeration over an LLL-reduced basis of Γ leaves 38 nonzero vectors (19 pairs) in it. For all of them the projection onto V_0 is certified over $\mathbb{Q}$; the minimum is attained at $\pm(2, 0, 1, -3, 0)$ (coordinates in the basis of Λ):

$$D_{\min}^2 = \frac{1634114770527727}{516898614620000},$$

$$D_{\min}^2 - \left(\frac{101}{100}\right)^2 4R^2 = \frac{1450500657237822255902962097875907780124229}{258447642807960215367421006872956903000000000} > 0,$$

so $D_{\min}/\text{diam } V_0 = 1.010897714548927\dots > 1.01$. □

An independent verifier that uses neither Qhull nor a floating-point enumerator reproduces the same R^2 , $D_{\min}^2$ and KKT certificates.

3.3 Dimension 7: $\chi(\mathbb{R}^7) \leq 1029$

Lamination builds a coloring of $\mathbb{R}^n$ from a coloring $(\Lambda', \Gamma' = \ker \psi)$ of $\mathbb{R}^{n-1}$: for a shift c , a height $t > 0$, a gluing $a \in \Lambda'/\Gamma'$ and a layer modulus m ,

$$\Lambda = \{(x + ic, it) : x \in \Lambda', i \in \mathbb{Z}\}, \quad \Gamma = \{(x + ic, it) : \psi(x) + ia = 0, m \mid i\}, \quad [\Lambda : \Gamma] = [\Lambda' : \Gamma'] m.$$

The squared diameter of the cell grows by at most t^2 , and the distances between “horizontal” same-colored cells (zero layer coordinate) do not decrease, so the excess width of the base pays for the layer height. The base here is the Eisenstein coloring $E_6^*/343$, $\Gamma' = (3 + \omega)E_6^*$, of width $\sqrt{7/6}$ (Corollary 4.5), and $m = 3$. In a $\mathbb{Z}$ -basis whose first six vectors span the base and whose seventh is the layer vector (c, t) , the rational point we verify is

$$Q = \begin{pmatrix} \frac{3}{4}M & g \\ g^\top & 17897/10^4 \end{pmatrix}, \quad M = \begin{pmatrix} 4 & 2 & 0 & -3 & 2 & 1 \\ 2 & 4 & 3 & 0 & 1 & 2 \\ 0 & 3 & 6 & 3 & 0 & 3 \\ -3 & 0 & 3 & 6 & -3 & 0 \\ 2 & 1 & 0 & -3 & 4 & 2 \\ 1 & 2 & 3 & 0 & 2 & 4 \end{pmatrix}, \quad (7)$$

$$10^4 g^\top = (-8632, -4197, 1601, 6617, -5275, 4434),$$

where $\frac{3}{4}M$ is a Gram matrix of E_6^* with $\lambda_1^2 = 3$, and $\Gamma = C\mathbb{Z}^7$ with C upper triangular, diagonal $(7, 1, 7, 1, 7, 1, 3)$, the only nonzero off-diagonal entries $C_{12} = C_{34} = C_{56} = -5$, and $\det C = 1029$ (the three blocks $\begin{pmatrix} 7 & -5 \\ 0 & 1 \end{pmatrix}$ realize the ideal of norm 7 in the three complex coordinates of the base).

Theorem 3.3. For $\Lambda = \mathbb{Z}^7$ with the form (7) and $\Gamma = C\mathbb{Z}^7$,

$$D(\Gamma)^2 = 7, \quad \text{diam}^2 V_0 = \frac{96858928789031597}{14761819462500000}, \quad d^2 = \frac{103332736237500000}{96858928789031597}, \quad d = 1.032878\dots,$$

so $\chi(\mathbb{R}^7, [1, \ell]) \leq 1029$ for all $1 \leq \ell \leq 103/100$; in particular, $\chi(\mathbb{R}^7) \leq 1029$.

Proof. Voronoi's theorem on $\Lambda/2\Lambda$ (with the complete bound $\max_{s \in \{0,1\}^7} Q(s, s)$) gives 127 pairs, i.e. 254 facets, the maximum in $\mathbb{R}^7$. The 30 368 vertices are obtained as exact solutions of integer systems; completeness is proved by exact closure along the 1-skeleton without assuming simplicity (1 600 vertices have nine active facets), giving $R^2 = 96858928789031597/59047277850000000$. The safe window $\|v\|^2 \leq 2(D_*^2 + \text{diam}^2 V_0) = 200191665026531597/7380909731250000$ contains 74 vectors, each with an exact KKT certificate. The minimum 7 is attained at 27 pairs $\pm v$ of horizontal vectors, the images of the 54 minimal vectors of E_6^* (the floor $\sqrt{7/3}\lambda_1$ of Proposition 4.1); the nearest layer vector gives $D^2 = 478385948533/68336903600 \approx 7.000404$. Margin for $\ell = 103/100$: $D(\Gamma)^2 - \ell^2 \text{diam}^2 V_0 = 5750986852163787427/147618194625000000 > 0$. $\square$

The inequality $\text{diam}^2 V_0 < 7$ was also obtained by a second route sharing no computation with the proof above: on the pieces of the common refinement of two shifted Voronoi partitions of the six-dimensional base (167 pieces, 103 provably empty) the covering radius is bounded by the maximum of an explicit convex quadratic, which equals $103310189182571717/59047277850000000 = 1.7496\dots < 7/4$, all in fractions.

3.4 Dimension 9: $\chi(\mathbb{R}^9) \leq 7203$

The same lamination over $E_8/2401$, $\Gamma' = (3 + \omega)E_8$ (width $\sqrt{7/6}$), with $m = 3$ and $t^2 = 8264707/6250000$: in a $\mathbb{Z}$ -basis whose first eight vectors span E_8 and whose ninth is the layer vector,

$$Q = \begin{pmatrix} \frac{3}{2}A & g \\ g^\top & 17982/10^4 \end{pmatrix}, \quad A = \begin{pmatrix} 2 & 1 & 1 & -1 & -1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 0 & -1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 2 & 0 & 0 & 0 & -1 \\ -1 & -1 & 0 & 0 & 2 & -1 & -1 & -1 \\ 1 & 1 & 0 & 0 & -1 & 2 & 1 & 1 \\ 1 & 1 & 0 & 0 & -1 & 1 & 2 & 1 \\ 1 & 1 & 0 & -1 & -1 & 1 & 1 & 2 \end{pmatrix}, \quad (8)$$

$$10^4 g^\top = (3329, 6390, 4411, -611, -3579, 5606, 4509, 8550),$$

$$C = \begin{pmatrix} -5 & -2 & -9 & -5 & -3 & -6 & -10 & -3 & 0 \\ 4 & 3 & -4 & -3 & 0 & 0 & -3 & 0 & 0 \\ 0 & 0 & 9 & 5 & 3 & 3 & 8 & 3 & 0 \\ 0 & 0 & -5 & -2 & 0 & -3 & -4 & 0 & 0 \\ 0 & 0 & -4 & -3 & 0 & -1 & -3 & -1 & 0 \\ 0 & 0 & 4 & 3 & 1 & 3 & 3 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & -3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 3 \end{pmatrix},$$

where A is a Gram matrix of E_8 ($\det A = 1$), $\Gamma = C\mathbb{Z}^9$, $|\det C| = 7203 = 3 \cdot 7^4$, and the Smith form of C is $\text{diag}(1, 1, 1, 1, 1, 7, 7, 7, 21)$. Unlike $\mathbb{R}^7$, the a priori bound $\text{diam}^2 V_0 \leq 6 + t^2$ gives only $R^2 \leq 1.8306 > 7/4$, so the claim rests on the complete list of vertices of the nine-dimensional cell; it became computable after an exact unimodular change of basis.

Theorem 3.4. For $\Lambda = \mathbb{Z}^9$ with the form (8) and $\Gamma = C\mathbb{Z}^9$,

$$D(\Gamma)^2 = 7, \quad \text{diam}^2 V_0 = \frac{1119552292542693}{165294140000000}, \quad d^2 = \frac{1157058980000000}{1119552292542693}, \quad d = 1.0166127\dots,$$

so $\chi(\mathbb{R}^9, [1, \ell]) \leq 7203$ for all $1 \leq \ell < d$; in particular, $\chi(\mathbb{R}^9) \leq 7203$.

Proof. Among 4504 vectors within the bound $\|v\|^2 \leq 6 + t^2$ one finds 376 pairs, i.e. 752 facets (of at most 1022; an independent enumeration over $\Lambda/2\Lambda$ without any bound gives the same list, and a missed facet could only weaken the bound). Exact closure along the 1-skeleton yields 1 654 230 vertices (1 649 640 simple, 4320 with fifteen and 270 with sixteen active facets) and $R^2 = 1119552292542693/661176560000000 = 1.69327281\dots < 7/4$. The safe window $\|v\|^2 \leq 2(D_*^2 + \text{diam}^2 V_0) = 2276611272542693/826470700000000$ contains 280 vectors

(140 pairs), each with an exact KKT certificate; the minimum 7 is attained at the 240 horizontal vectors (120 pairs), the images of the minimal vectors of E_8 , and no layer vector violates the threshold. Margin at $\ell = 1$: $D(\Gamma)^2 - \text{diam}^2 V_0 = 37506687457307/165294140000000 = 0.2269 \dots > 0$; at $\ell = 101/100$ it is still $150036863771988707/165294140000000000 > 0$. $\square$

The bound 7203 supersedes $\chi(\mathbb{R}^9) \leq 9604$ of Corollary 4.8 in both parameters (fewer colors, wider segment: $1.0166 \dots$ against $\sqrt{63/61} = 1.0162 \dots$).

4 The Eisenstein identity and the product rule

A lattice $\Lambda \subset \mathbb{R}^n$ is *Eisenstein* if it carries a multiplication by $\omega = e^{2\pi i/3}$ acting as an isometry without nonzero fixed vectors; then $\alpha\Lambda$ has index $|\alpha|^n$ for $\alpha \in \mathbb{Z}[\omega]$. The ABPR constructions with $7^{n/2}$ colors in dimensions 4, 6, 8, 24 are $\Gamma = (3+\omega)\Lambda$ ($|3+\omega|^2 = 7$) on D_4 , E_6^* , E_8 and the Leech lattice Λ_{24} [1]. Let $H_1 \subset \mathbb{C}$ be the Voronoi cell of $\mathbb{Z}[\omega]$: the regular hexagon with facets at distance $1/2$ and vertices at distance $1/\sqrt{3}$ from the origin.

Proposition 4.1 (lower bound via projection onto a plane). *Let Λ be Eisenstein, $w \in \Lambda \setminus \{0\}$, $P = \text{span}(w, \omega w)$ and $H = \|w\|H_1$ the Voronoi cell of $\mathbb{Z}[\omega]w \subset P$. Then $\pi_P(V_0) \subseteq H$ and $\text{dist}(\alpha w/2, V_0) \geq \|w\| \text{dist}(\alpha/2, H_1)$ for every $\alpha \in \mathbb{Z}[\omega]$. In particular $D((3+\omega)\Lambda) \geq \sqrt{7/3} \lambda_1(\Lambda)$.*

Proof. The six vectors $\pm w, \pm \omega w, \pm(1+\omega)w = \mp \omega^2 w$ lie in Λ and have length $\|w\|$; the half-spaces $\langle x, u \rangle \leq \|u\|^2/2$ over them contain V_0 and have normals in P , so $\pi_P(V_0)$ lies in the hexagon they cut out in P , which is H . The point $\alpha w/2$ lies in P and π_P is 1-Lipschitz. For $\alpha = 3 + \omega$, $\|w\| = 1$: the point $(5/4, \sqrt{3}/4)$ lies in the normal cone of the vertex $(1/2, 1/(2\sqrt{3}))$ of H_1 , at squared distance $9/16 + 1/48 = 7/12$; apply Lemma 2.2. $\square$

Lemma 4.2 (a point in the cell). *Let $u_1, \dots, u_s$ be shortest vectors of Λ , $q = \sum_i c_i u_i$ with $c_i \geq 0$, $\sum_i c_i \leq 1$, and $\langle q, u_j \rangle \leq \frac{1}{2} \lambda_1^2$ for all j . Then $q \in V_0$.*

Proof. $q \in V_0$ iff $\langle q, r \rangle \leq \frac{1}{2} \|r\|^2$ for all $r \in \Lambda$. For $r \neq u_i$, $\|r - u_i\|^2 \geq \lambda_1^2 = \|u_i\|^2$ gives $\langle r, u_i \rangle \leq \frac{1}{2} \|r\|^2$; hence $\langle q, r \rangle \leq (\sum_i c_i) \frac{1}{2} \|r\|^2 \leq \frac{1}{2} \|r\|^2$ for $r \notin \{u_i\}$, and for $r = u_j$ it is the hypothesis. $\square$

Proposition 4.3 (upper bound). *Let Λ be Eisenstein and w a shortest vector. Then the six vertices $\pm \omega^k q$, $q = \frac{1}{3}(2w + \omega w)$, of $H = \|w\|H_1$ lie in V_0 ; hence $V_0 \cap P = H$ and $\text{dist}(\alpha w/2, V_0) = \|w\| \text{dist}(\alpha/2, H_1)$ for all $\alpha \in \mathbb{Z}[\omega]$.*

Proof. From $w + \omega w = -\omega^2 w$ we get $\|w + \omega w\| = \|w\|$, i.e. $\langle w, \omega w \rangle = -\frac{1}{2} \lambda_1^2$. Then $\langle q, w \rangle = \frac{1}{3}(2 - \frac{1}{2}) \lambda_1^2 = \frac{1}{2} \lambda_1^2$ and $\langle q, \omega w \rangle = \frac{1}{3}(-1 + 1) \lambda_1^2 = 0$, so $q \in V_0$ by Lemma 4.2 ($q = \frac{2}{3}w + \frac{1}{3}\omega w$); the other vertices follow by replacing w with $\pm \omega^k w$. By convexity $H \subseteq V_0 \cap P \subseteq \pi_P(V_0) \subseteq H$ (Proposition 4.1), and the distance from $\alpha w/2 \in P$ to V_0 is squeezed between its distances to H from both sides. $\square$

Theorem 4.4 (Eisenstein identity). *For an Eisenstein lattice Λ and every $\alpha \in \mathbb{Z}[\omega] \setminus \{0\}$,*

$$D(\alpha\Lambda) = 2 \lambda_1(\Lambda) \cdot \text{dist}(\alpha/2, H_1), \quad (9)$$

and for $|\alpha| > 1$ the minimum is attained exactly at αw with w shortest. In particular

$$D((3+\omega)\Lambda)^2 = \frac{7}{3} \lambda_1(\Lambda)^2, \quad d(\Lambda, (3+\omega)\Lambda) = \frac{\sqrt{7/3}}{\rho(\Lambda)}, \quad \rho(\Lambda) = \frac{2R}{\lambda_1(\Lambda)}, \quad (10)$$

and $D(m\Lambda) = (m-1)\lambda_1(\Lambda)$ for every integer $m \geq 2$.

Proof. For $v = \alpha w$, Lemma 2.2 and Proposition 4.1 give $D(v) \geq 2\|w\| \text{dist}(\alpha/2, H_1) \geq 2\lambda_1 \text{dist}(\alpha/2, H_1)$, with equality for shortest w by Proposition 4.3. If $|\alpha| > 1$ then $|\alpha/2| \geq \sqrt{3}/2 > 1/\sqrt{3}$, so $\alpha/2 \notin H_1$ and the first inequality is strict for non-shortest w . For $\alpha = 3 + \omega$ the squared distance is $7/12$; for $\alpha = m$ the closest point of H_1 to $m/2$ is the facet midpoint $1/2$. $\square$

Corollary 4.5 (exact widths). *D_4 , E_6^* , E_8 and Λ_{24} have $\rho = \sqrt{2}$ [5], so each ABPR coloring with $7^{n/2}$ colors has width exactly $\sqrt{7/6}$: $\chi(\mathbb{R}^n, [1, \ell]) \leq 7^{n/2}$ for $\ell < \sqrt{7/6}$, $n = 4, 6, 8, 24$, and $\sqrt{7/6}$ cannot be improved within these constructions. The Coxeter–Todd lattice K_{12} has $\rho = \sqrt{8/3}$, so $d(K_{12}, (3+\omega)K_{12}) = \sqrt{7/8} < 1$ — no 7^6 -coloring of $\mathbb{R}^{12}$ this way (a partial negative answer to Open question 2 of [1]) — while $d(K_{12}, 3K_{12}) = 2/\rho = \sqrt{3/2}$, so $\chi(\mathbb{R}^{12}, [1, \ell]) \leq 3^{12}$ for $\ell < \sqrt{3/2}$.*

The threshold $\sqrt{7/3}$ appears in [1, Remark 9] as a sufficient condition $\rho \leq \sqrt{7/3}$ whose users were unknown; (10) shows it is also necessary and gives the exact width of every such lattice, and it turns the horizontal floors $D_{\min} = \sqrt{7}$ in Theorems 3.3 and 3.4 into theorems.

Proposition 4.6 (product rule). *Let $\Lambda = \bigoplus_{i=1}^s \Lambda_i$ be an orthogonal direct sum, $\Gamma = \bigoplus_i \Gamma_i$, $\Gamma_i \subseteq \Lambda_i$, $d_i = d(\Lambda_i, \Gamma_i)$, $k_i = [\Lambda_i : \Gamma_i]$. The maximum of $d(\Lambda, \Gamma)$ over independent rescalings of the blocks equals $(\sum_i d_i^{-2})^{-1/2}$, attained when $\text{diam}^2 V_0(\Lambda_i) \propto d_i^{-2}$; hence $\chi(\mathbb{R}^n, [1, \ell]) \leq \prod_i k_i$ for all $\ell < (\sum_i d_i^{-2})^{-1/2}$, $n = \sum_i n_i$.*

Proof. The cell of a direct sum is the product of the cells, so $\text{diam}^2 V_0 = \sum_i s_i$ with $s_i = \text{diam}^2 V_0(\Lambda_i)$, and $D(v)^2 = \sum_i D_i(v_i)^2$ by Lemma 2.2. Since $D_i(0) = 0$, the minimum over $\Gamma \setminus \{0\}$ is attained at a vector with one nonzero component, so $d^2 = \min_i (d_i^2 s_i) / \sum_j s_j$. With $t_i = d_i^2 s_i$, $\sum_j s_j = \sum_j t_j d_j^{-2} \geq \min_i t_i \sum_j d_j^{-2}$, with equality iff all t_i coincide, i.e. $s_i \propto d_i^{-2}$. Apply Proposition 2.1. $\square$

Width is thus an expendable resource: a block of width d_i costs $1/d_i^2$ out of a budget of 1, and blocks may have different indices.

Theorem 4.7. $\chi(\mathbb{R}^{10}, [1, \ell]) \leq 2401 \cdot 19 = 45619$ and $\chi(\mathbb{R}^{26}, [1, \ell]) \leq 19 \cdot 7^{12}$ for all $\ell < \sqrt{217/214}$; in particular $\chi(\mathbb{R}^{10}) \leq 45619$ and $\chi(\mathbb{R}^{26}) \leq 19 \cdot 7^{12}$.

Proof. The block $E_8/2401$ has $d_1^2 = 7/6$ (Corollary 4.5), cost $6/7$. The planar block $\Lambda_2 = \mathbb{Z}[\omega]$ ($V_0 = H_1$, $\text{diam}^2 V_0 = 4/3$), $\Gamma_2 = (5 + 2\omega)\mathbb{Z}[\omega]$ of index $N(5 + 2\omega) = 19$: by (9), $D(\Gamma_2) = 2 \text{dist}((5 + 2\omega)/2, H_1)$, and the point $(2, \sqrt{3}/2)$ lies in the normal cone of the vertex $(1/2, 1/(2\sqrt{3}))$ at squared distance $9/4 + 1/3 = 31/12$, so $D(\Gamma_2)^2 = 31/3$, $d_2^2 = 31/4$, cost $4/31$. Total $6/7 + 4/31 = 214/217 < 1$. The Leech lattice block $\Lambda_{24}/7^{12}$ has the same width $\sqrt{7/6}$. $\square$

Corollary 4.8. $\chi(\mathbb{R}^9, [1, \ell]) \leq 2401 \cdot 4 = 9604$ and $\chi(\mathbb{R}^{25}, [1, \ell]) \leq 4 \cdot 7^{12}$ for all $\ell < \sqrt{63/61}$.

Proof. The one-dimensional block $\mathbb{Z} \supset 4\mathbb{Z}$ has cells of length 1 at same-color distance 3, so $d = 3$ and cost $1/9$; $6/7 + 1/9 = 61/63 < 1$. $\square$

The previous bounds were 3^{10} , 3^{25} , 3^{26} and 17253 [1]; 45619 is the first bound in $\mathbb{R}^{10}$ below the classical 3^n tower. Three colors in the last block fail ($d = 2$, $6/7 + 1/4 > 1$): the budget $1/7$ left by $E_8/2401$ requires $d \geq \sqrt{7}$, which a planar lattice block first reaches (numerically) at index 19.

5 An index floor on E_8

Lemma 5.1 (inradius bound). $D(v) \leq \|v\| - \lambda_1(\Lambda)$ for $\|v\| \geq \lambda_1(\Lambda)$.

Proof. The ball of radius $\lambda_1/2$ lies in V_0 , so $\text{dist}(v/2, V_0) \leq \|v\|/2 - \lambda_1/2$. $\square$

Proposition 5.2 (shell floor). *If $D(v) < \text{diam} V_0$ for every $v \in \Lambda \setminus \{0\}$ with $\|v\|^2 < s^*$, then every $\Gamma \subset \Lambda$ defining a proper coloring satisfies $[\Lambda : \Gamma] \geq (s^*)^{n/2} / (\gamma_n^{n/2} \det \Lambda)$, γ_n the Hermite constant.*

Proof. Properness forces $\lambda_1(\Gamma)^2 \geq s^*$; Hermite's inequality gives $\lambda_1(\Gamma)^n \leq \gamma_n^{n/2} \det \Gamma = \gamma_n^{n/2} [\Lambda : \Gamma] \det \Lambda$. $\square$

Lemma 5.1 always allows $s^* = (\text{diam} V_0 + \lambda_1)^2$; since norms run over shells, s^* jumps to the next shell whenever the first shell above this threshold is forbidden entirely, which is certified by a one-line witness.

Lemma 5.3 (radial witness). *Let $\langle \Lambda, \Lambda \rangle \subseteq \frac{1}{e}\mathbb{Z}$, $\|v\|^2 = N$, and let a rational $s \in (0, \frac{1}{2})$ satisfy $s \lfloor e\sqrt{NM} \rfloor / e \leq M/2$ for every shell $M = \|u\|^2 < 4s^2 N$. Then $sv \in V_0$ and $D(v) \leq (1 - 2s)\sqrt{N}$.*

Proof. We need $s\langle v, u \rangle \leq \|u\|^2/2$ for all $u \in \Lambda$. For $M \geq 4s^2 N$ this is Cauchy-Schwarz, $s\sqrt{NM} \leq M/2$; for $M < 4s^2 N$, $\langle v, u \rangle$ is a multiple of $1/e$ not exceeding $\sqrt{NM}$, and the hypothesis applies. Then $D(v) \leq 2\|v/2 - sv\|$ by Lemma 2.2. $\square$

Theorem 5.4 (2401 is unimprovable on E_8). *If $\Gamma \subset E_8$ defines a proper coloring, then $[E_8 : \Gamma] \geq 2401$, with equality for $\Gamma = (3 + \omega)E_8$.*

Proof. For E_8 : $\lambda_1^2 = 2$, $R = 1$, $\text{diam} V_0 = 2$, $\det = 1$, integral and even. Lemma 5.1 forces $\|v\|^2 \geq (2 + \sqrt{2})^2 > 11$, so the first candidate shell is 12. For $\|v\|^2 = 12$ apply Lemma 5.3 with $e = 1$, $s = 1/4$: $4s^2 N = 3$, only $M = 2$ needs a check, and $\frac{1}{4}\lfloor \sqrt{24} \rfloor = 1 = M/2$; hence $D(v) \leq \frac{1}{2}\sqrt{12} = \sqrt{3} < 2$, the shell 12 is forbidden, $s^* = 14$, and Proposition 5.2 with $\gamma_8 = 2$ [2] gives $[E_8 : \Gamma] \geq 14^4/2^4 = 2401$. The sublattice $(3 + \omega)E_8$ has index 2401 and is proper by Corollary 4.5. $\square$

Coulson’s general bound $\chi_s \geq 2^{n+1} - 1$ [6, 1] gives only 511 for E_8 , so Theorem 5.4 answers Open question 8 of [1] for this lattice. The same argument (shells 4, 19/4, 5 of the bcc lattice, witness $e = 4$, $s = 1/4$) reproves Coulson’s floor 15 on the bcc lattice from its norm spectrum alone, and gives the floors 305 on E_6^* (record 343) and 5 688 009 064 on the Leech lattice (record 7¹²; $\gamma_{24} = 4$ [4]). On lattices in general position the norm spectrum is almost continuous and the screen is silent; whether 43 is minimal in $\mathbb{R}^4$ remains open.

6 Reproducibility and provenance

All code, exact certificates and data are at <https://github.com/ivaleo/Chromatic> (MIT license). Run from `audit-data/`, the command `python -m chromatic_research.campaigns.verify_main_results` recomputes the final inequalities of all five rows of Table 1 from the published fractions in rational arithmetic; with `-full` it also runs the complete verifiers of the rows 4, 5, 7 and 9 (the last enumerates 1 654 230 vertices and takes hours). The certificates are `results/dim4_k43_verify.json` and `results/cert43_eisenstein.json` (Theorem 3.1), `results/metric_deform_a5_132_refined_certificate.json` (Theorem 3.2), `results/dim7_1029_exact.json` (Theorem 3.3) and `results/dim9_7203_exact.json` (Theorem 3.4); each is self-contained (rational Gram form, sublattice, exact R^2 , complete window, KKT certificates) and can be re-verified by a short independent program. Every claim above is a theorem or an exact certificate; numerical results without proof strength are confined to the repository.

Contributions and the use of AI. The problem setting, the method and the computational pipeline, the index-132 construction, lamination, the product calculus and the lower bound via projection onto a plane are due to L.L. Ivanov. N.V. Glushkova found the index-43 lattice (the first point of the family (4)), proved Proposition 4.3, and wrote the exact verifiers of the laminations that turned 1029 and 7203 into Theorems 3.3 and 3.4 (with the decisive unimodular change of basis in $\mathbb{R}^9$). The symmetry-restricted search, the independent re-verifications and the shell floor are joint. The computational pipeline, campaign scripts and first drafts of the text were produced with systematic use of an AI assistant (general-purpose large language models in an agentic coding environment); problem statements, search directions, acceptance criteria and final verification belong to the authors, and the theorems do not depend on the provenance of the search code. Responsibility for all claims rests with the authors.

References

- [1] A. Arman, A. Bondarenko, A. Prymak, D. Radchenko, *Upper bounds on chromatic number of $\mathbb{E}^n$ in low dimensions*, Electron. J. Combin. **31** (2024), no. 2, #P2.35; arXiv:2112.13438.
- [2] H. F. Blichfeldt, *The minimum values of positive quadratic forms in six, seven and eight variables*, Math. Z. **39** (1935), 1–15.
- [3] L. I. Bogolubsky, A. M. Raigorodskii, *A remark on lower bounds for the chromatic numbers of spaces of small dimension with metrics ℓ_1 and ℓ_2* , Math. Notes **105** (2019), no. 2, 180–203.
- [4] H. Cohn, A. Kumar, *Optimality and uniqueness of the Leech lattice among lattices*, Ann. of Math. (2) **170** (2009), no. 3, 1003–1050.
- [5] J. H. Conway, N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Springer, New York, 1999.
- [6] D. Coulson, *A 15-colouring of 3-space omitting distance one*, Discrete Math. **256** (2002), 83–90.
- [7] A. D. N. J. de Grey, *The chromatic number of the plane is at least 5*, Geombinatorics **28** (2018), no. 1, 18–31.
- [8] G. Exoo, D. Ismailescu, M. Lim, *On the chromatic number of $\mathbb{R}^4$* , Discrete Comput. Geom. **52** (2014), no. 2, 416–423.
- [9] L. L. Ivanov, *An estimate for the chromatic number of the space $\mathbb{R}^4$* , Russian Math. Surveys **61** (2006), no. 5, 984–986.
- [10] L. L. Ivanov, *On the chromatic numbers of $\mathbb{R}^2$ and $\mathbb{R}^3$ with intervals of forbidden distances*, Bulletin of PFUR, Ser. Mathematics, Information Sciences, Physics (2011), no. 1, 14–29 (in Russian).
- [11] R. Radoičić, G. Tóth, *Note on the chromatic number of the space*, in: Discrete and Computational Geometry, Algorithms Combin. **25**, Springer, 2003, 695–698.

- [12] A. M. Raigorodskii, *Borsuk's problem and the chromatic numbers of some metric spaces*, Russian Math. Surveys **56** (2001), no. 1, 103–139.
- [13] A. M. Raigorodskii, *Coloring distance graphs and graphs of diameters*, in: Thirty Essays on Geometric Graph Theory (J. Pach, ed.), Springer, New York, 2013, 429–460.
- [14] A. M. Raigorodskii, *Cliques and cycles in distance graphs and graphs of diameters*, in: Discrete Geometry and Algebraic Combinatorics, Contemp. Math. **625**, Amer. Math. Soc., Providence, RI, 2014, 93–109.
- [15] A. Soifer, *The Mathematical Coloring Book*, Springer, New York, 2009.