

Geometric Root Calculators

Two fifth-order solvers and their construction cost

Ivan Besevic¹ · Independent research

Version 23 · 20 September 2026

<https://github.com/ivan-fr/pandrosion>

Abstract. We give two geometric realizations of fifth-order iterations for positive roots. A prepared fixed circle inverts a quadratic Padé model. Its selected real branch strictly reduces logarithmic error and alternates across the root; a short convexity argument proves convergence throughout the branch for every real $p > 2$. For integer degrees, a regular update uses $2p + 2$ joins, including the power stage. A second construction uses a decentered arc and converges monotonically from every positive start, also with exact order five. We compare their formulas with classical rational and quadratic approximation, derive explicit mobile-work bounds, and give a global circle–AD controller. Lean certificates and independent incidence computations support the analysis. The contribution is the specified constructions, their domains and their costs; fifth-order approximation itself is classical.

1 Two solvers, one construction problem

A formula specifies arithmetic; a geometric calculator must realize it. A product can be a pair of projections, while a square root can be a circle crossing. We ask how cheaply a positive-root correction can be realized with prepared lines and circles, and on which states the resulting iteration converges.

The paper has three main results, stated here with their precise roles:

1. **Fixed-circle solver.** A quadratic inverse model has exact order five and strictly contracts logarithmic error on its entire real branch (Theorems 4.1 and 4.3). The proof works uniformly for real $p > 2$.
2. **Decentered radical solver.** A moving arc realizes a different fifth-order correction, globally monotone on $(0, \infty)$ (Theorem 5.1). Its support uses the same mobile operations as the centered order-three AD support.
3. **Explicit geometric cost.** For integer $p \geq 3$, the prepared circle needs $2p + 2$ joins in its finite regular chart (Theorem 4.2). AD entry and chart fallback give a global algorithm with separately counted control work (Proposition 7.1).

The two dynamics are deliberately distinguished: circle steps cross the root with decreasing error; arc steps stay on the initial side. Rational Padé supplies a classical fifth-order comparator. Section 2 identifies the inherited approximation theory before the constructions are developed. The classical rectangle supports, historical reference and exploratory analog implementation are retained in the appendices.

¹Computational assistance: AI was used for derivations, proof development, figures and manuscript preparation. The author is responsible for the claims and their attribution. Section 8 distinguishes exact proofs, numerical checks and exploratory displays.

2 Position relative to existing methods

2.1 Classical formulas and geometric interpretation

High local order is not a novelty claim. Traub’s Padé organization of iteration formulas already unifies several classical schemes and their error terms [1, 2]. Gander treats both rational Halley and the radical Euler correction obtained by solving a quadratic Taylor model [3, pp. 131–133]. Scavo–Thoo develop the geometry of Halley’s method [4]; Melman connects geometric interpretation with global convergence [5]. Geometric constructions for nonlinear iteration also appear in Amat–Busquier–Gutiérrez [6]. Thus neither geometric iteration, global convergence via geometry, nor a radical correction is claimed here as a new general principle.

Our normalization is $a = X^{-1/p}$, $t = Xs^p$ and $s^+ = s\Phi(t)$. With $x = s/a$, the normalized iteration is $x^+ = x\Phi(x^p)$. Centered AD is exactly the classical h_{11} , projective AD is h_{12} , and the rational fifth-order baseline is h_{22} in the Padé family of Laszkiewicz–Ziętak [7, 8]; the latter two formulas are printed on slide 11 of the authors’ presentation. Karp–Lin prove convergence results for this rational family [9]. We use these maps as identified baselines, with new coordinates and explicit incidence inventories where applicable.

Table 1: Construction-by-construction attribution. The last column specifies what is analyzed here, rather than asserting absence from all prior literature.

Device	Closest classical object	Contribution evaluated here
AK / centered AD	Newton under inversion / Halley h_{11}	Shared rectangle support and counted readout; scalar formulas inherited.
Projective AD	Rational Padé h_{12}	Two-join support and its finite chart; scalar formula inherited.
Rational fifth order	Diagonal Padé h_{22}	Two factored four-join modules; explicit comparison protocol.
Fixed circle	Inverse of a $[2/2]$ Padé model; quadratic approximation	Prepared circle and branch selector; whole-branch dynamics and $2p + 2$ -join realization.
Decentered arc	Radical / quadratic approximation	Explicit displaced support and the global monotone theorem for (10).

The construction question also differs from existence under the Poncelet–Steiner theorem: a fixed circle and its center allow straightedge constructions [10], but a particular iterative protocol still needs its coordinates, admissible branch and work inventory. Our inventory follows the constructive concerns of géométrie and nomography [11, 12]; it is an upper bound under stated primitives, not a minimum-trace theorem.

2.2 Inverse Padé and quadratic approximation

The order-five rational comparator approximates $t^{-1/p}$ directly. The fixed circle instead approximates v^{-p} by $N(v)/D(v)$ and then solves $N(v) - tD(v) = 0$. These operations are not interchangeable. With the coefficients defined in (4), writing the circle equation as

$$(\gamma_p - t\alpha_p)v^2 + 2\beta_p(t-1)v + (\alpha_p - t\gamma_p) = 0$$

places it within the general algebraic form $A(t)v^2 + B(t)v + C(t) = 0$ associated with quadratic approximation. Shafer already develops quadratic approximation as an alternative to rational Padé when square roots are available [13]. The use of such an algebraic approximant therefore has a direct classical precedent.

The decentered correction belongs to the same broad algebraic class. Put

$$S(t) = (p-2)(t^2+1) + (10p+4)t.$$

Eliminating the positive radical from (10) gives

$$(p-1)^2 S(t)(v-1)^2 - 12p(tv-1)^2 = 0. \quad (1)$$

Squaring loses the sign information: the explicit positive radical in (10), rather than this polynomial alone, defines the iteration. In particular the equation has a double root at $t = 1$. We do not identify it with a particular normalized Shafer approximant solely from its quadratic form.

There is also a direct, checkable distinction between the three fifth-order maps in the same logarithmic error $e = \log(s/a)$:

Correction	Coefficient of e^5 in e^+
Fixed-circle inverse	$-(p^2-1)(p^2-4)/720$
Decentered arc	$(p-2)(p^2-1)(3p+1)/1440$
Rational diagonal Padé	$(p^2-1)(4p^2-1)/720$

The coefficients are pairwise distinct for $p > 2$. This proves nonidentity with the specified rational baseline in the stated normalization; it is not a classification of all algebraic iterations or coordinate conjugacies.

Comparison with tangent-circle methods. Kanwar–Singh–Bakshi include osculating-circle and ellipse constructions among their quadratic and cubic families [14]. Pakdemirli’s recent circular-arc algorithm uses the tangent circle with the function’s local curvature [15]. Our fixed circle is prepared once for the degree; its intersections encode a residual-dependent quadratic solve. Our decentered arc also has a fixed center, with radius prescribed by the residual. Neither is obtained by fitting the curvature of the graph at the current state. This construction-level distinction does not claim priority over every possible circle encoding.

Scope of the contribution. We claim the explicit incidence realizations and prove the stated branch/global dynamics and protocol bounds. Their formulas are positioned inside classical approximation theory. The source ledger (LITERATURE_REVIEW.md) records the full texts, formulas and abstracts consulted. A bounded literature comparison cannot establish historical priority, and no universal “first geometric solver”, optimal trace count or electrical speedup is asserted.

3 Coordinates as reusable computing elements

Let $X > 0$, let $p \geq 3$ be an integer, and write

$$a = X^{-1/p}, \quad t = Xs^p, \quad s^+ = s\Phi(t), \quad s > 0.$$

The state approaches the reciprocal root a ; inversion supplies the desired $X^{1/p}$. We use $e = \log(s/a)$ for local error. Points are capital letters, scalar iteration maps are $\mathcal{T}_{\text{label}}$, and polynomial coefficients use Greek letters. Thus a point and a correction function never share the same symbol.

In a rectangle with vertices $C = (0, 0)$, $B = (W, 0)$, $O = (0, H)$ and $A = (W, H)$, define two extended rails and a family of top centers:

$$R(q) = (W, H(1 - q)), \quad L(q) = (0, H(1 - q)), \quad \mathcal{U}_f = \left(\frac{Wf}{f-1}, H \right). \quad (2)$$

A line through the prepared center performs multiplication in one direction and division in the other:

$$\mathcal{U}_f R(q) \cap OC = L(fq), \quad \mathcal{U}_f L(b) \cap AB = R(b/f). \quad (3)$$

For example, substituting the first two points into a line equation and setting $x = 0$ gives ordinate $H(1 - fq)$. This is the whole arithmetic content of the rail device.

3.1 The power stage

Start at $P = R(s)$. The join CP meets the top line at $V = \mathcal{U}_{1/s}$. A leftward projection through $\mathcal{U}_{1/2}$ followed by a return through V changes a rail value q to $sq/2$. After $p - 1$ pairs, the endpoint is $Q = R(s^p/2^{p-1})$. Including the initial join, the stage uses $2p - 1$ lines. Its normalization factor is part of the design, not discarded arithmetic.

More generally, choosing the leftward factors $f_1, \dots, f_{p-1}$ gives an endpoint with value $s^p \prod_i f_i$. This allows the power stage to feed a device with its preferred residual scale. We use exactly this freedom for the fixed-circle calculator.

Power protocols, one state encoding. The later rectangle drawings use a different power stage: similar triangles and parallels produce $E = R(s^p)$ directly. They do not change the meaning of $R(s)$. The rail telescope instead produces a scaled power; its scale is absorbed into the prepared readout. Their costs must be kept separate.

Power stage	Geometric mechanism	Output and use
Rail telescope	$2p - 1$ joins; repeated projection	Scaled s^p ; pencils and fixed circle
Triangle chain	Horizontals and similar-triangle parallels	$R(s^p)$; original MA/AK/AD drawings
Binary products	Repeated squaring and multiplication	$R(s^p)$; large degrees and safe controllers

3.2 A precise notion of work

We count a join as J , a parallel-construction macro as $P_{\parallel}$, and a mobile compass arc as C_{arc} . Intersections of already drawn objects, fixed preparation and final inversion are excluded. When expanded into ruler-and-compass traces, our parallel macro costs $2J + 3C_{\text{arc}}$. These conventions are specified before any cost comparison.

The name Pandrosion refers to the addressee of the opening of Book III of Pappus' *Collection* [16]. Our rectangle follows a modern reconstruction [17]; the accelerated supports are modern constructions, not attributed to the ancient text.

4 A prepared circle as a quadratic solver

Suppose the desired correction factor is $v \approx 1$. Instead of evaluating a rational approximation to $t^{-1/p}$, approximate the inverse relation $t = v^{-p}$ by a quadratic rational model and solve it exactly. Put

$$\begin{aligned} \alpha_p &= (p+1)(p+2), & \beta_p &= p^2 - 4, & \gamma_p &= (p-1)(p-2), \\ N(v) &= \gamma_p v^2 - 2\beta_p v + \alpha_p, & D(v) &= \alpha_p v^2 - 2\beta_p v + \gamma_p. \end{aligned} \quad (4)$$

The model is $\mathcal{R}(v) = N(v)/D(v)$, the $[2/2]$ Padé approximant of v^{-p} at one. Its inverse branch $v(1) = 1$ defines $\mathcal{T}_{\text{circle}}(s) = sv(Xs^p)$. A circle can supply the quadratic root in a single crossing.

4.1 The eight-line cubic example

Use $W = 2, H = 4, p = 3$ and $X = 2$. Prepare a circle through B with center O_Γ , a projector Z , and constants

$$O_\Gamma = (-152/113, 126/113), \quad Z = (478/113, 396/113), \quad \rho = \frac{1}{2}, \quad k = \frac{5}{113}.$$

The first five joins form the rail power stage with left factors ρ and $\sigma = 20/113$, retaining $L_1 = L(\rho s)$ and ending at $T = R(kXs^3)$. The last three joins are the main device:

1. Draw ZT and select its prescribed intersection G with the fixed circle.
2. Draw BG to the top line, obtaining $U = \mathcal{U}_{\rho/v}$.
3. Draw UL_1 back to the right rail. Its value is $(\rho s)/(\rho/v) = sv$.




Figure 1: The fixed-circle device at $s = 3/4$. Left: the sixth join supplies G , and the seventh supplies the readout center U . Right: an independently enlarged view shows the eighth join and distinguishes P from P^+ . The circle is drawn once, before iteration. Equal horizontal and vertical scales are used within each view.

The computed next state is $0.793700551706633\dots$, with relative error about 3.24×10^{-8} . The circle is doing arithmetic: it replaces a longer rational readout by the solution of a quadratic model.

4.2 Why the inverse correction has order five

At $v = 1$, $N = D = 12$. The polynomial certificate for local order is

$$pND + v(N'D - ND') = p\alpha_p\gamma_p(v-1)^4. \quad (5)$$

Moreover $\gamma_p N = (\gamma_p v - \beta_p)^2 + 3(p^2 - 4)$ and $\alpha_p D = (\alpha_p v - \beta_p)^2 + 3(p^2 - 4)$, so the logarithms used below are well defined for $p > 2$.

Theorem 4.1 (Local inverse-circle order). *For real $p > 2$, the inverse branch through $v = 1$ exists near $t = 1$ and satisfies*

$$e^+ = -\frac{(p^2 - 1)(p^2 - 4)}{720}e^5 + O(e^7). \quad (6)$$

It therefore alternates sides locally and has exact order five.

Proof. Set $v = 1 + u$, $E(u) = \log \mathcal{R}(1 + u)/p$ and $K(u) = E(u) + \log(1 + u)$. Equation (5) gives

$$K'(u) = \frac{\alpha_p \gamma_p u^4}{(1 + u)N(1 + u)D(1 + u)}.$$

Thus $K(u)/u^5 \rightarrow \alpha_p \gamma_p / 720$, while $E(0) = 0$ and $E'(0) = -1$. The inverse function theorem identifies $E(u)$ with the input logarithmic error. Dividing the limits gives the coefficient in (6). Analyticity and $\mathcal{R}(1/v) = 1/\mathcal{R}(v)$ make the logarithmic update odd, giving the stated remainder. The coefficient is nonzero for $p > 2$. $\square$

Equation (5) specializes a classical Padé differential-defect identity [9, proof of Theorem 2.4]; integrating it proves contact with v^{-P} .

4.3 Which circle intersection?

The discriminant and the descending branch $\mathcal{R}'(v) < 0$ are

$$\begin{aligned} \Delta_p(t) &= (42p^2 - 24)t - 3(p^2 - 4)(1 + t^2), \\ \tau_{\pm} &= \frac{7p^2 - 4 \pm 4p\sqrt{3(p^2 - 1)}}{p^2 - 4}, \quad \tau_- \tau_+ = 1, \\ v_{\pm} &= \frac{p^2 + 2 \pm \sqrt{12(p^2 - 1)}}{p^2 - 4}. \end{aligned}$$

The transverse real interval is $\tau_- < t < \tau_+$; select its inverse with $v_- < v < v_+$. Positivity alone does not select this branch: the figure exhibits two positive roots at the same residual. At $\Delta_p(t) = 0$ they merge tangentially. Theorem 4.3 proves contraction in the open band; AD handles tangencies and starts outside it.




Figure 2: The fold of $\mathcal{R}$ for $p = 3$. Blue marks the descending branch through $(1, 1)$; gray marks the other branches. The level $t = 0.05$ has two positive preimages. Only $v \approx 3.1183$ belongs to the selected branch; $v \approx 6.3817$ must be rejected. Open circles mark the tangencies.

4.4 The general-degree construction

For each integer $p \geq 3$, prepare $O_\Gamma = (h, j)$, $Z = (z_x, z_y)$, $k > 0$ and $\rho > 0$, using the rational formulas in Appendix A. The circle passes through $B = (2, 0)$ and is independent of X and s . Put $\sigma = kX2^{p-3}/\rho$. The rail sequence uses

$$f_1 = \rho, \quad f_i = \frac{1}{2} \quad (2 \leq i \leq p-2), \quad f_{p-1} = \sigma.$$

It saves $L_1 = L(\rho s)$ and ends at $T = R(kt)$, because $\prod_i f_i = kX$. The same three final joins then give $R(sv)$.

Theorem 4.2 (Counted finite-circle protocol). *For the preparation in Appendix A, assume $X, s > 0$, $\tau_- < Xs^p < \tau_+$, and*

$$s \neq 1, \quad \sigma \neq 1, \quad v \neq \rho, \quad b(v) = (2-h)2v - 4j(\rho - v) \neq 0.$$

The selected branch is realized by $2p + 2$ mobile joins using the prepared circle.

Proof. The power stage costs $1 + 2(p-1) = 2p-1$ joins. To certify the remaining incidence, write

$$d = (2v, 4(\rho - v)), \quad n = d \cdot d, \quad b = (B - O_\Gamma) \cdot d, \quad G = B - \frac{2b}{n}d.$$

The pencil line through B has direction d . Substituting $B + \xi d$ into the circle equation gives $\xi(\xi n + 2b) = 0$: besides B , the other intersection is exactly the displayed G . Thus $b \neq 0$ keeps the two points distinct. The same line meets the top at $\mathcal{U}_{\rho/v}$ when $v \neq \rho$.

It remains to show that the projector Z and the residual point T select this G . Three points are collinear exactly when the determinant of their displacement vectors vanishes. Multiplying that determinant by n gives

$$I = (2 - z_x)(-2bd_y - z_y n) - [4(1 - kt) - z_y][(2 - z_x)n - 2bd_x].$$

The prepared parameters satisfy $I = \lambda_p[N(v) - tD(v)]$, with $\lambda_p \neq 0$. This identity converts the geometric incidence into exactly the inverse model, rather than an approximation to it. Since the selected root has $N(v) - tD(v) = 0$, it places G on ZT . Projecting G through B , then L_1 through the resulting center, gives $R(sv)$ by (3). These are three joins. The branch arc is parameterized by $v_- < v < v_+$ and can be marked during preparation using its endpoints and $G(1)$. The listed hypotheses exclude an infinite center, a coincident B, G , and tangency; the other rail intersections are transverse. $\square$

The exclusions concern this finite chart. In particular, $s = 1$ sends the initial center V to infinity even though the scalar correction can remain regular. A positive rescaling $\tilde{X} = c^p X$, $\tilde{s} = s/c$ preserves t and conjugates every correction. A dyadic choice can keep the limiting state away from one, controlling the distant initial center near convergence. The circle-specific exclusions still require their own branch and chart checks.

The result is a concrete upper bound on mobile work, not merely an existence theorem for straightedge construction. Its advantage over the rational fifth-order baseline will be quantified after the other devices have been introduced.

4.5 Uniform contraction on the whole real branch

Here “contraction” means strict error reduction relative to the root. No pairwise Lipschitz constant is required. A Cayley coordinate makes the whole-branch argument independent of integer-degree estimates.

Theorem 4.3 (Whole-branch contraction and alternation). *Let $p > 2$ be real and $\tau_- < t < \tau_+$, $t \neq 1$. On the descending inverse branch $v = v(t)$, put $e = \log t/p$ and $e^+ = e + \log v$. Then*

$$ee^+ < 0, \quad |e^+| < |e|.$$

Consequently the scalar circle iteration stays in this residual band and converges to $X^{-1/p}$ from every start in it. Its exact local order is five.

Proof. By reciprocity it suffices to consider $v > 1$, so $t < 1$. Set

$$x = \frac{v-1}{v+1}, \quad x_* = \sqrt{\frac{3}{p^2-1}}, \quad y = \frac{3px}{3+(p^2-1)x^2}.$$

The branch condition becomes $0 < x < x_* < 1$. Direct substitution gives

$$\frac{N(v)}{D(v)} = \frac{1-y}{1+y}, \quad \log v = 2 \operatorname{artanh} x, \quad -\log t = 2 \operatorname{artanh} y. \quad (7)$$

Positivity of N, D gives $0 < y < 1$. The branch inequality also gives $y > px/2$; hence $0 < x < px/2 < 1$ because $p > 2$.

The function $f(z) = \operatorname{artanh} z$ is strictly convex on $(0, 1)$ and $f(0) = 0$; indeed $f''(z) = 2z/(1-z^2)^2 > 0$. Thus $f(cz) > cf(z)$ whenever $c > 1$ and $0 < cz < 1$. Applying this with $c = p/2$ yields the entire upper-error estimate in one line:

$$\operatorname{artanh} y > \operatorname{artanh}(px/2) > \frac{p}{2} \operatorname{artanh} x, \quad p \log v < -2 \log t. \quad (8)$$

This proves $e^+ < -e$. To prove that the step crosses the root, use the already established defect certificate (5):

$$\frac{d}{dv} \log(v^p \mathcal{R}(v)) = \frac{p\alpha_p \gamma_p (v-1)^4}{vN(v)D(v)} > 0 \quad (v > 1).$$

The logarithm is zero at $v = 1$, so $p \log v + \log t > 0$ and $0 < e^+ < -e$. Replacing (t, v) by $(1/t, 1/v)$ proves the other half.

For convergence put $z = \log t$ and $F(z) = \log[tv(t)^p]$. The inverse branch is continuous, $F(0) = 0$, and $|F(z)| < |z|$ for $z \neq 0$. Reciprocity makes the band symmetric in z . An orbit starting at z_0 stays in the compact interval $[-|z_0|, |z_0|]$ inside that band. If its decreasing absolute errors had positive limit ℓ , a convergent subsequence $z_{n_k} \rightarrow z_*$ would satisfy $|z_*| = |F(z_*)| = \ell$, contradicting the strict inequality. Thus $z_n \rightarrow 0$. Theorem 4.1 supplies the local order. $\square$

4.6 Endpoint margin and proof coverage

The same argument settles the endpoint inequality without degree splitting. At $x = x_*$, one has $y_* = px_*/2 < 1$, $v_+ = (1 + x_*)/(1 - x_*)$ and $\tau_+ = (1 + y_*)/(1 - y_*)$. Strict convexity gives

$$2 \log \tau_+ - p \log v_+ = 4 \operatorname{artanh}(px_*/2) - 2p \operatorname{artanh} x_* > 0.$$

Thus $v_+^p < \tau_+^2$ for every real $p > 2$. No rational brackets for $\sqrt{12(p^2 - 1)}$, finite list of degrees or exponential numerical bound is needed. Scalar endpoint contraction does not make a tangent crossing a regular geometric step; the controller still uses AD there.




Figure 3: Sampled contraction ratios and endpoint margins illustrate the proved inequalities. The left curves use $p = 3, 5, 17$; the horizontal line is the strict-error threshold one. The right curve samples degrees 3–400. Its limiting margin is $2 \log(7 + 4\sqrt{3}) - 2\sqrt{3} \approx 1.80373$. These samples do not establish a universal numerical contraction factor.

Lean coverage. Theorem 4.3 is formalized for every real $p > 2$. The modules `RectangleFixedCircleUniform`, `RectangleFixedCircleDynamics` and `RectangleFixedCircleTheorem43` prove the uniform logarithmic inequality, alternation, existence and continuity of the selected inverse, the explicit residual band, orbit convergence and the exact fifth-order limit in the input-error variable. Their 69 declarations are audited. The earlier integer-degree proof in `RectangleFixedCircleBranch` remains unchanged and independently audited.

5 A decentered arc with global fifth-order convergence

The fixed circle solves for a correction externally. A second architecture keeps the power endpoint $E = R(s^p)$ and changes the interception support inside the rectangle. Prepare $M = (0, H(1 - X^{-1}))$. The slope of ME is $m = H(1 - t)/(WX)$. A line through A of slope $k_s = H\kappa/(WX)$ meets the parallel to ME through $P = R(s)$ at T_A . The horizontal through T_A gives P^+ , with

$$s^+ = s \frac{k_s}{k_s - m} = s \frac{\kappa}{\kappa - 1 + t}. \quad (9)$$

The decentered arc will supply $\kappa = 1 + g(t)$. The common triangle stage uses horizontals through $B_j = (Ws^j, H(1 - s^j))$ on OB and $L_j = L(s^j)$. Parallels to L_1B carry B_j to L_{j+1} ; the p th horizontal gives E . This is the triangle protocol of Section 3, not the scaled rail telescope. The classical AK, AD and projective supports are documented in Appendix B; MA remains in Appendix C.

5.1 Realizing the positive radical

For $p > 2$, define

$$q(t) = \sqrt{\frac{(p-2)(t^2+1) + (10p+4)t}{12p}}, \quad g(t) = (p-1)q(t), \quad \mathcal{T}_{\text{arc}}(s) = s \frac{1+g(t)}{t+g(t)}. \quad (10)$$

For integer $p \geq 3$, put $\eta = (p-1)\sqrt{(p-2)/(12p)}$, $\zeta = (5p+2)/(p-2)$, $d_0 = W/\eta$ and $\delta = (H/X)\sqrt{\zeta^2 - 1}$. Prepare

$$G_0 = (W, H + H\zeta/X), \quad O_{\text{arc}} = (W - d_0 + \delta, H - H/(X\eta)), \quad x = W - d_0.$$

Draw the arc centered at O_{arc} with radius $EG_0 = (H/X)(t + \zeta)$ and take the lower intersection D_{arc} with the prepared vertical. Its vertical distance below the center is

$$\sqrt{EG_0^2 - \delta^2} = \frac{H}{X} \sqrt{t^2 + 2\zeta t + 1}.$$

The support AD_{arc} consequently gives $\kappa = 1 + g(t)$ in (9). All preparation parameters use rational operations and square roots of positive quantities. The radicand stays positive for $t > 0$.




Figure 4: The decentered-arc device for $p = 7, X = W = 2, H = 4, s = 3/4$. The full circle geometry is shown separately from the local report, at independently chosen equal-axis scales. The displacement δ makes the vertical circle intersection perform a square root. The local panel retains the complete support and distinguishes the reported state.

5.2 Global monotone convergence of the decentered map

Unlike the inverse-circle correction, this map is defined for every positive residual and requires no choice between two inverse-model branches. It uses a mobile radius at every step. Unlike rational diagonal Padé, it realizes the correction with a single radical rather than two rational factors. These are practical differences in geometric realization, even though all three maps have fifth local order.

Centered AD and decentered AD use the same mobile support operations: one arc and one join. Their fixed preparations and coordinate scales differ. This is the precise sense in which order five is obtained at the mobile support cost of order three.

Theorem 5.1. *For real $p > 2$, $X > 0$ and every $s_0 > 0$, (10) converges to a monotonically without crossing it. Its logarithmic error satisfies*

$$e^+ = \frac{(p-2)(p^2-1)(3p+1)}{1440}e^5 + O(e^7). \quad (11)$$

Proof. The radicand is positive and $q(1/t) = q(t)/t$, so the correction is reciprocal. Write

$$\mathcal{E}(t) = \frac{\log t}{p} + \log(1 + g(t)) - \log(t + g(t)).$$

Define the auxiliary polynomials

$$\begin{aligned} A_*(t) &= (p-1)(t^2+1) + (10p+2)t, \\ B_*(t) &= (t+1)(6pt - (t-1)^2), \\ K_*(t) &= (p^2-5p-2)(t-1)^2 + 12p(3p+1)t. \end{aligned}$$

Differentiation and factorization give

$$\frac{d\mathcal{E}}{d \log t} = \frac{(p-2)(p-1)^2(qA_* - B_*)}{12p^2g(1+g)(t+g)}, \quad (12)$$

$$q^2A_*^2 - B_*^2 = \frac{p+1}{12p}(t-1)^4K_*. \quad (13)$$

Here $A_* > 0$. If $B_* \leq 0$, then $qA_* - B_* > 0$. Otherwise $(t-1)^2 < 6pt$. When $p^2 - 5p - 2 \geq 0$, $K_* > 0$ directly; otherwise

$$K_* = 6p^2(p+1)t + (p^2-5p-2)((t-1)^2 - 6pt) > 0.$$

Equation (13) now gives $qA_* > B_*$ for $t \neq 1$. Therefore $\mathcal{E}$ is strictly increasing with $\mathcal{E}(1) = 0$: the new logarithmic error has the old sign. Meanwhile $\Phi(t) - 1 = (1-t)/(t+g)$ moves the state toward the root. The orbit is monotone and bounded between s_0 and a . Its positive limit is a fixed point by continuity, and $\Phi(t) = 1$ implies $t = 1$.

Near one, rationalize $qA_* - B_*$ using (13). Since $q = 1$, $g = p-1$, $A_* = B_* = 12p$ and $K_* = 12p(3p+1)$ at $t = 1$, this gives

$$\lim_{t \rightarrow 1} \frac{d\mathcal{E}/d \log t}{(t-1)^4} = \frac{(p-2)(p^2-1)(3p+1)}{288p^5}.$$

Substituting $t = e^{pe}$ and integrating yields (11); reciprocity makes the analytic logarithmic map odd. $\square$

At $p = 3$, the leading coefficient is $+1/18$, whereas the fixed-circle inverse has $-1/18$. The former stays on the starting side; the latter alternates near the root. At $p = 2$, both scalar radical formulas reduce to the exact square-root correction, but the displayed decentered preparation must be replaced by a mean-proportional construction.

6 What the shorter construction saves

A useful comparison holds the state, requested accuracy and counting convention fixed. The rational competitors can be built on the same rail telescope. A prepared projector

$$Z(\mu, \nu) = \left(-\frac{W\mu}{1-\mu}, H\left(1 - \frac{\nu}{1-\mu}\right) \right)$$

sends $Q = R(q_p)$ to $\mathcal{U}_{f(q_p)}$, with $f(q) = \mu + \nu/q$. Two such projections and two rail joins multiply a stored value by f/g : a four-join module.

For Halley choose $f = (p-1)/(4p) + (p+1)/(4pt)$ and $g = (p+1)/(4p) + (p-1)/(4pt)$. Since $t = X2^{p-1}q_p$, each is of the required projector form. The cost is $2p-1+4 = 2p+3$ joins.

For rational diagonal Padé set $a_5 = (p-1)(2p-1)$, $b_5 = 8p^2-2$, $c_5 = (p+1)(2p+1)$. The correction is

$$\Phi_{\text{rat}}(t) = \frac{a_5 t^2 + b_5 t + c_5}{c_5 t^2 + b_5 t + a_5} = \prod_{r \in \{r_+, r_-\}} \frac{t+r}{rt+1}, \quad r_{\pm} = \frac{b_5 \pm \sqrt{12p^2(4p^2-1)}}{2a_5}.$$

Each factor uses the four-join module with $f_r = [1+r/t]/[2(1+r)]$ and $g_r = [r+1/t]/[2(1+r)]$. Both reuse Q , giving $2p+7$ joins. Its scalar map is globally monotone: its derivative is $a_5 c_5 (t-1)^4 / (c_5 t^2 + b_5 t + a_5)^2$, and numerator minus denominator is $6p(1-t)(1+t)$.

Table 2: Generic mobile protocol costs; fixed preparation and final inversion are excluded. A parallel macro is $2J + 3C_{\text{arc}}$. The scope column describes the exact scalar map.

Construction	Order	J	$P_{\parallel}$	C_{arc}	Scope
MA rectangle	1	2	$2p+1$	0	$X > 1$
AK rectangle	2	2	$2p+1$	0	Global
AD rectangle	3	3	$2p+1$	1	Global
Projective AD	4	4	$2p+1$	0	Global
Decentered AD	5	3	$2p+1$	1	Global
Pencil Halley	3	$2p+3$	0	0	Global
Pencil rational Padé	5	$2p+7$	0	0	Global
Fixed-circle inverse	5	$2p+2$	0	0	Real band
Circle-AD protocols	5	Section 7			Global

The fixed circle saves five joins per regular update relative to this rational fifth-order realization. A reproducible cubic example makes both the gain and the price of control visible.

Table 3: Steps / mobile joins to obtain $|s/a - 1| \leq 10^{-d}$ from $p = 3, X = 2, s_0 = 3/4$, computed at 250 digits. The last two rows include the 12 joins for initial binary residual caching in their totals; every circle trial here is regular and accepted.

Protocol	$d = 6$	$d = 12$	$d = 30$	$d = 60$
Pencil Halley	2 / 18	3 / 27	3 / 27	4 / 36
Pencil rational Padé	1 / 13	2 / 26	2 / 26	3 / 39
Circle, linear power	1 / 8	2 / 16	2 / 16	3 / 24
Circle, branch entry	1 / 30	2 / 48	2 / 48	3 / 66
Circle, residual gate	1 / 40	2 / 68	2 / 68	3 / 96

These are geometric work counts under the stated convention, not circuit time or energy measurements. The next section develops branch entry; the optional residual gate is documented in Appendix D.

7 From a convergence band to all positive starts

Use a regular descending circle readout whenever it exists; otherwise use centered AD (14). This classical Halley map is globally monotone toward a (Appendix B). The branch theorem makes an improvement gate unnecessary; Appendix D retains the independently formalized gate.

7.1 AD entry and geometric exceptions

Cache $E = R(s^p)$ and $R(t)$. Stop at $t = 1$; otherwise apply the chosen step and refresh its power and residual.

Proposition 7.1 (Global branch-driven protocol). *For every integer $p \geq 3$, $X > 0$ and $s_0 > 0$, this exact geometric protocol converges to $a = X^{-1/p}$. With the preparation specified below, it eventually uses the circle step unless already at the root, and has local order five.*

Proof. Outside the band, AD converges globally and therefore reaches the open band in finitely many steps. Inside, both maps strictly reduce $|\log t|$ and preserve the band. On the compact log-residual interval determined by entry, both scalar post-error functions are continuous. If the errors tended to $\ell > 0$, the larger post-error would be uniformly below the current error on $\ell \leq |\log t| \leq |\log t_{\text{entry}}|$. Every late step would decrease it by a fixed positive amount, a contradiction. Thus chart-driven switching also converges.

The finite chart is regular near the root. For the compact preparation ($p \neq 4$), $\rho = 1/2$ and

$$b(1) = \frac{48p(p+4)(p+1)}{(p-1)\Delta_p} > 0;$$

for $p = 4$, use the separate chart in Appendix A, where direct substitution also gives $\rho \neq 1$ and $b(1) \neq 0$. At the root $\Delta_p(1) = 36p^2 > 0$. Continuity therefore preserves the transverse crossing, distinct B, G and finite top intersection nearby; the eventual update is the order-five circle map. $\square$

Lean checks the complete scalar branch theorem and chart certificates; the switching assembly above remains a written proof.

7.2 Geometric arithmetic and complete trace counts

For positive a, b , reuse $R(a)$ when $b = 1$. Otherwise one join gives $V = CR(b) \cap OA = \mathcal{U}_{1/b}$, then four joins through $\mathcal{U}_{1/2}, V, \mathcal{U}_{1/2}, \mathcal{U}_{1/4}$ successively give $L(a/2), R(ab/2), L(ab/4), R(ab)$. This five-join product is finite, including its exact unit shortcut. The two centers $\mathcal{U}_{d/[2(1+d)]}$ and $\mathcal{U}_{1/[2(1+d)]}$ multiply a rail value by a fixed $d > 0$ in two joins.

With $m = \lfloor \log_2 p \rfloor + \text{popcount}(p) - 1$, binary powering and the fixed scale X cost $5m + 2$ joins. Each power makes m unit tests; branch and root-stop decisions are additional, uncharged decisions.

Table 4: Mobile upper bounds, including the next cache. Fixed preparation, existing intersections and decisions are excluded.

Operation	J	$P_{\parallel}$	C_{arc}
Initial power and residual cache	$5m + 2$	0	0
Branch-driven regular circle step	$5m + 8$	0	0
Branch-driven fallback after a full attempted readout	$5m + 10$	2	1

A trial costs six joins: two for $R(kt)$, one for $L(\rho s)$ and three for the circle readout. AD adds $2J + 2P_{\parallel} + C_{\text{arc}}$. For the cubic, initialization costs $12J$, a regular step $18J$ and full fallback 31 expanded traces. At $p = 10^6$, $m = 25$ gives $133J$ per regular step. These are complete protocol upper bounds.

8 Evidence and reproducibility

Lean now establishes all scalar claims of Theorem 4.3: real-parameter contraction, alternation, branch selection, invariant-band convergence and exact local order five. Branch-driven switching and the decentered map’s global radical argument remain written proofs. Appendix D has a separate formal switching theorem. The source map identifies these distinct scopes [18, 19].

Lean module	Checked component
<code>RectangleProjective</code>	Incidences and order-four dynamics.
<code>RectanglePencils*</code>	Rail/projector identities, dynamics and orders.
<code>RectangleFixedCircleUniform</code>	Uniform logarithmic inequality and alternation for every real $p > 2$, including scalar endpoints.
<code>RectangleFixedCircleDynamics</code>	Continuous inverse, explicit residual band, invariant orbits, convergence to $X^{-1/p}$ and exact fifth-order inverse-map limit.
<code>RectangleFixedCircleTheorem43</code>	
<code>RectangleFixedCircle</code>	Quadratic identities, charts, preparation, readout and a parameterized local limit.
<code>RectangleDecenteredArc</code>	Certificates for the written radical proof.
<code>RectangleCircleSafeguard</code>	Products, powers, residual gate, global switching and local transfer; 41 declarations audited.
<code>RectangleFastPower</code>	Product incidence and binary recursion.
<code>AnalogFastAD</code>	Six centered-arithmetic identities.

All three audits allow only the standard axioms `propext`, `Classical.choice` and `Quot.sound`. Physical trace counts are instrumented inventories, not a Lean cost semantics. Neither the proofs nor sampled contraction ratios certify uniform drawing size, browser accuracy or hardware performance.

The retained incidence campaigns check 162 circle configurations at 250 digits, 108 arc configurations at 160 digits, and two 144-trajectory controller campaigns at 180 digits. These finite tests supplement the proofs. The new uniform-proof verifier passes 25 symbolic checks, 240 branch/endpoint cases and 72 scalar trajectories at 400 digits, including noninteger parameters near $p = 2$ and large degrees. The accompanying `README.md` links the sources, audits, data and reproduction commands. The package includes a local PDF build and verifier; the [gallery](#) provides interactive views. The repository includes the MIT license and citation metadata; V20, V21 and analog V30 remain archived.

The numerical families are classical [7, 20, 21]. Section 2 records the exact identifications and limits of the novelty claim. Appendix E retains the exploratory analog material, which supplies no hardware evidence for either fifth-order solver.

9 Conclusion

Two geometric mechanisms realize distinct fifth-order corrections. The fixed circle alternates with decreasing logarithmic error throughout its real branch; a uniform convexity proof covers every real $p > 2$. The decentered arc converges monotonically from every positive state. For integer degrees their incidence protocols make the cost of each update explicit. Relative to the stated rational fifth-order realization, the regular fixed-circle protocol saves five joins; global entry and exceptional charts carry a separate, accounted cost. These are mathematical and constructive results, independent of the prospective analog application.

A Rational preparation for every integer degree

This appendix makes the general-circle protocol reproducible without searching through an earlier manuscript. Write $v = p - 3$ and use α_p, γ_p from (4). Define

$$\begin{aligned}\mathcal{D}_p &= 625v^5 + 10525v^4 + 56797v^3 + 88639v^2 + 14758v + 664, \\ \mathcal{K}_p &= 625v^5 + 1975v^4 + 4147v^3 + 2905v^2 + 448v + 340, \\ \chi_p &= 288p(p-2)(25p^2 - 244)/\mathcal{D}_p, \\ u_p &= 5 - 4\alpha_p/(9\gamma_p), \quad w_p = 8(5 - 2p)/(9(p-1)).\end{aligned}$$

Take

$$\begin{aligned}\rho &= \frac{1}{3}, & k &= \mathcal{K}_p/\mathcal{D}_p, & j &= -3\chi_p w_p/4, \\ h &= 2 + 2j - \chi_p u_p/2, & z_x &= 2 - \chi_p, & z_y &= \frac{384p(25p^3 - 50p^2 - 415p + 974)}{\mathcal{D}_p}.\end{aligned}$$

Then the incidence certificate in Theorem 4.2 has multiplier $\lambda_p = -64k\chi_p/(9\gamma_p)$. The positive coefficients imply $\mathcal{D}_p, \mathcal{K}_p > 0$ for $v \geq 0$; $25p^2 \neq 244$ for integer $p \geq 3$ makes χ_p, λ_p, j nonzero. All parameters are rational. The circle is drawn with center (h, j) through B . This uniform chart is checked symbolically; the Lean preparation certificates cover the compact chart below and the separate degree-four specialization. At $p = 4$ the formulas give the finite values

$$k = 145/2389, \quad O_\Gamma = (13502/7167, 3328/2389), \quad Z = (-214/2389, 2432/2389).$$

The cubic worked example uses a more compact chart. For $p \neq 4$, put $\Lambda_p = p^3 + 12p^2 + 39p - 26$ and take

$$\begin{aligned}\rho &= \frac{1}{2}, & k &= \frac{(p-2)(p^2 - 4p + 13)}{\Lambda_p}, \\ h &= 2 - \frac{72p(p+4)}{(p-1)\Lambda_p}, & j &= \frac{24p(p-2)(p+4)}{(p-1)\Lambda_p}, \\ z_x &= 2 - \frac{24p(p+4)(p-2)}{(p-4)\Lambda_p}, & z_y &= \frac{24p(p^2 + 2p - 26)}{(p-4)\Lambda_p}.\end{aligned}$$

Here $\lambda_p = -384p(p-2)(p+4)(p^2 - 4p + 13)/[(p-4)(p-1)\Lambda_p^2]$. Substitution checks both charts. For numerical evaluation the branch can be written without the removable cancellation at large t :

$$v(t) = \begin{cases} \frac{\alpha_p - t\gamma_p}{\beta_p(1-t) + \sqrt{\Delta_p(t)}}, & t \leq 1, \\ \frac{\sqrt{\Delta_p(t)} - \beta_p(1-t)}{t\alpha_p - \gamma_p}, & t > 1. \end{cases}$$

B Classical rectangle supports

B.1 AK and centered AD: orders two and three

AK: a fixed line. Prepare $K = (W(1 - X/p), 0)$. The support AK has $\kappa = p$, so $\mathcal{T}_{AK}(s) = ps/(p - 1 + t)$, Newton under inversion. It has order two and converges from every positive start.

AD: a centered length transfer. Prepare

$$O_{AD} = \left(W - \frac{2W}{p-1}, H - \frac{H(p+1)}{X(p-1)} \right).$$

Transfer $AE = Hs^p$ downward from this center to D_{AD} and join AD_{AD} . Then $\kappa = [p + 1 + (p - 1)t]/2$, giving

$$\mathcal{T}_{AD}(s) = s \frac{p + 1 + (p - 1)t}{p - 1 + (p + 1)t}. \quad (14)$$

This Halley map has cubic order and global monotone convergence. Earlier geometric interpretations appear in [3, 4]; here the rail encoding and readout are shared with AK.




Figure 5: Pandrosion AK, order two. The fixed support AK replaces MA while the triangle chain and green correction direction are retained. Here $p = 3, X = 2, s = 3/4$.




Figure 6: Pandrosion AD, order three. Transfer the orange length AE onto the prepared vertical through O_{AD} ; the lower arc point determines AD_{AD} . Here $O_{AD} = C$.

Both drawings retain the triangle stage, green transport and horizontal readout; only the support changes.

B.2 Projective AD: order four

Two joins supply the support. Prepare

$$\ell : y = H \left(1 + \frac{5p-1}{(p+1)X} \right), \quad Z_{\text{proj}} = \left(W \left(1 + \frac{5p-1}{2p(2p-1)} \right), H \left(1 + \frac{p+1}{(2p-1)X} \right) \right).$$

Set $D_{\text{proj}} = Z_{\text{proj}}E \cap \ell$ and join AD_{proj} . Substitution in (9) gives

$$\mathcal{T}_{\text{proj}}(s) = s \frac{2p((2p-1)t + p + 1)}{(p+1)t^2 + 2(2p-1)(p+1)t + (2p-1)(p-1)}. \quad (15)$$

The direct-root correction is Padé [2/1]; the reciprocal-state factor displayed here is [1/2]. Denote its positive denominator by $Q_{\text{proj}}(t)$. Direct differentiation gives

$$\mathcal{T}'_{\text{proj}}(s) = -\frac{2p(p^2-1)(2p-1)(t-1)^3}{Q_{\text{proj}}(t)^2}.$$

The map has its global maximum a at $s = a$. Also $\mathcal{T}_{\text{proj}}(s)/s - 1 = (1-t)((p+1)t + 5p-1)/Q_{\text{proj}}(t)$. Thus a first step lands at or below a , and subsequent steps increase to a . Integrating the derivative at a gives exact order four. The archival companion gives further dynamics and the original triangle-chain realization [22].




Figure 7: Projective AD [2/1], order four, at the same cubic state. The orange line $Z_{\text{proj}}E$ meets the prepared horizontal ℓ at D_{proj} . The blue support AD_{proj} receives the green correction. The inset separates P , T and P^+ .

C The linear MA reference

MA provides a baseline, not an additional accelerated method. Write its lower-right endpoint as $A_{\text{lin}} = B$. The parallel to ME through P meets MB ; a horizontal returns the next state. For $s \neq 1$,

$$\mathcal{T}_{\text{MA}}(s) = 1 - \frac{X-1}{XS_p(s)}, \quad S_p(s) = \sum_{j=0}^{p-1} s^j.$$

For $X > 1$ this scalar map is increasing and $\mathcal{T}_{\text{MA}}(s) - s = (1 - Xs^p)/(XS_p(s))$. It therefore converges monotonically to a from every positive start, with the removable scalar extension at $s = 1$. Its derivative at a is $1 - pa^{p-1}/S_p(a) \in (0, 1)$: convergence is linear.




Figure 8: Pandrosion MA, the linear reference construction, at $p = 3$, $X = 2$, $s = 3/4$. Thin lines construct the power by similar triangles. The green parallel meets the blue support MA_{lin} ; the horizontal gives P^+ . The historical endpoint A_{lin} is the lower-right vertex B in this paper. At $s = 1$ the geometric intersection is coincident despite the removable scalar extension.

The notation $S_p(s)$ denotes only this geometric sum.

D The independent residual-gate controller

This independently formalized V21 controller is retained for comparison. It is not required by the branch-driven protocol.

For a positive circle candidate c , construct $u = Xc^P$ and accept precisely when

$$(t \leq 1 \wedge t \leq u^2 \wedge tu^2 \leq 1) \quad \text{or} \quad (1 \leq t \wedge u^2 \leq t \wedge 1 \leq tu^2). \quad (16)$$

Otherwise use AD. Three rail comparisons implement this condition, equivalent to $|\log u| \leq |\log t|/2$. The errors decrease; if their limit were positive, eventually no halving step would be possible, leaving only AD and forcing the limit to zero. This is the V21 global switching theorem, including the incidence-based gate, formalized in `RectangleCircleSafeguard`. Strict branch contraction alone does not assert the stronger half-error inequality.

From the cached current state, a circle trial uses six joins and the candidate power/residual uses $5m + 2$. The gate's two products add at most ten joins and three ordered comparisons. Acceptance thus costs $5m + 18$ joins including the next cache. Full rejection adds AD ($2J + 2P_{\parallel} + C_{\text{arc}}$) and a fresh $5m + 2$ cache, for a total of $(10m + 22)J + 2P_{\parallel} + C_{\text{arc}}$. The common initialization costs $5m + 2$ joins.

Cubic protocol ($m = 2$)	Regular / accepted	Full fallback
Branch-driven	$18J$	$20J + 2P_{\parallel} + C_{\text{arc}}$
Residual gate	$28J$	$42J + 2P_{\parallel} + C_{\text{arc}}$

The expanded fallback bounds are respectively 31 and 53 traces. At $p = 10^6$, regular steps cost 133 versus 143 joins. A completed gate adds two product-identity tests to the m tests in each power evaluation. All these are protocol upper bounds; fixed preparation and decisions are not charged as traces.




Figure 9: The five-join product and the optional V21 residual gate. The branch-driven protocol reuses the product for powers and scales, but omits the two gate products and its three ordered comparisons. The geometric points and labels are unchanged from the independently checked V21 figure.

Its separate 144-trajectory campaign and 41-declaration axiom audit are reproduced in the repository.



E Binary powering and an exploratory analog extension

The linear telescope is attractive for a few degrees. At large p , binary powering gives a smaller arithmetic architecture. Define a top encoding $Z_{\times}(q) = (W + Hq, H)$ and the fixed unit $U_{\times} = (W + H, H)$. A parallel to $U_{\times}B$ copies $R(a)$ to $Z_{\times}(a)$. The parallel through $Z_{\times}(a)$ to $U_{\times}R(b)$ then meets the right rail at $R(ab)$: its slope is b and its horizontal displacement is $-Ha$.

Squaring the accumulator and multiplying by s at each binary one produces $E = R(s^p)$. With $L = \lfloor \log_2 p \rfloor$ and $h = \text{popcount}(p)$, there are $m = L + h - 1$ multiplications. Reusing the top copy of s , the power stage costs $mJ + (m + L)P_{\parallel}$. It feeds any of the four rectangle supports unchanged.

(b) Fast output E feeds the AD support

(a) One geometric multiplication



(c) Binary schedule: $13 = (1101)_2$; five multiplications instead of twelve




Figure 10: The fast-power construction and its AD readout, adapted from the V30 companion. The case $p = 13 = (1101)_2$ uses five products along exponents 1, 2, 3, 6, 12, 13. A product is an actual pair of line incidences, not a numerically inserted power. The output E enters the same moving support and horizontal readout.

The construction is an operation graph for a circuit: it identifies which signals must exist before a cell can act, and which intermediate values must survive a later correction. The implementation obligations are concrete:

Geometric object	Electrical role	Property to measure
Rail coordinate	Scaled voltage or current	Dynamic range and offset
Product incidence	Multiplier and weighted sums	Gain error and settling
Circle crossing	Quadratic solve plus branch choice	Stability and branch margin
Retained point	Sample-and-hold memory	Leakage and sampling error

We now separate the existing AD circuit from the quadratic cell suggested by the fixed-circle construction.

E.1 A behavioral baseline and a prospective quadratic cell

Existing order-three baseline. The V30 behavioral prototype implements centered AD with state $q = p(u - 1)$ after a digitally prepared scaling $s = cu$, $Y \approx Xc^p \in [1, 2]$. Its residual sensitivity is $\partial \log(Yu^p)/\partial q = 1/u$, rather than p/u for an uncentered state. Centered binary products feed the exact update

$$q^+ = q + \frac{2(1 + q/p)(1 - t)}{1 + t + (t - 1)/p}, \quad \hat{r} = \frac{1}{c(1 + q/p)}.$$

There are 25 power products at $p = 10^6$, and at most 37 over $3 \leq p \leq 10^6$. Netlists and calibration are in the companion [23].




Figure 11: The retained order-three AD behavioral topology. Memories, loading and switches are explicit; diamonds are arithmetic sources. Neither fifth-order solver has an electrical implementation in this study.

Prospective circle cell. A candidate feedback cell must enforce $N(v) - tD(v) = 0$ on the branch through $v = 1$ and form $s^+ = sv$. With $z = p(v - 1)$, division by 12 gives the exact specification

$$0 = (1 - t) - \frac{(1 - 2/p) + (1 + 2/p)t}{2} z + \frac{(1 - 3/p + 2/p^2) - t(1 + 3/p + 2/p^2)}{12} z^2.$$

Its derivative in z is -1 at $(t, z) = (1, 0)$, independently of the degree. It vanishes at a branch endpoint, so branch margin and fallback must be designed before this becomes a usable circuit.

Evidence retained from V30. Arithmetic tests cover 8,243 pairs. The calibrated behavioral SPICE campaign gives worst decoded-root error 0.279 ppm at 23.985 ms across 64 cases; a 4.785 ms policy meets one ppm in 18 new cases. At $p = 10^6$, $X = 500000$, digital preparation already places $1/c$ within 2.477×10^{-7} relative error. Reported readout times exclude preparation, programming and conversion. These are behavioral simulations, not silicon measurements.

A comparison of the proposed quadratic cell with AD must hold power, memory, loading and noise fixed and measure wrong-branch events, settling and decoded error. Neither faster electronics nor lower energy follows from the geometric join count. This appendix motivates that experiment; it supplies no electrical validation of the fifth-order methods.

References

- [1] J. F. Traub. On a class of iteration formulas and some historical notes. *Communications of the ACM*, 4(6):276–278, 1961. doi: 10.1145/366573.366606.
- [2] J. F. Traub. *Iterative Methods for the Solution of Equations*. Prentice-Hall, Englewood Cliffs, NJ, 1964.
- [3] Walter Gander. On halley’s iteration method. *The American Mathematical Monthly*, 92(2):131–134, 1985. doi: 10.2307/2322644.
- [4] T. R. Scavo and J. B. Thoo. On the geometry of halley’s method. *The American Mathematical Monthly*, 102(5):417–426, 1995. doi: 10.2307/2975033.
- [5] A. Melman. Geometry and convergence of euler’s and halley’s methods. *SIAM Review*, 39(4):728–735, 1997. doi: 10.1137/S0036144595301140.
- [6] S. Amat, S. Busquier, and J. M. Gutiérrez. Geometric constructions of iterative functions to solve nonlinear equations. *Journal of Computational and Applied Mathematics*, 157(1):197–205, 2003. doi: 10.1016/S0377-0427(03)00420-5.
- [7] Beata Laszkiewicz and Krystyna Ziętak. A Padé family of iterations for the matrix sector function and the matrix p th root. *Numerical Linear Algebra with Applications*, 16(11–12):951–970, 2009. doi: 10.1002/nla.656.
- [8] Krystyna Ziętak. A padé family of iterations for the matrix sector function and matrix p th root, 2009. URL https://cs.pwr.edu.pl/zietak/talks/GAMM_2009.pdf. GAMM presentation, joint work with Beata Laszkiewicz; slide 11 gives h_{12} and h_{22} .
- [9] Dmitrii B. Karp and Minghua Lin. Convergence analysis of a Padé family of iterations for the matrix sector function, 2011. URL <https://arxiv.org/abs/1102.3957>.
- [10] Jakob Steiner. *Die geometrischen Constructionen, ausgeführt mittelst der geraden Linie und eines festen Kreises*. Ferdinand Dümmler, Berlin, 1833. URL https://archive.org/details/bub_gb_m2wLAAAAYAAJ.
- [11] Émile Lemoine. *Géométrie ou Art des constructions géométriques*. Number 18 in Scientia, physique-mathématique. C. Naud, Paris, 1902. URL <https://catalogue.bnf.fr/ark:/12148/cb36049032t>.
- [12] Maurice d’Ocagne. *Esquisse d’ensemble de la nomographie*, volume 4 of *Mémoires des sciences mathématiques*. Gauthier-Villars, 1925. URL https://www.numdam.org/item/MSM_1925__4__1_0/.
- [13] R. E. Shafer. On quadratic approximation. *SIAM Journal on Numerical Analysis*, 11(2):447–460, 1974. doi: 10.1137/0711037.
- [14] Vinay Kanwar, Sukhjot Singh, and Suman Bakshi. Simple geometric constructions of quadratically and cubically convergent iterative functions to solve nonlinear equations. *Numerical Algorithms*, 47(1):95–107, 2008. doi: 10.1007/s11075-007-9149-4.
- [15] Mehmet Pakdemirli. Non-divergent circular arc root algorithm. *Mathematics and Computational Sciences*, 6(3):77–88, 2025. doi: 10.30511/mcs.2025.2028349.1171. URL <https://avesis.mcbu.edu.tr/yayin/3d95199c-1515-4b22-a182-0e86fa457b82/non-divergent-circular-arc-root-algorithm>.
- [16] Pappus of Alexandria. Book iii of the mathematical collection. College of the Holy Cross, Bookshelf 63, 2023. URL https://crossworks.holycross.edu/hc_books/63/. English translation by John B. Little, following Hultsch; opening address to Pandrosion.
- [17] Mickaël Launay. Pandrosion et la duplication du cube. Micmaths, video presentation, 2024. URL <https://www.youtube.com/watch?v=7T4neM660SM>. Rectangle construction and correction, approximately 16:37–19:10.
- [18] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction—CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021. doi: 10.1007/978-3-030-79876-5_37.
- [19] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs*, pages 367–381. ACM, 2020. doi: 10.1145/3372885.3373824. URL <https://leanprover-community.github.io/papers/mathlib-paper.pdf>.
- [20] NIST Digital Library of Mathematical Functions. Section 3.8: Nonlinear equations, 2026. URL <https://dlmf.nist.gov/3.8>. Accessed 16 September 2026.
- [21] George A. Baker, Jr. and Peter Graves-Morris. *Padé Approximants*, volume 59 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 2 edition, 1996. doi: 10.1017/CBO9780511530074. In particular, Section 1.5, Theorems 1.5.2–1.5.3, on argument and value homographic invariance.
- [22] Ivan Besevic. Reciprocal root geometry, version 20. Research preprint, 2026. URL https://github.com/ivan-fr/pandrosion/tree/main/paper/reciprocal_geometry_v20.
- [23] Ivan Besevic. Pandrosion geometry and analog computation, version 30. Research preprint, 2026. URL https://github.com/ivan-fr/pandrosion/tree/main/paper/pandrosion_analog_v30.