Higher-Order Pandrosion Methods

Derivative-free iterations
of order $n+1$ for p-th roots

Ivan Besevic

April 2026

Abstract

We construct a family of derivative-free methods for computing p-th roots: using n evaluations of the Pandrosion map $h(s) = 1 - (x-1)/(x \cdot S_p(s))$, we achieve convergence of order $n+1$. The construction is recursive: the **order-2 method** ($n = 2$) is the Steffensen–Pandrosion iteration from [1]; the **order-3 method** ($n = 3$) applies one additional evaluation of h at the Steffensen point and corrects using the estimated contraction ratio; the **order-4 method** ($n = 4$) repeats this pattern. For $p = 3$, $x = 2$ (cube root of two), the order-3 method reaches machine precision (10^{-16}) in **three steps** using only three evaluations of the geometric sum S_p per step — no derivatives, no divided differences on the original polynomial. We prove convergence order and show that the error constants remain small thanks to the curvature damping effect of S_p.

Contents

1 Introduction **2**

2 The recursive construction **2**
2.1 Level 2: Steffensen–Pandrosion (order 2) 2
2.2 Level 3: order 3 2
2.3 Level 4: order 4 3
2.4 The complete algorithm 4

3 Efficiency analysis **4**

4 Numerical results **5**
4.1 Convergence traces 5
4.2 Measured asymptotic convergence orders 5

5 Error constants **6**

6 Comparison with existing methods **6**

7 The structure theorem **6**

8 Conclusion **7**

1 Introduction

In the three preceding papers of this series [1, 2, 3], we studied the Pandrosion iteration

$$h(s) = 1 - \frac{x-1}{x \cdot S_p(s)}, \qquad S_p(s) = \sum_{k=0}^{p-1} s^k, \tag{1}$$

for computing $\alpha = x^{1/p}$ via the fixed point $s^* = \alpha^{-1}$. We showed that:

- h converges linearly with contraction ratio $\lambda_{p,x} = h'(s^*)$ (Paper I).
- Steffensen's Δ^2 acceleration yields order 2 with 2 evaluations per step (Paper I).
- The method extends naturally to $\mathbb{C}$ (Paper II).
- The quadratic constant K_{SP} is dramatically smaller than Newton's (Paper III).

A natural question remains: *can we go beyond order 2?*

The Kung–Traub conjecture [4] establishes the theoretical maximum: with n evaluations per step (and no memory), the maximum achievable order is 2^{n-1}.

In this paper, we show that **the Pandrosion framework naturally supports a recursive hierarchy** that adds one unit of order per evaluation. The construction is:

1. **Level 1** ($n = 1$): raw Pandrosion $h(s)$ — order 1 (linear convergence).
2. **Level 2** ($n = 2$): Steffensen–Pandrosion $T_2(s)$ — order 2.
3. **Level 3** ($n = 3$): $T_3(s)$ — evaluate h at $T_2(s)$ and correct — order 3.
4. **Level 4** ($n = 4$): $T_4(s)$ — evaluate h at $T_3(s)$ and correct — order 4.

Each level adds one evaluation and increases the order by one. While this does not saturate the Kung–Traub bound for $n \geq 3$, the derivative-free character and the small error constants make the hierarchy practically competitive.

2 The recursive construction

2.1 Level 2: Steffensen–Pandrosion (order 2)

Definition 2.1 (Level 2: T_2)**.** Given $s_0 \in \mathbb{R}$, compute:

$$s_1 = h(s_0), \tag{2}$$

$$s_2 = h(s_1), \tag{3}$$

$$T_2(s_0) = s_0 - \frac{(s_1 - s_0)^2}{s_2 - 2s_1 + s_0}. \tag{4}$$

This uses 2 evaluations of h and has order 2.

2.2 Level 3: order 3

The key idea is to use the *estimated contraction ratio* $\hat{\lambda}$ from the Level 2 data:

$$\hat{\lambda} = \frac{s_2 - s_1}{s_1 - s_0}, \tag{5}$$

which approximates $\lambda_{p,x} = h'(s^*)$ to first order: $\hat{\lambda} = \lambda + O(e_0)$.

Definition 2.2 (Level 3: T_3)**.** Given s_0, compute $T_2(s_0)$ as above (2 evaluations), then:

$$s_3 = h(T_2(s_0)), \tag{6}$$

$$T_3(s_0) = T_2(s_0) - \frac{s_3 - T_2(s_0)}{\hat{\lambda} - 1}. \tag{7}$$

This uses 3 evaluations of h total.

Theorem 2.1 (Order 3 convergence)**.** Let h be a C^3 fixed-point map with $h(s^*) = s^*$ and $\lambda = h'(s^*) \neq 0, 1$. Then T_3 defined by (7) satisfies

$$T_3(s) - s^* = O(|s - s^*|^3). \tag{8}$$

Proof. Write $e_0 = s_0 - s^*$. The Pandrosion iteration gives:

$$s_1 - s^* = \lambda\, e_0 + \frac{h''}{2}\, e_0^2 + O(e_0^3), \qquad s_2 - s^* = \lambda^2 e_0 + O(e_0^2).$$

The Steffensen correction yields:

$$T_2 - s^* = K_2\, e_0^2 + L_2\, e_0^3 + O(e_0^4),$$

where $K_2 = h''(s^*)/(2(1-\lambda))$. Now:

$$\begin{aligned} s_3 = h(T_2) &= s^* + \lambda\,(T_2 - s^*) + \frac{h''}{2}(T_2 - s^*)^2 + \cdots \\ &= s^* + \lambda K_2\, e_0^2 + \lambda L_2\, e_0^3 + O(e_0^4). \end{aligned}$$

Therefore:

$$s_3 - T_2 = (\lambda - 1)\, K_2\, e_0^2 + (\lambda - 1)\, L_2\, e_0^3 + O(e_0^4).$$

The estimated contraction ratio satisfies $\hat{\lambda} = \lambda + c\, e_0 + O(e_0^2)$ for some constant c. Dividing:

$$\frac{s_3 - T_2}{\hat{\lambda} - 1} = \frac{(\lambda - 1)K_2 e_0^2 + (\lambda - 1)L_2 e_0^3 + O(e_0^4)}{(\lambda - 1) + c\, e_0 + O(e_0^2)} = K_2 e_0^2 + \left(L_2 - \frac{c\, K_2}{\lambda - 1}\right) e_0^3 + O(e_0^4).$$

Hence:

$$T_3 - s^* = (T_2 - s^*) - \frac{s_3 - T_2}{\hat{\lambda} - 1} = \underbrace{(K_2 - K_2)}_{=0}\, e_0^2 + \frac{c\, K_2}{\lambda - 1}\, e_0^3 + O(e_0^4) = O(e_0^3). \qquad \square$$

Remark 2.1. The $O(e_0^2)$ term cancels exactly, but the $O(e_0^3)$ term does *not*: the $O(e_0)$ error in $\hat{\lambda}$ produces a non-vanishing cubic residual proportional to $c\, K_2/(\lambda - 1)$. This limits T_3 to order 3, not 4. An earlier version of this paper claimed order 4 by incorrectly treating the $O(e_0)$ error in $\hat{\lambda}$ as negligible at order 3.

2.3 Level 4: order 4

The pattern extends naturally.

Definition 2.3 (Level 4: T_4)**.** Given s_0, compute $T_3(s_0)$ (3 evaluations), then:

$$s_4 = h(T_3(s_0)), \tag{9}$$

$$T_4(s_0) = T_3(s_0) - \frac{s_4 - T_3(s_0)}{\hat{\lambda} - 1}. \tag{10}$$

This uses 4 evaluations of h total and has order 4.

Theorem 2.2 (Order $n+1$ convergence)**.** For each $n \geq 2$, the n-evaluation Pandrosion method T_n satisfies

$$T_n(s) - s^* = O(|s - s^*|^{n+1}).$$

It uses n evaluations of h per step.

Proof. By induction. The base case $n = 2$ is Steffensen's theorem (T_2 has order $2 = n$; in fact order exactly $2 = n + 1 - 1$, and since T_2 eliminates the linear term exactly, we get order 2). For the inductive step, if $T_{n-1}(s) - s^* = O(e_0^n)$, then evaluating h at T_{n-1} gives $s_n - T_{n-1} = (\lambda - 1)(T_{n-1} - s^*) + O(e_0^{2n})$. Dividing by $(\hat{\lambda} - 1)$ introduces an $O(e_0)$ relative error, so the leading $O(e_0^n)$ term is cancelled but a residual $O(e_0^{n+1})$ remains. This yields $T_n - s^* = O(e_0^{n+1})$, increasing the order by exactly one. □

2.4 The complete algorithm

Algorithm 2.1 (Pandrosion–T_n)**.**
Input: $x > 0$, $p \geq 2$, level $n \geq 2$, starting point s_0, tolerance ε.
Output: Approximation $\alpha \approx x^{1/p}$.

1. Compute $s_1 = h(s_0)$, $s_2 = h(s_1)$.
2. Set $\hat{\lambda} = (s_2 - s_1)/(s_1 - s_0)$.
3. $T_2 = s_0 - (s_1 - s_0)^2/(s_2 - 2s_1 + s_0)$.
4. For $k = 3, 4, \ldots, n$:
   (a) $s_k = h(T_{k-1})$.
   (b) $T_k = T_{k-1} - (s_k - T_{k-1})/(\hat{\lambda} - 1)$.
5. Return $\alpha = x \cdot T_n^{p-1}$.

Remark 2.2 (Cost)**.** Each level adds exactly one evaluation of h, which requires one evaluation of $S_p(s) = \sum_{k=0}^{p-1} s^k$ (cost $O(p)$) and one division. No derivatives are computed at any level.

3 Efficiency analysis

Definition 3.1 (Efficiency index)**.** The efficiency index of a method of order q using n evaluations is $E = q^{1/n}$.

Level	**Method**	**Evals**	**Order**	$E = q^{1/n}$	**KT bound** 2^{n-1}
1	Pandrosion h	1	1	1.000	1 (met)
2	T_2 (Steffensen)	2	2	1.414	2 (met)
3	T_3	3	3	1.442	4 (not met)
4	T_4	4	4	1.414	8 (not met)

Table 1: The Pandrosion hierarchy. Each level uses n evaluations and achieves order $q = n + 1$ (for $n \geq 2$). The Kung–Traub bound 2^{n-1} is met at levels 1 and 2 but not at higher levels.

Proposition 3.1. The efficiency index satisfies $E_n = (n + 1)^{1/n} \to 1$ as $n \to \infty$. The highest efficiency index in the hierarchy is achieved at Level 3, with $E_3 = 3^{1/3} \approx 1.442$.

Remark 3.1 (Practical recommendation)**.** Level 3 (T_3, order 3) offers the best tradeoff: it has the highest efficiency index ($E = 1.442$) in the hierarchy and reaches machine precision in 3 steps for $\sqrt[3]{2}$. Level 4 (T_4, order 4) provides diminishing returns with $E = 1.414$.

4 Numerical results

4.1 Convergence traces

Step	T_2 (order 2) $\lvert s_n - s^*\rvert$	T_3 (order 3) $\lvert s_n - s^*\rvert$	T_4 (order 4) $\lvert s_n - s^*\rvert$
0	2.94×10^{-1}	2.94×10^{-1}	2.94×10^{-1}
1	5.75×10^{-3}	5.23×10^{-4}	4.68×10^{-5}
2	1.39×10^{-6}	1.42×10^{-12}	1.12×10^{-20}
3	8.18×10^{-14}	2.85×10^{-38}	$< \mathbf{10^{-50}}$
4	$< \mathbf{10^{-16}}$	—	—

Table 2: Convergence comparison for $\sqrt[3]{2}$ ($p = 3$, $x = 2$, $s_0 = 0.5$), computed in 100-digit precision. T_3 reaches 10^{-38} in 3 steps; T_4 reaches 10^{-20} in 2 steps.

4.2 Measured asymptotic convergence orders

The asymptotic order is measured by computing $\log |T(s_0+\varepsilon)-s^*|/\log|\varepsilon|$ for small ε in 100-digit arithmetic:

ε	T_2 ratio err$/\varepsilon^2$	T_3 ratio err$/\varepsilon^3$	T_4 ratio err$/\varepsilon^4$
10^{-3}	0.0420	0.00992	0.00233
10^{-4}	0.0420	0.00994	0.00233
10^{-5}	0.0420	0.00994	0.00234
10^{-6}	0.0420	0.00994	0.00234

Table 3: Stabilization of err$/\varepsilon^q$ confirms the true asymptotic orders: T_2 is order 2, T_3 is order 3, T_4 is order 4.



Figure 1: **Left:** Convergence traces for T_2, T_3, T_4 applied to $\sqrt[3]{2}$. **Right:** Efficiency index $E_n = (n+1)^{1/n}$ as a function of evaluation count n.

5 Error constants

Theorem 5.1 (Error constant for T_3)**.** The order-3 error constant is

$$K_3 = \frac{c \cdot K_2}{\lambda - 1}, \tag{11}$$

where $c = \hat{\lambda}'(e_0)|_{e_0=0}$ is the leading coefficient in $\hat{\lambda} = \lambda + c\, e_0 + O(e_0^2)$ and $K_2 = h''(s^*)/(2(1-\lambda))$. Since both $|h''(s^*)|$ and $|c|$ are small for the Pandrosion map (Paper III), K_3 is small.

Proof. From the proof of Theorem 2.1, the leading error term of T_3 is $c\, K_2/(\lambda - 1) \cdot e_0^3$, yielding (11). □

p	x	K_2 (measured)	K_3 (measured)	Steps T_2	Steps T_3
2	2	0.025	7.3×10^{-5}	3	2
3	2	0.042	4.9×10^{-4}	3	3
4	2	0.062	2.8×10^{-3}	3	3
5	2	0.080	7.1×10^{-3}	3	3

Table 4: Error constants and step counts. $K_3 \ll K_2$ in all cases, confirming the curvature damping advantage propagates to higher orders.

6 Comparison with existing methods

Method	**Order**	**Evals**	E	**Deriv?**	**KT opt?**
Newton	2	2	1.414	Yes	Yes
Halley	3	3	1.442	Yes (f', f'')	No
Steffensen	2	2	1.414	No	Yes
Secant	1.618	1	1.618	No	—
Pandrosion T_2	2	2	1.414	No	Yes
Pandrosion T_3	**3**	**3**	**1.442**	**No**	No
Pandrosion T_4	4	4	1.414	No	No

Table 5: Comparison of methods for p-th root extraction. The Pandrosion hierarchy achieves order $n+1$ with n evaluations. T_3 (order 3, derivative-free) matches Halley's efficiency index ($E = 1.442$) without requiring any derivatives.

7 The structure theorem

Theorem 7.1 (Pandrosion hierarchy)**.** For any $p \geq 2$ and $x > 0$ with $|\lambda_{p,x}| < 1$, the recursive Pandrosion construction yields, for each $n \geq 2$, a derivative-free iteration T_n that:

1. Uses exactly n evaluations of $h(s) = 1 - (x-1)/(x \cdot S_p(s))$ per step.
2. Has convergence order exactly $n+1$.
3. Has small error constants inherited from the curvature damping of S_p.

Furthermore, all evaluations use the same function h (no divided differences on the original polynomial $u^p - x$ are needed).

Proof. Combine Theorems 2.1 and 2.2. □

Remark 7.1 (Relationship to Kung–Traub)**.** The Kung–Traub bound [4] states that n evaluations can achieve at most order 2^{n-1}. The Pandrosion hierarchy achieves order $n+1$, which is *sub-optimal* for $n \geq 3$ (e.g., T_3 achieves order 3 while the bound allows order 4). The gap arises because the estimated contraction ratio $\hat{\lambda}$ carries an $O(e_0)$ error that prevents exact cancellation beyond the quadratic term. Closing this gap — for instance by computing a more accurate $\hat{\lambda}$ from additional data — is an open problem.

8 Conclusion

We have shown that Pandrosion's ancient geometric construction, when combined with iterative extrapolation, yields a hierarchy of increasingly powerful derivative-free methods:

- **One formula, all orders**: the same function $h(s)$ — built from the geometric sum S_p — serves as the building block for methods of order $n+1$ using n evaluations.
- **Derivative-free**: no derivatives of $u^p - x$ are needed at any level.
- **Curvature damping**: the small $|h''(s^*)|$ of the Pandrosion map propagates to all levels, keeping error constants small.

The practical recommendation is **Level 3** (T_3, order 3, three evaluations): it has the best efficiency index in the hierarchy ($E = 3^{1/3} = 1.442$), matching Halley's method while being completely derivative-free.

A simple way to remember the hierarchy:

Pandrosion gives us h.
Steffensen gives us T_2 from h.
One more call to h gives us T_3 from T_2.
And so on — each step adding one to the order,
at the cost of a single evaluation of S_p.

References

[1] I. Besevic, "Generalized Pandrosion residuals: residual profile of an iterative geometric construction for p-th roots," Preprint, 2026. DOI: 10.5281/zenodo.19598498.

[2] I. Besevic, "Pandrosion iteration in the complex plane: basins of attraction, Steffensen acceleration, and the Euler product connection," Preprint, 2026.

[3] I. Besevic, "Kung–Traub optimality of Steffensen–Pandrosion: derivative-free p-th root extraction with near-minimal quadratic constant," Preprint, 2026.

[4] H. T. Kung and J. F. Traub, "Optimal order of one-point and multipoint iteration," *J. Assoc. Comput. Mach.*, vol. 21, no. 4, pp. 643–651, 1974.

[5] J. F. Steffensen, "Remarks on iteration," *Skandinavisk Aktuarietidskrift*, vol. 16, pp. 64–72, 1933.

[6] J. Stoer and R. Bulirsch, *Introduction to Numerical Analysis*, 3rd ed., Springer, New York, 2002.

[7] J. F. Traub and H. Woźniakowski, *A General Theory of Optimal Algorithms*, Academic Press, New York, 1980.