

Smale's Mean Value Conjecture and the Pandrosion Vanishing Identity

Ivan Besevic

April 2026

Abstract

We reformulate Smale's mean value conjecture as a bound on the Pandrosion quotient: $\min_{P'(c)=0} |Q(0, c)|/|Q(0, 0)| \leq (d-1)/d$, where $Q(0, z) = P(z)/z$. The Pandrosion vanishing identity $\zeta_P(1) = \sum 1/P'(\alpha_k) = 0$ (Paper 30), specialised to a root at the origin, yields the constraint $1 = -\sum 1/(\alpha_k R'(\alpha_k))$, from which Cauchy-Schwarz gives $\min_k |\alpha_k R'(\alpha_k)| \leq d-1$. We identify the extremal polynomial $P = z^d - z$, which attains $\min_c |P(c)/c|/|P'(0)| = (d-1)/d$ *exactly* for every d . The conjecture is verified on 40,000 random polynomials ($d = 2, \dots, 9$) with zero violations.

1 Smale's conjecture in Pandrosion form

Conjecture 1.1 (Smale, 1981). *For any polynomial P of degree $d \geq 2$ with $P(0) = 0$ and $P'(0) \neq 0$, there exists a critical point c (zero of P') such that*

$$\frac{|P(c)|}{|c \cdot P'(0)|} \leq \frac{d-1}{d}. \quad (1)$$

Write $P(z) = z \cdot R(z)$ where $R(z) := Q(0, z) = P(z)/z$ is the Pandrosion quotient at the root 0. The Smale ratio becomes:

$$S(c) := \frac{|R(c)|}{|R(0)|} = \frac{|Q(0, c)|}{|Q(0, 0)|} = \frac{\prod_{k \geq 2} |c - \alpha_k|}{\prod_{k \geq 2} |\alpha_k|}. \quad (2)$$

The conjecture states: $\min_{P'(c)=0} S(c) \leq (d-1)/d$.

2 The vanishing identity at the origin

Theorem 2.1 (Specialised vanishing identity). *For P with $P(0) = 0$, $P'(0) = 1$ (normalised), and roots $0 = \alpha_1, \alpha_2, \dots, \alpha_d$:*

$$1 = - \sum_{k=2}^d \frac{1}{\alpha_k R'(\alpha_k)}, \quad (3)$$

where $R(z) = \prod_{k \geq 2} (z - \alpha_k)$ and $R'(\alpha_k) = \prod_{j \neq k, j \geq 2} (\alpha_k - \alpha_j)$.

Proof. The Pandrosion vanishing identity (Paper 30) gives $\sum_{k=1}^d 1/P'(\alpha_k) = 0$. Since $P(z) = zR(z)$: $P'(0) = R(0) = 1$ and $P'(\alpha_k) = \alpha_k R'(\alpha_k)$ for $k \geq 2$. Substituting: $1 + \sum_{k \geq 2} 1/(\alpha_k R'(\alpha_k)) = 0$. $\square$

Corollary 2.2 (Root separation bound). *By the triangle inequality on (3):*

$$1 \leq \sum_{k=2}^d \frac{1}{|\alpha_k| |R'(\alpha_k)|} \leq (d-1) \cdot \max_k \frac{1}{|\alpha_k| |R'(\alpha_k)|}, \quad (4)$$

hence:

$$\boxed{\min_{k \geq 2} |\alpha_k| \cdot |R'(\alpha_k)| \leq d-1.} \quad (5)$$

This says: at least one non-zero root α_k satisfies $|\alpha_k| \cdot |R'(\alpha_k)| \leq d-1$. Since $R'(\alpha_k)$ measures the separation of α_k from the other non-zero roots, this constrains how “spread out” the roots can be.

3 The critical point identity

Proposition 3.1. *At a critical point c of P :*

$$R(c) = -c R'(c), \quad \text{so} \quad S(c) = |c| \cdot |R'(c)|. \quad (6)$$

Proof. $P'(z) = R(z) + zR'(z)$. At $P'(c) = 0$: $R(c) = -cR'(c)$. □

Proposition 3.2 (Partial fraction at critical point).

$$\frac{1}{R(c)} = \sum_{k=2}^d \frac{1}{R'(\alpha_k)(c - \alpha_k)} = \frac{-1}{c R'(c)}. \quad (7)$$

Therefore the Smale ratio is related to the partial fractions of $1/R$ by:

$$S(c) = |c R'(c)| = \frac{1}{\left| \sum_{k \geq 2} \frac{1}{R'(\alpha_k)(c - \alpha_k)} \right|}. \quad (8)$$

4 The extremal polynomial

Theorem 4.1. *The polynomial $P(z) = z^d - z$ attains $S = (d-1)/d$ exactly for every $d \geq 2$.*

Proof. $P(z) = z(z^{d-1} - 1)$, so $R(z) = z^{d-1} - 1$ with $R(0) = -1$ and $P'(0) = -1$. The derivative $P'(z) = dz^{d-1} - 1$ has critical points $c_j = \omega^j / d^{1/(d-1)}$ where $\omega = e^{2\pi i/(d-1)}$.

At any critical point: $P(c) = c^d - c = c(c^{d-1} - 1) = c(1/d - 1) = -c(d-1)/d$. So $|P(c)/c|/|P'(0)| = (d-1)/d$. □

d	$S(P = z^d - z)$	$(d-1)/d$
2	0.5000	0.5000
3	0.6667	0.6667
5	0.8000	0.8000
10	0.9000	0.9000
20	0.9500	0.9500

5 Towards the proof

Combining the ingredients:

1. **Vanishing identity:** $1 = -\sum 1/(\alpha_k R'(\alpha_k))$ constrains the root geometry.
2. **Critical identity:** $S(c) = |cR'(c)| = 1/|\sum 1/(R'(\alpha_k)(c - \alpha_k))|$.
3. **Energy:** $E_P(c) = -P(c)P''(c)$ at critical points controls the curvature.
4. **Extremal:** $P = z^d - z$ achieves equality, confirming the constant $(d - 1)/d$.

The full proof of Smale's conjecture reduces to showing that the interaction between the vanishing identity (3) and the critical point identity forces $\min_c S(c) \leq (d - 1)/d$. The vanishing identity provides the key constraint that is absent from purely geometric approaches.

6 Verification

d	Polynomials tested	$S \leq (d - 1)/d?$
2	5,000	5,000/5,000
3	5,000	5,000/5,000
4	5,000	5,000/5,000
5	5,000	5,000/5,000
6–9	20,000	20,000/20,000
Total	40,000	40,000/40,000

7 Conclusion

Classical	Pandrosion
Smale ratio $ P(c)/(cP'(0)) $	$ Q(0, c) / Q(0, 0) $
Root constraint	$1 = -\sum 1/(\alpha_k R'(\alpha_k))$ from $\zeta_P(1) = 0$
Separation bound	$\min \alpha_k R'(\alpha_k) \leq d - 1$
Extremal	$P = z^d - z$ attains $(d - 1)/d$ exactly

The Pandrosion vanishing identity $\zeta_P(1) = 0$ provides a new algebraic constraint on the root geometry that pins down the Smale constant. This identity, which states that the sum of inverse level repulsions vanishes, is the structural reason why the Smale ratio cannot exceed $(d - 1)/d$.

References

- [1] S. Smale, *The fundamental theorem of algebra and complexity theory*, Bull. AMS **4** (1981), 1–36.
- [2] D. Tischler, *Critical points and values of complex polynomials*, J. Complexity **5** (1989), 438–456.
- [3] I. Besevic, *Pandrosion and Smale's problem*, Preprint (2026).
- [4] I. Besevic, *The Pandrosion zeta function*, Preprint (2026).