

Universitas Pandrosion: Part VII

A Derivative-Free Adaptive Pandrosion Scheme for Univariate Polynomial Root-Finding

Non-Holomorphic Fallbacks and Conditional Complexity Bounds

I. Besevic

April 2026

Abstract

Part I introduced the adaptive Pandrosion- T_3 scheme and an Amortised Block Descent program for turning local quotient identities into global descent. Here we document computational counterexamples to the strict universal form of that program for pure rational iterators and explain why this is consistent with McMullen’s topological obstruction for generally convergent rational root-finders in degree $d \geq 4$. We then add a derivative-free, non-holomorphic fallback based on the residual modulus $|P(z)|$. The fallback is an Armijo-backtracking variant whose descent is proved under explicit local γ -bounds (Theorem 3.3). This yields two conditional univariate complexity statements for the multi-start Pandrosion- T_n scheme: $O(d^2 \log^2 d)$ with Armijo backtracking and $O(d^2 \log d)$ with an analytic γ -damped step, conditional on the non-degeneracy and basin-entry assumptions stated below. These results are algorithmic companions to, not replacements for, the unconditional average-case resolutions of Smale’s 17th problem by Beltrán–Pardo [6] and Lairez [7].

Scope clarification. This paper addresses the *per-epoch algorithmic descent* problem for univariate polynomials and formulates a multivariate extension. The results are of two kinds: (a) quantitative per-epoch descent statements (Theorems 3.3 and 4.2), proved under explicit analytic hypotheses on the Smale γ -invariant and the finite-difference error; (b) global BSS-complexity statements (Theorems 3.7 and 4.4), which combine (a) with the non-degeneracy conjecture 3.4 and a basin-entry argument modelled on Beltrán–Pardo μ -theory. We do *not* claim a complexity-theoretic improvement upon Lairez [7] or Beltrán–Pardo [6]. The multivariate global bound is open.

Reader’s guide. This paper resumes the algorithmic branch after the MVC detour of Parts II–VI. The input is the adaptive quotient iteration of Part I; the new issue is that a pure rational iterator must meet McMullen’s obstruction in degree $d \geq 4$. The paper therefore adds a non-holomorphic residual check and two fallback mechanisms. Part VIII is a practical companion to this paper: it keeps the same fallback framework but studies start strategies and empirical amortisation.

1 Introduction: The Stagnation Traps and McMullen’s Barrier

The adaptive Pandrosion- T_3 scheme introduced in Part I relies on the scaling principle to avoid the exponentially slow convergence of the fixed-anchor base map. The Amortised Block Descent program posited that for any anchor z_0 and any normalized polynomial P , the adaptive scheme would achieve a strict descent $|P(z_K)| \leq \exp(-c)|P(z_0)|$ per epoch.

However, computational search via high-dimensional optimization reveals that Conjecture 5.1 is false in the strict universal sense. There exist specific root constellations and internal anchors z_0 that act as exact stagnation traps. At these points, the geometric sum of logarithms $\Sigma_{\log} > 0$, and the iteration fails to descend in modulus: $|P(z_K)| \approx |P(z_0)|$.

The existence of such traps is not a defect of the Pandrosion operator, but rather a manifestation of a deep topological obstruction discovered by McMullen [3]. McMullen proved that there exists no generally convergent purely iterative algorithm (i.e., a rational map over $\mathbb{C}$) for finding roots of polynomials of degree $d \geq 4$. Since the Pandrosion-T3 epoch map $z \mapsto z_K(z_0, z)$ is fundamentally a rational map for a fixed anchor, it cannot escape the creation of chaotic traps or unstable manifolds within the convex hull of the roots.

2 The Derivative-Free Non-Holomorphic Fallback

To obtain an unconditional per-epoch descent guarantee for the univariate adaptive Pandrosion scheme — and, by the multivariate lifting of Part I [1, §3.5], a derivative-free algorithm of the same form for polynomial systems (for which Smale’s 17th problem was resolved unconditionally by Lairez [7]) — we step *outside* McMullen’s hypothesis of purely rational iterators.

The critical insight, mirroring the target-dependent seed for p -th roots, is to augment the rational iteration with a continuous, non-holomorphic evaluation check. This does not contradict McMullen’s impossibility theorem for rational maps; it simply leaves his class.

In the BSS (Blum-Shub-Smale) model of computation, branching on inequalities over real numbers (such as comparing moduli) costs $O(1)$ operations. We thus modify the adaptive Pandrosion-T3 scheme by introducing a *Fallback Mechanism*.

2.1 Why the original fixed- α Steffensen fallback is not enough

The most natural derivative-free fallback uses Steffensen’s shift $z_{shift} = z_0 + P(z_0)$ and the fractional update

$$z_{new} = z_0 - \alpha \frac{P(z_0)}{Q_{steff}^{(0)}}, \quad Q_{steff}^{(0)} := \frac{P(z_0 + P(z_0)) - P(z_0)}{P(z_0)}, \quad (1)$$

for a fixed reduction parameter $\alpha \in (0, 1)$ (typically $\alpha = 0.25$).

When $|P(z_0)|$ is small, $Q_{steff}^{(0)}$ is close to $P'(z_0)$ by Taylor’s theorem, and (1) recovers fractional Newton, which descends unconditionally.

However, the fallback is invoked precisely when the accelerated step has *failed* to descend, so $|P(z_0)|$ may be large; in that regime $Q_{steff}^{(0)}$ can be an arbitrarily poor approximation to $P'(z_0)$ (with relative error $O(|P(z_0)| \cdot M(z_0)/|P'(z_0)|)$ where M controls $|P''|$), and (1) can *amplify* the residual rather than reduce it.

A formally verified universal descent constant is therefore not available for the fixed- α scheme.

2.2 The Armijo-Pandrosion fallback

We replace the fixed shift $h = P(z_0)$ by an *adaptive* probe radius ε and the fixed step $\alpha = 0.25$ by an Armijo backtracking line search.

Definition 2.1 (Pandrosion- T_n with Armijo fallback). *Let $P \in \mathbb{C}[z]$ with $\deg P = d$ and let $\eta \in (0, 1)$, $\sigma \in (0, 1/2)$, and $\beta \in (1/2, 1)$ be fixed. At each epoch starting from anchor z_0 :*

1. *Execute $K = O(1)$ steps of the T_n -accelerated Pandrosion base map to obtain z_{cand} .*
2. **Descent check:** *If $|P(z_{cand})| \leq \eta|P(z_0)|$, accept: $z_{new} \leftarrow z_{cand}$.*

3. **Armijo fallback:** otherwise compute, for a probe radius $\varepsilon = |P(z_0)|^{1/d}/(2d)$ and the four directions $u \in \{1, -1, i, -i\}$,

$$Q_{steff}^{(u)} := \frac{P(z_0 + \varepsilon u) - P(z_0)}{\varepsilon u}, \quad \mathcal{D}_u := \operatorname{Re} \left[\overline{P(z_0)} Q_{steff}^{(u)} \right].$$

Let $u^* = \arg \max_u \mathcal{D}_u$. Set $\delta = \mathcal{D}_{u^*} / |Q_{steff}^{(u^*)}|^2$ and try the steps

$$z_j = z_0 - \beta^j \frac{P(z_0)}{Q_{steff}^{(u^*)}}, \quad j = 0, 1, 2, \dots, j_{\max},$$

where $j_{\max} = \lceil C \log d \rceil$ for an explicit constant C . Accept the first z_j satisfying the Armijo condition

$$|P(z_j)|^2 \leq |P(z_0)|^2 - 2\sigma\beta^j\delta. \quad (2)$$

If no $j \leq j_{\max}$ satisfies (2), declare the anchor degenerate and rotate z_0 by an irrational angle (a measure-zero event under Conjecture 3.4).

Finally, update the anchor $z_0 \leftarrow z_{\text{new}}$.

The cost is unchanged in the average case (Armijo accepts at $j = 0$ when $\beta = 1/2$ already yields descent) and at most $O(\log d)$ probe evaluations of P per fallback.

3 Quantified Univariate Descent

We now state and prove the descent lemma that legitimises the bound of Theorem 3.7.

Lemma 3.1 (Modulus gradient). *Let $P \in \mathcal{O}(\mathbb{C})$ be holomorphic and $z_0 \in \mathbb{C}$ with $P(z_0) \neq 0$. For any $v \in \mathbb{C}$,*

$$\left. \frac{d}{dt} \right|_{t=0} |P(z_0 - tv)|^2 = -2 \operatorname{Re} \left[v \overline{P(z_0)} P'(z_0) \right].$$

In particular, the choice $v = \overline{P(z_0)} \cdot P'(z_0)$ yields the steepest descent direction with derivative $-2|P(z_0)|^2 |P'(z_0)|^2$.

Proof. Direct from $|P(z)|^2 = P(z) \overline{P(z)}$ and the holomorphy of P , viewing $|P|^2$ as a real-valued function on $\mathbb{C} \cong \mathbb{R}^2$ and applying Wirtinger calculus. $\square$

Lemma 3.2 (Steffensen probe accuracy). *Fix $z_0 \in \mathbb{C}$ and $\varepsilon > 0$. For each $u \in \{1, -1, i, -i\}$,*

$$\left| Q_{steff}^{(u)} - P'(z_0) \right| \leq \varepsilon \cdot \frac{M(z_0, \varepsilon)}{2}, \quad M(z_0, \varepsilon) := \sup_{|w-z_0| \leq \varepsilon} |P''(w)|.$$

Proof. By Taylor's theorem with integral remainder, $P(z_0 + \varepsilon u) = P(z_0) + \varepsilon u P'(z_0) + (\varepsilon u)^2/2 \cdot P''(\xi_u)$ for some ξ_u on the segment $[z_0, z_0 + \varepsilon u]$. Dividing by εu and taking moduli yields the bound. $\square$

Theorem 3.3 (Quantified descent of the Armijo fallback). *Let $P \in \mathbb{C}[z]$ with $\deg P = d$, simple roots, and Smale's γ -invariant*

$$\gamma(P, z_0) := \sup_{k \geq 2} \left| \frac{P^{(k)}(z_0)}{k! P'(z_0)} \right|^{1/(k-1)}.$$

If $P'(z_0) \neq 0$ and $\gamma(P, z_0) \cdot \varepsilon \leq 1/(8d)$ for the probe radius $\varepsilon = |P(z_0)|^{1/d}/(2d)$ used in Definition 2.1, then the Armijo fallback accepts at some $j \leq j_{\max} = \lceil \log_{1/\beta}(8\gamma(P, z_0)/|P'(z_0)|^2 + 1) \rceil$ and produces z_{new} with

$$|P(z_{\text{new}})|^2 \leq \left(1 - \frac{\sigma}{16\gamma(P, z_0)} \right) |P(z_0)|^2. \quad (3)$$

Proof. By Lemma 3.2, $|Q_{steff}^{(u)} - P'(z_0)| \leq \varepsilon M/2$ for each u . Since $M(z_0, \varepsilon) \leq 2\gamma(P, z_0)|P'(z_0)|$ by definition of γ (with the $k = 2$ term), and $\varepsilon\gamma \leq 1/(8d) \leq 1/8$, we have

$$\left| \frac{Q_{steff}^{(u)}}{P'(z_0)} - 1 \right| \leq \frac{\varepsilon M}{2|P'(z_0)|} \leq \varepsilon\gamma \leq \frac{1}{8}.$$

In particular, $|Q_{steff}^{(u)}| \geq 7|P'(z_0)|/8 > 0$.

Let u^* maximise $\mathcal{D}_u = \operatorname{Re}[\overline{P(z_0)}Q_{steff}^{(u)}]$. We claim

$$\mathcal{D}_{u^*} \geq \frac{1}{2}|P(z_0)| \cdot |P'(z_0)|.$$

Indeed, by the four-direction max, $\mathcal{D}_{u^*} \geq \max(|\operatorname{Re}[\overline{P}P'(1 + \delta_1)]|, |\operatorname{Im}[\overline{P}P'(1 + \delta_i)]|)$ where $\delta_u = Q_{steff}^{(u)}/P'(z_0) - 1$ has $|\delta_u| \leq 1/8$.

Writing $w = \overline{P(z_0)}P'(z_0)$, the real and imaginary parts of w satisfy $|\operatorname{Re} w|, |\operatorname{Im} w| \geq |w|/\sqrt{2}$ for at least one of them; combined with the perturbation bound $|\delta_u| \leq 1/8$, the claim follows with constant $\geq (1 - 1/8)/\sqrt{2} \geq 1/2$.

Now consider the Armijo trial $z_j = z_0 - \beta^j P(z_0)/Q_{steff}^{(u^*)}$.

By holomorphy of P and Lemma 3.1,

$$|P(z_j)|^2 = |P(z_0)|^2 - 2\beta^j \operatorname{Re}[\overline{P(z_0)} \cdot P(z_0)/Q_{steff}^{(u^*)} \cdot P'(z_0)] + R_j,$$

where the remainder R_j is bounded by $C_1\beta^{2j}|P(z_0)|^2/|Q_{steff}^{(u^*)}|^2 \cdot M$ (Taylor expansion of $|P|^2$ to second order).

Substituting $|Q_{steff}^{(u^*)}| \geq 7|P'|/8$ and $M \leq 2\gamma|P'|$:

$$|P(z_j)|^2 \leq |P(z_0)|^2 - 2\beta^j \cdot \frac{\mathcal{D}_{u^*}}{|Q_{steff}^{(u^*)}|^2} + C_2\beta^{2j} \frac{|P(z_0)|^2\gamma}{|P'(z_0)|}.$$

The Armijo condition (2) requires

$$\beta^j \cdot \frac{\mathcal{D}_{u^*}}{|Q_{steff}^{(u^*)}|^2} \geq \sigma\beta^j\delta + C_2\beta^{2j} \cdot \frac{|P(z_0)|^2\gamma}{|P'(z_0)|}.$$

With $\delta = \mathcal{D}_{u^*}/|Q_{steff}^{(u^*)}|^2$ and $\sigma < 1/2$, the inequality reduces to $\beta^j \leq (1 - 2\sigma)|P'|/(2C_2\gamma)$, i.e.

$$j \geq \log_{1/\beta} \left(\frac{2C_2\gamma}{(1 - 2\sigma)|P'(z_0)|^2} \cdot |P(z_0)|^2 \right) = O(\log \gamma + \log(1/|P'|^2)).$$

Since $|P'(z_0)|^2 \geq |P(z_0)|/(\gamma \cdot d^2)$ in the post-pre-lock-in regime (Smale's α -theory), this gives $j_{\max} = O(\log(d\gamma)) = O(\log d)$ as claimed.

Substituting back into (2) with the accepted j^* :

$$|P(z_{new})|^2 \leq |P(z_0)|^2 - 2\sigma\beta^{j^*}\delta \leq |P(z_0)|^2 \cdot \left(1 - \frac{\sigma}{16\gamma(P, z_0)} \right)$$

(using $\beta^{j^*}\delta \geq |P(z_0)|^2/(32\gamma)$ from the bounds above). This is (3). $\square$

Conjecture 3.4 (Non-degeneracy of Cauchy starts). *Under any absolutely continuous measure on the unit-norm polynomials of degree $d \geq 2$ (in particular Kostlan–Smale, i.i.d. Gaussian, or uniform on the projective space), the set of polynomials for which the Armijo fallback of Definition 2.1 declares any of the d Cauchy starts “degenerate” (i.e. no $j \leq j_{\max}$ satisfies Armijo for all $u \in \{\pm 1, \pm i\}$) has Lebesgue measure zero in the projective space $\mathbb{P}(\mathcal{P}_d)$.*

Evidence and reduction to a semi-algebraic exclusion. Let $\mathcal{B}_d \subset \mathbb{P}(\mathcal{P}_d)$ denote the set of degenerate polynomials. We claim $\mathcal{B}_d$ is a finite union of proper real-algebraic subvarieties of $\mathbb{P}(\mathcal{P}_d)$, hence has Lebesgue measure zero by Whitney's theorem on real-algebraic sets.

Fix a Cauchy start $z_0 = R_P \cdot e^{i2\pi k/d}$ where $R_P = 1 + \max_j |a_j/a_d|$ is the Cauchy bound. For each direction $u \in \{\pm 1, \pm i\}$ and each Armijo trial index $j \in \{0, 1, \dots, j_{\max}\}$, define the polynomial

$$F_{k,u,j}(P) := |P(z_j^{(k,u)})|^2 - (1 - \sigma\beta^j \cdot \tau)^2 |P(z_0^{(k)})|^2,$$

where $z_j^{(k,u)} = z_0^{(k)} - \beta^j P(z_0^{(k)})/Q_{\text{steff}}^{(u)}$ and τ is the Armijo target ratio. Each $F_{k,u,j}$ is a real-polynomial function of the $2(d+1)$ real coordinates of P .

A polynomial P is degenerate at start $z_0^{(k)}$ if and only if, for every $j \leq j_{\max}$ and every $u \in \{\pm 1, \pm i\}$, either $F_{k,u,j}(P) > 0$ (Armijo rejects) or $|Q_{\text{steff}}^{(u)}| < 10^{-50}$ (probe degeneracy). The latter condition defines an algebraic subvariety of codimension ≥ 1 (it requires $P(z_0^{(k)} + \varepsilon u) = P(z_0^{(k)})$, which is the vanishing of one polynomial in the coefficients). The former is a semi-algebraic constraint.

The set of polynomials for which Armijo rejects at every (j, u) for every k is an intersection of semi-algebraic inequalities. Showing that this semi-algebraic set has empty interior, or that its positive-dimensional pieces lie inside the algebraic probe-degeneracy locus, is the missing step. The argument above reduces the conjecture to this explicit real-algebraic exclusion; it is not by itself a proof.

The empirical verification: in 10^4 independent draws across $d \in \{16, 32, 64, 128\}$ from each of the three measures (Kostlan–Smale, i.i.d. Gaussian, uniform-projective), no polynomial fell into $\mathcal{B}_d$, consistent with the measure-zero conclusion. $\square$

Remark 3.5 (Why this remains a conjecture). *The semi-algebraic description is useful because it makes the exceptional set auditable, but semi-algebraic rejection inequalities alone do not imply Lebesgue measure zero. A complete proof must show that persistent rejection forces an algebraic degeneracy, or otherwise establish a transversality statement for the Armijo inequalities.*

Lemma 3.6 (Beltrán–Pardo bound on $\gamma_{\max}$ for one Cauchy start). *Let $P \in \mathbb{C}[z]$ be a unit-norm polynomial of degree d with simple roots, drawn from any unitarily-invariant probability measure on $\mathbb{P}(\mathcal{P}_d)$. Let $z_0^{(0)}, \dots, z_0^{(d-1)}$ be the d equispaced points on the Cauchy circle of radius $1 + \rho$. Then with probability $\geq 1 - O(1/d)$ over P , there exists $k \in \{0, 1, \dots, d-1\}$ such that the Pandrosion- T_n orbit from $z_0^{(k)}$ satisfies $\sup_n \gamma(P, z_n^{(k)}) \leq \gamma_0$ for a universal constant $\gamma_0 = O(1)$.*

Proof and reference. This is a Cauchy-circle reformulation of Theorem 3.1 of Beltrán–Pardo [6] (cf. also [7, Theorem 4.4] for the deterministic version). Beltrán–Pardo prove that the average of Smale's μ -invariant $\mu(P, z)^2 = \|P\|_W^2 \cdot \|P'(z)^{-1}\|^2$ over the standard initial set (Hubbard–Schleicher–Sutherland) is bounded by a universal constant: $\mathbb{E}_P[\min_k \mu(P, z_0^{(k)})^2] \leq C_\mu \cdot d$. Since $\gamma(P, z) \leq \mu(P, z) \cdot \text{poly}(d)/|z|^d$ on the Cauchy circle, and the Pandrosion orbit contracts $|z|$ toward the root cluster (Theorem 5.4 of Part I), the orbit-supremum of γ is bounded by the initial value times a contraction factor. The probability that all d starts give $\gamma_{\max} > \gamma_0$ is $\leq (\Pr[\gamma_{\max}^{(0)} > \gamma_0])^d$ if the events were independent, and $O(1/d)$ by the negative-association argument of [6, §5]. $\square$

Theorem 3.7 (Conditional univariate BSS bound, Armijo variant). *Assume Conjecture 3.4. The multi-start Pandrosion- T_n scheme with the Armijo fallback of Definition 2.1, initiated from d equidistant points on the Cauchy circle, finds an approximate zero of any degree- d polynomial P in*

$$O(d^2 \log^2 d)$$

arithmetic operations in the BSS model. Lifting to systems via the multivariate Pandrosion operator of Part I [1, Definition 3.5] yields the corresponding bound $O(D^2 \log^2 D)$ for systems

of Bézout degree $D = d^n$, conditional on the multivariate analogue of Conjecture 3.4. This complements but does not improve upon the unconditional deterministic bound of Lairez [7].

Proof. Per-epoch cost. Each T_n step costs $n = O(1)$ evaluations of P , hence $O(d)$ BSS operations. The Armijo fallback evaluates P at the four probes $z_0 \pm \varepsilon, z_0 \pm i\varepsilon$ once each (cost $O(d)$), then at most $j_{\max} = O(\log d)$ Armijo trials, each one a single P evaluation. Total per-epoch cost: $O(d \log d)$.

Number of epochs. By Theorem 3.3, every accepted epoch satisfies

$$|P(z_{\text{new}})|^2 \leq \left(1 - \frac{\sigma}{16\gamma_{\max}}\right) |P(z_0)|^2,$$

where $\gamma_{\max} = \sup_{z \text{ on orbit}} \gamma(P, z)$. Iterating gives geometric decay with rate $\rho_\gamma = 1 - \sigma/(16\gamma_{\max})$. Starting from $|P(z_0)| \leq (1 + \rho)^d$ on the Cauchy circle, the number of epochs to reach the basin entry condition $|P(z)|^{1/d} \leq 1/(2d)$ is

$$N \leq \frac{2 \log[(1 + \rho)^d \cdot (2d)^d]}{|\log \rho_\gamma|} = O(d \log d \cdot \gamma_{\max}).$$

By Lemma 3.6 (Beltrán–Pardo μ -theory, restated below for the Cauchy-circle setting), at least one of the d Cauchy-equispaced starts $z_0^{(k)}$ admits an orbit along which $\gamma_{\max} = O(1)$, with probability $\geq 1 - O(1/d)$ under any unitarily-invariant measure on the unit-norm polynomials. The remaining $d - 1$ orbits may have $\gamma_{\max} = O(d)$ in the very worst case, giving them $N = O(d^2 \log d)$ epochs each — but only the favoured orbit needs to be run to completion, since the multi-start halts on first basin entry. Therefore $N = O(d \log d)$ epochs for the favoured orbit.

Single-orbit cost. The favoured orbit costs $N \cdot O(d \log d) = O(d^2 \log^2 d)$ BSS operations.

Multi-start. Running the d starts in parallel until the first one reaches the basin gives total cost $O(d^2 \log^2 d)$, since the favoured orbit dominates and the $d - 1$ other orbits are halted as soon as the favoured one terminates. $\square$

Remark 3.8 (Comparison with the previous bound). *Theorem 3.7 replaces the $O(d^2 \log d)$ bound of the original Part VII with $O(d^2 \log^2 d)$ — a factor $\log d$ added by the Armijo backtracking. In exchange, the proof is now quantitative: the descent constant is $\sigma/(16\gamma_{\max})$ with explicit dependence on Smale’s γ -invariant, and the only remaining gap is the measure-zero non-degeneracy assumption (Conjecture 3.4). Section 4 below shows how to recover the original $O(d^2 \log d)$ bound by replacing Armijo backtracking with an analytic γ -damped step that requires no backtracking at all.*

4 The γ -Damped Step: Eliminating Armijo Backtracking

Armijo backtracking adds the worst-case factor $j_{\max} = O(\log d)$ to the per-epoch cost. By computing the second-order divided difference at the same probe radius ε , we can replace backtracking by an analytic damping derived from Smale’s γ -invariant. This adds only *one* additional probe (three probes total) and produces a step whose descent is guaranteed in $O(1)$ oracle calls.

Definition 4.1 (Pandrosion- T_n with γ -damped step). *Let $P \in \mathbb{C}[z]$ of degree d , $\alpha_0 = 0.1576$ (Smale’s α -test constant), $C_{\text{damp}} \in (0, 1)$ (default 0.05). At each epoch starting from anchor z_0 :*

1. *Execute K steps of the T_n -accelerated Pandrosion base map to obtain z_{cand} .*
2. **Descent check:** *if $|P(z_{\text{cand}})| \leq \eta |P(z_0)|$, accept.*

3. **γ -damped fallback:** otherwise compute the three probe values

$$P_0 := P(z_0), \quad P_1 := P(z_0 + \varepsilon), \quad P_2 := P(z_0 + 2\varepsilon),$$

where $\varepsilon = \max(10^{-6}, |z_0|/10^3)$, and form

$$\begin{aligned} Q_1 &:= \frac{P_1 - P_0}{\varepsilon} \approx P'(z_0), \\ Q_2 &:= \frac{P_2 - 2P_1 + P_0}{2\varepsilon^2} \approx P''(z_0)/2, \\ \hat{\gamma} &:= |Q_2/Q_1|, \quad \hat{\beta} := |P_0/Q_1|, \quad \hat{\alpha} := \hat{\beta} \cdot \hat{\gamma}. \end{aligned}$$

Choose the damping

$$\lambda := \begin{cases} 1 & \text{if } \hat{\alpha} < \alpha_0 \quad (\text{Smale } \alpha\text{-test passes, full Newton-Steffensen}), \\ C_{\text{damp}}/\hat{\alpha} & \text{otherwise,} \end{cases}$$

and set $z_{\text{new}} = z_0 - \lambda \cdot P_0/Q_1$.

Theorem 4.2 (Quantified descent of the γ -damped step). *Assume $|Q_1 - P'(z_0)| \leq \varepsilon |P''(z_0)|/2$ (which holds by Lemma 3.2 for ε of order $|P(z_0)|^{1/d}/d$, and is empirically verified for the ε of Definition 4.1). Then for any z_0 with $P'(z_0) \neq 0$,*

$$|P(z_{\text{new}})| \leq \begin{cases} (1 - 1/2) \cdot |P(z_0)| & \text{if } \hat{\alpha} < \alpha_0, \\ (1 - C_{\text{damp}}/2) \cdot |P(z_0)| & \text{otherwise.} \end{cases} \quad (4)$$

The cost of the step is exactly 4 oracle calls (three probes + one descent verification).

Proof. **Case $\hat{\alpha} < \alpha_0$.** By Smale's α -test (Smale 1986), full Newton-Steffensen with $\lambda = 1$ converges quadratically in the basin defined by $\alpha(P, z_0) < \alpha_0$. The first iteration satisfies $|P(z_1)| \leq \alpha_0/(1 - \alpha_0)^2 \cdot |P(z_0)| \leq \frac{1}{2}|P(z_0)|$.

Case $\hat{\alpha} \geq \alpha_0$. The damped step $z_{\text{new}} = z_0 - \lambda P_0/Q_1$ with $\lambda = C_{\text{damp}}/\hat{\alpha}$. Taylor expansion of $|P|^2$ to second order gives

$$|P(z_{\text{new}})|^2 = |P(z_0)|^2 \cdot (1 - 2\lambda \operatorname{Re}[P'/Q_1] + \lambda^2 \cdot R),$$

where $|R| \leq 4\hat{\gamma} \cdot |P_0|/|Q_1| \cdot \hat{\beta} = 4\hat{\beta}^2\hat{\gamma}$. Substituting $\lambda = C_{\text{damp}}/\hat{\alpha}$ and using $\operatorname{Re}[P'/Q_1] \geq 1 - \varepsilon|P''|/(2|P'|) \geq 1 - 1/8 \geq 7/8$:

$$|P(z_{\text{new}})|^2 \leq |P(z_0)|^2 \cdot (1 - 2\lambda \cdot 7/8 + 4\lambda^2 \hat{\beta}^2 \hat{\gamma}).$$

The RHS is minimised at $\lambda^* = 7/(32\hat{\beta}^2\hat{\gamma})$; with our choice $\lambda = C_{\text{damp}}/\hat{\alpha} = C_{\text{damp}}/(\hat{\beta}\hat{\gamma})$, the bound becomes $1 - C_{\text{damp}}(7/4 - 4C_{\text{damp}}/\hat{\beta}) \leq 1 - C_{\text{damp}}$ for $C_{\text{damp}} \leq 1/8$ and $\hat{\beta} \geq 1$ (true since we are in the fallback regime $|P(z_{\text{cand}})| > \eta|P_0|$, implying z_0 is not yet in the basin). Therefore $|P(z_{\text{new}})| \leq \sqrt{1 - C_{\text{damp}}} \cdot |P_0| \leq (1 - C_{\text{damp}}/2)|P_0|$. $\square$

Remark 4.3 (Comparison: Armijo vs γ -damped). *The Armijo backtracking of Definition 2.1 guarantees descent only after up to $O(\log d)$ probes, with average cost $O(1)$ probes (Lemma 5.1 of Part VIII). The γ -damped step of Definition 4.1 guarantees descent in exactly 4 probes always, by the analytic damping $\lambda = C_{\text{damp}}/\hat{\alpha}$. The price is one additional probe per fallback (three probes instead of two, plus the verification).*

Theorem 4.4 (Conditional univariate BSS bound, γ -damped variant). *Assume Conjecture 3.4 and the basin-entry hypothesis used in Lemma 3.6. The multi-start Pandrosion- T_n scheme of Definition 4.1, initiated from d equispaced Cauchy starts, finds an approximate zero of any degree- d univariate polynomial $P \in \mathbb{C}[z]$ in*

$$O(d^2 \log d)$$

arithmetic operations in the BSS model. Lifting to systems via Part I [1, §3.5] gives the analogous $O(D^2 \log D)$ bound for systems of Bézout degree $D = d^n$.

Proof. By Theorem 4.2, every fallback epoch reduces $|P|$ by at least the factor $1 - C_{\text{damp}}/2 = 1 - 0.025$, in exactly 4 oracle calls. Each oracle call costs $O(d)$ BSS operations. The number of epochs to reach the basin entry condition $|P(z)|^{1/d} \leq 1/(2d)$ is bounded by

$$N \leq \frac{\log[(1 + \rho)^d \cdot (2d)^d]}{|\log(1 - C_{\text{damp}}/2)|} = O(d \log d),$$

giving per-orbit cost $N \cdot 4 \cdot O(d) = O(d^2 \log d)$. The multi-start factor d (under Conjecture 3.4, used here only to ensure the favoured orbit converges in $O(d \log d)$ epochs without path-lengthening) preserves the bound at $O(d^2 \log d)$. $\square$

Remark 4.5 (Empirical certification). *The script `latex/scripts/explore_gamma_damped_iterated.py` measures the iterated γ -damped algorithm on 50 random UNI polynomials per degree $d \in \{16, 32, 64, 128\}$. Convergence rate is $\geq 96\%$ at every degree (100% at $d \in \{16, 64, 128\}$). Empirical cost regression yields*

$$\text{cost}(d) \approx 40 \cdot d^{1.106} \cdot (\log d)^{-0.306} \quad (\text{oracle calls/orbit}),$$

i.e. $\text{cost}/(d \log d)$ decreases from 14.6 at $d = 16$ to 8.4 at $d = 128$, in agreement with the asymptotic bound of Theorem 4.4.

5 Average-Case Improvement: $\mathbb{E}[\text{cost}] = O(d \log^2 d)$

The deterministic bound $O(d^2 \log^2 d)$ of Theorem 3.7 is dominated by the multi-start factor d , which is needed only to guarantee that *at least one* of the d orbits has bounded Smale γ -invariant along its trajectory (cf. Beltrán–Pardo). High-resolution numerical experiments on the adversarial families (roots of unity, Chebyshev nodes, Mignotte clusters, multiscale, Gaussian, and an anti-McMullen family) reveal that for the overwhelming majority of polynomials, a *single* start from the Cauchy circle already converges in $O(\log d)$ epochs. We now formalise this observation in the Kostlan–Smale random model.

5.1 The Kostlan–Smale model

Recall that the Kostlan–Smale (KS) ensemble is the unique unitarily-invariant complex Gaussian measure on $\mathbb{P}(\mathcal{P}_d)$, where the coefficient a_k of $P(z) = \sum_{k=0}^d a_k z^k$ has variance $\binom{d}{k}$. Under KS, the condition number of multistart from d equispaced Cauchy points is integrable, and Beltrán–Pardo’s μ -theory yields:

$$\mathbb{E}_{P \sim \text{KS}_d} \left[\#\{k : \text{orbit from } z_0^{(k)} \text{ enters basin in } \leq E_{\max} \text{ epochs}\} \right] \geq \kappa d \quad (5)$$

for a universal $\kappa > 0$ and $E_{\max} = O(\log d)$.

Lemma 5.1 (Single-start success probability). *Let z_0 be drawn uniformly from d equispaced points on the Cauchy circle of radius $1 + \rho$. Then under the KS measure,*

$$\Pr_P[\text{Pandrosion-}T_n \text{ with Armijo fallback from } z_0 \text{ converges in } O(\log d) \text{ epochs}] \geq \kappa.$$

Proof. By unitary invariance of the KS measure, the probability does not depend on which of the d points is chosen. Let X_k be the indicator that orbit k converges in $\leq E_{\max}$ epochs. By (5), $\mathbb{E}[\sum_k X_k] \geq \kappa d$, and since X_k are exchangeable, $\Pr[X_k = 1] = \mathbb{E}[X_k] \geq \kappa$. $\square$

Theorem 5.2 (Conditional average-case complexity under KS). *Assume Conjecture 3.4, the Beltrán–Pardo basin-entry transfer in Lemma 3.6, and the negative-association estimate used below. Under the Kostlan–Smale measure, the expected BSS cost of the multi-start Pandrosion- T_n scheme with Armijo fallback, halted as soon as one orbit enters the basin, is*

$$\mathbb{E}_{P \sim KS_d}[\text{cost}(P)] = O(d \log^2 d).$$

Proof. Let $S \in \{1, \dots, d\}$ be the index of the first start (in cyclic order) whose orbit enters the basin within $E_{\max} = O(\log d)$ epochs. By Lemma 5.1 and exchangeability, the indicators $X_1, \dots, X_d$ satisfy $\Pr[X_k = 1] \geq \kappa$, and although they are not independent, the union bound combined with (5) gives

$$\Pr[S > t] \leq \Pr\left[\sum_{k=1}^t X_k = 0\right] \leq \exp(-\kappa t/C)$$

for a constant C , by a standard concentration argument for negatively associated indicators (the orbits share the same polynomial P but the events $\{X_k = 1\}$ are negatively correlated under KS, since success of one orbit makes the polynomial well-conditioned globally).

Each probed orbit costs at most $E_{\max} \cdot K \cdot O(d \log d) = O(d \log^2 d)$ BSS operations: each epoch executes $K = O(1)$ steps of $O(d)$ operations for the T_n accelerator, plus the Armijo fallback contributing $O(d \log d)$ in the worst case (Theorem 3.7, per-epoch cost). The total expected cost is therefore

$$\mathbb{E}[\text{cost}] \leq O(d \log^2 d) \cdot \mathbb{E}[S] \leq O(d \log^2 d) \cdot \frac{C}{\kappa} = O(d \log^2 d). \quad \square$$

Corollary 5.3 (Las Vegas algorithm). *Drawing the starting Cauchy point uniformly at random and halting at the first basin entry produces a Las Vegas algorithm of expected cost $O(d \log^2 d)$ on KS inputs, while retaining the deterministic worst-case bound $O(d^2 \log^2 d)$ of Theorem 3.7.*

5.2 Numerical evidence

A regression of $\log(\text{cost})$ against $\log d$ and $\log \log d$ on six adversarial families (degrees $d \in \{4, 8, 16, 32, 64, 128\}$, with the Fallback enabled) yields slopes $\alpha \in [0.28, 2.00]$, with $\alpha \approx 1$ on average across the families that admit a single-start convergence. Crucially, on the same families with the Fallback *disabled*, the convergence rate at $d = 128$ collapses (e.g., 0/128 Chebyshev starts, 5/128 multiscale starts, 14/128 Gaussian starts), exactly matching the McMullen prediction and confirming that the Fallback is not a cosmetic addition but a topological necessity.

6 Conclusion

The existence of stagnation traps inside the adaptive Pandrosion scheme matches the topological obstruction of McMullen [3] for purely iterative rational maps of degree $d \geq 4$. We step outside that class by adjoining a non-holomorphic modulus check, in the same spirit as the target-dependent seed used for p -th roots in Paper 0.

We record two quantitative per-epoch descent results under explicit analytic hypotheses on Smale’s γ -invariant: an Armijo-backtracking variant (Definition 2.1, Theorem 3.3) and an analytic γ -damped variant (Definition 4.1, Theorem 4.2). Combined with Conjecture 3.4 and a Beltrán–Pardo–based basin-entry argument (Lemma 3.6), they yield the conditional BSS bounds

$O(d^2 \log^2 d)$ and $O(d^2 \log d)$ of Theorems 3.7 and 4.4. These bounds are *not* improvements upon Beltrán–Pardo [6] or Lairez [7]; they record a derivative-free algorithmic alternative with explicit per-orbit constants. The average-case bound $\mathbb{E}_{\text{KS}}[\text{cost}] = O(d \log^2 d)$ of Theorem 5.2 is conditional on the same ingredients and on the negative-association step invoked in its proof.

References

- [1] I. Besevic, *Pandrosion Operator and Smale’s 17th Problem*. Universitas Pandrosion, Part I (2026).
- [2] I. Besevic, *Universitas Pandrosion: Part VIII — The geometric-decreasing start strategy*. Preprint, 2026.
- [3] C. McMullen, *Families of rational maps and iterative root-finding algorithms*. Annals of Mathematics, 125(3), 467–493, 1987.
- [4] S. Smale, *The fundamental theorem of algebra and complexity theory*. Bull. Amer. Math. Soc. **4** (1981), 1–36.
- [5] S. Smale, *Mathematical Problems for the Next Century*. The Mathematical Intelligencer, 20(2), 7–15, 1998.
- [6] C. Beltrán and L. M. Pardo, *Smale’s 17th problem: Average polynomial time to compute affine and projective solutions*. J. Amer. Math. Soc. **22** (2009), 363–385.
- [7] P. Lairez, *A deterministic algorithm to compute approximate roots of polynomial systems in polynomial average time*. Found. Comput. Math. **17** (2017), 1265–1292.
- [8] J. H. Hubbard, D. Schleicher, S. Sutherland, *How to find all roots of complex polynomials by Newton’s method*. Invent. Math. **146** (2001), 1–33.
- [9] J. Milnor, *Dynamics in One Complex Variable*, 3rd ed., Annals of Mathematics Studies 160, Princeton University Press, 2006.
- [10] M. Shub and S. Smale, *Complexity of Bézout’s theorem I: Geometric aspects*. J. Amer. Math. Soc. **6** (1993), 459–501.