

A Formally Verified Pandrosion–Steffensen Scheme for $z^p = x$ on $\mathbb{R}$ and $\mathbb{C}$

*Lean 4 / Mathlib v4.7.0 development, with closed-form real-line analysis
and a companion framework of finite cyclotomic-root identities*

Ivan Besevic

Independent Mathematical Research
github.com/ivan-fr/poussiere_de_x

April 2026

Abstract

We present a Lean 4 / Mathlib v4.7.0 development of a multi-start variant of the Pandrosion–Steffensen iteration for solving $z^p = x$ with $x \in \mathbb{C}^*$ and $p \geq 1$. The scheme combines the rational iteration $h_{p,x}(z) = 1 - (x - 1)/(x S_p(z))$ (where $S_p(z) = \sum_{k=0}^{p-1} z^k$), the Aitken–Steffensen Δ^2 accelerator $\sigma = \text{Aitken}(h)$, and a perturbative seed $\text{seed}_{\alpha,\varepsilon}(z) = \alpha + \varepsilon(z - \alpha)$ whose definition depends on the user-chosen cyclotomic target α .

The paper reports three kinds of content, explicitly separated: (a) *results that are new in this formalization*, concerning the universal convergence of the multi-start scheme with an explicit closed-form iteration count, the Borel-measurability and a.e. continuity of the resulting solution map, and a set of elementary algebraic identities for the finite series $\zeta_P(s) := \sum_{k=1}^p P'(\alpha_k)^{-s}$; (b) *Lean specializations of classical theorems* (Banach 1922, Aitken 1926, Schwarz 1890, Kronecker 1857, Aberth–Ehrlich 1967/1973) to the Pandrosion setting these are not new mathematics, only new formalizations. Three further modules (*PandrosionCardano*, *PandrosionVieta*, *PandrosionGalois*) record definitional-level identities for the cyclotomic anchors of $z^p = x$ and are not counted as specializations of the classical theorems; (c) *two short structural observations*: a compliance interface for the Kung–Traub upper bound for $c \geq 3$, and a statement-level recording of McMullen 1987. Two other classical themes (Lehmer’s Mahler-measure conjecture, Smale’s 17th problem) that earlier drafts discussed have been removed from scope, because the Pandrosion setting contributes nothing to either.

The Lean development comprises 121 modules with zero `sorry` and an audited dependency closure reducing to `{propext, Classical.choice, Quot.sound}`, verifiable per-theorem via `#print axioms` (see Section 10). The main contribution of the corpus is the formalization itself and the multi-start convergence analysis of (a), not any of the engagements in (c); readers primarily interested in the mathematics should focus on Sections 2–6.

Contents

1	Introduction and scope	3
1.1	Historical starting point: Pandrosion’s two sequences for $\sqrt[3]{2}$	3
1.2	The general iterator $h_{p,x}$	3
1.3	Contributions, separated by kind	4
1.4	Related work	4
1.5	On naming and scope	5
1.6	Reading order	5
2	The Pandrosion iteration on the real line	5
2.1	Closed-form contraction ratio	5
2.2	Optimal starting point: one-step preconditioning	6

2.3	Scaling preconditioning: exploiting monotonicity	6
2.4	Aitken Δ^2 : from linear to quadratic convergence	7
2.5	A quadratic local bound at the super-attractive fixed point	8
3	Cyclotomic anchors and Galois structure	8
3.1	Cyclotomic anchor family	8
3.2	Cyclic permutation of the anchors	9
4	Universal multi-start convergence	9
4.1	The multi-start seed	9
4.2	Universal multi-start theorem	10
4.3	Explicit iteration count	10
4.4	An a.e.-convergence statement for the plain iteration	11
4.5	Arbitrary target selector	11
5	Measurability and a.e. continuity of the solution map	11
6	Elementary identities on the derivative values at cyclotomic roots	13
7	Pandrosion specializations of classical theorems	13
8	Short structural observations	14
8.1	Kung–Traub upper bound for $c \geq 3$ (open since 1974)	14
8.2	McMullen 1987 (recorded but not formalized)	14
8.3	Two classical problems we deliberately do <i>not</i> engage with	14
8.4	Summary	15
9	Structure of the Lean corpus	15
10	Reproducibility	16
11	Conclusion	16

1 Introduction and scope

1.1 Historical starting point: Pandrosion's two sequences for $\sqrt[3]{2}$

Before stating the general iterator, we recall the two sequences that Pappus (*Collectio*, Book III) attributes to Pandrosion of Alexandria for approximating $\sqrt[3]{2}$. Given an initial parameter $u_0 \in (0, 4)$, the construction produces the pair

$$u_{n+1} = \frac{u_n}{2 - 2\left(\frac{4-u_n}{4}\right)^3}, \quad v_n = 2\left(\frac{4-u_n}{4}\right)^2, \quad u_0 = 0.5. \quad (1)$$

The sequence (u_n) is the internal parameter (a length in the Thales diagram); the sequence (v_n) is the *readout*, converging to $\sqrt[3]{2}$.

Normalization. The substitution $s_n := (4 - u_n)/4 \in (0, 1)$ eliminates all constants:

$$u_n = 4(1 - s_n), \quad v_n = 2s_n^2, \quad u_{n+1} = \frac{4(1 - s_n)}{2(1 - s_n^3)} = \frac{2}{1 + s_n + s_n^2},$$

which turns (1) into the single rational iteration

$$s_{n+1} = 1 - \frac{u_{n+1}}{4} = \frac{2s_n^2 + 2s_n + 1}{2s_n^2 + 2s_n + 2}. \quad (2)$$

The fixed point of (2) is $s^* = 2^{-1/3}$, giving $v^* = 2 \cdot 2^{-2/3} = 2^{1/3} = \sqrt[3]{2}$, and the local convergence is *linear* with ratio

$$\lambda_{3,2} = \frac{(\sqrt[3]{2} - 1)(\sqrt[3]{2} + 2)}{\sqrt[3]{4} + \sqrt[3]{2} + 1} \approx 0.2202.$$

Recognition of the general form. Writing $x = 2$, $p = 3$, and $S_3(s) = 1 + s + s^2$, the iteration (2) rearranges to

$$s_{n+1} = 1 - \frac{1}{2(1 + s_n + s_n^2)} = 1 - \frac{x - 1}{x \cdot S_p(s_n)} \Big|_{p=3, x=2}.$$

This is exactly the iterator $h_{p,x}$ introduced below, at $(p, x) = (3, 2)$. The fixed-point equation $s^* = 1 - (x - 1)/(x S_p(s^*))$ rearranges to $x \cdot (s^*)^p = 1$, i.e. $s^* = x^{-1/p}$, and the readout $v^* = x \cdot (s^*)^{p-1} = x^{1/p}$ recovers the target p -th root. The historical construction (1) is therefore the $(p, x) = (3, 2)$ instance of a one-parameter family defined in (3).

1.2 The general iterator $h_{p,x}$

On $\mathbb{C}^*$ with $p \in \mathbb{N}_{\geq 1}$, the Pandrosion iterator is

$$h_{p,x}(z) := 1 - \frac{x - 1}{x \cdot S_p(z)}, \quad S_p(z) := \sum_{k=0}^{p-1} z^k, \quad (3)$$

whose fixed points are exactly the p -th roots of $1/x$: $h_{p,x}(s) = s \iff s^p = 1/x$. The historical case (1) is the instance $(p, x) = (3, 2)$. Two constructions operate on top of $h_{p,x}$:

- an *algebraic acceleration*, the Aitken Δ^2 transform $\sigma_{p,x} := \text{Aitken}(h_{p,x})$, which upgrades linear to quadratic convergence without computing derivatives (Section 2);
- a *seeded multi-start* in $\mathbb{C}$, placing z_0 inside a controlled disk around a user-chosen cyclotomic anchor before iteration begins (Section 4).

Both are used below. We first analyze $h_{p,x}$ on the real line (Section 2), where the closed-form contraction ratio and preconditionings are cleanest, then extend to the complex case via multi-start.

1.3 Contributions, separated by kind

The paper makes explicit the difference between genuinely new formalized results, Pandrosion specializations of classical theorems, and honestly partial engagements with open problems.

Kind	What is in the corpus	Where
New real-line results for $h_{p,x}$	Closed-form linear contraction ratio $\lambda_{p,x} = h'_{p,x}(s^*)$, canonical derivative-free starting point $s_0^{\text{opt}} = h_{p,x}(1)$, scaling preconditioning $x^{1/p} = A^{1/p}(x/A)^{1/p}$ exploiting monotonicity of λ , Aitken Δ^2 upgrade to quadratic convergence with explicit constants.	Section 2.
New multi-start results in $\mathbb{C}$	Universal multi-start convergence, explicit closed-form iteration count, Borel-measurability and a.e. continuity of the multi-start solution map.	Sections 4, 5.
Elementary algebraic identities for ζ_P	Vanishing identity $\zeta_P(1) = 0$, $(p-1)$ higher-vanishing relations + normalization, closed-form product $\prod_k P'(\alpha_k) = (-1)^{p-1} p^p / x^{p-1}$.	Section 6. All of these reduce to the geometric-sum identity on roots of unity; we use a “finite Riemann-style” vocabulary only as a presentation device and claim no number-theoretic content.
Pandrosion specializations of classical theorems (not new mathematics, new Lean formalizations)	Banach 1922, FTA (Gauss 1799), Aitken 1926, Schwarz 1890, Kronecker 1857, Aberth–Ehrlich 1967/1973, Kung–Traub 1974 attainability at $c = 3$. Cardano, Vieta, and Galois modules exist in the corpus but are purely definitional for $z^p = x$ and not counted here.	Section 7.
Two short structural observations	Kung–Traub upper bound for $c \geq 3$ (compliance interface, not a proof); statement-level recording of McMullen 1987 without a formal proof.	Section 8. Lehmer’s conjecture and Smale’s 17th problem are deliberately outside scope (earlier drafts discussed them; they have been removed).

The paper does *not* claim to solve any of the four classical open problems in Section 8. Only restricted or structural results are proven there, each explicitly scoped.

1.4 Related work

Rational iterations for p -th roots are classical. Newton’s method applied to $f(u) = u^p - x$, Halley’s method, and the Schröder (1870) and König (1884) families supply iterations of arbitrary order with explicit asymptotic constants; Traub’s *Iterative Methods for the Solution of Equations* (1964) is the canonical systematic treatment of fixed-point iterations and their acceleration, and Ortega–Rheinboldt (1970) treats the multidimensional analogue. Aitken–Steffensen Δ^2 acceleration is standard textbook material [3]. The cyclotomic identities in Section 6 are elementary consequences of the geometric-sum vanishing $\sum_k \omega^{ks} = 0$ on roots of unity, which is used in every undergraduate treatment of cyclotomic polynomials.

What is specific to this paper is: (i) the rational iteration $h_{p,x}$ arising from the Pandrosion construction, which is *not* the Newton iteration of any polynomial and has order one rather than two; (ii) the closed-form ratio $\lambda_{p,x}$ and the preconditioning built on its monotonicity; (iii) the multi-start seed of §4.1 together with the measurability/a.e.-continuity analysis of the induced solution

map; (iv) the Lean 4 formalization itself, which is the main deliverable. We do not claim the iteration itself, or the ancillary identities, are novel mathematics; the contribution is to assemble them in a verified formal framework and to record the specific Pandrosion parameters.

1.5 On naming and scope

A remark on terminology. We define $\zeta_P(s) := \sum_{k=1}^p P'(\alpha_k)^{-s}$ as a finite series of p complex terms. All the identities we prove for it (including $\zeta_P(1) = 0$ and a closed-form for $\prod_k P'(\alpha_k)$) reduce to the geometric-sum identity on roots of unity and are elementary. We occasionally use a vocabulary reminiscent of the Riemann zeta function (“integer zeros of ζ_P ”) because the shape of the identities mirrors it, but we do *not* claim any number-theoretic content and the analogy should be read as typographic, not mathematical. In particular, nothing in this paper bears on the Riemann Hypothesis or on any transcendental question.

(The label “Niveau 5” used in Theorem 4.6 and in the Lean module naming is internal book-keeping for the corpus and has no external attestation; see the footnote in §4.)

1.6 Reading order

Section 2 develops the real-line theory of $h_{p,x}$: closed-form contraction ratio, optimal starting point, scaling preconditioning, and Aitken Δ^2 acceleration. Section 3 collects the cyclotomic identities for the complex setting. Sections 4–5 contain the multi-start results in $\mathbb{C}$ (universal convergence, iteration count, measurability and a.e. continuity). Section 6 collects the finite-sum identities for ζ_P . Section 7 gathers the Pandrosion specializations of classical theorems. Section 8 records the restricted / partial engagements with the four open problems. Sections 9 and 10 describe the Lean corpus structure and how to reproduce everything. A reader interested only in the new mathematical content can restrict attention to Sections 2–5 and 6.

2 The Pandrosion iteration on the real line

This section studies $h_{p,x}$ on $\mathbb{R}_{>0}$, where every quantity admits a closed form. The results of independent interest are three: a rational expression for the linear contraction ratio $\lambda_{p,x}$ depending only on the target (§2.1), an optimal derivative-free starting point (§2.2), and a scaling preconditioning that collapses the iteration count for large x (§2.3). Sections 2.4 and 2.5 then show that Aitken’s Δ^2 turns $h_{p,x}$ into the quadratically convergent $\sigma_{p,x}$ with a fully explicit local bound; this is the tool that drives the complex analysis of Section 4.

Throughout, $x > 1$ and $p \geq 2$; $\alpha := x^{1/p} > 1$ is the target p -th root and $s^* := 1/\alpha = x^{-1/p}$ is the unique fixed point of $h_{p,x}$ in $(0, 1)$. The readout is $v_n := x \cdot s_n^{p-1}$, converging to $v^* = \alpha$.

2.1 Closed-form contraction ratio

Theorem 2.1 (Closed-form contraction ratio). *For $x > 1$, $p \geq 2$, the iteration $s_{n+1} = h_{p,x}(s_n)$ is linearly convergent at s^* with ratio*

$$\lambda_{p,x} = h'_{p,x}(s^*) = \frac{(\alpha - 1) \sum_{k=1}^{p-1} k \alpha^{p-1-k}}{\sum_{k=0}^{p-1} \alpha^k}, \quad \alpha = x^{1/p}. \quad (4)$$

In particular, $\lambda_{p,x} \in (0, 1)$ is a rational function of the target α alone.

Direct computation: $h'_{p,x}(s) = (x - 1)S'_p(s)/(xS_p(s)^2)$, evaluated at $s^* = 1/\alpha$ and simplified using $x = \alpha^p$ and $xS_p(s^*) = \alpha/(\alpha - 1)$.

Why this matters. The number of iterations to reach precision δ is governed by

$$n_\delta \approx \left\lceil \frac{\log(\varepsilon_0/\delta)}{\log(1/\lambda_{p,x})} \right\rceil, \quad \varepsilon_0 := |v_0 - \alpha|. \quad (5)$$

Two levers for reducing n_δ are therefore available: reducing ε_0 (one-step preconditioning of the argument, §2.2) and reducing $\lambda_{p,x}$ (scaling of the target, §2.3). Thus $\lambda_{p,x}$ encodes the method-target interaction independently of the starting point.

Monotonicity and limits. A further computation (carried out in Lean in the corpus) shows that $\lambda_{p,x}$ is strictly increasing in both p and x on $[1, \infty)$. The limits are

$$\lim_{x \rightarrow \infty} \lambda_{p,x} = 1^- \quad (\text{for fixed } p), \quad \lim_{p \rightarrow \infty} \lambda_{p,x} = 1 - \frac{\ln x}{x-1} \in (0, 1) \quad (\text{for fixed } x > 1).$$

In particular, for fixed x the ratio does *not* tend to 1 as p grows – it saturates at $1 - \ln(x)/(x-1)$. The scaling preconditioning of §2.3 is built directly on the monotonicity in x .

2.2 Optimal starting point: one-step preconditioning

Proposition 2.2 (Optimal α -free starting point). *Among starting points $s_0 \in (0, 1)$ computable without knowledge of α , the choice*

$$s_0^{\text{opt}} := h_{p,x}(1) = 1 - \frac{x-1}{xp} \quad (6)$$

minimises the initial residual $|s_0 - s^|$ to leading order.*

Since $h_{p,x}$ is the method’s own one-step update and s^* is its fixed point, applying $h_{p,x}$ to the naive boundary value $s = 1$ gives the best single-step estimate of s^* that does not require α . Taylor expansion at $s = 1$ gives $s^* = 1 - (\ln x)/p + O(1/p^2)$, so $|s_0^{\text{opt}} - s^*| = |(x-1)/(xp) - (\ln x)/p| + O(1/p^2)$, substantially smaller than $|1 - s^*|$ or $|1/2 - s^*|$ in the relevant regime.

(p, x)	s^*	$s_0^{\text{opt}} = h_{p,x}(1)$	$ s_0^{\text{opt}} - s^* $	Iter. from s_0^{opt}	Iter. from $s_0 = 1/2$
(2, 2)	0.7071	0.7500	0.043	12	13
(3, 2)	0.7937	0.8333	0.040	14	16
(4, 2)	0.8409	0.8750	0.034	15	17
(3, 8)	0.5000	0.7083	0.208	42	*

Table 1: Iteration counts to reach $|v_n - \alpha| < 10^{-10}$. s_0^{opt} consistently reduces the count. (*For (3, 8), $s_0 = 1/2$ happens to coincide with s^* .)

2.3 Scaling preconditioning: exploiting monotonicity

For large x , the contraction ratio $\lambda_{p,x}$ is close to 1 and (5) explodes. The monotonicity $x \mapsto \lambda_{p,x}$ is however the basis of a simple remedy.

Proposition 2.3 (Scaling reduction). *Let $A \in \mathbb{N}_{\geq 1}$ satisfy $A^{1/p} \in \mathbb{N}$ and $x' := x/A \in [1, (1 + 1/A^{1/p})^p]$. Then*

$$x^{1/p} = A^{1/p} \cdot (x')^{1/p}, \quad (7)$$

so that computing $x^{1/p}$ reduces to computing $(x')^{1/p}$ and multiplying by the integer $A^{1/p}$. Taking $A = \lfloor x^{1/p} \rfloor^p$ makes x' close to 1 and the ratio $\lambda_{p,x'}$ close to 0.

The saving compounds over every iteration: with $\varepsilon'_0 \ll \varepsilon_0$ and $\log(1/\lambda_{p,x'}) \gg \log(1/\lambda_{p,x})$, (5) drops by a factor of several tens to several hundred (Table 2). This is the real-variable analogue of the complex multi-start seed of §4, which places z_0 close to an anchor by contracting towards it rather than by rescaling the argument.

x	$\lambda_{3,x}$	$A = \lfloor x^{1/3} \rfloor^3$	$x' = x/A$	$\lambda_{3,x'}$	Iterations (direct $\rightarrow$ scaled)
10	0.615	8	1.250	0.073	49 $\rightarrow$ 8
100	0.890	64	1.563	0.145	213 $\rightarrow$ 11
1000	0.973	729	1.372	0.103	$> 500 \rightarrow 9$
10000	0.994	9261	1.080	0.025	$> 500 \rightarrow 5$

Table 2: Scaling effect at $p = 3$, precision 10^{-10} , with $s_0 = s_0^{\text{opt}}$. The per-step ratio drops by one to two orders of magnitude and the iteration count collapses from hundreds to single digits.

Remark 2.4 (Double preconditioning). The two accelerations combine multiplicatively: (6) reduces ε_0 by a fixed factor, (7) reduces $1/\log(1/\lambda_{p,x})$ by an x -dependent factor, and (5) benefits from both simultaneously.

2.4 Aitken Δ^2 : from linear to quadratic convergence

The linear convergence of $h_{p,x}$ with explicit ratio $\lambda_{p,x}$ is exactly the input required by the Aitken Δ^2 transform.

Proposition 2.5 (Aitken Δ^2 on geometric sequences). *For any geometric sequence $s_n = s^* + C \cdot r^n$ with $r \neq 0, 1$ and $C \neq 0$,*

$$s_n - \frac{(s_{n+1} - s_n)^2}{s_{n+2} - 2s_{n+1} + s_n} = s^* \quad \text{for every } n.$$

Applied at each step to the Pandrosion iterates, this yields the derivative-free accelerator

$$\sigma_{p,x}(z) := z - \frac{(h_{p,x}(z) - z)^2}{h_{p,x}(h_{p,x}(z)) - 2h_{p,x}(z) + z} \quad (8)$$

(one Δ^2 step per iterate; compare [12, 13]).

Fig 7. Aitken-Steffensen acceleration (1926/1933, §113): linear h to quadratic σ




Figure 1: Left: a linearly convergent geometric sequence and its Δ^2 -accelerated version, which collapses to s^* at machine precision. Right: the h iteration (linear) vs $\sigma = \text{Aitken}(h)$ (quadratic) for $(x, p) = (2, 3)$, $z_0 = 1$.

2.5 A quadratic local bound at the super-attractive fixed point

Proposition 2.6 (Quadratic local bound). *The map $\sigma_{p,x}$ has a super-attractive fixed point at every p -th root of $1/x$: $\sigma_{p,x}(s^*) = s^*$ and $\sigma'_{p,x}(s^*) = 0$. On a neighbourhood $B(s^*, R_\sigma)$,*

$$|\sigma_{p,x}(s) - s^*| \leq K_\sigma \cdot |s - s^*|^2,$$

where the constants admit the explicit forms

$$K_\sigma = \left| \frac{h''_{p,x}(s^*)}{2(1 - \lambda_{p,x})} \right|, \quad R_\sigma = \min(1/(2K_\sigma), \text{dist}(s^*, \text{sing}(\sigma_{p,x}))),$$

so that $K_\sigma \cdot R_\sigma \leq 1/2$. The same bound holds in $\mathbb{C}$ on $B(\gamma_k, R_\sigma)$ for every cyclotomic anchor γ_k . The explicit formulas are formalized in `Pandrosion/Core/QuadraticBound.lean` and `SteffensenQuadraticBound.lean`.

This is the Schwarz-Pick-type estimate that the multi-start scheme in $\mathbb{C}$ exploits: once z is inside $B(\gamma_k, R/2)$, successive iterates contract quadratically and the log log iteration count of (4.5) follows.

3 Cyclotomic anchors and Galois structure

3.1 Cyclotomic anchor family

Definition 3.1 (Cyclotomic anchors). For $\alpha \in \mathbb{C}^*$ and $p \geq 1$, let

$$\gamma_k := \alpha \cdot e^{2\pi i k/p}, \quad k = 0, 1, \dots, p-1.$$

When $\alpha^p = 1/x$, the γ_k are the p distinct solutions of $z^p = 1/x$.

Fig 2. Cyclotomic anchor families $\gamma_k = \alpha \cdot e^{2\pi i k/p}$ for $z^p = 1/x$, $x = 2$ (§118-§119)




Figure 2: Cyclotomic anchor families for $p \in \{3, 4, 5, 7, 11, 13\}$ at $x = 2$. The anchors lie on the circle of radius $|\alpha| = 2^{-1/p}$, and the cyclic action $\omega \cdot \gamma_k = \gamma_{k+1 \bmod p}$ generates the splitting field.

3.2 Cyclic permutation of the anchors

By direct expansion of Definition 3.1, $\omega \cdot \gamma_k = \gamma_{(k+1) \bmod p}$ for every k (with $\omega^p = 1$). Likewise $\sum_k \gamma_k = 0$ and $\prod_k \gamma_k = (-1)^{p-1} \cdot x$. These are definitional consequences for $z^p = x$, not specializations of the Galois or Vieta theorems; we use them freely in the sequel.

4 Universal multi-start convergence

We now extend the iteration to $\mathbb{C}$. The linear-to-quadratic acceleration of Section 2.4 applies verbatim to the complex iterator (3), giving the Steffensen map $\sigma_{p,x}$ of (8) on all of $\mathbb{C}$. What is genuinely new over the real-line case is the need to control *which* cyclotomic anchor the iteration converges to: in $\mathbb{C}$, $\sigma_{p,x}$ has p attractive fixed points, the Julia set separating their basins is generically non-trivial, and a classical impossibility result of McMullen [18] rules out purely iterative universal root finders for $p \geq 3$ in a strong sense. The seed of §4.1 is designed precisely to exit the hypotheses of [18].

4.1 The multi-start seed

Definition 4.1 (Multi-start seed). For a cyclotomic anchor $\alpha \in \mathbb{C}^*$ with $\alpha^p = 1/x$ and $z \in \mathbb{C}$,

$$\text{seed}_{\alpha,\varepsilon}(z) := \alpha + \varepsilon \cdot (z - \alpha), \quad \varepsilon := \frac{R_\sigma(x, p, \alpha)}{2 \|z - \alpha\|} \quad (\varepsilon := 1 \text{ if } z = \alpha), \quad (9)$$

where $R_\sigma(x, p, \alpha)$ is the explicit quadratic-contraction radius from Proposition 2.6. The seed always lies in $\overline{B}(\alpha, R_\sigma/2)$.

The seed plays the role in $\mathbb{C}$ that the preconditionings of §§2.2–2.3 play on $\mathbb{R}$: it places the argument inside the quadratic-contraction disk of the target *before* the iteration begins. It is α -dependent by design, and this dependency is what takes the scheme out of the hypotheses of [18].

Fig 1. Newton on $z^5 = 1$ has fractal Julia set (McMullen 1987); Pandrosion multi-start circumvents via α -dependent seed — §95, §120




Figure 3: Structure of the multi-start seed. Each point $z \in \mathbb{C}$ is mapped to $\text{seed}_{\alpha,\varepsilon_{\text{seed}}}(z)$, which lies in $\overline{B}(\alpha, R_\sigma/2)$ regardless of z ; the iteration is then run from the seed. Colour indicates the number of iterations needed to reach precision 10^{-6} after the seed is applied. The figure illustrates how seed (9) places the scheme outside the hypotheses of [18] by making the iterate entry point α -dependent; we explicitly do *not* compare with Newton’s basin structure here, since the two schemes do not solve the same problem (Newton is parameter-free, multi-start is α -targeted).

4.2 Universal multi-start theorem

Theorem 4.2 (Universal multi-start convergence). *Let $x \in \mathbb{C}^*$, $p \geq 1$, $\alpha \in \mathbb{C}^*$ with $\alpha^p = 1/x$ and $S_p(\alpha) \neq 0$. Then for every $z \in \mathbb{C}$ and every $\varepsilon > 0$, there exist $\varepsilon_{\text{seed}} > 0$ and $N \in \mathbb{N}$ such that*

$$\forall n \geq N : \quad \|\sigma_{p,x}^n(\text{seed}_{\alpha, \varepsilon_{\text{seed}}}(z)) - \alpha\| \leq \varepsilon.$$

The construction is explicit: $\varepsilon_{\text{seed}} = R_\sigma(x, p, \alpha)/(2\|z - \alpha\|)$ when $z \neq \alpha$, and N is given by Theorem 4.5 below.

The content is: the seed is in $\overline{B}(\alpha, R_\sigma/2)$, the iteration stays in $B(\alpha, R_\sigma)$, and a quadratic-contraction argument gives the iteration count.

Remark 4.3 (The hypothesis $S_p(\alpha) \neq 0$). The non-vanishing condition $S_p(\alpha) \neq 0$ rules out the degenerate case where the denominator of $h_{p,x}$ vanishes at the anchor. Since $S_p(\alpha) = (\alpha^p - 1)/(\alpha - 1) = (1/x - 1)/(\alpha - 1)$ for $\alpha^p = 1/x$ and $\alpha \neq 1$, this is an event of codimension at least one in the (x, p) -parameter space and in particular holds generically; it fails only for specific arithmetic combinations (for instance when α is a non-trivial p -th root of unity, which requires $x = 1$). The Lean corpus carries this hypothesis explicitly in the statement.

4.3 Explicit iteration count

Definition 4.4 (Effective iteration count). For $K, R, \varepsilon > 0$ with $KR < 1$ and $K\varepsilon < 1$,

$$N_{\text{eff}}(K, R, \varepsilon) := \log_2 \lceil \log(K\varepsilon)/\log(KR) \rceil_+ + 1,$$

with $\lceil \cdot \rceil_+$ the natural-number ceiling clamped at 0. The hypotheses $KR < 1$ and $K\varepsilon < 1$ ensure both logarithms are negative and their ratio is well-defined and non-negative.

Theorem 4.5 (Explicit iteration count). *For any quadratically contracting sequence $e_{n+1} \leq K \cdot e_n^2$ with $e_0 \leq R$, $K \cdot R < 1$, and $\varepsilon < R$, every iterate $k \geq N_{\text{eff}}(K, R, \varepsilon)$ satisfies $e_k \leq \varepsilon$. Asymptotically $N_{\text{eff}} \sim \log_2 \log(1/\varepsilon)$.*

Fig 3. Effective Kung-Traub loglog complexity (§98, §101) beats Smale 17 for $z^p = x$ (§115)




Figure 4: Left: error $\|\sigma^n(\text{seed}) - \alpha\|$ vs iteration for $p \in \{3, 4, 5, 7\}$, $x = 2$, $z_0 = 1 + 0.5i$. Right: iteration count $N(\varepsilon)$ vs target precision for the same setup. The dashed line is $\log(1/\varepsilon)$; it is shown only for visual reference, not as an algorithmic comparison.

4.4 An a.e.-convergence statement for the plain iteration

It is natural to ask whether the plain Steffensen iteration $\sigma_{p,x}$ converges for almost every starting point to the principal anchor α_0 :

$$\forall^{\text{a.e.}} z \in \mathbb{C} : \quad \sigma_{p,x}^n(z) \xrightarrow{n \rightarrow \infty} \alpha_0.$$

Numerical experiments at $(x, p) = (2, 3)$ (Python sweep over $[-3, 3]^2$ at 200×200 resolution) show that roughly 0.015% of initial points converge instead to a non-principal anchor, so the plain-iteration version of the statement is *false*.¹

Theorem 4.6 (Multi-start version). *Under the standard hypotheses on (x, p, α) , the multi-start reformulation*

$$\forall z \in \mathbb{C}, \exists \varepsilon_{\text{seed}} > 0 : \quad \sigma_{p,x}^n(\alpha + \varepsilon_{\text{seed}} \cdot (z - \alpha)) \xrightarrow{n \rightarrow \infty} \alpha$$

holds unconditionally.

This is Theorem 4.2 restated. The content is not that the reformulation is “solved”: it is that by conditioning on the choice of target anchor α and adjusting the seed accordingly, the convergence problem decouples from the dynamics of $\sigma_{p,x}$ on all of $\mathbb{C}$. What the reformulation gives up, compared to the plain-iteration statement, is universality: the seed depends on α and can therefore not be interpreted as a “method that converges regardless of target”.

4.5 Arbitrary target selector

Theorem 4.7 (Target selection). *For every function $S : \mathbb{C} \rightarrow \text{Fin } p$, every $z \in \mathbb{C}$ and every $\varepsilon > 0$, the multi-start scheme with target $\gamma_{S(z)}$ converges to $\gamma_{S(z)}$ with an iteration count of the form of Theorem 4.5.*

Three selectors are implemented: constant (`principal_selector`), Voronoï-nearest (`nearest_anchor_selector`) and angular sector.

5 Measurability and a.e. continuity of the solution map

The `nearest_anchor_selector` of Section 4 uses `Classical.choose` on `Finset.exists_min_image`; this is not a priori measurable.

Definition 5.1 (Borel selector).

$$S_{\text{Borel}}(\alpha, p, z) := \min'(\text{argmin}_{k \in \text{Fin } p} \|\gamma_k - z\|).$$

Theorem 5.2 (Borel measurability). *The map S_{Borel} is Borel-measurable. Consequently the solution map $z \mapsto \gamma_{S_{\text{Borel}}(z)}$ is Borel-measurable $\mathbb{C} \rightarrow \mathbb{C}$.*

The proof shows that each level set is a finite Boolean combination of closed half-planes $\{z : \|z - \gamma_s\| \leq \|z - \gamma_t\|\}$.

Definition 5.3 (Voronoi boundary and its complement). $\text{VorBdy}(\alpha, p) := \text{VoronoiBoundary}(\gamma_0, \dots, \gamma_{p-1})$, $\text{VorInt}(\alpha, p) := \mathbb{C} \setminus \text{VorBdy}(\alpha, p)$.

These are Voronoï-geometric objects, not the classical Fatou–Julia sets of $\sigma_{p,x}$. Earlier drafts of this corpus used the names “Pandrosion Julia / Fatou set” for them; this was a poor choice of vocabulary, because it suggests a relationship to Fatou–Julia dynamics of $\sigma_{p,x}$ that we do not establish. The theorems below are statements about Voronoï geometry of the cyclotomic anchor set, nothing more.

¹In the Lean corpus this a.e.-statement is tracked internally under the label “Niveau 5” (modules `Niveau5Master`, `Niveau5StrictMultiStart`, etc.). The label is ours; it does not correspond to any externally attested conjecture, and there is no “Niveau 4” hidden elsewhere – it is simply internal bookkeeping.

Theorem 5.4 (Voronoi boundary is Lebesgue-null). $\text{vol}(\text{VorBdy}(\alpha, p)) = 0$.

Theorem 5.5 (A.e. continuity). *The Borel solution map is continuous at every $z \in \text{VorInt}(\alpha, p)$. Combined with Theorem 5.4, it is continuous at Lebesgue-a.e. z .*

Fig 4. Pandrosion Julia-null theorem (§103): ∂V is Lebesgue-null, multi-start deterministic a.e.

$p = 4$: Voronoi basins V_k + Julia frontier (red, Lebesgue-null) = 7: Voronoi basins V_k + Julia frontier (red, Lebesgue-null)




Figure 5: Voronoi decomposition of $\mathbb{C}$ for cyclotomic anchors at $p = 4$ and $p = 7$, $x = 2$. Each colored region V_k is the basin of attraction of γ_k under the selector. The red frontier is the Voronoi boundary of the cyclotomic anchor set; it is Lebesgue-null.

Fig 8. Voronoi basins (left, structural) + multi-start speed (right, ~ 3 -5 iter) — §102 measurable, §103 Julia-null, §105 a.e. continuous




Figure 6: Left: Voronoi basins as the structural reference, computed from the nearest-anchor criterion only (no dynamics). Right: number of multi-start iterations needed to reach precision 10^{-8} ; it is bounded and essentially flat over the window, in line with Theorem 4.5.

Together, Theorems 4.2, 5.2, 5.4 and 5.5 give the analytic character of the multi-start solution map in our framework: measurable everywhere, continuous outside the Voronoi boundary of the cyclotomic anchor set, with that boundary Lebesgue-null.

6 Elementary identities on the derivative values at cyclotomic roots

For $P(z) = z^p - 1/x$ with $\alpha^p = 1/x$ and cyclotomic anchors $\alpha_k = \gamma_k$ (Definition 3.1), one has $P'(\alpha_k) = p\alpha_k^{p-1} = p/(x\alpha_k)$. Define

$$\zeta_P(s) := \sum_{k=1}^p P'(\alpha_k)^{-s} = (x/p)^s \sum_{k=1}^p \alpha_k^s. \quad (10)$$

Every identity used elsewhere in the corpus about the values $P'(\alpha_k)$ reduces to the geometric-sum vanishing $\sum_{k=0}^{p-1} \omega^{ks} = 0$ for $\omega = e^{2\pi i/p}$ and $p \nmid s$. We record the three forms that Lean modules cite directly.

Proposition 6.1 (Three elementary identities). *Let $p \geq 2$ and $x \neq 0$.*

(i) *For every $s \in \mathbb{Z}$ with $1 \leq s \leq p-1$,*

$$\zeta_P(s) = \sum_{k=1}^p P'(\alpha_k)^{-s} = 0.$$

In particular $\zeta_P(1) = 0$.

(ii) *For every $m \in \mathbb{Z}$ with $0 \leq m \leq p-2$, $\sum_{k=1}^p \alpha_k^m / P'(\alpha_k) = 0$; for $m = p-1$, $\sum_{k=1}^p \alpha_k^{p-1} / P'(\alpha_k) = 1$.*

(iii) $\prod_{k=1}^p P'(\alpha_k) = (-1)^{p-1} p^p / x^{p-1}$.

Items (i) and (ii) are the same statement under $s \leftrightarrow m+1$ via $1/P'(\alpha_k) = x\alpha_k/p$; item (iii) is the product counterpart. All three follow from $\sum_k \omega^{ks} = 0$ for $p \nmid s$.

We keep this section deliberately short: the identities are logged because the Lean modules `PandrosionZeta.lean`, `PandrosionHigherVanishing.lean`, and `PandrosionSpectralDeterminant.lean` cite them by name, not because there is substantial mathematics here. The shape of (i)–(ii) is reminiscent of vanishing identities for classical arithmetic zeta functions, but the analogy is typographic only and no number-theoretic claim is made.

7 Pandrosion specializations of classical theorems

The corpus contains Lean formalizations of a number of classical theorems, *specialized* to the Pandrosion setting. The purpose is to have everything in one formal framework; the mathematical statements are classical. The following table summarizes what is formalized where.

Module	Classical theorem (specialized)	Role in the Pandrosion framework
§112	Banach fixed point (1922); FTA (Gauss 1799)	No-cycle on multi-start orbits; $z^p = x$ has p explicit roots.
§113	Aitken-Steffensen (1926/1933); Schwarz (1890)	Acceleration identity; quadratic local bound.
§114	Kung-Traub 1974 attainability at $c = 3$	Efficiency $E(4, 3) = 2^{2/3}$ is reached.
§115	(See Section 8.)	Restricted to $z^p = x$.
§117	Aberth-Ehrlich (1967/1973)	Multi-start realizes their scheme for $z^p = x$.
§118	Kronecker (1857)	Applied to cyclotomic anchors with $ \alpha = 1$.
§120	McMullen (1987) (See Section 8.)	Multi-start avoids the obstruction by α -dependent seed.

We stress: none of these are new results. Their place in this paper is to make the full Pandrosion framework self-contained in Lean. The corresponding Lean modules are short specializations, not independent developments of the classical theorems.

Definitional-level items (not listed above). Three further modules, `PandrosionCardano` (§121), `PandrosionVieta` (§122), and `PandrosionGalois` (§119), record the identities $\gamma_k = \alpha\omega^k$ for $p = 3$, $e_1 = 0$ and $e_p = (-1)^{p-1}x$, and $\omega \cdot \gamma_k = \gamma_{(k+1) \bmod p}$ respectively. For the monomial family $z^p = x$ all three are direct expansions of Definition 3.1 and do not rely on the classical Cardano, Vieta, or Galois theorems as such. We therefore do not count them as specializations of those theorems; they remain in the Lean corpus as named lemmas for downstream convenience only.

A representative example.

Theorem 7.1 (Multi-start orbit has no non-trivial cycle). *For every (x, p, α) in the hypotheses and every seed $\text{seed} \in B(\alpha, R_\alpha/2)$, the orbit $(\sigma_{p,x}^n(\text{seed}))_n$ has no non-trivial cycle: if $\sigma_{p,x}^n(\text{seed}) = \sigma_{p,x}^m(\text{seed})$ with $n < m$, then $\sigma_{p,x}^n(\text{seed}) = \alpha$.*

This is Banach’s fixed-point theorem applied to $\sigma_{p,x}$ seen as a $1/4$ -contraction on $B(\alpha, R_\alpha/2)$, itself a consequence of $K_\alpha \cdot R_\alpha \leq 1/2$.

8 Short structural observations

We record two places (Kung–Traub, McMullen) where the Pandrosion setting intersects a classical or open problem, and one explicit out-of-scope note for two further classical themes (Lehmer, Smale 17). None of the material below constitutes progress on the underlying problems; each subsection states what the corpus contributes and, just as importantly, what it does not.

8.1 Kung–Traub upper bound for $c \geq 3$ (open since 1974)

The conjecture $q \leq 2^{c-1}$ is proven for $c = 2$. Attainability $q = 4$ at $c = 3$ is witnessed classically (Jarratt 1969) and in our formal setting.

What the corpus provides. An interface: every `DerivativeFreeMethod` in the corpus carries a *compliance field* asserting $q \leq 2^{c-1}$ for its own instance. This is not a proof of the Kung–Traub conjecture, only a structural assertion that any instance we construct satisfies the bound. The conjecture itself is recorded as a named `Prop`; the general case for $c \geq 3$ remains open.

8.2 McMullen 1987 (recorded but not formalized)

McMullen’s barrier is a classical theorem. A full formal proof in Lean would require Fatou–Julia and post-critically-finite complex-dynamic machinery that is not present in Mathlib v4.7.0.

What the corpus provides. The statement is recorded as a named `Prop`; it is *not* formally proven. Independently, Theorem 4.2 shows that our multi-start scheme has an α -dependent seed and therefore does not fall under McMullen’s hypotheses on purely iterative universal root finders. This observation places our scheme outside the hypothesis of [18]; it does not reproduce or challenge the theorem.

8.3 Two classical problems we deliberately do *not* engage with

We close this section by noting two classical problems that earlier drafts of the corpus were tempted to discuss: Lehmer’s 1933 Mahler-measure conjecture, and Smale’s 17th problem on deterministic polynomial root-finding. In both cases, the only non-vacuous things the Pandrosion setting offers are a trivial observation (for Lehmer: that $\beta = x \cdot \alpha$ has Mahler measure x^{p-1} , which is ≥ 2 for $x \geq 2$, far from the conjecture’s extremal regime near 1) and a scope-restricted algorithm (for Smale: the $\mathcal{O}(\log \log(1/\varepsilon))$ deterministic complexity for the one-parameter family $z^p = x$, which is not the coefficient-uniform problem Smale asks about). Since in neither case is there progress worth reporting, we refrain from a dedicated subsection and simply record that these two problems lie outside the scope of the corpus.

The Lean module `Pandrosion/Core/PandrosionOpenProblems.lean` nevertheless carries statement-level Props (`lehmer_conjecture_statement`, `smale17_deterministic_statement`, `pandrosion_mahler_observation`) for historical reasons. These are not referenced anywhere else in the corpus, produce no theorems, and should be read as frozen records of the discussion above rather than as active content. A future cleanup pass will either delete them or move them into a separate `archive` namespace.

8.4 Summary

No claim is made that any classical problem is solved, or even substantively advanced. The material in this section consists of a compliance interface (Kung–Traub) and a recording of a classical statement (McMullen). Readers primarily interested in the mathematics should focus on Sections 2–6.

9 Structure of the Lean corpus

The 121-module corpus is organised in dependency layers:

- **Foundations** (§1–§30): geometric sums, fixed-point characterization, abstract Aitken–Steffensen framework, Voronoï basins, cyclotomic anchor definitions, Kung–Traub interface.
- **Core spine** (§31–§94): explicit super-attractive rate, global loglog convergence, real-line $p = 2$ unconditional discharge.
- **Multi-start** (§95–§101): universal convergence, target selection, the multi-start a.e.-convergence reformulation of Theorem 4.6 (the plain-iteration version is numerically *false*; what the module establishes is the multi-start theorem, not a proof of the original a.e. statement), explicit iteration count. Module names prefixed `Niveau5*` are historical bookkeeping and will be renamed in a future cleanup pass.
- **Measure/topology** (§102–§105): Borel measurability, Voronoï frontier Lebesgue-null, a.e. continuity.
- **Finite-zeta identities** (§106–§111): the identities of Section 6.
- **Classical specializations** (§112–§122): as listed in Section 7.
- **Short structural observations** (§123, `PandrosionOpenProblems`): named Props for Kung–Traub ($c \geq 3$) and McMullen 1987, matching Section 8. The module also contains statement-level Props for Lehmer and Smale 17 that are retained for historical reasons; these are not used by any other module and should be read as out-of-scope notes, consistent with the paper text.

Fig 10. Pandrosion-Steffensen Lean 4 corpus structure — from foundations to publication




Figure 7: Dependency structure of the Lean corpus. The dependency closure of every load-bearing theorem reduces to the Lean whitelist `{propxt, Classical.choice, Quot.sound}`.

10 Reproducibility

The Lean source lives at github.com/ivan-fr/poussiere_de_x. It targets Lean 4 with Mathlib v4.7.0 and is built with

Listing 1: Incremental build command

```
1 bash scripts/lean-incremental.sh
```

The 121-module corpus compiles in about 60–90 seconds on a modern laptop. Per-theorem axiom auditing is available via `#print axioms`: for every load-bearing theorem referenced in this paper (Theorems 2.1, 4.2, 4.5, 5.2, 5.4, 5.5, and the ζ_P identities of Section 6), the output is the standard Lean whitelist `{propxt, Classical.choice, Quot.sound}`; this can be reproduced by running `#print axioms <theorem>` at the end of the relevant module.

Every figure in this paper is reproducible from `latex/figures_pandrosion.py` (dependencies: `numpy`, `matplotlib`, `scipy`).

11 Conclusion

What this paper reports, concretely, is a Lean 4 development built around the Pandrosion iterator $h_{p,x}$ and its multi-start Steffensen extension for $z^p = x$, together with the following ingredients:

1. real-line analysis of $h_{p,x}$: closed-form contraction ratio $\lambda_{p,x} = h'_{p,x}(s^*)$ (Theorem 2.1), optimal derivative-free starting point $s_0^{\text{opt}} = h_{p,x}(1)$ (Proposition 2.2), scaling preconditioning exploiting monotonicity of λ (Proposition 2.3), and Aitken Δ^2 upgrade to quadratic convergence with explicit local bound (Propositions 2.5–2.6);

2. a universal convergence theorem for the complex multi-start scheme with an explicit closed-form iteration count (Theorems 4.2, 4.5);
3. a Borel-measurability and a.e.-continuity analysis of the induced solution map (Theorems 5.2–5.5);
4. a set of elementary finite-sum identities for the series $\zeta_P(s) := \sum_k P'(\alpha_k)^{-s}$ (Section 6), all reducing to roots-of-unity arithmetic;
5. Lean specializations of classical theorems to the Pandrosion setting (Section 7);
6. two short structural observations (Kung–Traub compliance interface, McMullen 1987 statement-level recording) in Section 8; two further classical themes that earlier drafts discussed (Lehmer, Smale 17) have been removed as being outside scope.

The paper does not claim progress on any classical open problem. The main value of the corpus is as a self-contained Lean development of a multi-start root-finding scheme and its ancillary algebraic identities.

Future directions

Natural extensions are: non-cyclotomic anchor families (general polynomials), which would require dropping the closed-form cyclotomic identities that drive Sections 3 and 6; and a formalization of Fatou–Julia in Mathlib, which would make a proper in-Lean proof of McMullen’s 1987 theorem accessible. We make no conjecture regarding the other three problems discussed in Section 8.

References

- [1] de Moura, L., Ullrich, S., *The Lean 4 Theorem Prover and Programming Language*, CADE-28, Springer LNCS (2021).
- [2] The Mathlib Community, *Mathlib: A unified library of mathematics formalized*, github.com/leanprover-community/mathlib4.
- [3] W. H. Press, S. A. Teukolsky, W. T. Vetterling, B. P. Flannery, *Numerical Recipes: The Art of Scientific Computing*, 3rd ed., Cambridge University Press, 2007.
- [4] Cardano, G., *Ars Magna* (1545), Cap. XI.
- [5] Viète, F., *Opera Mathematica* (1591).
- [6] Gauss, C.F., Thesis on the fundamental theorem of algebra (1799).
- [7] Galois, É., *Mémoire sur les conditions de résolubilité des équations par radicaux* (1832, posth. Liouville 1846).
- [8] Kronecker, L., Berlin Akademie (1857).
- [9] Riemann, B., *Über die Anzahl der Primzahlen unter einer gegebenen Grösse* (1859).
- [10] Schwarz, H.A., J. Reine Angew. Math. **70** (1869); Pick, G. (1916).
- [11] Banach, S., Fund. Math. **3** (1922).
- [12] Aitken, A.C., Proc. Royal Soc. Edinburgh **46** (1926), 289–305.
- [13] Steffensen, J.F., Skandinavisk Aktuarietidskrift **16** (1933), 64–72.

- [14] Lehmer, D.H., *Annals of Math.* **34** (1933), 461–479.
- [15] Ehrlich, L.W., *Comm. ACM* **10** (1967), 107–108.
- [16] Aberth, O., *Math. Comp.* **27** (1973), 339–344.
- [17] Kung, H.T., Traub, J.F., *J. ACM* **21** (1974), 643–651.
- [18] McMullen, C., *Annals of Math.* **125** (1987), 467–493.
- [19] Smale, S., *Mathematical Problems for the Next Century*, *Math. Intelligencer* **20** (1998), 7–15.
- [20] Bürgisser, P., Cucker, F., *Annals of Math.* **174** (2011), 1785–1836.
- [21] Besevic, I., *The Pandrosion zeta function and the discriminant as spectral determinant*, Preprint (2026).