

Pandrosion Operator and Smale’s 17th Problem

Universal polynomial evaluation, non-local geometry, and the Euler product connection

Ivan Besevic

April 2026

Abstract

We extend the generalized Pandrosion iteration — a geometric fixed-point method originally designed for computing p -th roots via the geometric progression polynomial $S_p(z)$ — from the real line to the complex plane and, crucially, to *polynomial systems* $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$, the proper setting of Smale’s 17th problem (1998). The univariate Pandrosion sequence is the special case $n = 1$ of a *universal evaluation generator*: by anchoring the matrix-valued difference quotient $\mathbf{Q}_F(\mathbf{z}_0, \mathbf{z})$ defined by the telescoping identity $F(\mathbf{z}) - F(\mathbf{z}_0) = \mathbf{Q}_F(\mathbf{z}_0, \mathbf{z})(\mathbf{z} - \mathbf{z}_0)$, we define a non-local rational iteration $\mathbf{z}_{n+1} = \mathbf{z}_0 - \mathbf{Q}_F(\mathbf{z}_0, \mathbf{z}_n)^{-1}F(\mathbf{z}_0)$ that recovers the exact p -th root formula in $n = 1$, the multivariate Newton iteration as the limiting case where the anchor tracks the iterate, and applies uniformly to any polynomial map F . For univariate $P(z) = xz^p - 1$ we characterize the complex contraction ratio.

Our main rigorous contributions, in both univariate ($n = 1$) and multivariate ($n \geq 2$) settings, are threefold: (i) a universal local convergence theorem establishing an explicit contraction basin of radius $\eta_F(\zeta)$ around every regular zero ζ of F ; (ii) exact polynomial factorisations avoiding the singular Σ barriers of Newton’s method completely; (iii) a macroscopic kinematic identity yielding a bounded telescopic product. We identify a *far-anchor obstruction* for fixed-anchor Cauchy-circle starts and resolve it via the **scaling principle**: adaptive anchor reanchoring every $K = O(1)$ steps keeps the effective argument $\|\mathbf{Q}_F^{-1}F\| \approx 1$. The resulting adaptive Pandrosion- T_3 scheme is verified numerically on four adversarial univariate families up to degree $d = 500$, and on multivariate Smale-17-style polynomial systems with Bézout degree d^n up to $(n, d) \in \{(2, 3), (3, 3), (4, 2)\}$ (cf. `benchmark_smale17_v4.py`). Conditionally on Conjecture 5.19, this yields a deterministic, derivative-free $O(d^2 \log d)$ -operation BSS algorithm for univariate root-finding; the multivariate Smale-17 average-case $O(\text{poly}(d, n))$ bound is established unconditionally only by Beltrán–Pardo [4] and Lairez [5], and our scheme provides a derivative-free *algorithmic alternative* with comparable empirical complexity, not a competing complexity-theoretic resolution. We close with a formal Pandrosion-language rewriting of the Euler product for $\zeta(s)$.

Distinction from Smale’s Mean Value Conjecture. The present work and the companion Parts II–VI [16] address *two distinct conjectures* of Smale, often confused in the literature:

- **Smale’s Mean Value Conjecture (MVC, 1981, [17]):** a univariate inequality $\min_{P'(\zeta)=0} |P(\zeta)/(\zeta P'(0)| \geq (d-1)/d$ for monic polynomials P of degree d with $P(0) = 0$. Treated in Parts II–VI.
- **Smale’s 17th Problem (1998, [18]):** a complexity-theoretic problem about the average-case cost of finding an approximate zero of a *system* $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ of n polynomials in n unknowns, with Bézout degree d^n . Resolved positively by Beltrán–Pardo (2009, [4]) probabilistically and Lairez (2017, [5]) deterministically. Treated in this Part I and in Parts VII–VIII.

The present paper develops the algorithmic framework for the latter; we do *not* claim to improve upon the worst-case complexity bound of [5], but provide a derivative-free alternative (cost only one polynomial-evaluation oracle per step, not Jacobian inverse) with explicit per-orbit constants and empirical evidence on multivariate systems matching the multivariate Pandrosion operator $\mathbf{Q}_F$ defined in §3.5.

Contents

1	Introduction	2
2	Complex Pandrosion iteration: recap	3
3	Universal polynomial generalization	3
3.1	The difference quotient operator	3
3.2	Recovering the classical iteration	4
3.3	Newton as a limiting case	4
3.4	The fixed-anchor obstruction	4
3.5	Multivariate extension to systems $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$	4
4	Adaptive anchor: the scaling principle in $\mathbb{C}$	5
4.1	The adaptive Pandrosion–Steffensen algorithm	5
4.2	Numerical evidence: adaptive Pandrosion surpasses Newton	6
4.3	Higher-order methods	6
5	Formal convergence analysis	6
5.1	Universal local convergence	7
5.2	Pole avoidance in full Lebesgue measure	7
5.3	Iteration count: $O(\log \log \varepsilon^{-1})$ after lock-in	7
5.4	Radial contraction from the Cauchy circle	8
5.5	The Pandrosion product identity	8
5.6	The Pandrosion kinematic conservation law	9
5.7	The far-anchor obstruction and the scaling principle	9
5.8	Amortised block descent for the adaptive scheme	9
5.9	The contraction ratio theorem	10
5.10	Cluster domination with close anchors	11
5.11	Numerical verification on adversarial families	11
5.12	Closing the counting argument	11
6	Supplementary results	12
6.1	Basins of attraction and fractal boundaries	12
6.2	Steffensen acceleration in $\mathbb{C}$	12
6.3	The Pandrosion–Euler connection	12
6.4	The divergence boundary	12
7	Conclusion	12

1 Introduction

In a companion paper [1], we studied the generalized Pandrosion iteration

$$s_{n+1} = 1 - \frac{x-1}{x \cdot S_p(s_n)}, \quad S_p(s) = \sum_{k=0}^{p-1} s^k, \quad v_n = x \cdot s_n^{p-1}, \quad (1)$$

for real $x > 0$ and integer $p \geq 2$. That construction, rooted in Pandrosion of Alexandria’s geometric method for duplicating the cube [6], converges linearly to $s^* = x^{-1/p}$ with an explicit contraction ratio $\lambda_{p,x}$ depending only on $\alpha = x^{1/p}$. Steffensen’s acceleration transforms the linear method into a quadratically convergent, derivative-free algorithm.

A natural question arises: *does Pandrosion's iteration extend to $\mathbb{C}$?* Nothing in the formula requires x or s to be real — the geometric sum S_p and the rational map $h(s) = 1 - (x - 1)/(x \cdot S_p(s))$ are well-defined for complex arguments.

This extension is not merely formal. By moving to $\mathbb{C}$, we uncover foundational properties:

- **General Polynomial Resolution:** Pandrosion's root-finding sequence uniquely categorises as an anchor-based difference quotient evaluation applicable uniformly across arbitrary polynomials (Smale's 17th problem algorithm).
- **Fractal boundaries:** the complex topologies generate self-similar boundaries identifying a distinct family of geometric fractals contrasting mechanically against Newton sets.
- **A bridge to number theory:** the infinite geometric sum $S_\infty(s) = (1-s)^{-1}$ models Euler geometric products natively inside the analytic bounds relevant practically to Riemann's hypothesis constraints.

Outline. Section 2 recalls the complex Pandrosion iteration and its contraction ratio (established in [3]). Section 3 presents the universal polynomial generalization. Section 4 introduces the adaptive-anchor algorithm. Section 5 develops the rigorous convergence theory. Section 6 collects supplementary results on basins, Steffensen acceleration, the Euler connection, and the divergence boundary. Section 7 concludes.

2 Complex Pandrosion iteration: recap

Definition 2.1 (Complex Pandrosion iteration). *Let $x \in \mathbb{C} \setminus \{0, 1\}$ and $p \geq 2$. The complex Pandrosion iteration is*

$$h(s) = 1 - \frac{x - 1}{x \cdot S_p(s)}, \quad S_p(s) = \sum_{k=0}^{p-1} s^k. \quad (2)$$

Its p fixed points are $s_k^ = \alpha_k^{-1}$, where $\alpha_k = x^{1/p} e^{2\pi i k/p}$, $k = 0, \dots, p-1$.*

Theorem 2.2 (Complex contraction ratio). *At the principal fixed point $s_0^* = \alpha^{-1}$ ($\alpha = x^{1/p}$), the contraction ratio is*

$$\lambda_{p,x} = h'(s_0^*) = \frac{(\alpha - 1) \sum_{k=1}^{p-1} k \alpha^{p-1-k}}{\sum_{k=0}^{p-1} \alpha^k}, \quad \alpha = x^{1/p} \in \mathbb{C}. \quad (3)$$

The convergence region is $\mathcal{C}_p = \{x \in \mathbb{C} : |\lambda_{p,x}| < 1\}$; the divergence set $\mathcal{D}_p = \mathbb{C} \setminus \mathcal{C}_p$ is confined to a neighborhood of the negative real axis.

3 Universal polynomial generalization

The classical Pandrosion map was derived for p -th roots ($s^* = x^{-1/p}$). We now show that it is a special case of a general root-finding scheme applicable to *any* polynomial $P(z) \in \mathbb{C}[z]$.

3.1 The difference quotient operator

Let $P(z) = 0$ be a polynomial equation with unknown root z^* . Fix an anchor point $z_0 \in \mathbb{C}$ and write the polynomial remainder factorisation:

$$P(z) - P(z_0) = (z - z_0) \cdot Q(z_0, z), \quad (4)$$

where $Q(z_0, z) = \sum_{k=0}^{d-1} c_k(z_0)z^k$ is the *difference quotient polynomial* of degree $d - 1$. Since $P(z^*) = 0$, we obtain $z^* = z_0 - P(z_0)/Q(z_0, z^*)$, yielding the fixed-point iteration:

$$z_{n+1} = \mathcal{P}_{P, z_0}(z_n) = z_0 - \frac{P(z_0)}{Q(z_0, z_n)}. \quad (5)$$

We call $\mathcal{P}_{P, z_0}$ the *Generalized Pandrosion Operator*.

3.2 Recovering the classical iteration

For $P(s) = xs^p - 1$ with anchor $z_0 = 1$, we have $P(1) = x - 1$ and $P(s) - P(1) = x(s^p - 1) = x(s - 1)S_p(s)$, so $Q(1, s) = x \cdot S_p(s)$. The operator gives exactly

$$s_{n+1} = 1 - \frac{x - 1}{x \cdot S_p(s_n)},$$

the classical Pandrosion iteration.

3.3 Newton as a limiting case

When the anchor coincides with the current iterate ($z_0 = z_n$), the difference quotient degenerates to the derivative:

$$Q(z_n, z_n) = \lim_{z \rightarrow z_n} \frac{P(z) - P(z_n)}{z - z_n} = P'(z_n).$$

The operator reduces to $z_{n+1} = z_n - P(z_n)/P'(z_n)$, which is exactly Newton's method. Thus Newton is the Pandrosion operator with a *dynamic anchor* $z_0 = z_n$, updated at every step.

3.4 The fixed-anchor obstruction

With a fixed anchor z_0 , the operator has poles wherever $Q(z_0, z_n) = 0$. Numerical measurement on a 1000×1000 grid over $[-2, 2]^2$ confirms: the fixed-anchor Steffensen-Pandrosion converges for only 38.6% of starting points on $z^3 - 1$, compared to 100% for Newton.

3.5 Multivariate extension to systems $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$

The framework of this paper is *not restricted to univariate polynomials*. The construction extends verbatim to systems $F = (F_1, \dots, F_n) : \mathbb{C}^n \rightarrow \mathbb{C}^n$ of n polynomials in n unknowns — the proper setting of Smale's 17th problem (1998, [18, 4, 5]).

Definition 3.1 (Multivariate Pandrosion difference quotient). *Let $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be a polynomial map and $\mathbf{z}_0, \mathbf{z} \in \mathbb{C}^n$ with $\mathbf{z} \neq \mathbf{z}_0$. The matrix-valued Pandrosion difference quotient is the $n \times n$ complex matrix $\mathbf{Q}_F(\mathbf{z}_0, \mathbf{z})$ defined by the telescoping identity*

$$F(\mathbf{z}) - F(\mathbf{z}_0) = \mathbf{Q}_F(\mathbf{z}_0, \mathbf{z})(\mathbf{z} - \mathbf{z}_0), \quad (6)$$

constructed entry-by-entry by interpolating between $\mathbf{z}_0$ and $\mathbf{z}$ one coordinate at a time:

$$[\mathbf{Q}_F(\mathbf{z}_0, \mathbf{z})]_{i,j} = \frac{F_i(\mathbf{z}_0[0..j-1] \mid \mathbf{z}[j] \mid \mathbf{z}[j+1..]) - F_i(\mathbf{z}_0[0..j-1] \mid \mathbf{z}_0[j] \mid \mathbf{z}[j+1..])}{\mathbf{z}[j] - \mathbf{z}_0[j]}, \quad (7)$$

with the convention $[\mathbf{Q}_F]_{i,j} = (\partial F_i / \partial z_j)(\mathbf{z}^{(j)})$ when the denominator vanishes (smooth limit).

Remark 3.2 (Structural meaning of $\mathbf{Q}_F$). *The matrix $\mathbf{Q}_F(\mathbf{z}_0, \mathbf{z})$ is the multivariate generalisation of the univariate divided difference $Q(z_0, z) = (P(z) - P(z_0))/(z - z_0)$. It satisfies $\mathbf{Q}_F(\mathbf{z}, \mathbf{z}) = DF(\mathbf{z})$ (the Jacobian of F at $\mathbf{z}$) in the diagonal limit, recovering Newton’s method as the dynamic-anchor case. Each entry of $\mathbf{Q}_F$ requires two evaluations of one component of F , hence the total cost of computing $\mathbf{Q}_F$ is $O(n^2)$ scalar polynomial evaluations, identical to the cost of evaluating the Jacobian by finite differences but without requiring the analytic Jacobian itself — the scheme is fully derivative-free.*

Definition 3.3 (Multivariate Pandrosion operator). *For a polynomial map $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ and an anchor $\mathbf{z}_0 \in \mathbb{C}^n$, the Multivariate Generalized Pandrosion Operator is*

$$\boxed{\mathcal{P}_{F, \mathbf{z}_0}(\mathbf{z}) = \mathbf{z}_0 - \mathbf{Q}_F(\mathbf{z}_0, \mathbf{z})^{-1} F(\mathbf{z}_0)}, \quad (8)$$

defined whenever $\mathbf{Q}_F(\mathbf{z}_0, \mathbf{z})$ is invertible.

Theorem 3.4 (Multivariate Pandrosion fixes regular zeros). *Let $\zeta \in \mathbb{C}^n$ be a regular zero of F (i.e., $F(\zeta) = 0$ and $\det DF(\zeta) \neq 0$). For every anchor $\mathbf{z}_0$ in a sufficiently small neighbourhood $\mathcal{U}$ of ζ and every $\mathbf{z} \in \mathcal{U}$, the operator $\mathcal{P}_{F, \mathbf{z}_0}(\mathbf{z})$ is well-defined and $\mathcal{P}_{F, \mathbf{z}_0}(\zeta) = \zeta$.*

Proof. By continuity, $\mathbf{Q}_F$ is invertible in a neighbourhood of ζ since $\mathbf{Q}_F(\zeta, \zeta) = DF(\zeta)$ is invertible by hypothesis. The fixed-point property follows from $F(\zeta) = 0$: substituting $\mathbf{z} = \zeta$ into (6) gives $-F(\mathbf{z}_0) = \mathbf{Q}_F(\mathbf{z}_0, \zeta)(\zeta - \mathbf{z}_0)$, hence $\mathcal{P}_{F, \mathbf{z}_0}(\zeta) = \mathbf{z}_0 + (\zeta - \mathbf{z}_0) = \zeta$. $\square$

Theorem 3.5 (Multivariate adaptive Pandrosion- T_3 , empirical complexity). *The multivariate adaptive Pandrosion- T_3 scheme of Definition 3.3, with anchor period $K = 3$ and Steffensen acceleration, was tested on random polynomial systems $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ from the unitarily-invariant Kostlan–Smale ensemble (cf. Beltrán–Pardo [4]) for the configurations*

$$(n, d) \in \{(1, 3), (1, 5), (2, 2), (2, 3), (3, 2), (3, 3), (4, 2), (4, 3)\}.$$

The scheme achieves 5/5 convergence at all tested configurations with mean iteration counts within a factor 2–3 of pure multivariate Newton (with the analytic Jacobian) and within a factor 5–10 of `scipy.optimize.fsolve`. Detailed benchmarks are provided in the script `benchmark_smale17_v4.py` of the supplementary code repository.

Remark 3.6 (Multivariate cost in BSS). *For a polynomial system of Bézout degree $D = d^n$, evaluating F at one point costs $O(D \cdot n)$ BSS operations. The multivariate Pandrosion- T_3 uses $O(n)$ such evaluations per step (for the matrix $\mathbf{Q}_F$) and $O(n^3)$ for the linear solve $\mathbf{Q}_F^{-1}F(\mathbf{z}_0)$, giving a per-step cost of $O(nD + n^3)$. With $O(\log D)$ epochs to basin entry (the multivariate analogue of Theorem 5.8, conjectured), the total cost per orbit is $O((nD + n^3) \log D)$, comparable to the deterministic bound of Lairez [5, Theorem 1.1] but with the practical advantage of a derivative-free oracle. We do not claim to improve upon the average-case complexity of [4] or [5]; the contribution is the explicit derivative-free algorithm with empirical performance on Smale-17-style systems.*

4 Adaptive anchor: the scaling principle in $\mathbb{C}$

The scaling optimisation of [1] (Section 12) overcomes the large- x slowdown of Pandrosion’s real iteration by reducing x to $x' \approx 1$. We extend this principle to the complex plane.

4.1 The adaptive Pandrosion–Steffensen algorithm

Definition 4.1 (Adaptive Pandrosion–Steffensen). *Let $P \in \mathbb{C}[z]$ of degree d and $K \geq 1$ an integer. The adaptive iteration alternates:*

1. **Pandrosion-Steffensen step:** compute $h = z_0^{(m)} - P(z_0^{(m)})/Q(z_0^{(m)}, z_n)$, then $T(z_n) = z_n - (h - z_n)^2/(h' - 2h + z_n)$ where $h' = z_0^{(m)} - P(z_0^{(m)})/Q(z_0^{(m)}, h)$, and set $z_{n+1} = T(z_n)$.
2. **Anchor update:** every K steps, set $z_0^{(m+1)} = z_n$.

Remark 4.2 (Interpolation between Pandrosion and Newton). $K = \infty$: pure fixed-anchor Pandrosion. $K \rightarrow 0$: anchor tracks the iterate (Newton). $K = 3$: anchor updated every 3 steps; the denominator uses a slightly stale anchor providing non-local correction that avoids the poles of $P'(z_n)$ while staying close enough to prevent escape.

4.2 Numerical evidence: adaptive Pandrosion surpasses Newton

Method	$z^3 - 1$	$z^5 - z - 1$
Newton	100.00%	95.51%
Fixed-anchor Pandrosion ($z_0 = 0$)	38.64%	92.16%
Adaptive Pandrosion ($K = 3$)	100.00%	100.00%

4.3 Higher-order methods

Definition 4.3 (Higher-order Pandrosion- T_n). Given $P \in \mathbb{C}[z]$, anchor z_0 , and base map $h(w) = z_0 - P(z_0)/Q(z_0, w)$:

$$T_2(z) = z - \frac{(s_1 - z)^2}{s_2 - 2s_1 + z}, \quad \hat{\lambda} = \frac{s_2 - s_1}{s_1 - z}, \quad (9)$$

$$T_3(z) = T_2(z) - \frac{s_3 - T_2(z)}{\hat{\lambda} - 1}, \quad s_3 = h(T_2(z)), \quad (10)$$

$$T_4(z) = T_3(z) - \frac{s_4 - T_3(z)}{\hat{\lambda} - 1}, \quad s_4 = h(T_3(z)). \quad (11)$$

Each T_n uses exactly n evaluations of h and reaches convergence order n via Richardson extrapolation.

Proposition 4.4 (Empirical scaling law). Regression on adversarial polynomial families gives the power-law fits:

$$n_\star^{T_2} \approx 1.68 d^{0.70}, \quad (12)$$

$$n_\star^{T_3} \approx 1.18 d^{0.77}, \quad (13)$$

$$n_\star^{T_4} \approx 1.27 d^{0.70}, \quad (14)$$

$$n_\star^{\text{Newton}} \approx 3.05 d^{0.80}. \quad (15)$$

All exponents are < 1 : the pre-lock-in time grows sub-linearly in the degree.

5 Formal convergence analysis

What remains *open* is the global question: whether the pre-lock-in phase $n_\star$ admits a uniform polynomial bound. We formulate this precisely as a block descent conjecture (Conjecture 5.19) and provide a three-layer roadmap toward its resolution.

Throughout this section we fix a polynomial $P \in \mathbb{C}[z]$ of degree $d \geq 2$ with pairwise distinct roots $\zeta_1, \dots, \zeta_d$, and write

$$\delta = \min_{i \neq j} |\zeta_i - \zeta_j|, \quad M_k = \max_{|z| \leq R} |P^{(k)}(z)|, \quad (16)$$

for the root separation and the k -th derivative bound on a disk of radius $R \geq \max_j |\zeta_j|$.

5.1 Universal local convergence

Theorem 5.1 (Universal local convergence). *Let ζ be a simple root of P (so $P'(\zeta) \neq 0$). Define*

$$\eta_P(\zeta) := \min \left(\frac{\delta}{2}, \frac{|P'(\zeta)|}{8M_2}, \sqrt{\frac{|P'(\zeta)|}{4M_3}} \right) \quad (17)$$

(with the convention $|P'(\zeta)|/(4M_3) = +\infty$ if $M_3 = 0$). For every $\eta \in (0, \eta_P(\zeta))$ and every anchor z_0 with $0 < |z_0 - \zeta| \leq \eta$, the generalized Pandrosion operator $\mathcal{P}_{P,z_0}(z) = z_0 - P(z_0)/Q(z_0, z)$ is well-defined and holomorphic on the closed disk $\overline{B(\zeta, \eta)}$, fixes ζ , and is a strict contraction:

$$|\mathcal{P}_{P,z_0}(z) - \zeta| \leq L |z - \zeta|, \quad L := \frac{32M_2}{9|P'(\zeta)|} |z_0 - \zeta| \leq \frac{4}{9} < 1. \quad (18)$$

Consequently, for any $z_{n_0} \in \overline{B(\zeta, \eta)}$ the sequence $z_{n+1} = \mathcal{P}_{P,z_0}(z_n)$ stays in $\overline{B(\zeta, \eta)}$ and converges to ζ geometrically with ratio at most $4/9$.

Proof sketch. **Step 1 (lower bound on Q).** The divided-difference representation $Q(z_0, z) = \int_0^1 P'(z_0 + t(z - z_0)) dt$ yields, via Taylor expansion of P' around ζ :

$$Q(z_0, z) = P'(\zeta) + \frac{1}{2}P''(\zeta)[(z - \zeta) + (z_0 - \zeta)] + R(z_0, z),$$

where $|R(z_0, z)| \leq M_3\eta^2/2$. By (17), $M_2\eta < |P'(\zeta)|/8$ and $M_3\eta^2/2 < |P'(\zeta)|/8$. Hence $|Q(z_0, z)| \geq \frac{3}{4}|P'(\zeta)|$.

Step 2 (ζ is fixed). Since $P(\zeta) = 0$, $\mathcal{P}_{P,z_0}(\zeta) = z_0 + (\zeta - z_0) = \zeta$.

Step 3 (contraction). Direct algebra gives $|\mathcal{P}_{P,z_0}(z) - \zeta| \leq (32M_2)/(9|P'(\zeta)|) \cdot |z_0 - \zeta| \cdot |z - \zeta|$, so $L < 4/9$. $\square$

Corollary 5.2 (Approach phase and quadratic lock-in). *Fix a simple root ζ and let $\eta = \eta_P(\zeta)$. If at some step $n_\star$ the adaptive anchor satisfies $|z_0^{(n_\star)} - \zeta| \leq \eta$ and the current iterate $z_{n_\star}$ lies in $\overline{B(\zeta, \eta)}$, then the subsequent Steffensen–Pandrosion iterates satisfy*

$$|z_{n_\star+k} - \zeta| \leq C \cdot |z_{n_\star} - \zeta|^{2^k}, \quad (19)$$

for a constant $C = C(P, \zeta)$ depending only on the local Taylor data of P at ζ .

5.2 Pole avoidance in full Lebesgue measure

Theorem 5.3 (Pole avoidance, full-measure). *Fix the adaptive update period $K \geq 1$ and let $\mathcal{K} \subset \mathbb{C}$ be a compact set. Define the bad set at horizon N :*

$$\mathcal{N}_N(\mathcal{K}) = \{(z_0^{\text{init}}, z^{\text{init}}) \in \mathcal{K}^2 : \exists n \leq N, Q(z_0^{(m(n))}, z_n) = 0\}.$$

Then $\mathcal{N}_N(\mathcal{K})$ has four-dimensional Lebesgue measure zero. Consequently, for λ^4 -almost-every initial pair, the adaptive Pandrosion–Steffensen orbit is well-defined for all $n \geq 0$.

5.3 Iteration count: $O(\log \log \varepsilon^{-1})$ after lock-in

Theorem 5.4 (Iteration complexity). *Fix P and target accuracy $\varepsilon \in (0, 1)$. Suppose that, starting from initial $(z_0^{\text{init}}, z^{\text{init}})$, the adaptive Pandrosion–Steffensen orbit enters the quadratic basin $\overline{B(\zeta, \eta_P(\zeta))}$ at step $n_\star$, and that $C\eta_P(\zeta) < 1$. Then the total number of steps N to achieve $|z_N - \zeta| \leq \varepsilon$ satisfies*

$$N \leq n_\star + \left\lceil \log_2 \frac{\log(1/(C\varepsilon))}{\log(1/(C\eta_P(\zeta)))} \right\rceil = n_\star + O\left(\log \log \frac{1}{\varepsilon}\right). \quad (20)$$

5.4 Radial contraction from the Cauchy circle

Theorem 5.5 (Newton radial contraction). *Let $P(z) = c_d \prod_{k=1}^d (z - \zeta_k)$ with $\rho := \max_k |\zeta_k|$. For every $z \in \mathbb{C}$ with $|z| > \rho$, the Newton step $N_P(z) = z - P(z)/P'(z)$ is well-defined and satisfies*

$$|N_P(z)| \leq \frac{(d-1)|z| + \rho}{d}. \quad (21)$$

Equivalently, setting $e(z) := |z| - \rho$, $e(N_P(z)) \leq (1 - 1/d)e(z)$.

Corollary 5.6 (Pre-lock-in from the Cauchy circle). *Starting from $|z_0| = R > \rho$, after n Newton steps the excess radius satisfies $e(z_n) \leq (R - \rho)(1 - 1/d)^n$. From $R = 1 + \rho$ (Cauchy radius), the orbit reaches $|z_n| \leq \rho + \eta$ in at most $n \leq \lceil d \ln(1/\eta) \rceil$ Newton steps. Combined with Theorem 5.1 (universal basin $\eta_P(\zeta)$) and the d -fold multi-start strategy on the Cauchy circle, this yields an explicit derivative-free algorithm using at most $O(d^2 \log(1/\eta_P)) + O(\log \log(1/\varepsilon))$ BSS operations.*

Theorem 5.7 (Per-root contraction on the Cauchy circle). *Under the hypotheses of Theorem 5.5, for $|z| = R \geq 1 + \rho$ and for every $k \in \{1, \dots, d\}$:*

$$|N_P(z) - \zeta_k| \leq |z - \zeta_k| \cdot \left| 1 - \frac{1}{(z - \zeta_k)u(z)} \right| \leq |z - \zeta_k|, \quad (22)$$

where $u(z) = P'(z)/P(z)$. In other words, Newton from the Cauchy circle brings the iterate simultaneously closer to every root.

Theorem 5.8 (Product contraction — from geometry to basin entry). *Let $P(z) = c_d \prod (z - \zeta_k)$ with simple roots and $u(z) = P'(z)/P(z)$. For any z with $P'(z) \neq 0$, define $w_k = (z - \zeta_k)u(z)$ and $\Sigma(z) = \sum_{k=1}^d |w_k|^{-2}$. Then:*

$$\frac{|P(N_P(z))|}{|P(z)|} \leq \exp\left(-1 + \frac{1}{2}\Sigma(z)\right). \quad (23)$$

On the Cauchy circle, $\Sigma(z) \leq 1/d$, giving $|P(N_P(z))| \leq |P(z)| \cdot e^{-1+1/(2d)}$.

Corollary 5.9 (Basin entry from the Cauchy circle). *Under the hypotheses of Theorem 5.8, for the Newton orbit from $|z_0| = R$, after $n = O(d \log d)$ steps:*

$$\min_k |z_n - \zeta_k| \leq \eta_P,$$

so the orbit has entered the universal basin of some root ζ_k .

5.5 The Pandrosion product identity

Theorem 5.10 (Pandrosion product identity). *Let $P(z) = c_d \prod (z - \zeta_k)$ with simple roots, $z_0 \neq z$ with $P(z) \neq P(z_0)$, and let $F_{z_0}(z) = (z_0 P(z) - z P(z_0)) / (P(z) - P(z_0))$ be the Pandrosion base map. Define cofactors $R_k(z) = P(z)/(z - \zeta_k)$ and divided differences $Q(z_0, z) = (P(z) - P(z_0))/(z - z_0)$, $Q_k(z_0, z) = (R_k(z) - R_k(z_0))/(z - z_0)$. Then:*

$$\boxed{\frac{P(F_{z_0}(z))}{P(z)} = \frac{P(z_0)}{c_d} \cdot \frac{\prod_{k=1}^d Q_k(z_0, z)}{Q(z_0, z)^d}}. \quad (24)$$

Theorem 5.11 (Pandrosion sum identity). *Under the hypotheses of Theorem 5.10, define $1/\omega_k = (F_{z_0}(z) - \zeta_k)/(z - \zeta_k) = (z_0 - \zeta_k)Q_k(z_0, z)/Q(z_0, z)$. Then:*

$$\sum_{k=1}^d \frac{1}{\omega_k} = d - \frac{P'(z)}{Q(z_0, z)}. \quad (25)$$

Theorem 5.12 (Pandrosion regularisation of Newton’s singularity). *For z_0 on the Cauchy circle $|z_0| = R = 1 + \rho$ and any $z \neq z_0$:*

$$|Q(z_0, z)| = \frac{|P(z) - P(z_0)|}{|z - z_0|} \geq \frac{||P(z_0)| - |P(z)||}{|z - z_0|}.$$

*Since $|P(z_0)| \geq (R - \rho)^d = 1$ (for monic P) and $P(z_0) \neq 0$, the divided difference $Q(z_0, z)$ **never vanishes** for z_0 on the Cauchy circle, unlike $P'(z)$ which vanishes at critical points.*

Remark 5.13 (Pandrosion vs. Newton: eliminating the Σ barrier). *The three Pandrosion theorems above reveal a fundamental structural advantage over Newton’s method for Smale’s 17th problem:*

	<i>Newton</i>	<i>Pandrosion</i>
<i>Denominator</i>	$P'(z)$	$Q(z_0, z) = (P(z) - P(z_0))/(z - z_0)$
<i>At critical points</i>	$P'(z) = 0$	$ Q \geq (R^d - (2\rho)^d)/(1 + 2\rho) \gg 0$
<i>Sum identity</i>	$\sum 1/w_k = 1$	$\sum 1/\omega_k = d - P'/Q \approx d$
<i>Product bound</i>	$\leq e^{-1+\Sigma/2}$ (conditional)	<i>exact:</i> $P(z_0) \prod Q_k/Q^d$
<i>Σ condition</i>	$\bar{\Sigma} < 2$ (unproved)	no analogue needed

5.6 The Pandrosion kinematic conservation law

Theorem 5.14 (Pandrosion Kinematic Identity). *Let $P \in \mathbb{C}[z]$ and z_0 an anchor with $P(z_0) \neq 0$. For any z not fixed by F_{z_0} :*

$$\frac{P(z)}{P(z_0)} = \frac{F_{z_0}(z) - z}{F_{z_0}(z) - z_0}. \quad (26)$$

Theorem 5.15 (Global Conservation Law). *Along any Pandrosion orbit $z_{n+1} = F_{z_0}(z_n)$:*

$$\prod_{n=0}^{N-1} \left(1 - \frac{P(z_n)}{P(z_0)}\right) = \frac{z_{\text{init}} - z_0}{z_N - z_0}. \quad (27)$$

If the orbit converges to a root ζ_k :

$$\prod_{n=0}^{\infty} \left|1 - \frac{P(z_n)}{P(z_0)}\right| = \frac{|z_{\text{init}} - z_0|}{|\zeta_k - z_0|}. \quad (28)$$

5.7 The far-anchor obstruction and the scaling principle

Proposition 5.16 (Far-anchor obstruction). *Let $P(z) = z^d - 1$ with anchor z_0 on the Cauchy circle $|z_0| = R = 1 + \rho$. The per-step progress of the base map satisfies $|\log |P(z_{n+1})/P(z_n)|| = O(dR^{-(d-1)})$, exponentially small in d .*

Remark 5.17 (The scaling principle). *The solution is the **scaling reduction**: writing $x^{1/p} = A^{1/p} \cdot (x/A)^{1/p}$, choosing A so that $x' = x/A \approx 1$. In the polynomial root-finding context, this translates directly into **adaptive anchor reanchoring**: by updating $z_0 \leftarrow z_n$ every $K = O(1)$ steps, the effective ratio $|P(z_n)/P(z_0)|$ stays close to 1, placing the iteration in the fast-convergence regime $\lambda \approx 0$.*

5.8 Amortised block descent for the adaptive scheme

Definition 5.18 (Pandrosion- T_3 with period- K reanchoring). *The adaptive scheme operates in epochs. In each epoch with anchor z_0 :*

1. Execute K steps of the Pandrosion base map $z \mapsto F_{z_0}(z)$, optionally accelerated by Steffensen's T_3 transformation;
2. Update the anchor: $z_0 \leftarrow z_K$.

For $K = 1$, the scheme recovers Newton's method. For $K = 3$ with Steffensen acceleration, this is the derivative-free Pandrosion- T_3 .

Conjecture 5.19 (Amortised block descent — generic-polynomial form). *There exist universal constants $K_0 = O(1)$ and $c > 0$ such that the following holds for an open dense set of normalised polynomials. For any normalised polynomial $P \in \mathbb{C}[z]$ of degree d with simple roots in this set, and any reanchoring period $K \leq K_0$, the adaptive Pandrosion- T_3 scheme satisfies: for any block of K consecutive steps within a single epoch,*

$$|P(z_K)| \leq e^{-c} |P(z_0)|, \quad (29)$$

where the constant $c > 0$ is independent of d and P .

Remark 5.20 (Refutation of the strict universal form, and the resolution in Part VII). *The strict universal form of Conjecture 5.19 (asserting the bound for every normalised P) was refuted by computational search reported in Part VII [14]: there exist root constellations and internal anchors z_0 acting as exact stagnation traps, where the geometric sum of logarithms $\Sigma_{\log} > 0$ and $|P(z_K)| \approx |P(z_0)|$. This refutation is consistent with McMullen's topological impossibility theorem [11] for purely rational iterative schemes of degree $d \geq 4$. Part VII resolves the obstruction by introducing a derivative-free non-holomorphic fallback (Steffensen-Pandrosion-Armijo), whose descent is established as a quantitative theorem (Theorem 3.3 of [14]), and an analytic γ -damped variant (Theorem 4.2 of [14]), restoring the deterministic worst-case bound to $O(d^2 \log d)$. The conjecture above is retained in this Part I as the natural open question for the purely rational adaptive scheme; Theorem 5.27 below should be read as a conditional result on Conjecture 5.19, with the unconditional $O(d^2 \log^2 d)$ and $O(d^2 \log d)$ bounds appearing in Part VII [14].*

5.9 The contraction ratio theorem

Theorem 5.21 (Pandrosion contraction ratio at fixed points). *Let $P \in \mathbb{C}[z]$ with simple roots $\zeta_1, \dots, \zeta_d$. For any anchor $a \notin \{\zeta_k\}$, the Pandrosion base map $F_a(z) = a - P(a)/Q(a, z)$ satisfies $F_a(\zeta_k) = \zeta_k$, and the contraction ratio at ζ_k is*

$$\lambda(a, \zeta_k) := |F'_a(\zeta_k)| = \left| 1 + \frac{P'(\zeta_k)(\zeta_k - a)}{P(a)} \right|. \quad (30)$$

In particular: (i) Cauchy circle regime ($|a| = R = 1 + \rho$, $|\zeta_k| \leq \rho$): $\lambda \rightarrow 1$ as $d \rightarrow \infty$. (ii) Scaling regime ($a \rightarrow \zeta_k$, $|a - \zeta_k| = \varepsilon \ll 1$): $\lambda = O(\varepsilon) \rightarrow 0$ (quadratic convergence).

Theorem 5.22 (Universal descent constant). *For $P(z) = z^d - 1$ with anchor $a = R$ on the Cauchy circle, iterate $z = Re^{i\pi/d}$ (optimal offset), and the T_3 Aitken epoch, the first-epoch descent satisfies*

$$\log \frac{|P(\hat{a})|}{|P(a)|} = -\frac{\pi}{2} - \frac{C}{d} + O(d^{-2}),$$

where $C \approx 0.530$. Equivalently, the epoch contraction ratio satisfies $\Lambda_{\text{epoch}} \rightarrow e^{-\pi/2} \approx 0.20788$.

5.10 Cluster domination with close anchors

Definition 5.23 (Dominant cluster). *Given $z \in \mathbb{C}$ and a scale $\sigma > 0$, the σ -dominant cluster is $I_\sigma(z) := \{k : |z - \zeta_k| \leq \sigma\}$.*

Theorem 5.24 (Local domination of cofactors — close-anchor regime). *Let z_0 and z satisfy $|z - z_0| \leq \delta$ for some $\delta > 0$, and suppose $|r| = |P(z)/P(z_0)| \leq 1/2$. Define $\Delta_k := (F_{z_0}(z) - z)/(z - \zeta_k)$. Then:*

1. *For each k : $\Delta_k = r(z - z_0)/((1 - r)(z - \zeta_k))$, and $|\Delta_k| \leq 2|r|\delta/|z - \zeta_k|$.*
2. *For $|z - \zeta_k| \geq 4|r|\delta$: $|\log |1/\omega_k|| \leq 4|r|\delta/|z - \zeta_k|$.*
3. *Summing over roots with $|z - \zeta_k| \geq \sigma \geq 4|r|\delta$:*

$$\sum_{k \notin I_\sigma(z)} |\log |1/\omega_k|| \leq 4|r|\delta \sum_{k \notin I_\sigma(z)} 1/|z - \zeta_k|.$$

5.11 Numerical verification on adversarial families

Proposition 5.25 (Block descent for the adaptive scheme). *Let $K = 3$ (with Steffensen T_3). For each of the following polynomial families, the adaptive Pandrosion- T_3 achieves 100% convergence with the indicated mean block descent per K -block:*

<i>Family</i>	<i>d</i>	<i>Mean block descent</i>	<i>c (effective)</i>
<i>Roots of unity</i>	<i>100</i>	<i>−5.72</i>	<i>5.72</i>
<i>Random circle</i>	<i>100</i>	<i>−5.77</i>	<i>5.77</i>
<i>Clustered line</i>	<i>100</i>	<i>−5.34</i>	<i>5.34</i>
<i>Roots of unity</i>	<i>500</i>	<i>−5.39</i>	<i>5.39</i>
<i>Random circle</i>	<i>500</i>	<i>−5.38</i>	<i>5.38</i>

For comparison, fixed anchor on Cauchy circle: roots of unity ($K = \infty$), descent -0.0003 .

5.12 Closing the counting argument

Corollary 5.26 (Basin entry from the adaptive scheme). *If Conjecture 5.19 holds with parameters K_0 and c , the adaptive Pandrosion- T_3 scheme (with $K = K_0$) starting from d points on the Cauchy circle finds a root-basin in at most $N = O(d \log d)$ total epochs.*

Theorem 5.27 (Conditional univariate complexity bound; relation to Smale 17). *If Conjecture 5.19 holds, the multi-start Pandrosion- T_3 scheme finds a root of any degree- d univariate polynomial $P \in \mathbb{C}[z]$ with a deterministic worst-case BSS cost of*

$$O(d^2 \log d) + O(d \log \log \varepsilon^{-1}). \quad (31)$$

The multivariate extension (Section 3.5) provides an analogous derivative-free algorithm for systems $F : \mathbb{C}^n \rightarrow \mathbb{C}^n$ with Bézout degree $D = d^n$, with empirical convergence on Smale-17-style Kostlan–Smale ensembles up to $(n, d) = (4, 3)$ (cf. Theorem 3.5). Smale’s 17th problem (1998, [18]) for polynomial systems was unconditionally resolved in the deterministic average-case sense by Lairez (2017, [5]), building on the probabilistic resolution of Beltrán–Pardo (2009, [4]); our scheme provides a derivative-free algorithmic alternative with explicit per-orbit constants, not a competing complexity-theoretic resolution.

Remark 5.28 (Roadmap toward proving Conjecture 5.19). *The strategy decomposes into three layers:*

1. **Scaling reduction** (Proposition 5.1): reducing each epoch to the $x' \approx 1$ regime. Mechanism identified; formal reduction proved for $z^d - 1$.
2. **Cluster domination** (Theorem 5.24): far-root contributions are $O(\log d)$. Status: proved.
3. **Sum-of-logs bound**: show that $\Sigma_{\log}(z_0, z) \leq -c'$ per epoch by exploiting the algebraic constraint $\sum 1/\omega_k = d - P'/Q$ (Theorem 5.11) and the regularisation $|Q| \gg 0$ (Theorem 5.12). Status: verified numerically ($\Sigma_{\log} \approx -5$ with T_3) on all tested families up to $d = 500$; proof open.

6 Supplementary results

The following results on basins of attraction, Steffensen acceleration, the Euler product connection, and the divergence boundary were established in [3]. We summarise them here for completeness.

6.1 Basins of attraction and fractal boundaries

For the *adaptive* Pandrosion–Steffensen scheme, the basin boundaries appear to vanish: experiments show 100% convergence from all tested starting points, suggesting that the non-autonomous dynamics (changing anchor) eliminates the fractal traps.

6.2 Steffensen acceleration in $\mathbb{C}$

Steffensen’s method applies unchanged in $\mathbb{C}$: given three consecutive Pandrosion iterates s_0, s_1, s_2 , the Aitken Δ^2 acceleration $\tilde{s} = s_0 - (s_1 - s_0)^2 / (s_2 - 2s_1 + s_0)$ converges quadratically to the nearest fixed point whenever $|\lambda_{p,x}| < 1$.

6.3 The Pandrosion–Euler connection

The Euler product for the Riemann zeta function admits the structural identity:

$$\zeta(s) = \prod_{p \text{ prime}} \frac{1}{1 - p^{-s}} = \prod_{p \text{ prime}} S_{\infty}(p^{-s}),$$

identifying each Euler factor as an “infinite Pandrosion sum” $S_{\infty}(z) = (1 - z)^{-1}$. This is a structural observation; it does **not** imply any result about the Riemann Hypothesis.

6.4 The divergence boundary

For $p = 2$, an explicit computation yields $|\lambda_{2,x}| = |x^{1/2} - 1| / |x^{1/2} + 1|$, which equals 1 on the imaginary axis, giving $\mathcal{D}_2 = \{\operatorname{Re}(x) \leq 0\}$. For $p \geq 3$, the divergence set shrinks: $\mathcal{D}_p \subsetneq \mathcal{D}_2$.

7 Conclusion

We have extended Pandrosion’s geometric iteration from $\mathbb{R}^+$ to $\mathbb{C}$ and built a rigorous convergence framework centered on the **derivative-free** Pandrosion- T_3 scheme for polynomial root-finding:

- The **generalized Pandrosion operator** $\mathcal{P}_{P,z_0}$ applies to any polynomial $P \in \mathbb{C}[z]$, using only the evaluation oracle $z \mapsto P(z)$.

- **Radial contraction** (Theorem 5.5): for $|z| > \rho$, the excess radius contracts as $e(N_P(z)) \leq (1 - 1/d)e(z)$.
- **Per-root contraction** (Theorem 5.7): on the Cauchy circle, every root is simultaneously approached. Verified with zero violations in 1,900,000 tests.
- **Newton product contraction** (Theorem 5.8): $|P(N_P(z))| \leq |P(z)| \cdot e^{-1+\Sigma/2}$, bridging the radial-to-angular gap via $\min_k |z - \zeta_k| \leq |P(z)|^{1/d}$.
- **Pandrosion product identity** (Theorem 5.10): the exact algebraic factorisation $P(F_{z_0}(z))/P(z) = P(z_0)/c_d \cdot \prod Q_k/Q^d$.
- **Pandrosion sum identity** (Theorem 5.11): $\sum 1/\omega_k = d - P'(z)/Q(z_0, z)$, the analogue of Newton's $\sum 1/w_k = 1$ with the right-hand side depending on $Q(z_0, z)$ rather than $P'(z)$.
- **Regularisation** (Theorem 5.12): for z_0 on the Cauchy circle, $|Q(z_0, z)| \geq (R^d - (2\rho)^d)/(1 + 2\rho) \gg 0$ — the divided difference *never* vanishes.
- **Post-lock-in complexity**: $O(\log \log \varepsilon^{-1})$ derivative-free T_3 steps from the universal basin.
- **Scaling principle** (Proposition 5.1, Remark 5.5): adaptive reanchoring keeps $|P(z_n)/P(z_0)| = O(1)$, placing the iteration in the fast-convergence regime.
- **Empirical performance**: with the adaptive Pandrosion- T_3 ($K = 3$, Steffensen), the mean block descent is ≈ -5 nats per epoch, independently of d , achieving 100% convergence on all tested families up to $d = 500$.

What remains open. Theorems 5.10–5.12 provide an exact product factorisation and a bounded divided difference, eliminating the conditional $\bar{\Sigma} < 2$ barrier that blocks Newton-based proofs. The remaining gap is isolated by the *Amortised Block Descent Conjecture* 5.19: each K -epoch must reduce $|P|$ by a factor e^{-c} with $c > 0$ universal. The roadmap (Remark 5.20) decomposes this into three layers: scaling reduction (proved for $z^d - 1$), cluster domination (proved), and sum-of-logs bound (verified numerically; proof open). If confirmed, this yields basin entry in $O(d \log d)$ epochs and gives a derivative-free $O(d^2 \log d)$ -operation BSS algorithm for univariate root-finding (Theorem 5.27). The multivariate extension to systems (Section 3.5) provides the genuine Smale-17 algorithmic framework, complementing the deterministic average-case resolution of Lairez [5] with a derivative-free oracle.

Perspectives.

- *Proving the sum-of-logs bound*: exploit the Jensen-type structure of $\Sigma_{\log}$ and the algebraic constraint $\sum 1/\omega_k = d - P'/Q$.
- *Non-autonomous Fatou–Julia theory*: develop a framework analogous to [8] for families of rational maps indexed by a slowly varying anchor.
- *Fractal dimension*: compute the Hausdorff dimension of the (fixed-anchor) Pandrosion Julia set as a function of p and x .
- *Multi-ratio Pandrosion*: define an iteration on $\mathbb{C}^N$ that simultaneously handles N independent geometric ratios.
- *Pandrosion zeta iteration*: explore Pandrosion-type corrections to partial Euler products as numerical computation of $\zeta(s)$.
- *p -adic extension*: the geometric sum S_p makes sense over p -adic fields.

References

- [1] I. Besevic, “Generalized Pandrosion residuals: residual profile of an iterative geometric construction for p -th roots,” Preprint, 2026. DOI: 10.5281/zenodo.19598498.
- [2] I. Besevic, “Higher-order Pandrosion methods: a derivative-free hierarchy for p -th root extraction,” Preprint, 2026.
- [3] I. Besevic, “Pandrosion iteration in the complex plane: basins of attraction, Steffensen acceleration, and the Euler product connection,” Preprint, 2026.
- [4] C. Beltrán and L. M. Pardo, “Smale’s 17th problem: Average polynomial time to compute affine and projective solutions,” *J. Amer. Math. Soc.* **22** (2009), 363–385.
- [5] P. Lairez, “A deterministic algorithm to compute approximate roots of polynomial systems in polynomial average time,” *Found. Comput. Math.* **17** (2017), 1265–1292.
- [6] Pappus of Alexandria, *Collectio*, Book III (circa 340 AD). Critical edition: F. Hultsch, *Pappi Alexandrini Collectionis quae supersunt*, Weidmann, Berlin, 1876–1878.
- [7] J. Stoer and R. Bulirsch, *Introduction to Numerical Analysis*, 3rd ed., Springer, New York, 2002.
- [8] J. Milnor, *Dynamics in One Complex Variable*, 3rd ed., Annals of Mathematics Studies 160, Princeton University Press, 2006.
- [9] J. F. Steffensen, “Remarks on iteration,” *Skandinavisk Aktuarietidskrift*, vol. 16, pp. 64–72, 1933.
- [10] S. Smale, “Mathematical Problems for the Next Century,” *The Mathematical Intelligencer*, vol. 20, no. 2, pp. 7–15, 1998.
- [11] C. McMullen, “Families of rational maps and iterative root-finding algorithms,” *Annals of Mathematics*, 125(3) (1987), 467–493.
- [12] J. H. Hubbard, D. Schleicher, S. Sutherland, “How to find all roots of complex polynomials by Newton’s method,” *Invent. Math.* **146** (2001), 1–33.
- [13] M. Shub and S. Smale, “Complexity of Bézout’s theorem I: Geometric aspects,” *J. Amer. Math. Soc.* **6** (1993), 459–501.
- [14] I. Besevic, *Universitas Pandrosion: Part VII — Breaking McMullen’s Barrier and the Resolution of Smale’s 17th Problem*. Preprint, 2026.
- [15] I. Besevic, *Universitas Pandrosion: Part VIII — The Geometric-Decreasing Start Strategy*. Preprint, 2026.
- [16] I. Besevic, *Universitas Pandrosion: Parts II–VI on Smale’s Mean Value Conjecture (1981)*. Preprint series, 2026.
- [17] S. Smale, “The fundamental theorem of algebra and complexity theory,” *Bull. Amer. Math. Soc.* **4** (1981), 1–36. (*The Mean Value Conjecture, often abbreviated MVC, originates here.*)
- [18] S. Smale, “Mathematical Problems for the Next Century,” *The Mathematical Intelligencer*, vol. 20, no. 2, pp. 7–15, 1998. (*Problem 17 originates here, on the average-case complexity of finding zeros of polynomial systems.*)

- [19] P. Bürgisser and F. Cucker, *On a problem posed by Steve Smale*. Annals of Mathematics **174** (2011), 1785–1836.
- [20] M. Shub and S. Smale, “Complexity of Bézout’s theorem II–V,” SIAM J. Comput. **22** (1993), Computational Algebraic Geometry, J. Complexity **9** (1993), J. Amer. Math. Soc. **6** (1993), Theoretical Computer Science **133** (1994).