11/12/2019 Project 5

Ivana Zdravevska

MATH333-017 Probability and Statistics

Prof. Matthew Michal


Project 5

```
In [47]: from numpy import mean
    from scipy.stats import norm
    import matplotlib.pyplot as plt
    import statistics

array1_norm = norm.rvs(loc=58,scale=10, size=100)
    mean_norm = array1_norm.mean()
    print("The mean of a normal distribution array with a mean of 58, standa
    rd deviation of 10, and a size of 100 is " + str(mean_norm))

means = [mean(norm.rvs(loc=58,scale=10, size=100)) for _i in range(1000
)]
    plt.hist(means)
    plt.show()
    print('The mean of the sample means is {}'.format(mean(means)))
    print('The standard deviation of the sample means is {}'.format(statistics.stdev(means)))
```

The mean of a normal distribution array with a mean of 58, standard deviation of 10, and a size of 100 is 57.693801920609765



The mean of the sample means is 57.953916060320765

The standard deviation of the sample means is 0.9803550195810288


11/12/2019 Project 5

```
In [10]: from numpy import mean
    from scipy.stats import geom
    import matplotlib.pyplot as plt
    import statistics

arrayl_geom = geom.rvs(p=0.3, size=100)
    mean_geom = arrayl_geom.mean()
    print("The mean of a geometric distribution array with a probability of
        0.3 and a size of 100 is " + str(mean_geom))

means2 = [mean(geom.rvs(p=0.3, size=100)) for _i in range(1000)]
    plt.hist(means2)
    plt.show()
    print('The mean of the sample means is {}'.format(mean(means2)))
    print('The standard deviation of the sample means is {}'.format(statistics.stdev(means2)))
```

The mean of a geometric distribution array with a probability of 0.3 and a size of 100 is 3.92



The mean of the sample means is 3.32452
The standard deviation of the sample means is 0.2767134617974701

When calculating the theoretical standard deviation of a sample, we should use 1000 for the value of n. Since the variance of a sample mean is σ^2 , the standard deviation is σ and increases, the variance shrinks. Therefore, as the size increases we get more precise estimates of the group means. The standard error decreases leading to an increase in higher accuracy. The smaller the standard error, the more powerful the statistical test is since it brings us closer to the true value and away from an error. With n being bigger, the variance shrinks which means that the measure of the spread in the sample also decreases giving us data that is less spread out and more accurate.