

## Solutions to Question Sheet 3

MATH40003 Linear Algebra and Groups

Term 2, 2019/20

Problem sheet released on Wednesday of week 4. All questions can be attempted before the problem class on Monday Week 5. Question 3 is suitable for tutorials. Solutions will be released on Wednesday of week 5.

**Question 1** For each of the following matrices  $A \in M_3(\mathbb{R})$ , find the eigenvalues and eigenvectors. Then diagonalise  $A$ , or prove it cannot be diagonalised.

$$(i) \begin{pmatrix} -1 & -2 \\ 4 & 5 \end{pmatrix} \quad (ii) \begin{pmatrix} 1 & 1 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{pmatrix} \quad (iii) \begin{pmatrix} 1 & 2 & 2 \\ 1 & 2 & -1 \\ -1 & 1 & 4 \end{pmatrix} \quad (iv) \begin{pmatrix} 2 & 1 & 0 \\ -1 & 0 & 1 \\ 1 & 3 & 1 \end{pmatrix}.$$

**Solution:** (i) Eigenvalues 1, 3 with eigenvectors  $a \begin{pmatrix} 1 \\ -1 \end{pmatrix}$  and  $b \begin{pmatrix} 1 \\ -2 \end{pmatrix}$  respectively, for any non-zero real numbers  $a, b$ . So if we set (for instance)  $P = \begin{pmatrix} 1 & 1 \\ -1 & -2 \end{pmatrix}$  then  $P^{-1}AP = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}$ .

(ii) Eigenvalues 1, 2, 3 with eigenvectors  $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ . So setting  $P = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$  gives  $P^{-1}AP = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}$ .

(iii) Characteristic polynomial is  $(x-1)(x-3)^2$ , so eigenvalues are 1, 3. For  $\lambda = 1$ , eigenvectors are scalar multiples of  $\begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$ . For  $\lambda = 3$  eigenvectors are  $\begin{pmatrix} a+b \\ a \\ b \end{pmatrix}$  for any  $a, b$  (not both zero). So taking  $P = \begin{pmatrix} 2 & 1 & 1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$  for instance gives  $P^{-1}AP = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix}$ .

(iv) Eigenvalues 2, -1. For  $\lambda = 2$  eigenvectors are multiples of  $(1, 0, 1)^T$ ; for  $\lambda = -1$  eigenvectors are multiples of  $(1, -3, 4)^T$ . So there is no basis of  $\mathbb{R}^3$  consisting of eigenvectors and therefore the matrix is not diagonalisable.

**Question 2** For which values of  $c$  is the matrix  $\begin{pmatrix} 1-2c & 4c & -c \\ -c & 2c+1 & -c \\ 0 & 0 & -1 \end{pmatrix} \in M_3(\mathbb{R})$  diagonalisable?

**Solution:** The characteristic polynomial is  $(1-x)^2(1+x)$ , so eigenvalues are  $1, -1$  with  $1$  repeated. For  $c \neq 0$ , the eigenvectors for  $\lambda = 1$  are scalar multiples of  $(2, 1, 0)^T$ , so we cannot form a  $3 \times 3$  matrix  $P$  with linearly independent eigenvectors as its columns. But for  $c = 0$  the eigenvectors for  $\lambda = 1$  are  $(a, b, 0)^T$ , so we can find an invertible  $P$  such as  $\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$  such that  $P^{-1}AP$  is  $\text{diag}(-1, 1, 1)$ . So the matrix is diagonalisable if and only if  $c = 0$ .

**Question 3** Let  $A = \begin{pmatrix} -10 & -18 \\ 9 & 17 \end{pmatrix} \in M_2(\mathbb{R})$ .

- (a) Find an invertible  $2 \times 2$  matrix  $P$  such that  $P^{-1}AP$  is diagonal.
- (b) Find  $A^n$ , where  $n$  is an arbitrary positive integer.
- (c) Find a matrix  $B \in M_2(\mathbb{R})$  such that  $B^3 = A$ .
- (d) Find a matrix  $C \in M_2(\mathbb{C})$  such that  $C^2 = A$ .
- (e) Prove that there is no  $C \in M_2(\mathbb{R})$  such that  $C^2 = A$ .

**Solution:** (a) Solving for the eigenvalues  $-1, 8$ , let  $P$  be the matrix with columns the eigenvectors, i.e.  $P = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$ . Then  $P^{-1}AP = D = \begin{pmatrix} -1 & 0 \\ 0 & 8 \end{pmatrix}$ .

(b) As seen in lectures,  $(P^{-1}AP)^n = P^{-1}A^nP$ , hence  $P^{-1}A^nP = D^n$ , giving  $A^n = PD^nP^{-1} = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} (-1)^n & 0 \\ 0 & 8^n \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ . This works out as  $\begin{pmatrix} 2 \cdot (-1)^n - 8^n & 2 \cdot (-1)^n - 2 \cdot 8^n \\ (-1)^{n+1} + 8^n & (-1)^{n+1} + 2 \cdot 8^n \end{pmatrix}$ .

(c) If  $E = \begin{pmatrix} -1 & 0 \\ 0 & 2 \end{pmatrix}$  then  $E^3 = D$ , so  $(PEP^{-1})^3 = PE^3P^{-1} = PDP^{-1} = A$ . So take  $B = PEP^{-1} = \begin{pmatrix} -4 & -6 \\ 3 & 5 \end{pmatrix}$ .

(d) If  $F = \begin{pmatrix} i & 0 \\ 0 & \sqrt{8} \end{pmatrix}$  then  $F^2 = D$ , so  $C = PFP^{-1} = \begin{pmatrix} -\sqrt{8} + 2i & -2\sqrt{8} + 2i \\ \sqrt{8} - i & 2\sqrt{8} - i \end{pmatrix}$  satisfies  $C^2 = A$  just as in (c).

(e) Suppose  $C^2 = A$  with all entries of  $C$  real. Then  $\det(C)^2 = \det(C^2) = \det(A) = -8$ . This is impossible as  $\det(C)$  is real.

**Question 4** Suppose  $V$  is a vector space over a field  $F$  and  $T : V \rightarrow V$  is linear. If  $\lambda \in F$ , let  $E_\lambda = \{v \in V : T(v) = \lambda v\}$ . Prove that this is a subspace of  $V$  and  $\lambda$  is an eigenvalue of  $T$  if and only if  $E_\lambda \neq \{0\}$ .

**Solution:** Either use the test for a subspace, or note that  $E_\lambda$  is the kernel of the linear map  $(T - \lambda Id) : V \rightarrow V$ , and is therefore a subspace.

**Question 5** For each of the linear maps  $\theta_i$  below, write down the matrix representing  $\theta_i$  with respect to the standard basis. Hence find the eigenvalues of  $\theta_i$  and for each eigenvalue  $\lambda$ , find the eigenspace  $E_\lambda$ . Determine whether  $\theta_i$  is diagonalizable.

- i)  $\theta_1 : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  given by

$$\theta_1 : \begin{pmatrix} a \\ b \\ c \end{pmatrix} \mapsto \begin{pmatrix} c - b \\ a - c \\ c \end{pmatrix}.$$

ii)  $\theta_2 : \mathbb{C}^3 \rightarrow \mathbb{C}^3$  given by

$$\theta_2 : \begin{pmatrix} a \\ b \\ c \end{pmatrix} \mapsto \begin{pmatrix} c-b \\ a-c \\ c \end{pmatrix}.$$

**Solution:**

i) The characteristic polynomial is

$$\det \begin{pmatrix} x & 1 & -1 \\ -1 & x & 1 \\ 0 & 0 & x-1 \end{pmatrix} = \dots = (x-1)(x^2+1)$$

So the only eigenvalue is  $\lambda = 1$ .

$$E_1 = \left\{ \begin{pmatrix} a \\ b \\ c \end{pmatrix} : \begin{pmatrix} c-b \\ a-c \\ c \end{pmatrix} = \begin{pmatrix} a \\ b \\ c \end{pmatrix} \right\} = \left\{ \begin{pmatrix} a \\ b \\ c \end{pmatrix} : b=0, a=c \right\}.$$

So this is a 1-dimensional space, with basis  $\{(1, 0, 1)^T\}$ . The only eigenvectors of  $\theta$  are multiples of  $(1, 0, 1)^T$ . The map is not diagonalizable.

ii) The characteristic polynomial is again  $x(x^2+1)$ , but this time the roots are  $1, \pm i$ . Since there are 3 distinct eigenvalues, the map is diagonalizable. The eigenspaces are given by

$$E_1 = \left\{ \begin{pmatrix} a \\ 0 \\ a \end{pmatrix} : a \in \mathbb{C} \right\}, \quad E_i = \left\{ \begin{pmatrix} ib \\ b \\ 0 \end{pmatrix} : b \in \mathbb{C} \right\}, \quad E_{-i} = \left\{ \begin{pmatrix} a \\ ia \\ 0 \end{pmatrix} : a \in \mathbb{C} \right\}.$$

A basis of eigenvectors would therefore be  $(1, 0, 1)^T, (i, 1, 0)^T, (1, i, 0)^T$ .

**Question 6** For each of the linear maps  $T$  in Question 2 of Sheet 2, compute the eigenvalues and eigenvectors of  $T$  and determine whether or not  $T$  is diagonalisable.

**Solution:** (i) The matrix of  $T$  with respect to the standard basis is  $\begin{pmatrix} -1 & 1 & -1 \\ 0 & -4 & 6 \\ 0 & -3 & 5 \end{pmatrix}$ .

So the char poly is  $(x+1)^2(x-2)$ . The eigenvalues are  $-1, 2$ . The eigenspace  $E_{-1}$  is spanned by  $(-1, 2, 1)$ ; the eigenspace  $E_2$  is spanned by  $(0, 1, 1)$ . There is no basis of eigenvectors, so  $T$  is not diagonalisable.

(ii) Matrix of  $T$  w.r.t. basis  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$  is  $A = \begin{pmatrix} 1 & -2 & 0 & 0 \\ 1 & 4 & 0 & 0 \\ 0 & 0 & 1 & -2 \\ 0 & 0 & 1 & 4 \end{pmatrix}$ .

So the char poly is  $(x-2)^2(x-3)^2$ , so the eigenvalues are  $2, 3$ . The eigenspace  $E_2$  of  $A$  is spanned by  $(2, -1, 0, 0)^T$  and  $(0, 0, 2, -1)^T$ . So the eigenspace  $E_2$  of  $T$  is spanned by  $\begin{pmatrix} 2 & 0 \\ -1 & 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 & 2 \\ 0 & -1 \end{pmatrix}$ . Likewise the eigenspace  $E_3$  of  $T$  is spanned by  $\begin{pmatrix} 1 & 0 \\ -1 & 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 & 1 \\ 0 & -1 \end{pmatrix}$ . So  $T$  is diagonalisable.

(iii)  $T$  sends  $1 \mapsto 0$ ,  $x \mapsto 3x$ ,  $x^2 \mapsto x + 6x^2$ , so matrix of  $T$  wrt basis  $1, x, x^2$  is  $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 3 & 1 \\ 0 & 0 & 6 \end{pmatrix}$ . The eigenvalues are 0, 3, 6. Corresponding eigenvectors are 1 (the constant function),  $x$  and  $x + 3x^2$  (and their non-zero scalar multiples).  $T$  is diagonalisable.

**Question 7** As in Question 9 of Sheet 1, let  $A$  be the  $n \times n$  matrix

$$A = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & -a_0 \\ 1 & 0 & 0 & \cdots & 0 & -a_1 \\ 0 & 1 & 0 & \cdots & 0 & -a_2 \\ & & & \cdots & & \\ 0 & 0 & 0 & \cdots & 1 & -a_{n-1} \end{pmatrix}$$

where the  $a_i$  are in the field  $F$ . Let  $e_1, \dots, e_n$  be the standard basis of  $F^n$ .

- i) Prove that  $F^n$  is spanned by the vectors  $e_1, Ae_1, \dots, A^{n-1}e_1$ . What is  $A^n e_1$  as a linear combination of these?
- ii) Show that for every  $v \in F^n$  there is a polynomial  $q(x)$  (over  $F$ ) of degree at most  $n - 1$  such that  $v = q(A)e_1$  (where  $q(A)$  is the result of substituting  $A$  for  $x$  into the polynomial  $q$ ).
- iii) Deduce that  $\chi_A(A)$  is the zero matrix (this is a special case of the Cayley - Hamilton Theorem).

**Solution:** (i) Note that as the columns of  $A$  are the images of the standard basis vectors,  $Ae_i = e_{i+1}$  for  $1 \leq i < n$ . Thus  $A^i e_1 = e_{i+1}$  for  $1 \leq i < n$ . We also have

$$A^n e_1 = A(A^{n-1} e_1) = Ae_n = -a_0 e_1 - a_1 e_2 - \dots - a_{n-1} e_n = -a_0 e_1 - a_1 Ae_1 - \dots - a_{n-1} A^{n-1} e_1.$$

(ii) By (i) each  $e_i$  is in (the span of)  $e_1, Ae_1, \dots, A^{n-1} e_1$ , so the span of these is the whole of  $F^n$ . So if  $v \in F^n$  there are  $b_0, \dots, b_{n-1} \in F$  with

$$v = b_0 e_1 + b_1 Ae_1 + \dots + b_{n-1} A^{n-1} e_1 = (b_0 I_n + b_1 A + \dots + b_{n-1} A^{n-1}) e_1.$$

The result follows.

(iii) By (i) we have  $p(A)e_1 = 0$ , where  $p(x) = a_0 + a_1 x + \dots + a_{n-1} x^{n-1} + x^n$ . Let  $v, q$  be as in (ii). Then  $p(A)v = p(A)q(A)e_1 = q(A)p(A)e_1 = 0$  (as  $A$  commutes with powers of itself). So  $p(A) = 0$ . But by Qu 9, Sheet 1,  $\chi_A(x) = p(x)$ .

**Question 8** In this question you can use Q7. Unless stated otherwise, you can choose which field to use.

- (a) Find a  $3 \times 3$  matrix which has characteristic polynomial  $x^3 - 7x^2 + 2x - 3$ .
- (b) Find a  $3 \times 3$  matrix  $A$  such that  $A^3 - 2A^2 = I_3$ .
- (c) Find a  $4 \times 4$  invertible matrix  $B$  such that  $B^{-1} = B^3 + I_4$ .
- (d) Find a  $5 \times 5$  invertible matrix  $B$  such that  $B^{-1} = B^3 + I_5$ .
- (e) Find a real  $4 \times 4$  matrix  $C$  such that  $C^2 + C + I_4 = 0$ .
- (f) For each  $n \geq 2$  find an  $n \times n$  matrix  $D$  such that  $C^n = I_n$  but  $C \neq I_n$ .

**Solution:** (a)  $\begin{pmatrix} 0 & 0 & 3 \\ 1 & 0 & -2 \\ 0 & 1 & 7 \end{pmatrix}$  works (by Q9, Sh 1)

(b) Take  $A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 2 \end{pmatrix}$ . This works, by Qu8.

(c) Multiplying through by  $B$ , the equation is  $B^4 + B - I = 0$ . So we can use Qu8 to find such a matrix  $4 \times 4$  matrix  $B$ . Note that as the constant term of the char poly is non-zero,  $B$  is indeed invertible.

(d) Take  $B_0$  as in (c), and let  $B = \begin{pmatrix} B_0 & 0 \\ 0 & \lambda \end{pmatrix}$ , where  $\lambda$  is a complex root of  $x^4 + x - 1$ .

(e) By Qu7 the  $2 \times 2$  matrix  $A = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}$  satisfies  $A^2 + A + I = 0$ . So take  $C = \begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix}$ .

(f) Use Qu7 to get a non-identity  $n \times n$  matrix with char poly  $x^n - 1$ .