

Factorization systems

Ivan Kobe

November 20, 2024

A factorization system in a category \mathcal{C} consists of two classes of morphisms (L, R) , such that both L and R contain all isomorphisms and are closed under composition, and every morphism $f : C \rightarrow D$ in \mathcal{C} admits a factorization into a morphism $l \in L$ followed by a morphism $r' \in R$, which is unique up to unique isomorphism among such factorizations.

$$\begin{array}{ccccc}
 & & E & & \\
 & \nearrow l & & \nwarrow r & \\
 C & & & & D \\
 & \searrow l' & & \nearrow r' & \\
 & & E' & &
 \end{array}$$

$i \cong$

Plan of work:

- show that factorization systems descend to slices
- construct a few elementary examples (in the **Set**, **Grp**, **CRing** etc.)
- show that the left class in a f.s. has the right cancellation property, and, dually, the right class has the left cancellation property
- define the orthogonality relation and show various closure properties of orthogonal complements (under (co)limits, (co)base change, etc.)
- either of the classes in a f.s. determines the other via orthogonal complements
- recognition principle for factorization systems: (L, R) is a factorization system iff every morphism $f : C \rightarrow D$ in \mathcal{C} admits an (apriori non-unique) (L, R) -factorization, both classes are replete, and $L \perp R$

If time permits it, we would also like to use the general theory of factorization systems to define (left-exact) modalities and study some of their properties. In particular, we would like to show that every (left-exact) modality determines a (left-exact) reflective subcategory and vice versa.