

Discrete Random Variables

David Armstrong

UCI

Discrete Random Variables

- *Variable*: A quantity that may take different values.
- *Random variable*: A variable that may assume different values with certain probabilities.
 - One way to think of it as a function that assigns a real number to each outcome in the sample space.
 - A *discrete* random variable is one who can only be discrete values.
 - For now, we will focus on bounded discrete random variables.
- Example of a discrete random variable:
 - Flip a coin 3 times.
Then $S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$.
 - Let X be a random variable that is equal to the number of heads in 3 flips of a coin.
 - X can be 0,1,2, or 3.

The Support

- Denote the *support* of X as \mathbb{S}_X .
- The *support* of X is the space of values which X has a positive probability of occurring.
 - Notationally, $X : S \rightarrow \mathbb{S}_X$
- An example is flipping a coin once.
- $S = \{T, H\}$.
 - If we let X be equal to the number of heads (this is the same as setting heads to equal 1 and tails to equal 0).
 - Then $\mathbb{S}_X = \{0, 1\}$.
- In the example on the previous slide with 3 coin flips, $\mathbb{S}_X = \{0, 1, 2, 3\}$.
- The concept of the support of a random variable is an important one.
- Once the appropriate random variable is specified, we can focus only on the support of it as opposed to the entire sample space.
- Example: If we flip a coin 100 times and interested in the number of heads seen.
 - What does X represent?
 - How many elements are in the sample space?
 - What is the support of the random variable X ?

Example of the Support

Assume we flip a coin 3 times and let X be the number of heads.

- What does $X = 0$ represent?
- What is the probability all three of the flips land in tails?
- What is the probability that $X = 1$
- What is the probability that $X = 2$
- What is the probability that $X = 3$
- Is this a valid distribution?

Probability Mass Function (pdf)

The *probability distribution* of X assigns a number to all values x in \mathbb{S}_X such that:

- $0 \leq P(X = x) \leq 1$
- $\sum_{x \in \mathbb{S}_x} P(X = x) = 1$

Notationally we state $f(x) = P(X = x)$.

With discrete random variables, $f(x)$ is termed the *probability mass function (p.m.f.)*.

- From here on out, we will refer to X as the random variable.
- We will denote x as the values that X can be.
- For example flipping a coin 3 times, and setting X to be the number of heads.
 - X is the random variable.
 - X can be set equal to x where $x = 0, 1, 2$, or 3 .

Cumulative Distribution Function (cdf)

- With the p.m.f. of a discrete random variable, we can compute quantities such as $P(X < a)$ or $P(a \leq X < b)$ for some set constants of a and b .
- The *cumulative distribution function* (cdf) of a random variable at value X is $P(X \leq x)$.
- Notationally this is $F(x) = P(X \leq x)$
 - It is the sum of all probabilities which have $X \leq x$.

Properties of discrete random variables

Let a and b be a values of x

- $P(X \leq a) = 1 - P(X > a)$.
 - $P(X > a) = P(X \geq a + 1)$
 - $P(X < a) = P(X \leq a - 1)$.
 - Something that looks counter intuitive, but holds true for discrete distributions.
 - $P(a \leq X \leq b) = P(X \leq b) - P(X < a)$.
-
- $P(a < X \leq b) = P(X \leq b) - P(X \leq a)$.

Expectation - Discrete Random Variables

The p.m.f. completely determines the probability distribution of a discrete random variable.

- The *expectation* of X can be viewed as the mean or average of X .
- Within a frequentist framework, it can be seen as the average of X across many trials of the experiment.
- The *expectation* of X is denoted as $E(X)$.
- $E(X) = \sum_{x \in \mathbb{S}_X} xP(X = x) = \sum_{x \in \mathbb{S}_X} xf(x)$.
- Can be viewed as averaging over all possible X values while weighting each possible value by its probability.

Properties of expectations.

- If a and b are constants and X is a random variables, then:

$$E(a + bX) = E(a) + E(bX) = a + bE(X).$$

- If X and Y are random variables, then $E(X + Y) = E(X) + E(Y)$.
 - As a result if a and b are constants, then $E(aX + bY) = aE(X) + bE(Y)$
 - As a further result, let X_i be random variables and a_i 's be constants.
 - $E\left(\sum_i a_i X_i\right) = \sum_i a_i E(X_i)$

Expectation - Discrete Random Variables

Functions of X are also random variables.

- We can set $h(X)$ to be a function of X .
 - Example: $h(X) = X^2$.
 - In the number of heads in 3 coin flips, $\mathbb{S}_X = \{0, 1, 2, 3\}$
 - The support of X^2 will be $\mathbb{S}_{X^2} = \{0, 1, 4, 9\}$
- We can take expectation of these functions without having to first find the distribution of $h(X)$ first.
 - $$E(h(X)) = \sum_{x \in \mathbb{S}_X} h(x)P(X = x) = \sum_{x \in \mathbb{S}_X} h(x)f(x).$$
- Just like with X , $E(h(X))$ can be viewed as averaging over all possible $h(X)$ values while weighting each possible value by the probability of X .

Variance - Discrete Random Variables

Now we come to another quantity that describes the probability distribution of X , known as the *variance*.

- The *variance* of X is defined to be the average squared deviation from the mean.
 - $\text{Var}(X) = E[(X - E(X))^2] = E[(X - \mu)^2]$.
 - $\text{Var}(X) = E(X^2) - [E(X)]^2$
 - We denote the variance of X as σ^2 .
 - Since it is the expected value of a squared random variable, $\sigma^2 > 0$.
 - $\sigma = \sqrt{\text{Var}(X)}$ is known as the *standard deviation* of X .
- σ - The typical distance of the datapoints to the mean.

Properties

- If c is a constant, then $\text{Var}(c) = 0$
- If c is a constant, then $\text{Var}(cX) = c^2 \text{Var}(X)$

If X and Y are independent random variables.

- $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$

As a result, if X_i 's are independent random variables and c_i 's are constants, then:

- $\text{Var}(\sum_i c_i X_i) = \sum_i c_i^2 \text{Var}(X_i)$

Discrete Random Variables: Distribution

Returning to the example of flipping a coin 3 times. A distribution table for X , the number of heads, can be constructed below.

	$x = 0$	$x = 1$	$x = 2$	$x = 3$	Total
$P(X = x)$					

- Calculate $f(2)$
- Calculate $F(2)$.
- What is the expectation of X ?

$$E(X) = \sum_{x \in \mathbb{S}_X} xf(x)$$

- What is the expectation of $5 - 3X$?

Discrete Random Variables: Distribution

Returning to the example of flipping a coin 3 times. A distribution table for X , the number of heads, can be constructed below.

	$x = 0$	$x = 1$	$x = 2$	$x = 3$	Total
$P(X = x)$					

- Let $h(X) = X^2$. What is the expectation of $h(x)$?

- Calculate the Variance of X

- What is the Variance of $5 - 3X$?

- What is the standard deviation of X ?

Discrete Random Variables: Classes

Example: Say X is the number of days a student is registered to take classes at University of California Irvine.

x	1	2	3	4	5
$f(x)$	0.10	0.30	0.25	0.25	0.10

- What is the expectation of X ?
- What is the expectation of $5 + 3X$?
- Let $h(X) = X^2$. What is the expectation of $h(x)$?
- What is the Variance of X ?