

point estimate = \bar{x} or \hat{p} , critical value: z-score, standard error: $\frac{\sigma}{\sqrt{n}}$, $\frac{s}{\sqrt{n}} = \frac{1 - \frac{0.01}{2}}{\frac{0.01}{2}}$, sample size = $n = (\frac{z \cdot \sigma}{m})^2$, margin of error = (critical value) * (standard error)

C interval: $\bar{x} - m < \mu < \bar{x} + m$
 $\hat{p} - m < p < \hat{p} + m$

construct confidence interval

known θ z-method: (Z interval) $z_{\alpha/2} = \text{invNorm}(\text{C-level}, 0, 1, \text{center})$ $m = z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}$	unknown θ T-method: (T interval) $t_{\alpha/2} = \text{invT}(\frac{1 - \frac{\alpha}{2}}{2}, n - 1)$ $m = t_{\alpha/2} \cdot \frac{s}{\sqrt{n}}$	C-interval for proportion: (1 prop Z Int) $\hat{p} = \frac{x}{n}$ $z_{\alpha/2} = \text{invNorm}(\text{C-level}, 0, 1, \text{center})$ $m = z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$ $n = \hat{p}(1 - \hat{p}) \left(\frac{z_{\alpha/2}}{m}\right)^2$ or $0.25 \left(\frac{z_{\alpha/2}}{m}\right)^2$
critical values using chi-square: $\chi^2_{1 - \frac{\alpha}{2}} = (1 - \frac{\alpha}{2}, \text{df})$ $\chi^2_{\frac{\alpha}{2}} = (\frac{\alpha}{2}, \text{df})$	C-interval for population θ is $\sqrt{\frac{(n-1)S^2}{\chi^2_{\alpha/2}}} < \theta < \sqrt{\frac{(n-1)S^2}{\chi^2_{1 - \alpha/2}}}$	-

null hypothesis: $H_0: \mu = \mu_0$

If $p < \alpha$, reject H_0 . Not enough evidence

alternate hypothesis: $H_1: \mu < \mu_0, \mu > \mu_0, \mu \neq \mu_0$

If $p > \alpha$, do not reject H_0 . Enough evidence

level of significance: $\alpha = 0.05$ (if not mentioned)

smaller p is, stronger against H_0

Type I error: reject H_0 when H_0 is true ($p > \alpha$)

Type II error: do not reject H_0 when H_0 is false ($p < \alpha$)

Hypothesis test

known θ z-test: Test statistic: $z = \frac{\bar{x} - \mu_0}{\frac{\sigma}{\sqrt{n}}}$ p-value: $p = \text{normalcdf}(\frac{-\infty}{z}, \frac{z}{\infty}, 0, 1)$ left-tailed $p = \text{normalcdf}(\frac{z}{\infty}, \frac{\infty}{z}, 0, 1)$ right-tailed $p = 2 \cdot \text{normalcdf}(\frac{z}{\infty}, \frac{\infty}{z}, 0, 1)$ two-tailed	unknown θ T-test: Test statistic: $t = \frac{\bar{x} - \mu_0}{\frac{s}{\sqrt{n}}}$ p-value: $p = \text{tcdf}(\frac{-\infty}{t}, \frac{t}{\infty}, 0, n - 1)$ left-tailed $p = \text{tcdf}(\frac{t}{\infty}, \frac{\infty}{t}, 0, n - 1)$ right-tailed $p = 2 \cdot \text{tcdf}(\frac{t}{\infty}, \frac{\infty}{t}, 0, n - 1)$ two-tailed	Hypothesis test for proportion $H_0: p = p_0$ $H_1: p < p_0, p > p_0, p \neq p_0$ 1 prop Z-test Test statistic: $z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1 - p_0)}{n}}}$ p-value: $p = \text{normalcdf}(\frac{-\infty}{z}, \frac{z}{\infty}, 0, 1)$ left-tailed $p = \text{normalcdf}(\frac{z}{\infty}, \frac{\infty}{z}, 0, 1)$ right-tailed $p = 2 \cdot \text{normalcdf}(\frac{z}{\infty}, \frac{\infty}{z}, 0, 1)$ two-tailed
---	---	--

$H_0: \mu_1 = \mu_2$

$H_1: \mu_1 < \mu_2, \mu_1 > \mu_2, \mu_1 \neq \mu_2$

standard error of $\bar{x}_1 - \bar{x}_2 = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$

degree of freedom: smaller of $n_1 - 1$ and $n_2 - 1$

Two means: Independent samples

z-sampTTest:

Test statistic: $t = \frac{(\bar{x}_1 - \bar{x}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$

p-value: $p = \text{tcdf}(\frac{-\infty}{z}, \frac{z}{\infty}, 0, \text{df})$ left-tailed
 $p = \text{tcdf}(\frac{z}{\infty}, \frac{\infty}{z}, 0, \text{df})$ right-tailed
 $p = 2 \cdot \text{tcdf}(\frac{z}{\infty}, \frac{\infty}{z}, 0, \text{df})$ two-tailed

$H_0: p_1 = p_2$

$H_1: p_1 < p_2, p_1 > p_2, p_1 \neq p_2$

mean = $p_1 - p_2$, standard deviation = $\sqrt{\frac{p_1(1 - p_1)}{n_1} + \frac{p_2(1 - p_2)}{n_2}}$

$\hat{p} = \frac{x_1 + x_2}{n_1 + n_2}$, standard error = $\sqrt{\frac{\hat{p}(1 - \hat{p})}{n_1} + \frac{\hat{p}(1 - \hat{p})}{n_2}} = \sqrt{\hat{p}(1 - \hat{p}) \left(\frac{1}{n_1} + \frac{1}{n_2}\right)}$

Two proportions

z-propZTest:

Test statistic: $z = \frac{(\hat{p}_1 - \hat{p}_2)}{\sqrt{\hat{p}(1 - \hat{p}) \left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$

p-value: $p = \text{normalcdf}(\frac{-\infty}{z}, \frac{z}{\infty}, 0, 1)$ left-tailed
 $p = \text{normalcdf}(\frac{z}{\infty}, \frac{\infty}{z}, 0, 1)$ right-tailed

matched pairs: dependent samples

$d = \bar{x}_1 - \bar{x}_2$

\bar{d} = mean of d

$H_0: \mu_d = 0$

$H_1: \mu_d < 0, \mu_d > 0, \mu_d \neq 0$

Two means: paired samples

T-Test:

Test statistic: $t = \frac{\bar{d} - \mu_0}{\left(\frac{s_d}{\sqrt{n_d}}\right)}$

p-value: $p = \text{tcdf}(\frac{-\infty}{z}, \frac{z}{\infty}, 0, n_d - 1)$

Assumptions: SRS and $n > 30$ or normally distributed

Assumptions for proportion: SRS, population $\geq 20 \cdot n$, categories = 2
 and each categories > 10