

## Lecture 2

### 0) Introduction

1) Formal quantizations

2) Microlocalization.

### 0) Introduction

Let  $A$  be  $\mathbb{Z}$ -graded Poisson algebra (where, for simplicity,  $\deg \{ \cdot, \cdot \} = -1$ ),  $A = \bigoplus_{i \in \mathbb{Z}} A_i$ .

We have defined the notion of a filtered quantization,  $\mathcal{A}$  of  $A$ . This is a filtered associative algebra  $\mathcal{A} = \bigcup_{i \in \mathbb{Z}} \mathcal{A}_{\leq i}$  w.

$[\mathcal{A}_{\leq i}, \mathcal{A}_{\leq j}] \subset \mathcal{A}_{\leq i+j-1}$  ( $\leadsto \text{gr } \mathcal{A}$  comes w.  $\{ \cdot, \cdot \}$  of  $\deg -1$ )

& a fixed graded Poisson algebra isomorphism  $\text{gr } \mathcal{A} \xrightarrow{\sim} A$ .

If  $\mathcal{A}_{\leq -1} \neq \{0\}$  we also need to impose the technical condition:  $\mathcal{A} \xrightarrow{\sim} \varprojlim_{i \rightarrow -\infty} \mathcal{A} / \mathcal{A}_{\leq i}$ .

In this lecture we'll define some kind of a localization of  $\mathcal{A}$  by an element of  $A$ . And this requires a connection with a related class of quantizations: formal ones.

## 1) Formal quantizations

Let  $A$  be a commutative algebra. By its **formal deformation** (over  $\mathbb{C}[[\hbar]]$ ) we mean an associative  $\mathbb{C}[[\hbar]]$ -algebra  $\mathcal{A}_\hbar$  together w. an algebra isomorphism  $\iota: \mathcal{A}_\hbar/(\hbar) \xrightarrow{\sim} A$  s.t.

(i)  $\hbar$  is not a zero divisor in  $\mathcal{A}_\hbar$

(ii)  $\mathcal{A}_\hbar$  is complete & separated in the  $\hbar$ -adic topology i.e.

$$\mathcal{A}_\hbar \xrightarrow{\sim} \varprojlim_n \mathcal{A}_\hbar/(\hbar^n).$$

Note that  $\mathcal{A}_\hbar/(\hbar)$  carries a Poisson bracket. Indeed, since  $\mathcal{A}_\hbar/(\hbar)$  is commutative,  $[a, b] \in \hbar \mathcal{A}_\hbar \forall a, b \in \mathcal{A}_\hbar$ . Thx to (i),  $\frac{1}{\hbar}[a, b] \in \mathcal{A}_\hbar$  is uniquely defined. We set

$$\{a + (\hbar), b + (\hbar)\} = \frac{1}{\hbar}[a, b] + (\hbar)$$

getting a Poisson bracket on  $\mathcal{A}_\hbar/(\hbar)$  (**exercise**). If  $A$  was equipped with a Poisson bracket &  $\iota$  is a Poisson isomorphism, then we say that  $\mathcal{A}_\hbar$  is a **formal quantization** of  $A$ .

### 1.1) From filtered quantizations to graded formal.

By a **grading** on  $\mathcal{A}_\hbar$  we mean a collection of gradings on each  $\mathcal{A}_\hbar/(\hbar^n)$  s.t.

- $\deg \hbar = 1$

- $\mathcal{A}_\hbar / (\hbar^{n+1}) \longrightarrow \mathcal{A}_\hbar / (\hbar^n)$  is graded  $\forall n$ .

( $\mathcal{A}_\hbar$  itself cannot be graded b/c of completeness)

Let's explain how to get from a filtered quantization to a graded formal quantization. We'll do this in two steps. Let  $\mathcal{A}$  be a filtered quantization of graded  $A$ .

1) From  $\mathcal{A} = \bigcup_{i \in \mathbb{Z}} \mathcal{A}_{\leq i}$  we form the **Rees algebra**

$$R_\hbar(\mathcal{A}) := \bigoplus_{i \in \mathbb{Z}} \mathcal{A}_{\leq i} \hbar^i \subset \mathcal{A}[\hbar^{\pm 1}], \text{ a graded subalgebra}$$

Note that  $R_\hbar(\mathcal{A})/(\hbar) \cong \text{gr } \mathcal{A}$ ,  $R_\hbar(\mathcal{A})/(\hbar-1) \xrightarrow{\sim} \mathcal{A}$  (exercise)

2) Set  $\mathcal{A}_\hbar := \varprojlim_{n \rightarrow \infty} R_\hbar(\mathcal{A})/(\hbar^n)$

**Exercise:** Prove that  $\mathcal{A}_\hbar$  is a graded formal quantization of  $A$ . Hint: prove first that  $\mathcal{A}_\hbar / (\hbar^n) \xrightarrow{\sim} R_\hbar(\mathcal{A})/(\hbar^n)$ .

## 1.2) From graded formal to filtered

Now let  $\mathcal{A}_\hbar$  be a graded formal quantization of  $A$ . We want to produce a filtered quantization. Recall that

$$\mathcal{A}_\hbar(n) := \mathcal{A}_\hbar / (\hbar^n) \text{ is graded: } \mathcal{A}_\hbar(n) = \bigoplus_{i \in \mathbb{Z}} \mathcal{A}_\hbar(n)_i. \text{ The projection}$$

$\mathcal{A}_{\hbar}^{(n+1)} \rightarrow \mathcal{A}_{\hbar}^{(n)}$  is a graded algebra homomorphism. Set

$$\mathcal{A}_{\hbar,i} := \varprojlim_{n \rightarrow \infty} \mathcal{A}_{\hbar}^{(n)}{}_i, \quad \mathcal{A}_{\hbar}^{\text{fin}} := \bigoplus_{i \in \mathbb{Z}} \mathcal{A}_{\hbar,i}.$$

**Exercise:** 1) Show that  $\mathcal{A}_{\hbar}^{\text{fin}}$  is a  $\mathbb{C}[\hbar]$ -subalgebra of  $\mathcal{A}_{\hbar}$  & a graded  $\mathbb{C}[\hbar]$ -algebra.

2) Show that  $\mathcal{A}_{\hbar}^{\text{fin}}/(\hbar) \xrightarrow{\sim} A$  &  $\mathcal{A}_{\hbar}^{\text{fin}}/(\hbar-1)$  is a filtered quantization of  $A$  (w.r.t. the filtration induced by the grading on  $\mathcal{A}_{\hbar}^{\text{fin}}$ ).

3) Show that the assignments  $\mathcal{A} \mapsto \varprojlim_{n \rightarrow \infty} R_{\hbar}(\mathcal{A})/(\hbar^n)$  &  $\mathcal{A}_{\hbar} \mapsto \mathcal{A}_{\hbar}^{\text{fin}}/(\hbar-1)$  are mutually inverse bijections between filtered & graded formal quantizations of  $A$ .

**Rem:**  $\mathcal{A}_{\hbar}^{\text{fin}}/(\hbar-1)$  is complete w.r.t. the filtration b/c  $\mathcal{A}_{\hbar}$  is complete in the  $\hbar$ -adic topology. This kind of explains that condition in the general definition of a filtered quantization.

## 2) Microlocalization.

Setting: Let  $A$  be a Poisson algebra &  $\mathcal{A}_{\hbar}$  be its formal

quantization. For  $f \in A$  we want to construct from  $\mathcal{H}_\hbar$  a formal quantization of  $A[f^{-1}]$ , a kind of localization, to be denoted by  $\mathcal{H}_\hbar[f^{-1}]$ .

Then we will explain a modification for filtered quantizations.

## 2.1) Localization in noncommutative rings

Let  $R$  be an associative (unital) ring &  $S \subset R$  be a subset closed under products. We want to have the localization  $R_S$  i.e. a ring with a homomorphism  $\eta: R \rightarrow R_S$  s.t.  $\forall$  ring  $R'$ :

$$\eta^*: \text{Hom}_{\text{rings}}(R_S, R') \xrightarrow{\sim} \{\varphi: R \rightarrow R' \mid \varphi(s) \text{ is invertible } \forall s \in S\}$$

Moreover, we want it to consist of "fractions"  $rs^{-1}$ . Sufficient conditions for the latter were found by Ore:

$$(01) \quad \forall a \in R, s \in S \Rightarrow \exists b \in R, t \in S \mid at = sb \Leftrightarrow s^{-1}a = bt^{-1}$$

(a.k.a. every left fraction is also a right action)

$$(02) \quad \forall a \in R, s \in S \text{ s.t. } sa = 0 \Rightarrow \exists t \in S \mid at = 0$$

Note that these conditions are vacuous if  $R$  is commutative.

Example/exercise: Suppose  $s \in R$  satisfies  $\text{ad}(s)^n = 0$  for some  $n \in \mathbb{Z}_{>0}$ , where  $\text{ad}(s): R \rightarrow R$  is given by  $r \mapsto [s, r]$ . Then  $\{s^m \mid m \in \mathbb{Z}_{\geq 0}\}$  satisfies (01) & (02).

Assume (01) & (02). Consider the following relation on  $R \times S$ :  $(a, s) \sim (a', s')$  if  $\exists t, t' \in S$  w.  $at = a't'$  &  $st = s't'$ .

It turns out that:

- this is an equivalence relation, let  $a/s$  denote the equivalence class of  $(a, s)$
- and the set of equivalence classes has a ring structure making it the localization  $R_S$ .

## 2.2) Construction of $\mathcal{A}_\hbar[f^{-1}]$

Recall that any formal quantization is the inverse limit of its quotients by  $(\hbar^n)$ . Set  $\mathcal{A}_\hbar(n) = \mathcal{A}_\hbar / (\hbar^n)$ . We are going to define an inverse system  $\mathcal{A}_\hbar(n)[f^{-1}]$  of flat  $\mathbb{C}[\hbar]/(\hbar^n)$ -algebras.

Let  $\hat{f}$  denote a lift of  $f$  to  $\mathcal{A}_\hbar(n)$ . Since  $[\mathcal{A}_\hbar(n), \mathcal{A}_\hbar(n)] \subset \hbar \mathcal{A}_\hbar(n)$ , we see that  $S := \{\hat{f}^m\}$  satisfies (01) & (02). Then we set  $\mathcal{A}_\hbar(n)[f^{-1}] := \mathcal{A}_\hbar(n)_S$ . Here are the properties:

1)  $\mathcal{A}_\hbar(n)[f^{-1}]$  is independent of the choice of  $\hat{f}$ :  
for a different lift  $\hat{f}' \exists n > 0$  s.t.

$$\hat{f}' \hat{f}^{n-1} = \hat{f}^n (1 + \hbar a) \text{ for some } a \in \mathcal{A}_\hbar(n)$$

2)  $\mathcal{A}_\hbar(n)[f^{-1}]$  is flat over  $\mathbb{C}[\hbar]/(\hbar^n)$

3) We have natural epimorphism

$$\mathcal{A}_\hbar(n+1)[f^{-1}] \twoheadrightarrow \mathcal{A}_\hbar(n)[f^{-1}]$$

The proofs are *exercises*. Now thx to these properties we can set  $\mathcal{A}_\hbar[f^{-1}] := \varprojlim_n \mathcal{A}_\hbar(n)[f^{-1}]$ .

*Exercise:*  $\mathcal{A}_\hbar[f^{-1}]$  is a formal quantization of  $A[f^{-1}]$ .

### 2.3) Filtered setting

One can ask when  $\mathcal{A}_\hbar[f^{-1}]$  has a grading. The answer: when  $f$  is homogeneous: each  $\mathcal{A}_\hbar(n)[f^{-1}]$  is graded & the homomorphism in 3) is graded. So using the correspondence

7

between filtered & graded formal quantizations in Sec 1 we define  $\mathcal{H}[f^{-1}]$ . Note that if  $\mathcal{H}$  is  $\mathbb{Z}_{\geq 0}$ -filtered and  $f \in \Lambda$ ; for  $i > 0$ , then  $\mathcal{H}[f^{-1}]$  is only  $\mathbb{Z}$ -filtered (in essentially all cases).