

Emergent Time from Quantum Variability

(Tiempo emergente de la variabilidad cuántica)

Iván Mauricio Burbano Aldana^{*}

Code: 201423205

Advised by: Dr. Andrés Fernando Reyes Lega^{**}

Departamento de Física, Universidad de los Andes, A.A.
4976-12340, Bogotá, Colombia

October 22, 2018

Abstract

Operator algebras, as general frameworks for the description of the observables of physical systems, have been an important tool towards rigorous investigations in information theory, statistical physics, and quantum field theory[1, 2, 3, 4]. In this project we will familiarize with these techniques and explore Connes' proposal for the emergence of time as a quantum phenomenon[5]. By building from [6], we will begin through the study of a generalization of the Radon-Nikodým theorem to von Neumann algebras[7]. This central result will yield a canonical dynamical mapping $\mathbb{R} \rightarrow \text{Out}(\mathcal{M})$ due to the noncommutativity of the von Neumann algebra \mathcal{M} . For type III von Neumann algebras, which are intimately related to the physics of systems with an infinite number of degrees of freedom[8], this prescription provides a class of dynamical evolutions which differ only locally from each other. It is in this sense that \mathcal{M} should be regarded by itself as a dynamical object. After understanding the theoretical and mathematical details of this construction, we will aim to present in great detail physically meaningful examples. We hope that this exercise will shed light into the physical meaning of Connes' mathematical proposal and guide research towards more concrete formulations.

Resumen

Las álgebras de operadores, como marcos generales para la descripción de observables en sistemas físicos, han sido una herramienta importante para investigaciones rigurosas en teoría de la información, física

^{*}im.burbano10@uniandes.edu.co

^{**}anreyes@uniandes.edu.co

estadística y teoría cuántica de campos[1, 2, 3, 4]. En este proyecto vamos a familiarizarnos con estas técnicas y exploraremos la propuesta de Connes sobre la emergencia del tiempo como un fenómeno cuántico[5]. Partiendo de [6], empezaremos estudiando una generalización del teorema de Radon-Nykodym para álgebras de von Neumann[7]. Este resultado central presentará un mapa dinámico canónico $\mathbb{R} \rightarrow \text{Out}(\mathcal{M})$ debido a la no-conmutatividad del álgebra de von Neumann \mathcal{M} . Para álgebras de von Neumann tipo III, las cuales están íntimamente relacionadas con la física de sistemas con un número infinito de grados de libertad[8], esta prescripción provee una clase de evoluciones dinámicas que solo difieren localmente entre si. Es en este sentido que \mathcal{M} debe de ser entendido como un objeto dinámico por si mismo. Después de entender los detalles teóricos y matemáticos de esta construcción, pretendemos mostrar en detalle ejemplos de importancia física. Esperamos que este ejercicio esclarezca el significado físico de la propuesta matemática de Connes y guíe la investigación hacia formulaciones más concretas.

1. State of the Art

2. Objectives

3. Theoretical Framework

In quantum mechanics observables are identified with operators on a separable Hilbert space \mathcal{H} . The spectrum of such an operator is then interpreted as the possible set of values the observation can yield. Therefore, the operators representing observables must be selfadjoint. Given that real observations are represented by concrete measurement devices whose range of outcomes is always bounded, we conclude that the observables must be in the bounded operators $\mathcal{B}(\mathcal{H})$ ¹. The following mathematical structures implement these requirements and are thus candidate structures for the algebra of observables of physical systems.

Definition 3.1 *A C^* -algebra \mathcal{A} is a Banach $*$ -algebra which satisfies the C^* -condition, that is, for every $a \in \mathcal{A}$ we have*

$$\|a^*a\| = \|a\|^2. \quad (1)$$

A von Neumann algebra (W^ -algebra for short) \mathcal{M} is a C^* -algebra for which there exists a predual, that is, a Banach space \mathcal{M}_* such that $\mathcal{M} = (\mathcal{M}_*)^*$ as Banach spaces.*

¹One can alternatively claim that, in light of the spectral theorem[9], every unbounded operator can be understood in terms of its spectral projections. These are in particular bounded operators. This point of view corresponds to adopting the position that every measurement can be ultimately broken down into a (possibly uncountably infinite) set of propositions.

On the other hand, physical states, which constitute the physical information required to evaluate the expected value of observables, can also be defined in this setting.

Definition 3.2 *Let \mathcal{A} be a C^* -algebra. A state on \mathcal{A} is a normalized positive linear functional $\omega : \mathcal{A} \rightarrow \mathbb{C}$. If \mathcal{M} is a von Neumann algebra, a state ω on \mathcal{M} is said to be normal if $\omega \in \mathcal{M}_*$ under the canonical embedding $\mathcal{M}_* \subseteq \mathcal{M}^*$.*

Note that in the context of von Neumann algebras and through the restriction to normal states, the structure of observables and states is complementary. This is evidence of the fact that each of them is meaningless without the other and, in fact, determines it. Finally, we can also consider dynamical evolutions in this framework. For definiteness we will work in the Heisenberg picture.

Definition 3.3 *A dynamical evolution on a C^* -algebra \mathcal{A} is a strongly continuous representation $\alpha : \mathbb{R} \rightarrow \text{Aut } \mathcal{A}$. A dynamical evolution on a W^* -algebra \mathcal{M} is a weakly continuous representation $\alpha : \mathbb{R} \rightarrow \text{Aut } \mathcal{M}$.*

The relationship between this framework and the one usually used in classical and quantum formulations is given by the Gelfand-Naimark theorem and its commutative version[?], Sakai's theorem on the relation between abstract and concrete W^* -algebras[?], its commutative analogue, Riesz theorem[?, ?, 10], Gleason's theorem[?].

4. Methodology

5. Timetable

6. Expected Results

Recommended Jury

Universidad de los Andes

1. Alexander Cardona Guio

Associate Professor of the Mathematics Department

acardona@uniandes.edu.co

CV: <https://pentagono.uniandes.edu.co/~acardona/ACcurriculumvitae.pdf>

2. Nelson Javier Buitrago Aza

Postdoctoral Researcher of the Physics Department

nj.buitragoa@uniandes.edu.co

3. Pedro Bargueño de Retes

Faculty Professor of the Physics Department

p.bargueno@uniandes.edu.co

CV: http://scienti.colciencias.gov.co:8081/cv lac/visualizador/generarCurriculoCv.do?cod_rh=0001572334

Other

1. Aiyalam Parameswaran Balachandran

Professor Emeritus at Syracuse University

balachandran38@gmail.com

2. Jorge Andrés Plazas Vargas

Assistant Professor at Universidad Javeriana

jorge.plazas@javeriana.edu.co

CV: http://scienti.colciencias.gov.co:8081/cv lac/visualizador/generarCurriculoCv.do?cod_rh=0001544588

3. Leonardo Arturo Cano García

Professor at Univesidad Nacional

lcanog@unal.edu.co

CV: http://scienti.colciencias.gov.co:8081/cv lac/visualizador/generarCurriculoCv.do?cod_rh=0001450991

Signatures

Iván Mauricio Burbano Aldana

Andrés Fernando Reyes Lega

References

- [1] Masanori Ohya and Dénes Petz. *Quantum Entropy and its Use*. Springer-Verlag, 1993.
- [2] Ola Bratteli and Derek W Robinson. *Operator Algebras and Quantum Statistical Mechanics*, volume 1. Springer, second edition, 1987.
- [3] Ola Bratteli and Derek W. Robinson. *Operator Algebras and Quantum Statistical Mechanics*, volume 2. Springer, second edition, 1997.

- [4] Alain Connes and Matilde Marcolli. *Noncommutative Geometry, Quantum Fields and Motives*. American Mathematical Society and Hindustan Book Agency, 2008. URL: <http://www.alainconnes.org/docs/bookwebfinal.pdf>.
- [5] Alain Connes. Temps et aléa du quantique, 2015. URL: <https://www.youtube.com/watch?v=ODAngTW8deg{&}t=3047s>.
- [6] Iván Mauricio Burbano. *KMS States and Tomita-Takesaki Theory*. B.sc. thesis, Universidad de los Andes, 2017. URL: <https://github.com/ivanmbur/Monografia/blob/master/monografia.pdf>.
- [7] Alain Connes. Une classification des facteurs de type III. *Ann. scient. Éc. Norm. Sup.*, 6(4):133–252, 1973.
- [8] Jakob Yngvason. The Role of Type III Factors in Quantum Field Theory. *Reports on Mathematical Physics*, 55(1):135–147, 2004. URL: <https://arxiv.org/pdf/math-ph/0411058.pdf>, arXiv:0411058, doi: 10.1016/S0034-4877(05)80009-6.
- [9] Brian C. Hall. *Quantum Theory for Mathematicians*. Springer, 2013. URL: <http://link.springer.com/10.1007/978-1-4614-7116-5>, doi: 10.1007/978-1-4614-7116-5.
- [10] Edwin Hewitt and Karl Stromberg. *Real and Abstract Analysis*. Springer, New York, 1975.