

KMS states and Tomita-Takesaki Theory


Iván Mauricio Burbano Aldana

Advised by: Prof. Andrés Fernando Reyes Lega

Universidad de los Andes


May 13, 2018

Motivation



Can we obtain the equations of motion from the equilibrium state?

Motivation



Can we obtain the equations of motion from the equilibrium state?

Maybe in quantum thermal systems.

$$e^{-\beta H} \circlearrowright e^{-iHt}$$

$$\text{temperature} \iff i \times \text{time}$$

Outline

- 1 Classical and Quantum Theories
- 2 Algebraic Quantum Mechanics
- 3 KMS States
- 4 Tomita-Takesaki Theory
- 5 The Canonical Time Evolution

Elements of Classical and Quantum Theories

Classical theories

- Auxiliary space: locally compact Hausdorff space X ;

Quantum theories

- Auxiliary space: separable Hilbert space \mathcal{H}

Elements of Classical and Quantum Theories

Classical theories

- Auxiliary space: locally compact Hausdorff space X ;
- Observables: continuous functions $C(X)$ on X ;

Quantum theories

- Auxiliary space: separable Hilbert space \mathcal{H}
- Observables: self-adjoint operators on \mathcal{H}

Elements of Classical and Quantum Theories

Classical theories

- Auxiliary space: locally compact Hausdorff space X ;
- Observables: continuous functions $C(X)$ on X ;
- States: probability measures P on X ;

Quantum theories

- Auxiliary space: separable Hilbert space \mathcal{H}
- Observables: self-adjoint operators on \mathcal{H}
- States: positive, self-adjoint, normalized and trace-class operators ρ on \mathcal{H} ;

Elements of Classical and Quantum Theories

Classical theories

- Auxiliary space: locally compact Hausdorff space X ;
- Observables: continuous functions $C(X)$ on X ;
- States: probability measures P on X ;
- Expected values: $\int f dP$.

Quantum theories

- Auxiliary space: separable Hilbert space \mathcal{H}
- Observables: self-adjoint operators on \mathcal{H}
- States: positive, self-adjoint, normalized and trace-class operators ρ on \mathcal{H} ;
- Expected values: $\text{tr}(A\rho)$.

Algebraic Quantum Mechanics

- Observables: A C^* -algebra \mathcal{A} :
 - ▶ Complete normed vector space with product and involution;
 - ▶ C^* property: $\|A^*A\| = \|A\|^2$;
 - ▶ A C^* -algebra can always be realized as a uniformly closed subset of the bounded operators on a Hilbert space [Bratteli and Robinson, 1987]. It is called a von Neumann algebra or W^* -algebra if $\mathcal{A}'' = \mathcal{A}$ where the commutant \mathcal{A}' of a set \mathcal{A} of bounded operators on a Hilbert space is defined as the set of all bounded operators which commute with every element of \mathcal{A} .
- States: Positive normalized linear functionals $\omega : \mathcal{A} \rightarrow \mathbb{C}$.

GNS Construction

Start with a C^* -algebra \mathcal{A} and a state ω .

- $\mathcal{N}_\omega := \{A \in \mathcal{A} \mid \omega(AA^*) = 0\}$
- Hilbert space $\mathcal{H}_\omega := \overline{\mathcal{A}/\mathcal{N}_\omega}$ with $\langle [A], [B] \rangle := \omega(A^*B)$
- Define the representation extending

$$\begin{aligned}\pi_\omega : \mathcal{A} &\rightarrow \mathcal{B}(\mathcal{H}_\omega) \\ A &\mapsto \pi_\omega(A) : \mathcal{H}_\omega \rightarrow \mathcal{H}_\omega \\ [B] &\mapsto [AB]\end{aligned}$$

- Cyclic vector $\Omega_\omega := [1]$, that is, $\overline{\mathcal{A}\Omega_\omega} = \mathcal{H}_\omega$
- This is the unique $*$ -representation of \mathcal{A} with a cyclic vector Ω_ω such that $\omega(A) = \langle \Omega_\omega, \pi_\omega(A)\Omega_\omega \rangle$.

Cyclic representations of W^* -algebras

Theorem (★)

If \mathfrak{M} is a W^* -algebra and ω is a faithful ($\omega(A^*A) = 0 \rightarrow A = 0$) normal ($\omega(A) = \text{tr}(\rho A)$) state then its cyclic representation $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega)$ satisfies

- π_ω is faithful (injective);
- $\pi_\omega(\mathfrak{M})$ is a von Neumann algebra;
- Ω_ω is separating for $\pi_\omega(\mathfrak{M})$ ($\pi_\omega(A)\Omega_\omega = 0 \rightarrow \pi_\omega(A) = 0$).

Dynamical Systems

Time evolution is represented by a one-parameter group of automorphisms

$$\begin{aligned}\tau : \mathbb{R} &\rightarrow \text{Aut}(\mathcal{A}) \\ t &\mapsto \tau_t.\end{aligned}$$

Dynamical systems consist of an $C^*(W^*)$ -algebra with a time evolutions which satisfy certain continuity properties.

KMS States

Definition

Let (\mathcal{A}, τ) be a dynamical system. We say that a state ω is a (τ, β) -KMS state if for all $A, B \in \mathcal{A}$ there exists a continuous bounded function $F_{A,B} : \overline{\mathfrak{D}_\beta} \rightarrow \mathbb{C}$ analytic on \mathfrak{D}_β (the strip of the complex plane bounded by $\operatorname{Im} z = 0$ and $\operatorname{Im} z = \beta$) such that

$$\begin{aligned}F_{A,B}(t) &= \omega(A\tau_t(B)) \\ F_{A,B}(t + i\beta) &= \omega(\tau_t(B)A)\end{aligned}$$

for all $t \in \mathbb{R}$.

KMS states as Equilibrium states

KMS states are a candidate for a general definition of thermodynamic equilibrium in quantum systems[Haag, 1992][Duvenhage, 1999]:

- KMS states are invariant under the dynamics $\omega(\tau_t(A)) = \omega(A)$;
- In finite dimensional Hilbert spaces with Schrödinger's time evolution τ , the only possible (τ, β) -KMS states are the β -Gibbs states

$$\mathcal{B}(\mathcal{H}) \rightarrow \mathbb{C}$$

$$A \mapsto \frac{\text{tr}(Ae^{-\beta H})}{\text{tr}(e^{-\beta H})}.$$

Tomita-Takesaki Theory

For a W^* -algebra \mathfrak{M} equipped with a cyclic and separating vector Ω Tomita-Takesaki theory yields:

- a one-parameter unitary group $t \mapsto \Delta^{it}$;
- a modular conjugation J .

Theorem (Tomita-Takesaki)

- $J\mathfrak{M}J = \mathfrak{M}'$;
- $\Delta^{it}\mathfrak{M}\Delta^{-it} = \mathfrak{M}$ for all $t \in \mathbb{R}$.

Proof.

[Duvenhage, 1999]



Tomita-Takesaki, Time Evolution and KMS States

Theorem (★)

$t \mapsto \Delta^{it}$ is the unique strongly continuous one-parameter unitary group on \mathcal{H} that satisfies the KMS condition with respect to \mathcal{K} such that $\Delta^{it}\mathcal{K} \subseteq \mathcal{K}$ for all $t \in \mathbb{R}$.

Theorem (★)

Let \mathfrak{M} be a von Neuman algebra and ω a faithful normal state. Consider the unitary group $t \mapsto \Delta^{it}$ associated to the pair $(\pi_\omega(\mathfrak{M}), \Omega_\omega)$. Then the one-parameter group of automorphisms given by $\alpha_t = \pi_\omega^{-1}(\Delta^{it}\pi_\omega(A)\Delta^{-it})$ makes (\mathfrak{M}, α) a W^ -dynamical system.*

Proof.

[Duvenhage, 1999]



The Canonical Time Evolution

Theorem (★★★)

Let \mathfrak{M} be a von Neumann algebra and ω be a faithful normal state. Then (\mathfrak{M}, τ) with $\tau_t(A) = \alpha_{-t/\beta}(A)$ and α the modular group of (\mathfrak{M}, ω) is the unique W^ -dynamical system such that ω is a (τ, β) -KMS state.*

Proof.

[Duvenhage, 1999]



Further work

- Classical KMS states and Tomita-Takesaki theory.
- Understanding KMS states from "first principles":
 - ▶ stability;
 - ▶ passivity.
- Relativistic generalization of KMS states.
- Entropy ambiguities.

References I



Bratteli, O. and Robinson, D. W. (1987).
Operator Algebras and Quantum Statistical Mechanics 1.
Springer, 2nd edition.



Duvenhage, R. D. V. (1999).
Quantum statistical mechanics , KMS states and Tomita-Takesaki theory.
Msc, University of Pretoria.



Haag, R. (1992).
Local Quantum Physics: Fields, Particles, Algebras.
Springer, 2nd edition.