

KMS states and Tomita-Takesaki Theory


Iván Mauricio Burbano Aldana

Advised by: Prof. Andrés Fernando Reyes Lega

Universidad de los Andes


November 20, 2017

Motivation



Can we obtain the equations of motion from the equilibrium state?

Motivation



Can we obtain the equations of motion from the equilibrium state?

Maybe in quantum thermal systems.

$$e^{-iHt} \circ e^{-\beta H}$$

Outline

- 1 Classical and Quantum Theories
- 2 Algebraic Quantum Mechanics
- 3 KMS States
- 4 Tomita-Takesaki Theory
- 5 The Canonical Time Evolution

Elements of Classical and Quantum Theories

Classical theories

- Auxiliary space: locally compact Hausdorff space X ;

Quantum theories

- Auxiliary space: separable Hilbert space \mathcal{H}

Elements of Classical and Quantum Theories

Classical theories

- Auxiliary space: locally compact Hausdorff space X ;
- Observables: continuous functions on a locally compact Hausdorff space $C(X)$;

Quantum theories

- Auxiliary space: separable Hilbert space \mathcal{H}
- Observables: self-adjoint operators on \mathcal{H}

Elements of Classical and Quantum Theories

Classical theories

- Auxiliary space: locally compact Hausdorff space X ;
- Observables: continuous functions on a locally compact Hausdorff space $C(X)$;
- States: probability measures P on X ;

Quantum theories

- Auxiliary space: separable Hilbert space \mathcal{H}
- Observables: self-adjoint operators on \mathcal{H}
- States: positive, self-adjoint, normalized and trace-class operators ρ on \mathcal{H} ;

Elements of Classical and Quantum Theories


Classical theories

- Auxiliary space: locally compact Hausdorff space X ;
- Observables: continuous functions on a locally compact Hausdorff space $C(X)$;
- States: probability measures P on X ;
- Expected values: $\int f dP$.


Quantum theories

- Auxiliary space: separable Hilbert space \mathcal{H}
- Observables: self-adjoint operators on \mathcal{H}
- States: positive, self-adjoint, normalized and trace-class operators ρ on \mathcal{H} ;
- Expected values: $\text{tr}(A\rho)$.


Difference between Classical and Quantum theories




Difference between Classical and Quantum theories



Difference between Classical and Quantum theories



Difference between Classical and Quantum theories



Algebraic Quantum Mechanics

- Observables: A $C^*(W^*)$ -algebra \mathcal{A} :
 - ▶ Complete normed vector space with product and involution;
 - ▶ C^* property: $\|A^*A\| = \|A\|^2$;
 - ▶ If $W^*: \mathcal{A}'' = \mathcal{A}$.
- States: Positive normalized linear functionals $\omega : \mathcal{A} \rightarrow \mathbb{C}$.

GNS Construction

Start with a C^* -algebra \mathcal{A} and a state ω .

- $\mathcal{N}_\omega := \{A \in \mathcal{A} \mid \omega(AA^*) = 0\}$
- Hilbert space $\mathcal{H}_\omega := \overline{\mathcal{A}/\mathcal{N}_\omega}$ with $\langle [A], [B] \rangle := \omega(A^*B)$
- Define the representation extending

$$\begin{aligned}\pi_\omega : \mathcal{A} &\rightarrow \mathcal{B}(\mathcal{H}_\omega) \\ A &\mapsto \pi_\omega(A) : \mathcal{H}_\omega \rightarrow \mathcal{H}_\omega \\ [B] &\mapsto [AB]\end{aligned}$$

- Cyclic vector $\Omega_\omega := [1]$
- Important facts:
 - ▶ $\omega(A) = \langle \Omega_\omega, A\Omega_\omega \rangle$
 - ▶ If \mathcal{A} is a W^* -algebra, π_ω is a $*$ -isomorphism, $\pi_\omega(\mathcal{A})$ is a W^* -algebra, and Ω_ω is cyclic and separating ★.

Dynamical Systems

Time evolution is represented by a one-parameter group of automorphisms

$$\begin{aligned}\tau : \mathbb{R} &\rightarrow \text{Aut}(\mathcal{A}) \\ t &\mapsto \tau_t.\end{aligned}$$

Dynamical systems consist of an $C^*(W^*)$ -algebra with a time evolution which satisfy certain continuity properties.

KMS States

Definition

Let (\mathcal{A}, τ) be a dynamical system. We say that a state ω is a (τ, β) -KMS state if for all $A, B \in \mathcal{A}$ there exists a continuous bounded function $F_{A,B} : \overline{\mathfrak{D}_\beta} \rightarrow \mathbb{C}$ analytic on \mathfrak{D}_β (the strip of the complex plane bounded by $\text{Im } z = 0$ and $\text{Im } z = \beta$) such that

$$\begin{aligned}F_{A,B}(t) &= \omega(A\tau_t(B)) \\ F_{A,B}(t + i\beta) &= \omega(\tau_t(B)A)\end{aligned}$$

for all $t \in \mathbb{R}$.

KMS states as Equilibrium states

KMS states are a candidate for a general definition of thermodynamic equilibrium in quantum systems[Haag, 1992][Duvenhage, 1999]:

- KMS states are invariant under the dynamics $\omega(\tau_t(A)) = \omega(A)$;
- In finite dimensional Hilbert spaces with Schrödinger's time evolution τ , the only possible (τ, β) -KMS states are the β -Gibbs states

$$\mathcal{B}(\mathcal{H}) \rightarrow \mathbb{C}$$

$$A \mapsto \frac{\text{tr}(Ae^{-\beta H})}{\text{tr}(e^{-\beta H})}.$$

Tomita-Takesaki Theory

For a W^* -algebra \mathfrak{M} equipped with a cyclic and separating vector Ω Tomita-Takesaki theory yields:

- a one-parameter unitary group $t \mapsto \Delta^{it}$;
- a modular conjugation J .

Theorem (Tomita-Takesaki)

- $J\mathfrak{M}J = \mathfrak{M}'$;
- $\Delta^{it}\mathfrak{M}\Delta^{-it} = \mathfrak{M}$ for all $t \in \mathbb{R}$.

Proof.

[Duvenhage, 1999]



Tomita-Takesaki, time evolution and KMS states

Theorem (★)

$t \mapsto \Delta^{it}$ is the unique strongly continuous one-parameter unitary group on \mathcal{H} that satisfies the KMS condition with respect to \mathcal{K} such that $\Delta^{it}\mathcal{K} \subseteq \mathcal{K}$ for all $t \in \mathbb{R}$.

Theorem (★)

Let \mathfrak{M} be a von Neuman algebra and ω a faithful normal state. Consider the unitary group $t \mapsto \Delta^{it}$ associated to the pair $(\pi_\omega(\mathfrak{M}), \Omega_\omega)$. Then the one-parameter group of automorphisms given by $\alpha_t = \pi_\omega^{-1}(\Delta^{it}\pi_\omega(A)\Delta^{-it})$ makes (\mathfrak{M}, α) a W^ -dynamical system.*

Proof.

[Duvenhage, 1999]



The Canonical Time Evolution

Theorem (★★★)

Let \mathfrak{M} be a von Neumann algebra and ω be a faithful normal state. Then (\mathfrak{M}, τ) with $\tau_t(A) = \alpha_{-t/\beta}(A)$ and α the modular group of (\mathfrak{M}, ω) is the unique W^ -dynamical system such that ω is an (α, β) -KMS state.*

Proof.

[Duvenhage, 1999]



Further work

- Classical KMS states and Tomita-Takesaki theory.
- Understanding KMS states from "first principles":
 - ▶ stability;
 - ▶ passivity.
- Relativistic generalization of KMS states.
- Entropy ambiguities.

References I



Bell, J. S. (1964).
The Einstein Podolsky Rosen Paradox.
Physics, 1(3):195–200.



Duvenhage, R. D. V. (1999).
Quantum statistical mechanics , KMS states and Tomita-Takesaki theory.
Msc, University of Pretoria.



Einstein, A., Podolsky, B., and Rosen, N. (1935).
Can Quantum-Mechanical Description of Reality Be Considered Complete?
Physical Review, 47.



Haag, R. (1992).
Local Quantum Physics: Fields, Particles, Algebras.
Springer, 2nd edition.

References II



Reyes-Lega, A. F. (2013).

Some Aspects of Operator Algebras in Quantum Physics.

In Cano, L., Cardona, A., Ocampo, H., and Reyes-Lega, A. F., editors, *Geometric, Algebraic and Topological Methods for Quantum Field Theory*, pages 1–74. World Scientific.



Wilce, A. (2012).

Quantum Logic and Probability Theory.

Stanford Encyclopedia of Philosophy.