

Probability paradoxes

Ivan Murashko

Contents

1	Base definitions of probability theory	1
2	Monty Hall problem	2
3	Waiting time on a bus stop	2

Introduction

The goal for the article is to demonstrate several paradoxes that are related to probability theory and how can they can be solved.

1 Base definitions of probability theory

I am going to provide several definitions. I will give the both formal and informal definitions and show how they are related each other.

We will start with the simplest example.




Figure 1: Probability example

Example 1.

TBD [1]



Figure 2: Probability space. It consists of elementary events: a , b , c and d , each of them has equal probability $p_{a,b,c,d} = \frac{1}{4}$




Figure 3: Condition probability. Original probability space. $P(R) = \frac{5}{6}$, $P(B) = \frac{3}{6}$, $P(G) = \frac{4}{6}$




Figure 4: Condition probability. $P(G|R) = \frac{3}{5}$, $P(B|R) = \frac{2}{5}$

2 Monty Hall problem

TBD

3 Waiting time on a bus stop

TBD

References

- [1] А. Н. Колмогоров. Основные понятия теории вероятностей / А. Н. Колмогоров. — Москва: Наука, 1974.



Figure 5: Condition probability. $P(R|B) = \frac{2}{3}$, $P(G|B) = \frac{1}{3}$




Figure 6: Condition probability. $P(B|G) = \frac{1}{4}$, $P(R|G) = \frac{3}{4}$