



VRIJE
UNIVERSITEIT
BRUSSEL

¹ **A search for flavour changing neutral currents
2 involving a top quark and a Z boson, using the
3 data collected by the CMS collaboration at a
4 centre of mass of 13 TeV**

⁵ Van Parijs, Isis

⁶ Proefschrift ingediend met het oog op het behalen van de academische graad
⁷ Doctor in de Wetenschappen.

Published in Faculteit Wetenschappen & Bio-ingenieurswetenschappen
Vrije Universiteit Brussel
At 1. June 2017.

⁸

Responsible Contact: I. Van Parijs
Institute for High Energy Physics
Promotor: Prof. Jorgen D'Hondt

9 First Referee: Prof. Dr. J. D'Hondt
 Date of Hand-in: 10 November 2017
 Date of Defense: 10 December 2017

Contents

11	1 An introduction to the theory	1
12	1.1 Elementary particles and forces	1
13	1.2 Standard Model Lagrangian	2
14	1.3 Flavour changing currents in the SM	5
15	1.4 Top quark physics in the SM	6
16	1.5 Experimental results on the SM top quark	8
17	1.6 Motivations for new physics	12
18	1.7 An effective approach beyond the SM: FCNC involving a top quark	13
19	1.8 Experimental constraints on top-FCNC	15
20	2 Experimental set-up	17
21	2.1 The Large Hadron Collider	17
22	2.2 The Compact Muon Solenoid	20
23	2.2.1 CMS coordinate system	21
24	2.2.2 Towards the heart of CMS	22
25	2.2.3 Data acquisition	30
26	2.2.4 Phase 1 upgrades	31
27	2.2.5 CMS computing model	32
28	3 Analysis techniques	33
29	3.1 Hadron collisions at high energies	33
30	3.2 Event generation	36
31	3.2.1 Fundamentals of simulating a proton collision	36
32	3.2.2 Programs for event generation	36
33	3.2.3 Generating FCNC top-Z interactions	38
34	3.2.4 Generating SM background events	39
35	3.3 Multivariate analysis techniques: Boosted Decision Trees	42
36	3.4 Statistical methodology	44
37	3.4.1 The absence of signal: limits	45
38	3.4.2 Adding sources of uncertainty	47
39	3.4.3 Asymptotic approximation of the CL method	48
40	3.4.4 Extracting the signal model parameters	48
41	4 Event reconstruction and selection	49

42	4.1 Event reconstruction	49
43	4.2 Event selection	51
44	4.3 Regions and channels	51
45	4.4 Data driven background simulation	51
46	5 The search for FCNC involving a top quark and a Z boson	53
47	5.1 Construction of template distributions	53
48	5.2 Systematic uncertainties	53
49	5.3 Limit setting procedure	53
50	5.4 Result and discussion	53
51	6 Conclusion and outlook	55
52	Bibliography	57

Theoretical basis

1

54 The Standard Model (SM) [1] is a name given in 1970s to a theory describing the fundamental
 55 particles and their interactions. This quantum field theory describes the particles and their
 56 interactions as fields and has successfully incorporated three of the four fundamental forces in
 57 the universe. In [Section 1.1](#), the particle content of the SM is summarised, while [Section 1.2](#)
 58 describes the SM Lagrangian and its symmetries. In [Section 1.3](#), the flavour content of the SM
 59 is highlighted, while [Section 1.4](#) focusses on the SM top quark. The latest experimental results
 60 of the top quark are given in [Section 1.5](#).

61 The successful theory of the SM has some shortcomings which are discussed in [Section 1.6](#)
 62 and lead to searches for a more general theory. One of such a search is using effective field
 63 theory (EFT) [2] to search for new physics in a model independent way. In [Section 1.7](#) an EFT
 64 model focussing on flavour changing neutral currents (FCNC) involving a top quark is presented.
 65 Its current experimental constraints are given in [Section 1.8](#).

66 1.1 Elementary particles and forces

67 The interactions in nature can be described by four forces, the strong force, the electromagnetic
 68 (EM) force, the weak force and the gravitational force. These interactions happen via particles
 69 with an integer spin known as bosons. The strong interaction is mediated by eight gluons g ,
 70 while the electromagnetic force is mediated by photons γ , and the weak force by Z and W^\pm
 71 bosons. In [Table 1.1](#), the forces and their characteristics are shown. The gravitational force is
 72 the only force not included in the SM and can be neglected for energies lower than the Planck
 scale ($1.22 \cdot 10^{19}$ GeV).

Table 1.1: The four forces of nature and their characteristics.

	Range	Mediator
Strong force	$10^{\text{-}e} - 15$ m	8 gluons
Electromagnetic force	∞	photon
Weak force	$10^{\text{-}18}$ m	W^\pm , Z bosons
Gravitational force	∞	unknown

The fermions are the particles that make up the visible matter in the universe. They carry half integer spin and can be subdivided into leptons and quarks, where leptons don't interact strongly. Each fermion has a corresponding anti-fermion which has the same mass and is oppositely charged. The electron e^- is the first elementary particle discovered [3] and belongs to the first generation of leptons together with the electron neutrino ν_e . The second generation compromises the muon μ^- and muon neutrino ν_μ , whereas the third generation consists of the tau τ and tau neutrino ν_τ . The neutrino's are neutral particles, while the other leptons have charge $\pm q_e$ where q_e represents the elementary charge of $1.602 \cdot 10^{-19}$ C. The masses of charged leptons differ by four orders of magnitude between the first and third generations. In the SM the neutrino's are assumed to be massless, nonetheless it is experimentally established that neutrino do have a tiny non-zero mass. In Table 1.2, the leptons and their properties in the SM are summarised.

Table 1.2: The properties of the leptons in the three generations of the SM [4], where q_e represents the elementary charge.

Generation	Particle	Mass	Charge
First	e^-	0.511 MeV	$-q_e$
	ν_e	≈ 0	0
Second	μ^-	106 MeV	$-q_e$
	ν_μ	≈ 0	0
Third	τ	1 777 MeV	$-q_e$
	ν_τ	≈ 0	0

85

The quarks can also be divided into three generations. Unlike the leptons, they carry colour charge and can interact via the strong interaction. The top quark, discovered in 1995 at the Tevatron [5, 6], is the heaviest SM particle with a mass close to 173.1 ± 0.6 GeV¹ [4]. The quarks and their properties are summarized in Table 1.3. In nature, only colour neutral objects can exist. This has as consequence that quarks are bound through gluons into mesons (quark+anti-quark) and baryons (three quarks). These mesons and baryons are mostly short-lived and unstable particle that rapidly decay through W^\pm and Z bosons, associated with a fermion. The only known stable baryon is the proton, made up of two up quarks and one down quark.

The scalar boson, commonly known as the Higgs boson, is the last piece of the SM and is discovered in 2012 [7, 8]. It is responsible for the masses of the W^\pm and Z boson, and that of the fermions.

97 1.2 Standard Model Lagrangian

The SM is a quantum field theory and thus describes the dynamics and kinematics of particles and forces by a Lagrangian \mathcal{L} . The theory is based on the $SU_C(3) \times SU_L(2) \times U_Y(1)$ gauge symmetry, where $SU_L(2) \times U_Y(1)$ describes the electroweak interaction and $SU_C(3)$ the strong

¹In this thesis all masses and energies are expressed in natural units, where the speed of light and \hbar are taken to be equal to one.

Table 1.3: The properties of the quarks in the three generations of the SM [4], where q_e represents the elementary charge.

	Generation	Particle	Mass	Charge
First	up u	$2.2^{+0.6}_{-0.4}$ MeV	$\frac{2}{3} q_e$	
	down d	$4.7^{+0.5}_{-0.4}$ MeV	$\frac{-1}{3} q_e$	
Second	charm c	1.28 ± 0.03 GeV	$\frac{2}{3} q_e$	
	strange s	96^{+8}_{-4} MeV	$\frac{-1}{3} q_e$	
Third	top t	173.1 ± 0.6 GeV	$\frac{2}{3} q_e$	
	bottom b	$4.18^{+0.04}_{-0.03}$ GeV	$\frac{-1}{3} q_e$	

101 coupling. The indices refer to colour C, the left chiral nature of the $SU_L(2)$ coupling L, and the
 102 weak hypercharge Y. Its Lagrangian is constructed such that contains symmetries representing
 103 physics conservation laws such as conservation of energy, momentum and angular momentum.
 104 The symmetries under local group transformations are sustained by demanding gauge invariance
 105 .

The $U_Y(1)$ group has one generator Y with an associated gauge field B_μ . The three gauge fields W_μ^1 , W_μ^2 , and W_μ^3 , are associated to $SU_L(2)$ with three generators that can be written as half of the Pauli matrices:

$$T_1 = \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad T_2 = \frac{1}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \text{and} \quad T_3 = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1.1)$$

The generators T^a satisfy the Lie algebra:

$$[T^a, T^b] = i\epsilon^{abc} T_c \text{ and } [T^a, Y] = 0, \quad (1.2)$$

106 where ϵ^{abc} is an antisymmetric tensor. The gauge fields of $SU_L(2)$ only couple to left-handed
 107 fermions as required by the observed parity violating nature of the weak force. The $SU_C(3)$
 108 group represents quantum chromodynamics (QCD). It has eight generators corresponding to
 109 eight gluon fields $G_\mu^{1..8}$. Unlike $SU_L(2) \times U_Y(1)$, $SU_C(3)$ is not chiral.

Under $SU_C(3)$ quarks are colour triplets while leptons are colour singlets. This implies that the quarks carry a colour index ranging between one and three, whereas leptons do not take part in strong interactions. Based on the chirality, the quarks and leptons are organized in doublets or singlets. Each generation i of fermions consists of left-handed doublets and right-handed singlets:

$$l_{L,i} = \begin{pmatrix} e^-_{L,i} \\ \nu_{L,i} \end{pmatrix}, \quad e^-_{R,i}, \quad q_{L,i} = \begin{pmatrix} u_{L,i} \\ d_{L,i} \end{pmatrix}, \quad u_{R,i}, \quad \text{and} \quad d_{R,i} \quad (1.3)$$

The SM Lagrangian can be decomposed as a sum of four terms

$$\mathcal{L}_{SM} = \mathcal{L}_{gauge} + \mathcal{L}_f + \mathcal{L}_{Yuk} + \mathcal{L}_\phi, \quad (1.4)$$

NOTE:
should I explain gauge invariance or is a reference enough?

that are related to the gauge, fermion, Yukawa and scalar sectors. The gauge Lagrangian regroups the gauge fields of all three symmetry groups, and the fermionic part consists of kinetic energy terms for quarks and leptons. The interaction between fermions and the scalar doublet ϕ gives rise to fermion masses and is described by the Yukawa Lagrangian. The scalar part of the Lagrangian is composed of a kinematic and potential component related to the scalar boson.

For the electroweak theory, two coupling constants are introduced, namely g' for $U_Y(1)$ and g for $SU_L(2)$. The physically observable gauge bosons of this theory are the photon field A_μ , the Z boson field Z_μ^0 , and the W field W_μ^\pm . These are a superposition of the four gauge fields of $SU_L(2) \times U_Y(1)$:

$$A_\mu = \sin\theta_W W_\mu^1 + \cos\theta_W B_\mu, \quad Z_\mu^0 = \cos\theta_W W_\mu^3 - \sin\theta_W B_\mu, \quad \text{and} \quad W_\mu^\pm = \sqrt{\frac{1}{2}} (W_\mu^1 \mp W_\mu^2), \quad (1.5)$$

where θ_W represents the weak mixing angle defined as $\tan\theta_W = \frac{g'}{g}$.

The coupling constant representing the strength of the QCD interactions is denoted as g_s . In QCD there is asymptotic freedom whereby the strong coupling constant becomes weaker as the energy with which the interaction between strongly interacting particles is probed increases, and stronger as the distance between the particles increases. A consequence of this is known as colour confinement. The quarks and gluons can not exist on their own and are not observed individually. They are bound in colour neutral states called hadrons, this process is known as hadronisation.

123 Electroweak symmetry breaking

In $\mathcal{L}_{\text{gauge}}$ and \mathcal{L}_f are no mass terms for fermions present because only singlets under $SU_C(3) \times SU_L(2) \times U_Y(1)$ can acquire a mass with an interaction of the type $m^2 \phi^\dagger \phi$ without breaking the gauge invariance. In order to accommodate mass terms for fermions and gauge fields, electroweak symmetry breaking, leading to \mathcal{L}_ϕ is introduced.

The scalar doublet is introduced in the SM as

$$\phi = \frac{1}{\sqrt{2}} \begin{pmatrix} \varphi_1 + i\varphi_2 \\ \varphi_3 + i\varphi_4 \end{pmatrix}. \quad (1.6)$$

NOTE:
check if I
need to add
constants
here

Its field potential is of the form

$$V(\phi) = \mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2, \quad (1.7)$$

with $\mu^2 < 0$ and λ a positive integer. This choice of parameters gives the potential a "Mexican hat" shape. It has an infinite set of minima (ground states) and by expanding the field around an arbitrary choice of ground state, the electroweak symmetry is broken (EW):

$$\phi = \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix} + \hat{\phi}, \quad (1.8)$$

where v is the vacuum expectation value (vev), measured to be around 245 GeV and corresponds to $\sqrt{\frac{-\mu}{\lambda}}$. The scalar doublet's four degrees of freedom is reduced to three degrees of freedom

that couple to the gauge fields and mix with the W^+ , W^- and Z bosons. The remaining fourth degree of freedom has given rise to a physically observable particle , called the Brout-Englert-Higgs (BEH) boson. This spontaneous symmetry breaking leaves the gauge invariance intact and gives masses to the W^\pm and Z bosons as:

$$m_W = \frac{1}{2}v|g| \quad \text{and} \quad m_Z = \frac{1}{2}v\sqrt{g'^2 + g^2}. \quad (1.9)$$

- 128 The Brout-Englert-Higgs field couples universally fermions with a strength proportional to their
129 masses, and to gauge bosons with a strength proportional to the square of their masses.

130 1.3 Flavour changing currents in the SM

Flavour changing charged currents are introduced in 1963 by Nicola Cabibbo [9]. Via interaction with a W boson the flavour of the quarks is changed. At the time of the postulation only up, down, and strange quarks were known and the charged weak current was described as a coupling between the up quark and d_{weak} , where d_{weak} is a linear combination of the down and strange quarks, $d_{\text{weak}} = \cos\theta_c d + \sin\theta_c s$. This linear combination is a direct consequence of the chosen rotation

$$\begin{pmatrix} d_{\text{weak}} \\ s_{\text{weak}} \end{pmatrix} = \begin{pmatrix} \cos\theta_c & \sin\theta_c \\ -\sin\theta_c & \cos\theta_c \end{pmatrix} \begin{pmatrix} d \\ s \end{pmatrix} = \mathcal{R} \begin{pmatrix} d \\ s \end{pmatrix}, \quad (1.10)$$

where the rotation angle θ_c is known as the Cabibbo angle. This provides a definition for the charged weak current between u and d quarks,

$$J_\mu = \bar{u} \gamma_\mu (1 + \gamma_5) d_{\text{weak}}. \quad (1.11)$$

A consequence of Cabibbo's approach is that the s_{weak} is left uncoupled, leading to Glashow, Iliopoulos and Maiani (GIM) [10–12] to require the existence of a fourth quark with charge $\frac{2}{3}q_e$. This quark, known as the charm quark, couples to s_{weak} and a new definition of the charged weak current is modified to

$$J_\mu = (u \ c) \gamma_\mu (1 + \gamma_5) \mathcal{R} \begin{pmatrix} d \\ s \end{pmatrix} = \bar{U} \gamma_\mu (1 + \gamma_5) \mathcal{R} D. \quad (1.12)$$

The neutral weak current is defined as

$$J_3 = \bar{U} \gamma_\mu (1 + \gamma_5) [\mathcal{R}, \mathcal{R}^\dagger] D, \quad (1.13)$$

- 131 and is diagonal in flavour space. This has as consequence that no flavour changing neutral
132 currents occur at tree-level Feynmann diagrams².

Kobayashi and Maskawa generalised the Cabibbo rotation matrix to accommodate for a third generation of quarks. The result is a 3×3 unitary matrix known as the CKM matrix, responsible

NOTE:
should I
explain
feynmann
diagrams?

²Feynmann diagrams are physical representation of interaction between particles. They are based on Feynmann rules [1].

for the mixing of weak interaction states of down-type quarks:

$$\begin{pmatrix} d_{\text{weak}} \\ s_{\text{weak}} \\ b_{\text{weak}} \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix} = \mathcal{V}_{\text{CKM}} \begin{pmatrix} d \\ s \\ b \end{pmatrix}. \quad (1.14)$$

The unitarity of the matrix ($\mathcal{V}_{\text{CKM}}^\dagger \mathcal{V}_{\text{CKM}} = \mathbb{1}$). A general 3×3 unitary matrix depends on three real angles and six phases. For the CKM matrix, the freedom to redefine the phases of the quark eigenstates can remove five of the phases, leaving a single physical phase known as the Kobayashi-Maskawa phase. This phase is responsible for the charge parity violation in the SM [13]. Each element V_{ij} of \mathcal{V}_{CKM} represents the transition probability of a quark i going to a quark j , and is experimentally determined to be [4]

$$\mathcal{V}_{\text{CKM}} = \begin{pmatrix} 0.97425 \pm 0.00022 & 0.2253 \pm 0.0008 & (4.13 \pm 0.49) 10^{-3} \\ 0.225 \pm 0.008 & 0.986 \pm 0.016 & (41.1 \pm 1.3) 10^{-3} \\ (8.4 \pm 0.6) 10^{-3} & (40.0 \pm 2.7) 10^{-3} & 1.021 \pm 0.032 \end{pmatrix}. \quad (1.15)$$

133 From Equation 1.15 follows that top quarks predominantly decay via charged weak currents to
 134 bottom quarks, with a probability consist with unity. In the SM, FCNC can only occur via higher
 135 loop Feynmann diagrams which are highly suppressed. The expected transition probabilities for
 136 a top quark decaying via a FCNC interaction in the SM are given in Table 1.4, where it is clear
 137 that the FCNC sector of the SM is still beyond the reach of the sensitivity of current experiments.

Table 1.4: The predicted branching ratios \mathcal{B} for FCNC interactions involving the top quark in the SM [14]

Process	\mathcal{B} in the SM	Process	\mathcal{B} in the SM
$t \rightarrow uZ$	$8 \cdot 10^{-17}$	$t \rightarrow cZ$	$1 \cdot 10^{-14}$
$t \rightarrow u\gamma$	$4 \cdot 10^{-16}$	$t \rightarrow c\gamma$	$5 \cdot 10^{-14}$
$t \rightarrow ug$	$4 \cdot 10^{-14}$	$t \rightarrow cg$	$5 \cdot 10^{-12}$
$t \rightarrow uH$	$2 \cdot 10^{-17}$	$t \rightarrow cH$	$3 \cdot 10^{-15}$

138

1.4 Top quark physics in the SM

NOTE: Add source

Discovered in 1995 by the CDF and D0 collaborations at Tevatron with proton-antiproton data, the top quark plays an important role in studying high energy physics. Its Yukawa interaction is given by

$$\mathcal{L}_{\text{top-Yukawa}} = -\frac{\lambda_t}{\sqrt{2}} \bar{t}_L t_R - \frac{\lambda_t}{\sqrt{2}} \bar{H} \bar{t}_L t_R + \text{h.c.}, \quad (1.16)$$

yielding a Yukawa coupling of

$$\lambda_t = \frac{\sqrt{2} m_t}{v} = 0.991 \pm 0.003, \quad (1.17)$$

140 with the top mass m_t equal to 172.44 ± 0.49 GeV [4]. This Yukawa coupling is very large
 141 compared to the other Yukawa couplings in the SM $\mathcal{O}(10^{-2})$, leading to the belief that the top
 142 quark may have an important role in understanding the mechanism of electroweak symmetry
 143 breaking. On top of this, the very short lifetime of the top quark makes it an excellent candidate
 144 for property studies. Its high mass, almost 40 times higher than the mass of the closest particle
 145 in mass, leads to a large coupling with the Higgs boson and makes the top quark an interesting
 146 candidate for the understanding of how particles acquire mass.

147 The CKM matrix element V_{tb} , given in [Equation 1.15](#), is experimentally found to be much
 148 larger than V_{ts} , V_{td} , and close to unity. The top quark decays through electroweak interactions
 149 since the W boson mass is smaller than the top mass and the W boson can be on shell. A
 150 consequence of this is that the top quark has a very short lifetime of only $1/\Gamma_t \approx 5 \cdot 10^{-25}$ s leading
 151 to the fact that the formation of bound states involving top quarks are not allowed. This lifetime
 152 is even shorter than the typical hadronisation timescale of $1/\Lambda_{QCD} \approx 10^{-23}$ s, prohibiting gluons
 153 to radiate from the top quark and keeping its spin coherent. Since the electroweak interactions
 154 have a V-A coupling structure, the top quark spin orientation can be derived from the angular
 155 distributions of its decay products. This makes it possible to study the polarisation of top quarks
 156 from the angular distributions in various processes.

NOTE: Add source

NOTE: Add source

157 The massiveness of the top quark leads to the fact that a large amount of energy is needed
 158 to create one. This is only the case for high energy collisions such as those in the Earth's
 159 upper atmosphere as cosmic rays collide with particles in air, or by particle accelerators. The
 160 production of top quarks happens in two ways: single via the electroweak interaction or in
 161 pairs via the strong interaction. At hadron colliders, the dominant production mechanism is top
 162 quark production via gluon ($gg \rightarrow t\bar{t}$) or quark fusion ($q\bar{q} \rightarrow t\bar{t}$). In [Figure 1.1](#), the different top
 163 pair production mechanisms are shown. The production channel of gluon fusion is the main
 164 contributor to the top pair cross section at the LHC compared to quark fusion at Tevatron. The
 165 $gg \rightarrow t\bar{t}$ process contributes 80-90% to the total top pair cross section in the LHC centre-of-mass
 166 energy regime of 7-14 TeV [4]. In [Table 1.5](#) the predicted top pair production cross sections are
 given for the LHC and Tevatron.

Figure 1.1: Leading order diagrams of the top pair production. Gluon fusion (right and middle) are the dominant processes at the LHC, while quark fusion (left) is the dominant one at Tevatron.

Table 1.5: Predictions on the top quark pair production cross sections at next-to-next-to-leading order with next-to-next-to-leading log soft gluon resummation per centre-of-mass energy [4]. The first uncertainty is from scale dependence, while the second uncertainty originates from parton density functions.

Experiment	Top mass	Centre-of-mass energy	Cross section (pb)
Tevatron	$m_t = 173.3$ GeV	$\sqrt{s} = 1.96$ TeV	$\sigma_{t\bar{t}} = 7.16^{+0.11+0.17}_{-0.20-0.12}$
LHC	$m_t = 173.2$ GeV	$\sqrt{s} = 7$ TeV	$\sigma_{t\bar{t}} = 173.6^{+4.5+8.9}_{-5.9-8.9}$
LHC	$m_t = 173.2$ GeV	$\sqrt{s} = 8$ TeV	$\sigma_{t\bar{t}} = 247.7^{+6.3+11.5}_{-8.5-11.5}$
LHC	$m_t = 173.2$ GeV	$\sqrt{s} = 13$ TeV	$\sigma_{t\bar{t}} = 816.0^{+19.4+34.4}_{-28.6-34.4}$

168 The singly produced top quarks are produced via the electroweak interaction. These production
 169 mechanisms are subdivided at leading order into three main channels based on the virtuality
 170 ($Q^2 = -p_\mu p^\mu$) of the exchanged W boson. In Figure 1.2, the corresponding Feynman diagrams
 171 are shown. The single top quark production cross section, given in Table 1.6, are smaller than
 172 the top pair production cross sections since the electroweak coupling strength is smaller than
 173 the strong coupling strength. In addition, for the single top production, there the need of sea
 174 quarks (b, \bar{q}) in the initial states for which the parton density functions increase less steeply at
 low momentum fractions compared to the gluon parton density functions.




Figure 1.2: Leading order Feynman diagrams of the electroweak production of single top quarks in the s -channel (left), t -channel (middle), and for the tW associated production. Figure taken from [15].

175

176 The production via the t -channel has a virtuality of the W boson $Q^2 > 0$, making it space-like.
 177 It is produced via the scattering of the W boson of a bottom quark coming from a proton or
 178 from gluon splitting ($g \rightarrow b\bar{b}$). This process is also known as W-gluon fusion production. It has
 179 the highest single top quark cross section in proton collisions and the top quark production is
 180 roughly twice more than the antitop quarks. This is a consequence of the up-down valence
 181 quark composition of the proton. This feature makes the t -channel sensitive to the parton
 182 density functions of the proton. The s -channel is the production mechanism with the smallest
 183 cross section. Here the W boson is time-like ($Q^2 < 0$) which requires the W boson to have a
 184 large virtuality to produce the heavier top quark. It is produced from two quarks belonging
 185 to the same isodoublet (e.g. $u \bar{d}$) and subsequently decays to $t \bar{b}$. This process get enhanced
 186 by many beyond the Standard Model scenarios via the addition of new heavy particles such
 187 as W' . The tW -channel has a top quark produced in association with a W boson produced on
 188 shell $Q^2 = -m_W^2$. This mode is negligible at Tevatron, but of relevant size at the LHC. The
 189 tW -channel is sensitive to new physics affecting the Wtb vertex.

190 1.5 Experimental results on the SM top quark

191 In this section a selection of experimental results of measurements on the SM are presented.
 192 In Figure 1.3, a summary plot of the CMS cross section measurements can be found. The
 193 estimations by the CMS and ATLAS collaborations of the CKM matrix element V_{tb} from single
 194 top quark measurement is given in Figure 1.4. The most precise estimation of V_{tb} originates from
 195 a combination of t -channel cross section measurements at 7 and 8 TeV by the CMS collaboration
 196 resulting in $|f_L V_{tb}| = 0.998 \pm 0.038$ (exp.) ± 0.016 (theo.). Assuming the $f_L = 1$ and $|V_{tb}| < 1$,
 197 this result yields a limit of $|V_{tb}| > 0.92$ at 95% confidence level. The most recent top mass
 198 measurements are given in Figure 1.5. The CMS combined amounts to is $m_t = 172.44 \pm 0.48$
 199 GeV from 7+8 TeV data.

Table 1.6: Predictions on the single top quark production cross sections at next-to-leading order per centre-of-mass energy [4]. The uncertainties from scale dependence and from parton density functions are combined in quadrature or given separately (scale + PDF). For the t -channel the relative proportions to t and \bar{t} are 65% and 35%. For the s -channel this respectively 69% and 31%. The tW -channel has an equal proportion of top and antitop quarks. For Tevatron, the top mass is assumed to be 173.3 GeV, while for the LHC predictions $m_t = 172.5$ GeV [4, 16].

Experiment	Centre-of-mass energy	Cross section $\sigma_{t+\bar{t}}$ (pb)		
		t -channel	s -channel	tW -channel
Tevatron	$\sqrt{s} = 1.96$ TeV	$2.06^{+0.13}_{-0.13}$	$1.03^{+0.05}_{-0.05}$	-
LHC	$\sqrt{s} = 7$ TeV	$63.89^{+2.91}_{-2.52}$	$4.29^{+0.19}_{-0.17}$	$15.74^{+0.40+1.10}_{-0.40-1.14}$
LHC	$\sqrt{s} = 8$ TeV	$84.69^{+3.76}_{-3.23}$	$5.24^{+0.22}_{-0.20}$	$22.37^{+0.60+1.40}_{-0.60-1.40}$
LHC	$\sqrt{s} = 13$ TeV	$216.99^{+9.04}_{-7.71}$	$10.32^{+0.40}_{-0.36}$	$71.7^{+1.80+3.40}_{-1.80-3.40}$

200 In general the various measurements show a good agreement with the SM predictions and by
 201 lack of deviations of the SM, limits on the anomalous couplings can be derived. The estimated
 202 coupling strengths per operator contributing to single top quark production obtained from
 203 various measurements at the LHC and Tevatron are shown in Figure 1.6. These results are
 204 consistent with the SM expectation for which those operators vanish.




Figure 1.3: Summary of the SM cross section measurements performed by the CMS collaboration. Figure taken from [17]




Figure 1.4: Estimations of the SM V_{tb} CKM element from single top cross section measurements. Figure taken from [18].




Figure 1.5: Summary of the top mass direct measurements performed by CMS and ATLAS, and compared with the LHC and LHC+Tevatron combinations. The results below the line are produced after the LHC and LHC+Tevatron combinations. Figure taken from [18].




Figure 1.6: Global fit results of top quark effective field theory to experimental data including all constrainable operators at dimension six. For the operators, the Warsaw basis of [19] is used. The bounds are set on the Wilson coefficients of various operators contributing to top quark production and decay in two cases (red) all other coefficients set to zero, or (blue) all other coefficient are marginalised over. Figure taken from [20].

205 1.6 Motivations for new physics

206 Many high energy experiments confirm the success of the SM. In particular the scalar boson,
 207 the cornerstone of the SM, has consecrated the theory. Unfortunately there are also strong
 208 indications that the SM ought to be a lower energy expression of a more global theory. The
 209 existence of physics beyond the SM (BSM) [21] is strongly motivated. These motivations are
 210 based on direct evidence from observation such as the existence of neutrino masses, the existence
 211 of dark matter and dark energy, or the matter-antimatter asymmetry, and also from theoretical
 212 problems such as the hierarchy problem, the coupling unification or the large numbers of free
 213 parameters in the SM.

214 In the SM, the neutrino is assumed to be massless, whilst experiments with solar, atmospheric,
 215 reactor and accelerator neutrinos have established that neutrinos can oscillate and change
 216 flavour during flight [22, 23]. These oscillations are only possible when neutrino's have masses.
 217 The flavour neutrinos (ν_e , ν_μ , ν_τ) are then linear expressions of the fields of at least three mass
 218 eigenstate neutrinos ν_1 , ν_2 , and ν_3 .

219 The ordinary or baryonic matter described by the SM describes only 5% of the mass (energy)
 220 content of the universe. Astrophysical evidence indicated that dark matter is contributing
 221 to approximately 27%, and dark energy to 68% of the content of the universe. From the
 222 measurements of the temperature and polarizations anisotropies of the cosmic microwave
 223 background by the Planck experiment [24], the density of cold non baryonic matter is determined.
 224 Cold dark matter is assumed to be only sensitive to the weak and gravitational force, leading
 225 to only one possible SM candidate: the neutrino. However, these are too light to account for
 226 the vast amount of dark matter and other models are needed. Dark energy is assumed to be
 227 responsible for the acceleration in the expansion of the universe [25].

228 At the Big Bang matter and antimatter is assumed to be produced in equal quantities. However,
 229 it is clear that we are surrounded by matter. So where did all the antimatter go? In 1967,
 230 Sakharov identified three mechanisms that are necessary to obtain a global matter antimatter
 231 asymmetry [26]. These mechanisms are those of baryon and lepton number violation, that at a
 232 given moment in time there was a thermal imbalance for the interactions in the universe, and
 233 there is charge C and charge parity CP violation³.

234 The large numbers of free parameters in the SM are taken as nine fermion masses, three CKM
 235 mixing angles and one CP violating phase, one EM coupling constant g' , one weak coupling
 236 constant g , one strong coupling constant g_s , one QCD vacuum angle, one vacuum expectation
 237 value, and one mass of the scalar boson. This large number of free parameters lead to the
 238 expectation of a more elegant, general theory beyond the SM.

239 The hierarchy problem [27] is related to the huge difference in energy between the weak
 240 scale and the Planck scale. The vev of the Brout-Englert-Higgs field determines the weak scale
 241 that is approximately 246 GeV. The radiative corrections to the scalar boson squared mass m_H^2 ,
 242 coming from its self couplings and couplings to fermions and gauge bosons, are quadratically

³The rate of a process $i \rightarrow f$ can be different from the CP-conjugate process: $\tilde{i} \rightarrow \tilde{f}$. The SM includes sources of CP-violation through the residual phase of the CKM matrix. However, these could not account for the magnitude of the asymmetry observed.

243 proportional to the ultraviolet momentum cut-off Λ_{UV} . This cut-off is at least equal to the energy
 244 to which the SM is valid without the need of new physics. The SM is valid up to the Planck mass
 245 making the correction to m_H^2 about thirty orders of magnitude larger than m_H^2 . This implies that
 246 an extraordinary cancellation of terms should happen. This is also known as the naturalness
 247 problem of the H boson mass.

The correction to the squared mass of the scalar boson coming from a fermion f , coupling to the scalar field ϕ with a coupling λ_f is given by

$$\Delta m_H^2 = -\frac{|\lambda_f|^2}{8\pi^2} \Lambda_{\text{UV}}^2, \quad (1.18)$$

while the correction to the mass from a scalar particle S with a mass m_S , coupling to the scalar field with a Lagrangian term $-\lambda_{\text{mathrm}{S}} |\phi|^2 |S|^2$ is

$$\Delta m_H^2 = -\frac{|\lambda_S|^2}{16\pi^2} \left(\Lambda_{\text{UV}}^2 - 2m_S^2 \ln \left(\frac{\Lambda_{\text{UV}}}{m_S} \right) + \dots \right). \quad (1.19)$$

248 As one can see the correction term to m_H^2 is much larger than m_H^2 itself. By introducing BSM
 249 physic models that introduce new scalar particles at TeV scale that couple to the scalar boson
 250 can cancel the Λ_{UV}^2 divergence and avoid this fine-tuning.

251 The choice of the $SU_C(3) \times SU_L(2) \times U_Y(1)$ symmetry group itself as well as the separate
 252 treatment of the three forces included in the SM raises concern. The intensity of the forces
 253 show a large disparity around the electroweak scale, but have comparable strengths at higher
 254 energies. The electromagnetic and weak forces are unified in a electroweak interaction, but the
 255 strong coupling constant does not encounter the other coupling constants at high energies. In
 256 order to reach a grand unification, the running of couplings can be modified by the addition of
 257 new particles in BSM models.

258 1.7 An effective approach beyond the SM: FCNC involving a top 259 quark

260 The closeness of the top mass to the electroweak scale led physicist to believe that it is a sensitive
 261 probe for new physics. Its property study is therefore an important topic of the experimental
 262 program at the LHC. Several extensions of the SM enhance the FCNC branching ratios and can
 263 be probed at the LHC [14], from which some of them are shown in Table 1.7. Previous searches
 264 have been performed at the Fermilab Tevatron by the CDF [28] and D0 [29] collaborations,
 265 and at the LHC by the ATLAS [30–33] and CMS [34–38] collaborations.

266 The impact of BSM models can be written in a model independent way by means of an effective
 267 field theory valid up to an energy scale Λ . The leading effects are parametrized by a set of
 268 fully gauge symmetric dimension-6 operators that are added to the SM Lagrangian and can be
 269 reduced to a minimal set of operators as discussed in [39, 40]. The full Lagrangian, neglecting
 270 neutrino physics, in the fully gauge symmetric case is given by

$$\mathcal{L}_{\text{SM+EFT}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{\bar{c}_i}{\Lambda^2} \mathcal{O}_i + \mathcal{O}\left(\frac{1}{\Lambda^3}\right), \quad (1.20)$$

Table 1.7: The predicted branching ratios \mathcal{B} for FCNC interactions involving the top quark in some BSM models [14]: quark singlet (QS), generic two Higgs doublet model (2HDM) and the minimal supersymmetric extensions to the SM (MSSM);

Process	QS	2HDM	MSSM	Process	QS	2HDM	MSSM
$t \rightarrow uZ$	$\leq 1.1 \cdot 10^{-4}$	—	$\leq 2 \cdot 10^{-6}$	$t \rightarrow cZ$	$\leq 1.1 \cdot 10^{-4}$	$\leq 10^{-7}$	$\leq 2 \cdot 10^{-6}$
$t \rightarrow u\gamma$	$\leq 7.5 \cdot 10^{-9}$	—	$\leq 2 \cdot 10^{-6}$	$t \rightarrow c\gamma$	$\leq 7.5 \cdot 10^{-9}$	$\leq 10^{-6}$	$\leq 2 \cdot 10^{-6}$
$t \rightarrow ug$	$\leq 1.5 \cdot 10^{-7}$	—	$\leq 8 \cdot 10^{-5}$	$t \rightarrow cg$	$\leq 1.5 \cdot 10^{-7}$	$\leq 10^{-4}$	$\leq 8 \cdot 10^{-5}$
$t \rightarrow uH$	$\leq 4.1 \cdot 10^{-5}$	$\leq 5.5 \cdot 10^{-6}$	$\leq 10^{-5}$	$t \rightarrow cH$	$\leq 4.1 \cdot 10^{-5}$	$\leq 10^{-3}$	$\leq 10^{-5}$

where the Wilson coefficients \bar{c}_i depend on the considered theory and on the way that new physics couples to the SM particles. Considering that Λ is large, contributions suppressed by powers of Λ greater than two are neglected. Moreover, all four fermion operators are omitted for the rest of this thesis. After electroweak symmetry breaking the operators induce [14, 41] both corrections to the SM couplings and new interactions at tree level such as FCNC interactions. The FCNC interactions of the top quark that are not present in the SM are given by

$$\mathcal{L}_{\text{EFT}}^t = \frac{\sqrt{2}}{2} \sum_{q=u,c} \left[g' \frac{\kappa_{t\gamma q}}{\Lambda} A_{\mu\nu} \bar{t} \sigma^{\mu\nu} (f_{\gamma q}^L P_L + f_{\gamma q}^R P_R) q \right. \quad (1.21)$$

$$+ \frac{g}{2\cos\theta_W} \frac{\kappa_{tZq}}{\Lambda} Z_{\mu\nu} \bar{t} \sigma^{\mu\nu} (f_{Zq}^L P_L + f_{Zq}^R P_R) q \quad (1.22)$$

$$+ \frac{\sqrt{2}g}{4\cos\theta_W} \zeta_{tZq} \bar{t} \gamma^\mu (\tilde{f}_q^L P_L + \tilde{f}_q^R P_R) q Z_\mu \quad (1.23)$$

$$+ g_S \frac{\kappa_{gqt}}{\Lambda} Z_{\mu\nu} \bar{t} \sigma^{\mu\nu} (f_{gq}^L P_L + f_{gq}^R P_R) q G_{\mu\nu}^a \quad (1.24)$$

$$+ \eta_{Hqt} \bar{t} (\hat{f}_q^L P_L + \hat{f}_q^R P_R) q H + \text{h.c.} \Big], \quad (1.25)$$

where the value of the FCNC couplings at scale Λ are represented by $\kappa_{tZq}, \kappa_{gqt}, \kappa_{t\gamma q}, \zeta_{tZq}$, and η_{Hqt} . These are assumed to be real and positive, with the unit of GeV^{-1} for κ_{tXq}/Λ and no unit for ζ_{xqt} and η_{xqt} . In the equation $\sigma^{\mu\nu}$ equals to $\frac{i}{2} [\gamma^\mu, \gamma^\nu]$, and the left- and right-handed chirality projector operators are denoted by P_L and P_R . The electromagnetic coupling constant is denoted by g' , the strong interaction coupling is denoted as g_s , while the electroweak interaction is parametrised by the coupling constant g and the electroweak mixing angle θ_W . The complex chiral parameters are normalized according to $|f_{xq}^L|^2 + |f_{xq}^R|^2 = 1$, $|\tilde{f}_q^L|^2 + |\tilde{f}_q^R|^2 = 1$, and $|\hat{f}_q^L|^2 + |\hat{f}_q^R|^2 = 1$. In the expression for $\mathcal{L}_{\text{EFT}}^t$, the unitary gauge is adopted and the scalar field is expanded around its vacuum expectation value with H being the SM scalar boson, and the field strength tensors of the photon A_μ , the gluon field $G_\mu^{1\dots 8}$, and the Z boson Z_μ^0 are defined as

$$A_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu, \quad Z_{\mu\nu} = \partial_\mu Z_\nu - \partial_\nu Z_\mu, \quad \text{and} \quad G_{\mu\nu} = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_S f_{bc}^a G_\mu^b G_\nu^c. \quad (1.26)$$

271 Denoting the structure constant of the $SU_C(3)$ group as f_{bc}^a . Note that there are two coupling
 272 constants arising in $\mathcal{L}_{\text{EFT}}^t$, which is a residue of electroweak symmetry breaking. The massive Z
 boson will appear in both the Z_μ^0 field as well as the covariant derivative, leading to an extra
 Z-vertex.

275 1.8 Experimental constraints on top-FCNC

Experiments commonly put limits on the branching ratio's which allow an easier interpretation across different EFT models by use of the branching ratio \mathcal{B}

$$\mathcal{B}(t \rightarrow qX) = \frac{\delta_{tXq}^2 \Gamma_{t \rightarrow qX}}{\Gamma_t}, \quad (1.27)$$

276 where $\Gamma_{t \rightarrow qX}$ represents the FCNC decay width⁴ for a coupling strength $\delta_{tXq}^2 = 1$, and Γ_t the full
277 decay width of the top quark. In the SM, supposing a top quark mass of 172.5 GeV, the full
278 width becomes $\Gamma_t^{\text{SM}} = 1.32 \text{ GeV}$ [42].

279 Searches for top-FCNC usually adopt a search strategy depending on the experimental set-up
280 and the FCNC interaction of interest, looking either for FCNC interactions in the production of
281 a single top quark or in its decay for top pair interactions. In Figure 1.7, these two cases are
282 shown for the tZq vertex.



Figure 1.7: Feynman diagrams for the tZq FCNC interaction, where the FCNC interaction is indicated with the shaded dot. (a) Single top production through an FCNC interaction. (b) Top pair production with an FCNC induced decay.

283

284 The observation of top-FCNC interactions has yet to come and experiments have so far only
285 been able to put upper bounds on the branching ratios. An overview of the best current limits is
286 given in Table 1.8 . In Figure 1.8 a comparison is shown between the current best limits set by
287 ATLAS and CMS with respect to several BSM model benchmark predictions. From there one can
288 see that FCNC searches involving a Z or H boson are close to excluding or confirming several
289 BSM theories.

NOTE:
Check atlases result
for tZq from
top2017
proceedings
when they
appear

⁴The decay width of a certain process represents the probability per unit time that a particle will decay. The total decay width, defined as all possible decay widths of a particle, is inversely proportional to its lifetime.

Table 1.8: Overview of the most stringent observed and expected experimental limits on top-FCNC branching ratios \mathcal{B} at 95% confidence level.

Process	Search mode	Observed \mathcal{B}	Expected \mathcal{B}	Experiment	[Reference]
$t \rightarrow uZ$	top pair decay and single top production	$2.2 \cdot 10^{-4}$	$2.7 \cdot 10^{-4}$	CMS	[34]
$t \rightarrow u\gamma$	single top production	$1.3 \cdot 10^{-4}$	$1.9 \cdot 10^{-4}$	CMS	[36]
$t \rightarrow ug$	single top production	$4.0 \cdot 10^{-5}$	$3.5 \cdot 10^{-5}$	ATLAS	[31]
$t \rightarrow uH$	top pair decay	$2.4 \cdot 10^{-3}$	$1.7 \cdot 10^{-3}$	ATLAS	[33]
$t \rightarrow cZ$	top pair decay and single top production	$4.9 \cdot 10^{-4}$	$12 \cdot 10^{-4}$	CMS	[34]
$t \rightarrow c\gamma$	single top production	$2.0 \cdot 10^{-3}$	$1.7 \cdot 10^{-3}$	CMS	[36]
$t \rightarrow cg$	single top production	$2.0 \cdot 10^{-4}$	$1.8 \cdot 10^{-4}$	ATLAS	[31]
$t \rightarrow cH$	top pair decay	$2.2 \cdot 10^{-3}$	$1.6 \cdot 10^{-3}$	CMS	[33]




Figure 1.8: Current best limits set by CMS and ATLAS for top-FCNC interactions.

Experimental set-up

2

291 The main objective of the Large Hadron Collider (LHC) was the search for the Brout-Englert-
 292 Higgs boson. The Large Electron Positron (LEP) [43] and Tevatron [44] experiments had
 293 established that the mass of the scalar boson has to be larger than 114 GeV [45, 46], and smaller
 294 than approximate 1 TeV due to unitarity and perturbativity constraints [47]. On top of this,
 295 the search for new physics such as supersymmetry or the understanding of dark matter were
 296 part of the motivation for building the LHC. Since the start of its operation, the LHC is pushing
 297 the boundaries of the Standard Model, putting the most stringent limits on physics beyond the
 298 Standard Model as well as precision measurements of the parameters of the Standard Model. A
 299 milestone of the LHC is the discovery the scalar boson in 2012 by the two largest experiments
 300 at the LHC [7, 8].

301 This chapter is dedicated to the experimental set up of the LHC and the Compact Muon
 302 Solenoid (CMS) experiment. [Section 2.1](#) describes the LHC and its acceleration process for
 303 protons to reach their design energies. The CMS experiment and its components are presented
 304 in [Section 2.2](#).

305 2.1 The Large Hadron Collider

306 The LHC has started its era of cutting edge science on 10 September 2008 [48] after approval by
 307 the European Organisation of Nuclear Research (CERN) in 1995 [49]. Installed in the previous
 308 LEP tunnels, the LHC consists of a 26.7 km ring, that is installed between 45 and 170 m under
 309 the French-Swiss border amidst Cessy (France) and Meyrin (Switzerland). Built to study rare
 310 physics phenomena at high energies, the LHC can accelerate two type of particles, protons or
 311 ions Pb^{45+} , and provides collisions at four interaction points, where the particle bunches are
 312 crossing. Experiments for studying the collisions are installed on each interaction point.

313 As can be seen in [Figure 2.1](#), the LHC is last element in a chain that creates, injects and
 314 accelerates protons. The starting point is the ionisation of hydrogen, creating protons that are
 315 injected in a linear accelerator (LINAC 2). Here, the protons obtain an energy of 50 MeV. They
 316 continue to the proton synchrotron booster (PSB or Booster), where the packs of protons are
 317 accelerated to 1.4 GeV and each pack is split up in twelve bunches with 25 ns spacing for Run 2
 318 (50 ns for Run 1). The proton synchrotron (PS) then increases their energy to 25 GeV before the

super proton synchrotron (SPS) increases the proton energy up to 450 GeV. Each accelerator ring expands in radius in order to reduce the energy loss of the protons by synchrotron radiation¹. Furthermore, the magnets responsible for the bending of the proton trajectories have to be strong enough to sustain to higher proton energy. Ultimately, the protons are injected into opposite directions into the LHC, where they are accelerated to 3.5 TeV (in 2010 and 2011), 4 TeV (in 2012 and 2013) or 6.5 TeV (in 2015 and 2016) [50]. Before the start of the LHC in 2010, the previous energy record was held by the Tevatron collider at Fermilab, colliding proton with antiprotons at $\sqrt{s} = 1.96$ TeV. When completely filled, the LHC nominally contains 2220 bunches in Run 2, compared to 1380 in Run 1 (design: 2200).




Figure 2.1: Schematic representation of the accelerator complex at CERN [51]. The LHC (dark blue) is the last element in chain of accelerators. Protons are successively accelerated by LINAC 2, the Booster, the Proton Synchrotron (PS) and the Super Proton Synchrotron (SPS) before entering the LHC.

327

Inside the LHC ring [52], the protons are accelerated by the means of radio frequency cavities, while 1232 dipole magnets of approximately 15 m long, each weighing 35 t ensure the deflection of the beams. The two proton beams circulate in opposite direction in separate pipes inside of the magnet. Through the use of a strong electric current in the coils around the beam pipe, magnetic fields are generated and cause the protons to bend in the required orbits. In order to get the coil to become superconducting and able to produce - with the aid of an iron return yoke - a strong magnetic field of 8.3 T, the magnet structure is surrounded by a vessel. This vessel is filled with liquid Helium making it possible to cool down the magnet to 1.9 K. In order to get more focussed and stabilised proton beams, additional higher-order multipole and corrector magnets are placed along the LHC beam line.

¹This energy loss is proportional to the fourth power of the proton energy and inversely proportional to the bending radius.

338 The LHC is home to seven experiments, each located on an interaction point:

- 339 • A Toroidal LHC ApparatuS (ATLAS) [53] and the Compact Muon Solenoid (CMS) [54]
 340 experiments are the two general purpose detectors at the LHC. They both have a hermetic,
 341 cylindrical structure and were designed to search for new physics phenomena along with
 342 precision measurements of the Standard Model. The existence of two distinct experiments
 343 allows cross-confirmation of any discovery.
- 344 • A Large Ion Collider Experiment (ALICE) [55] and the LHC Beauty (LHCb) [56] exper-
 345 iments are focusing on specific phenomena. ALICE studies strongly interacting matter
 346 at extreme energy densities where a quark-gluon plasma forms in heavy ion collisions
 347 (Pb-Pb or p-Pb). LHCb searches for differences between matter and antimatter with the
 348 focus on b physics..
- 349 • The forward LHC (LHCf) [57] and the TOTal cross section, Elastic scattering and diffraction
 350 dissociation Measurement (TOTEM) [58] experiments are two smaller experiments that
 351 focus on head on collisions. LHCf consists of two parts placed before and after ATLAS
 352 and studies particles created at very small angles. TOTEM is placed in the same cavern as
 353 CMS and measures the total proton-proton cross section and studies elastic and diffractive
 354 scattering.
- 355 • The Monopoles and Exotics Detector At the LHC (MoEDAL) [59] experiment is situated
 356 near LHCb and tries to find magnetic monopoles.

For the enhancement of the exploration of rare events and thus enhancing the number of collisions, high beam energies as well as high beam intensities are required. The luminosity [60] is a measurement of the number of collisions that can be produced in a detector per square meter and per second and is the key role player in this enhancement. The LHC collisions create a number of events per second given by

$$N_{\text{event}} = L\sigma_{\text{event}}, \quad (2.1)$$

where σ_{event} is the cross section of the event of interest and L the machine luminosity. This luminosity depends only on the beam parameters and is for a Gaussian beam expressed as

$$L = \frac{1}{4\pi} N_b n_b f_{\text{rev}} \frac{N_b}{\epsilon_n} \left(1 + \left(\frac{\theta_c \sigma_z}{2\sigma^*} \right)^2 \right)^{-\frac{1}{2}} \frac{\gamma_r}{\beta^*}. \quad (2.2)$$

- 357 The number of particles per bunch is expressed by N_b , while n_b is the number of bunches
 358 per beam, f_{rev} the revolution frequency, γ_r the relativistic gamma factor, ϵ_n the normalized
 359 transverse beam emittance - a quality for the confinement of the beam , β^* the beta function at
 360 the collision point - a measurement for the width of the beam, θ_c the angle between two beams
 361 at the interaction point, σ_z the mean length of one bunch, and σ^* the mean height of one bunch.
 362 In Equation 2.2, the blue part represents the stream of particles, the red part the brilliance, and
 363 the green part the geometric reduction factor due to the crossing angle at the interaction point.

The peak design luminosity for the LHC reached in 2016 is $10^{34} \text{ m}^{-2}\text{s}^{-1}$, which leads to about 1 billion proton interactions per second. In 2016, the LHC was around 10% above this design luminosity [61]. The luminosity is not a constant in time since it diminishes due to collisions between the beams, and the interaction of the protons and the particle gas that is trapped in the centre of the vacuum tubes due to the magnetic field. The diffusion of the beam degrades the emittance and therefore also the luminosity. For this reason, the mean lifetime of a beam inside the LHC is around 15 h. The integrated luminosity - the luminosity provided in a certain time range - recorded by CMS and ATLAS over the year 2016 is given in Figure 2.2. In Run 2, the peak luminosity is $13\text{-}17 \cdot 10^{33} \text{ cm}^{-2}\text{s}^{-1}$ compared to $7.7 \cdot 10^{33} \text{ cm}^{-2}\text{s}^{-1}$ in Run 1.




Figure 2.2: Cumulative offline luminosity measured versus day delivered to (blue), and recorded by CMS (orange) during stable beams and for proton collisions at 13 TeV centre-of-mass energy in 2016. The delivered luminosity accounts for the luminosity delivered from the start of stable beams until the LHC requests CMS to turn off the sensitive detectors to allow a beam dump or beam studies.

372

Multiple proton-proton interactions can occur during one bunch crossing, referred to as pileup. On average, the number of pileup events is proportional to the luminosity times the total inelastic proton-proton cross section. In 2016, an average of about 27 of pileup interactions has been observed in 13 TeV proton collisions at the interaction point of CMS. For 2012, this number was about 21 pileup interactions for 8 TeV collisions.

2.2 The Compact Muon Solenoid

At one of the collision points of the LHC, the CMS detector[62–64] is placed. Weighing 14 000 t, this cylindrical detector is about 28.7 m long and 15 m in diameter. It has an onion like structure of several specialised detectors and contains a superconducting solenoid with a magnetic field of 3.8 T. Living in a hadronic environment, multi-jet processes produced by the strong interaction are a main source of background for rare physics processes. Therefore, good identification,

momentum resolution, and charge determination of muons, electrons and photons are one of the main goals of the CMS detector. Additionally, a good charged particle momentum resolution and reconstruction efficiency in the inner tracker provides identification for jets coming from b quarks or tau particles can be identified. Also the electromagnetic resolution for an efficient photon and lepton isolation as well as a good hadronic calorimeter for the missing transverse energy² were kept into account while designing CMS. In Figure 2.3, an overview of the CMS detector is shown.




Figure 2.3: Mechanical layout of the CMS detector. Figure taken from [65].

390

2.2.1 CMS coordinate system

The coordinate system used by CMS can be found in Figure 2.4. The origin of the right handed orthogonal coordinate system is chosen to be the point of collisions. The x-axis points towards the centre of the LHC ring such that the y-axis points towards the sky, and the z-axis lies tangent to the beam axis. Since the experiment has a cylindrical shape, customary coordinates are used to describe the momentum \vec{p} : the distance $p = |\vec{p}|$, the azimuthal angle³ $\phi \in [-\pi, \pi]$, the

²The missing transverse energy comes from an imbalance in the transverse plane. This will be discussed in Chapter 4.

³The azimuthal angle is the angle between the x-axis and the projection in the transverse plane of the momentum \vec{p} , denoted as \vec{p}_T .

pseudo-rapidity⁴ η :

$$\eta = -\ln \left(\tan \left(\frac{\theta}{2} \right) \right). \quad (2.3)$$

For the energies considered at the LHC, where $E \gg m$, the pseudo-rapidity is a good approximation of the rapidity y

$$y = \frac{1}{2} \ln \left(\frac{E + p_z}{E - p_z} \right), \quad (2.4)$$

- 392 where the difference of rapidities of two particles is invariant under a Lorentz boost in the z-direction.




Figure 2.4: Representation of the coordinate system used by CMS. The point of origin is put at the collision point. The x-axis points towards the centre of the LHC ring such that the z-axis lies tangent to the beam axis.

393

394 2.2.2 Towards the heart of CMS

- 395 The CMS detector can be divided into two parts. A central barrel is placed around the beam
 396 pipe ($|\eta| < 1.4$), and two plugs (end caps) ensure the hermeticity of the detector. In [Figure 2.3](#)
 397 and [Figure 2.5](#) the onion like structure of the CMS detector is visible. The choice of a solenoid of
 398 12.9 m long and 5.9 m diameter gives the advantage of bending the particle trajectories in the
 399 transverse plane. The hadronic calorimeter ([Section 2.2.2.3](#)), the electromagnetic calorimeter
 400 ([Section 2.2.2.4](#)) and the tracker ([Section 2.2.2.5](#)) are within the solenoid ([Section 2.2.2.2](#)),
 401 while the muon chambers ([Section 2.2.2.1](#)) are placed outside the solenoid. The data used for
 402 the search presented in this thesis is collected after the long shutdown 1. After discussing each
 403 part of CMS in their Run 1 configuration, [Section 2.2.4](#) elaborates on their different upgrades
 404 for the data collected in Run 2.

⁴The pseudo rapidity is expressed by the polar angle θ between the direction of \vec{p} and the beam.




Figure 2.5: Schematic view of the CMS detector in the Run I configuration. The longitudinal view of one quarter of the detector is given on top, while the transversal view is shown on the bottom. The muon system barrel elements are denoted as $MBZ/N/S$, where $z = -2 \dots +2$ is the barrel wheel number, $n = 1 \dots 4$ the station number and $S = 1 \dots 12$ the sector number. Similarly, the steel return yokes are denoted as $YBZ/N/S$. The solenoid is denoted as $CB0$, while the hadronic calorimeter is denoted as HE (end cap)/ HB (barrel)/ HF (forward) and the electromagnetic calorimeter as EE (end cap)/ EB (barrel). The green part represents the tracking system (tracker + pixel). Figure taken from [66]

2.2.2.1 Muon system

The outermost part of CMS consists of the muon system. The magnet return yoke is interleaved with gaseous detector chambers for muon identification and momentum measurement. The barrel contains muon stations arranged in five separate iron wheels, while in the end cap four muon stations are mounted onto three independent iron discs on each side. Each barrel wheel has 12 sectors in the azimuthal angle.

The muon system is divided into three parts, shown in Figure 2.6. The muon rate and neutron induced backgrounds are small and the magnetic field is very low for the barrel, thus CMS can use drift tube (DT) chambers. For the end caps however, the muon and background flux is much higher and there is a need to use cathode strip chambers (CSC) which are able to provide a faster response, higher granularity and have a better resistance against radiation. In order to form a redundant trigger system, resistive plate chambers (RPC) are added. This makes a total of 250 DT, 540 CSC and 610 RPC chambers. In Figure 2.5 the arrangement is shown.




Figure 2.6: (Left) Schematic view of one quarter of the CMS muon system in the Run I configuration. The cathode strip chambers (CSC) are shown in green, the drift tubes (DT) are shown in yellow, while the resistive plate chambers (RPC) are shown in blue. Figure taken from [66]. (Right) Cathode strip chambers (ME+4/2 chambers on YE+3). Photo taken from [67].

417

Providing a measurement for $|\eta| < 1.2$, the DT chambers in the barrel are on average $2 \times 2.5 \text{ m}^2$ in size and consist of 12 layers of DT cells⁵ arranged in three groups of four. The $r\phi$ coordinate is provided by the two outside groups, while the middle group measures the z coordinate. For the outer muon station, the DT chambers contain only 8 layers of DT cells, providing a muon position in the $r\phi$ plane. There are four CSC stations in each end cap, providing muon measurements for $0.9 < |\eta| < 2.4$ (Run I configuration). These CSCs are multi-wired proportional chambers that consist of 6 anode wire planes crossed by 7 copper strips cathode panels in a gas volume. The r coordinate is provided by the copper strips, while the ϕ coordinate comes from the anode wires, giving a two dimensional position measurement. There are six

⁵The DT cells are 4 cm wide gas tubes with positively charged stretched wires inside.

427 layers of RPCs in the barrel muon system and one layer into each of the first three stations
 428 of the end cap. They are made from two high resistive plastic plates with an applied voltage
 429 and separated by a gas volume. Read out strips mounted on top of the plastic plates detect the
 430 signal generated by a muon passing through the gas volume. The RPCs provide a fast response
 431 with a time resolution of 1 ns and cover a range of $|\eta| < 1.8$ for the Run 1 configuration.

432 The muon system provides triggering on muons, identifying muons and improves the momentum
 433 measurement and charge determination of high p_T muons. On top of the muon system,
 434 the muon energy is deposited in the electromagnetic calorimeter, the hadronic calorimeter, and
 435 outer calorimeter. The high magnetic field enables an efficient first level trigger and allows a
 436 good momentum resolution of $\Delta p/p \approx 1\%$ for a p_T of 100 GeV and $\approx 10\%$ for a p_T of 1 TeV.
 437 There is an efficient muon measurement up to $|\eta| < 2.4$.

NOTE:
check numbers for run
2

438 2.2.2.2 Solenoid

439 Making use of the knowledge of previous experiments like ALEPH and DELPHI at LEP and H1
 440 at HERA, CMS choose for a large super conducting solenoid with a length of 12.9 m and a
 441 inner bore of 5.9 m [64]. With 2 168 turns, a current of 19.5 kA and a total energy of 2.7 GJ, a
 442 large bending power can be obtained for a modestly-sized solenoid. In order to ensure a good
 443 momentum resolution in the forward regions, a favourable length/radius was necessary. In
 444 [Figure 2.7](#), a photo of the CMS solenoid is shown.

445 The solenoid uses a high-purity aluminium stabilised conductor with indirect cooling from
 446 liquid helium, together with fully epoxy impregnation. A four-layer winding is implemented that
 447 can withstand an outward pressure of 64 atm. The NbTi cable is co-extruded by pure aluminium
 448 that acts as a thermal stabilizer and has an aluminium alloy for mechanical reinforcement. The
 449 return of the magnetic field is done by fives wheels, noted by YB in [Figure 2.5](#).

450 2.2.2.3 Hadronic calorimeter

451 The hadronic calorimeter (HCAL) is dedicated to precisely measure the energy of charged and
 452 neutral hadrons via a succession of absorbers and scintillators. This makes it crucial for physics
 453 analyses with hadronic jets or missing transverse energy. The HCAL barrel extends between
 454 $1.77 < r < 2.95$ m, where r is the radius in the transverse plane with respect to the beam. Due
 455 to space limitations, the HCAL needs to be as small as possible and is made from materials
 456 with short interaction lengths⁶. On top of this, the HCAL should be as hermetic as possible and
 457 extend to large absolute pseudo rapidities such that it can proved a good measurement of the
 458 missing transverse energy.

459 The quality of the energy measurements is dependent on the fraction of the hadronic shower
 460 that can be detected. Therefore, the HCAL barrel (HB) inside the solenoid is reinforced by an
 461 outer hadronic calorimeter between the solenoid and muon detectors (HO, see [Figure 2.8](#)),
 462 using the solenoid as extra absorber. This increases the thickness to 12 interaction lengths.

⁶Here the interaction length is the nuclear interaction length and this is the length needed for absorbing 36.7% of the relativistic charged particles. For the electromagnetic calorimeter this is defined in radiation length X_0 . The radiation length is the mean distance over which a high energy electron loses all but $1/e$ of its energy by bremsstrahlung.




Figure 2.7: (Left) CMS solenoid during the long shutdown in 2013. (Right) An impression of the solenoid magnet taken from [68].

463 The HB and HO provide measurements for $|\eta| < 1.3$, while an end cap on each side (HE,
464 $1.3 < |\eta| < 3$) and a forward calorimeter (HF, $3.0 < |\eta| < 5.2$) extend the pseudo rapidity
465 range.




Figure 2.8: (Left) Longitudinal view of the CMS detector showing the locations of the HB, HE, HO, and HF calorimeters. Figure taken from [54]. (Right) CMS barrel calorimeter. Photo taken from [70].

466 The HB is made of 16 absorber plates where most of them are built from brass and others are
467 made from stainless steel and is about five to ten interaction lengths thick. It is divided in $\eta \times \phi$
468 towers and contains 2592 read out channels. The HO complements the HB and extends the
469 reach up to twelve interaction lengths. This subsystem contains 2160 read out channels. The HE

470 is also composed of brass absorber plates and has a thickness corresponding to approximately
 471 ten interaction lengths, with 2592 read out channels.. The HF experiences intense particle
 472 fluxes with an expected energy of 760 GeV deposited on average in a proton interaction at a
 473 centre-of-mass of 14 TeV, compared to 100 GeV in the rest of the detector. Therefore, these are
 474 Cherenkov light detectors made of radiation hard quartz fibers. The main causes of such large
 475 energy events are high energy muons, cosmic particles and charged particles from late showering
 476 hadrons. During Run 1, it became clear that the glass windows of the photon multiplier tubes
 477 (PMTs) had to be replaced which was done during LS1 [71]. The HF represents 1728 read out
 478 channels.

479 The HCAL and electromagnetic calorimeter combined can measure the hadron energy with a
 480 resolution $\Delta E/E \approx 100\% \sqrt{E[\text{GeV}]} + 5\%$.

481 2.2.2.4 Electromagnetic calorimeter

482 The electromagnetic calorimeter (ECAL) is designed to measure the energy of photons and
 483 electrons and covers $|\eta| < 3$. It is an hermetic, homogeneous detector and consists of 75 848
 484 lead tungstate (PbWO_4) crystals. These crystals have a fast response time - 80% of the light
 485 is emitted within 25 ns - and are radiation hard. The electromagnetic showers produced by
 486 passing electrons or photons ionize the crystal atoms which emit a blue-green scintillation light,
 487 that is collected by silicon avalanche photodiodes (APDs) in the barrel and vacuum phototriodes
 488 (VPTs) in the end caps. The crystals and the APD response is sensitive to temperature changes
 489 and require a stable temperature.




Figure 2.9: (Left) Schematic cross section of the electromagnetic calorimeter taken from [54]. (Right top) The ECAL barrel during construction [72]. (Right bottom) One half of an EE [73].

490 There are three regions: a central barrel (EB), an endcap region (EE) and a preshower (ES)
 491 (Figure 2.9). The EB has an inner radius of 129 cm and corresponds to a pseudo rapidity of $0 <$
 492 $|\eta| < 1.479$. At a distance of 314 cm from the vertex and covering a pseudo rapidity of $1.479 <$

493 $|\eta| < 3.0$, are the EE. They consist of semi-circular aluminium plates from which structural
 494 units of 5×5 crystals (super crystals) are supported. The ES is placed in front of the crystal
 495 calorimeter over the end cap pseudo rapidity range with two planes of silicon strip detectors as
 496 active elements.

The electromagnetic shower will typically involve more than one channel. More than 90% of the energy of a 35 GeV electron or photon is contained in a 5×5 matrix of crystals. Therefore, a clustering algorithm is performed in order to associate the energy deposits to the particles impinging the calorimeter. The achieved precision [74] for the barrel is 2.10^{-3} rad in ϕ and 10^{-3} in η . For the end caps this is 5.10^{-3} rad in ϕ and 2.10^{-3} in η . The energy is reconstructed by a supercluster algorithm, taking into account energy radiated via bremsstrahlung or conversion [54]. The energy resolution is given by

$$\frac{\sigma(E)}{E} = \frac{2.8\%}{\sqrt{E}} \oplus \frac{0.128}{E(\text{GeV})} \oplus 0.3\%, \quad (2.5)$$

497 in the absence of a magnetic field, where the contributions come from the stochastic, noise and
 498 constant terms respectively. The dominating term is the constant term ($E_{\text{shower}} \approx 100$ GeV)
 499 and thus the performance is highly dependent on the quality of calibration and monitoring .

500 2.2.2.5 Inner tracking system and operations

501 The tracking system (tracker) [76] is the detecting unit closest to the point of interaction.
 502 Responsible for the reconstruction of trajectories from charged particles with $|\eta| < 2.5$ that are
 503 bent by the magnetic field, it provides a measurement of the momentum. The tracker is also
 504 responsible for the determination of the interaction point or vertex. It should be able to provide
 505 high granularity as well as speed, and be able to endure high radiation. For this reason, the
 506 CMS collaboration choose silicon detector technology.

507 The tracking system consists of a cylinder of 5.8 m long and 2.5 m in diameter. It is immersed
 508 in a co-axial magnetic field of 3.8 T due to the solenoid. As shown Figure 2.10, the tracker
 509 is built up from a large silicon strip tracker with a small silicon pixel inside. The inner region,
 510 pixel ($4.4 < r < 10.2$ cm), gets the highest flux of particles. Therefore, pixel silicon sensors
 511 of 100×150 μm area used. It consists of three cylindrical barrels that are complemented by
 512 two discs of pixel modules at each side. The silicon strip tracker ($20 < r < 116$ cm) has three
 513 subdivisions. The Tracker Inner Barrel and Discs (TIB, TID, see Figure 2.12) are composed
 514 of four barrel layers accompanied by three discs at each end. The outer part of the tracker -
 515 Tracker Outer Barrel (TOB) - consists of 6 barrel layers. In the outer discs, there are nine discs
 516 of silicon sensors, referred to as Tracker End Caps (TEC).

517 The pixel, shown in Figure 2.11 has 1440 modules that cover an area of about 1 m^2 and have
 518 66 million pixels. It provides a three-dimensional position measurement of the hits arising from
 519 the interaction from charged particles with the sensors. In transverse coordinate ($r\phi$), the hit
 520 position resolution is about $10 \mu\text{m}$, while $20\text{-}40 \mu\text{m}$ is obtained in the longitudinal coordinate
 521 (z). The sensor plane position provides the third coordinate. The silicon strip trackers consists
 522 of 15 148 single sided modules placed in the TIB, TID and the first four rings of the TEC.
 523 They provide 9.3 million readout channels. In the TOB and the outer three rings of the TEC,
 524 double sided modules are used. These modules are constructed from two back-to-back single




Figure 2.10: Schematic cross section of the top half of the CMS tracking system in the $r\phi$ plane. The centre of tracker is shown with a star and corresponds to the approximate position of the proton collision point. The green dashed lines are an indication for each named tracker subsystem. The strip tracker modules that provide two-dimensional hits are shown by thin, black lines, while those able to reconstruct three-dimensional hit positions are shown by thick, blue lines. The pixel modules, shown in red, also provide three-dimensional hits. [63]

525 sided modules, where one module is rotated through a stereo angle. This covers an active area
 526 of about 198 m^2 . The TIB and TID provide position measurements in $r\phi$ with a resolution
 527 of approximately $13\text{-}38 \mu\text{m}$, while the TOB provides a resolution of about $18\text{-}47 \mu\text{m}$. The
 528 resolution in the z direction is approximately $230 \mu\text{m}$ in the TIB/TID and $530 \mu\text{m}$ in the TOB.
 529 To allow overlay and avoid gaps in acceptance, each module is shifted slightly in r or z with
 530 respect to its neighbouring modules within a layer. With this detector lay out, at least nine
 531 points per charged particle trajectory can be measured in an $|\eta|$ range up to 2.4.




Figure 2.11: The pixel barrel being re-installed after the Long Shutdown in 2015, around the beam pipe at CMS [77].



Figure 2.12: First half of the inner tracker barrel, consisting of three layers of silicon modules [78].

532 **2.2.3 Data acquisition**

533 At a design luminosity of $10^{34} \text{ 1/(m}^2 \text{ s)}$, the proton interaction rate exceeds 1 GHz. This makes
 534 it impossible for the CMS experiment to store all the data generated. For this, a two level trigger
 535 system has been put in place. The first level (Level-1) is a custom hardware system, while a
 536 second level (HLT) is software based running on a large farm of computers. In run II, with the
 537 increase in centre of mass energy and a higher luminosity, a larger number of simultaneous
 538 inelastic collisions per crossing is expected with respect to run I. For this, the CMS Level-1 has
 539 been upgraded [81].

540 **CMS Level-1 trigger**

541 The Level-1 trigger has to be a flexible, maintainable system, capable of adapting to the evolving
 542 physics programme of CMS [82]. Its output rate is restricted to 100 kHz imposed by the CMS
 543 readout electronics. It is implemented by custom hardware and selects events containing candi-
 544 date objects - eg ionization deposits consistent with a muon, or energy clusters corresponding
 545 to an electron / photon / tau lepton / missing transverse energy / jet. Collisions with large
 546 momenta can be selected by using scalar sum of the transverse momenta of the jets.

547 By buffering the raw data from the CMS subdetectors in front-end drivers, the level-1 trigger
 548 has a pipeline memory of 3.2 μs to decide whether to keep an event or reject it. The trigger
 549 primitives (TP) from the calorimeters and muon detectors are processed in several steps and
 550 combined into a global trigger. This information is then combined with the input from the other
 551 subsystems for the HLT. The separate inputs are synchronized to each other and the LHC orbit
 552 clock and sent to the global trigger module. Here, level-1 trigger algorithms are performed
 553 within 1 μs to decide whether to keep the event.

554 For run II, all hardware, software, databases and the timing control system have been replaced.
 555 The main changes are that the muon system now uses the redundancy of three muon detector
 556 system earlier to make a high resolution muon trigger. Other upgrades are that the calorimeter
 557 system isn't bound any more for streaming data the data and the global trigger has more level-1
 558 trigger algorithms.

559 **CMS HLT trigger**

560 The HLT is an array of commercially available computers with programmable menu that has
 561 output rate of on average 400 Hz for off-line event storage. The data processing is based on a
 562 HLT path. This is a set of algorithmic steps to reconstruct objects and make selections on them.
 563 Here, the information of all sub detectors can be used to perform algorithms on higher level
 564 reconstructed objects.

565 **2.2.4 Phase 1 upgrades**

566 Before the start of taking collision data for 13 TeV operations on 3 June 2015, CMS had a long
 567 shutdown (LS1)[80]. During this shutdown, the section of the beryllium beam pipe within CMS
 568 was replaced by a narrower one. This operation required the pixel to be removed and reinserted
 569 into CMS. In Run 2, higher particle fluxes with respect to Run 1 are expected. To avoid long
 570 damage caused by the intense particle flux at the heart of CMS, the tracker is been made ready

571 to operate at much lower temperature than during Run I. The electromagnetic calorimeter
 572 preshower system had been damaged during Run 1, therefore the preshower discs were removed,
 573 repaired and reinstalled successfully inside CMS in 2014. To help the discrimination between
 574 interesting low momentum muons coming from collisions and muons caused by backgrounds, a
 575 fourth triggering and measurement station for muons was added in each of the end caps. Several
 576 new detectors were installed into CMS for measuring the collision rate within the detector and
 577 monitors beam related backgrounds.

578 During the LS1, the muon system underwent major upgrades [83, 84]. In the fourth station
 579 of each end cap, the outermost rings of CSC and RPC chambers were completed, providing an
 580 angular coverage of $1.2 < |\eta| < 1.8$ for Run 2, increasing the system redundancy, and allowing
 581 tighter cuts on the trigger quality. In order to reduce the environmental noise, outer yoke discs
 582 have been placed on both sides for the end caps. At the innermost rings of the first station,
 583 the CSCs have been upgraded by refurbishing the readout electronics to make use of the full
 584 detector granularity instead of groups of three as was the case for Run 1. In Figure 2.6 (right),
 585 the refurbishing of the CSCs is shown.

586 Since the HF experiences intense particle fluxes, it became clear during Run 1 that the glass
 587 windows of the PMTs need replacing. For the ECAL in Run 1, the energy reconstruction happened
 588 via a weighted sum of the digitized samples [75]. For Run 2 however, the reconstruction had
 589 to be made more resistant for out of time pile up and a multi-fit approach has been set into
 590 place. In this approach, the pulse shape is modelled as a sum of one in-time pulse plus the out
 591 of time pulses [74]. The energy resolution is better than 2% in the central barrel region and
 592 2-5 % elsewhere.

During the first data taking period of the LHC (2010 to 2013), the tracker operated at +4°C. With the higher LHC beam intensities from 2015 onwards, the tracker needs to be operated at much lower temperatures. This is due to the fact with intense irradiation, the doping concentration changes, the leakage current increases proportional to the fluence and the charge collection efficiency decreases due to charge trapping. Mostly the leakage current (I) is affected by the temperature change:

$$I \propto T^2 e^{-\frac{E_g}{2kT}}, \quad (2.6)$$

593 where T is the operating temperature, E_g the band gap and k the Boltzmann constant. There is
 594 approximately a factor 15 between the leakage currents at room temperatures and at -10 °C.

595 During the LS1, the CMS cooling plant was refurbished[79](Figure 2.14) and the fluorocarbon
 596 cooling system overhauled. To help to suppress the humidity inside the tracker, new methods
 597 for vapour sealing and insulation were applied (Figure 2.13). Furthermore, several hundred
 598 high-precision sensors are used to monitor the humidity and temperature. In order to get as
 599 dry air as possible, a new dry-gas plant provides eight times more dry gas (air or nitrogen)
 600 than during the first run, and allows regulation if the flow. As final addition, the cooling
 601 bundles outside the tracker are equipped with heater wires and temperature sensors in order to
 602 maintain safe operations above the cavern dew point For the data taking in 2015-2016, the
 603 tracker operated at -15°C.




Figure 2.13: Tracker bulkhead being put into closed state with insulation pieces installed during an early trial in fall 2013



Figure 2.14: New Tracker high-capacity dry-gas plant with membrane separation system [80].

604 After the first half of Run 2, the innermost part of detection material in CMS (pixel) was
 605 upgraded by adding a fourth layer , enhancing the particle tracking capabilities of CMS. The
 606 data used in the framework of this thesis however is from before this upgrade.

607 2.2.5 CMS computing model

608 The selected data is stored, processed and dispersed via the Worldwide Large Hadron Collider
 609 GRID (WLCG)[85, 86]. This has a tiered structure that function as a single, coherent system:.

610 At CERN, a single Tier-0 is located. The raw data collected by CMS is archived here, and
 611 a first reconstruction of the data is done. This data is then already in a file format usable for
 612 physics analysis. Furthermore, it is able to reprocess data when new calibrations are made
 613 available. The Tier-0 site distributes this data to a total of seven Tier-1 centres. They carry out
 614 data reprocessing and store real data as well as simulated data. The Tier-1 further distribute
 615 the data to over 50 Tier-2 centres. These make the data accessible for physics analysis and are
 616 also being used for the production of simulated data. The data is made accessible for physicists
 617 around the world.

Analysis techniques

3

619 In order to disentangle the collisions coming from high energy experiments, the hadron collisions
 620 are disentangled. In [Section 3.1](#), the predictions behind hadron-hadron collision at high energies are
 621 presented. These are used to generate events via Monte Carlo event generators, explained
 622 in [Section 3.2](#). Machine learning helps to disentangle signal- and background like events.
 623 In [Section 3.3](#), the multivariate technique of boosted decision trees is explained. This yields
 624 powerful discriminants for separating signal and background events and provides distributions
 625 that go through template-based maximum likelihood fits. The fitting method used in the search
 626 presented in this thesis is discussed in [Section 3.4](#).

627 3.1 Hadron collisions at high energies

In hadron collisions at sufficiently high momentum transfer, all partons can be approximated as free making it possible to treat hadron-hadron scattering as a single parton-parton interaction. The momentum of the parton can then be expressed as a fraction of the hadron momentum

$$\vec{p}_{\text{parton}} = x \vec{p}_{\text{hadron}}, \quad (3.1)$$

where x is referred to as the Björken scaling variable. The interaction $p_A p_B \rightarrow X$ can then be factorised in terms of partonic cross sections $\hat{\sigma}_{ij \rightarrow X}$ [87]

$$\sigma_{p_A p_B \rightarrow X} = \sum_{ij} \iint dx_1 dx_2 f_i^A(x_1, Q^2) f_j^B(x_2, Q^2) d\hat{\sigma}_{ij \rightarrow X}, \quad (3.2)$$

628 where i and j are the partons resolved from protons A and B, $f_i(x_i, Q^2)$ the parton density
 629 functions (PDF), and Q^2 the factorisation scale more commonly denoted as μ_F . The factorisation
 630 scale is the scale at which the hadronic interaction can be expressed as a product of the partonic
 631 cross section and the process independent PDF. In [Figure 3.1](#), the kinematic regions in x and
 632 μ_F are shown for fixed target and collider experiments.

633 The parton density functions (PDF) [88–90] give the momentum distribution of the proton
 634 amongst its partons at an energy scale μ_F . These functions can not be determined from first
 635 principles and have to be obtained from global fits to data. The PDFs are obtained from measure-
 636 ments on deep inelastic scattering using lepton-proton collision by the HERA collider [91],




Figure 3.1: Kinematic regions in momentum fraction x and factorisation scale Q^2 probed by fixed-target and collider experiments. Some of the final states accessible at the LHC are indicated in the appropriate regions, where y is the rapidity. In this figure, the incoming partons have $x_{1,2} = (M/14\text{TeV})e^{\pm y}$ with $Q = M$ where M is the mass of the state shown in blue in the figure. For example, exclusive J/ψ and Υ production at high $|y|$ at the LHC may probe the gluon PDF down to $x \sim 10^{-5}$. Figure taken from [4].

637 supplemented with proton-antiproton collisions from Tevatron at Fermi lab [92], and proton
 638 collision data from the ATLAS, CMS and LHCb collaborations at the LHC (Run 1) [93]. These
 639 measurements are included in global PDF sets known as the PDF4LHC recommendation [90].
 640 From their measurement at scale μ_F these PDFs can be extrapolated using the DGLAP equations
 . The PDFs are used to calculate the cross section of a certain process and are therefore used
 as input for the Monte Carlo generators used to make the simulated data samples at the LHC.

NOTE: A 641
 source 642
 643 In the framework of this thesis, the NLO PDF4LHC15_100 set is used. This set is an envelope
 644 of three sets, CT14, MMHT2014 and NNPDF3.0 [90]. In Figure 3.2 the dependency of the PDFs
 645 on the momentum fraction x is shown for the NNPDF3.0 set on hadronic scale ($\mu_F^2 = (10\text{GeV})^2$)
 646 and LHC scale ($\mu_F^2 = (10^4\text{GeV})^2$). For most values of the momentum fraction, the gluon density
 647 dominates, meaning that it is easier to probe muons than the quarks. For x close to one, the
 648 parton densities of the up and down quarks (the valence quarks of the proton) dominate over
 649 the gluon density. The charm, anti-up, and anti-down quarks have lower densities in general
 650 since those are sea quarks which originate in the proton only through gluon splitting. The
 651 resolution scale Q^2 is typically taken to be the energy scale of the collision. For the top quark pair
 652 production a scale of $Q^2 = (350\text{GeV})^2$ is chosen, meaning that the centre-of-mass energy of the
 653 hard interaction is about twice the top quark mass. The uncertainty on the parton distributions
 654 is evaluated using the Hessian technique [94], where a matrix with a dimension identical to the
 655 number of free parameters needs to be diagonalised. In the case of PDF4LHC15_100 set, this
 656 translates into 100 orthonormal eigenvectors and 200 variations of the PDF parameters in the
 657 plus and minus direction.

At high energies divergences can appear from quantum fluctuations. For the theory still to be




Figure 3.2: The momentum fraction x times the parton distribution functions $f(x)$, where $f = u_v, d_v, \bar{u}, \bar{d}, s, c$, or g as function of the momentum fraction obtained in the NNLO NNPDF3.0 global analysis at factorisation scales $\mu^2 = 10 \text{ GeV}^2$ (left) and $\mu^2 = 10^4 \text{ GeV}^2$ (right), with $\alpha_s(M_Z^2) = 0.118$. The gluon PDF has been scaled down by a factor of 0.1. Figure taken from [4].

able to describe the experimental regime, a renormalization scale μ_R is used to redefine physical quantities. A consequence of this method is that the coupling constants will run as function of μ_R . Beyond this scale, the high energy effects such as the loop corrections to propagators (self energy) are absorbed in the physical quantities through a renormalization of the fields. In particular the running behaviour of the strong coupling constant¹ α_s is found to be

$$\alpha_s = \frac{\alpha_s(\mu_0^2)}{1 + \alpha_s(\mu_0^2) \frac{33-2n_f}{12\pi} \ln\left(\frac{|\mu_R^2|}{\mu_0^2}\right)}, \quad (3.3)$$

with n_f the number of quarks and μ_0 the reference scale on which the coupling is known. The current world average of the strong coupling constant at the Z boson mass is $\alpha_s(\mu_F = m_Z) = 0.1181 \pm 0.0011$ [4]. From Equation 3.3 one can see easily that the coupling strength decreases with increasing renormalization scale, this known as asymptotic freedom. Additionally, following the behaviour of $\alpha_s(\mu_R^2)$, a limit $\Lambda_{\text{QCD}} \approx 200 \text{ MeV}$ is found for which α_s becomes larger than one. Under this limit, the perturbative calculations of observables can no longer be done.

Cross sections be written in terms of interacting vertices contributing to the matrix element (ME) originating from elements of a perturbative series [95], allowing them to be expanded as

¹The strong coupling constant is defined as $\alpha_s = \frac{g_s^2}{4\pi}$.

a power series of the coupling constant α

$$\sigma = \sigma_{\text{LO}} \left(1 + \left(\frac{\alpha}{2\pi} \right) \sigma_1 + \left(\frac{\alpha}{2\pi} \right)^2 \sigma_2 + \dots \right). \quad (3.4)$$

664 Leading order (LO) accuracy contains the minimal amount of vertices in the process, then
 665 depending on where the series is cut off one speaks of next-to-leading order (NLO), or next-
 666 to-next-to-leading order (NNLO) accuracy in α . Predictions including higher order correction
 667 tend to be less affected by theoretical uncertainties originating from a variation of the chosen
 668 renormalization and factorisation scales.

669 3.2 Event generation

670 In order to compare reconstructed data with theoretical predictions, collision events are gen-
 671 erated and passed through a simulation of the CMS detector and an emulation of its readout.
 672 For the detector simulation, a so-called Full Simulation package [96, 97] based on the Geant4
 673 toolkit [98] is employed. It allows a detailed simulation of the interactions of the particles with
 674 the detector material.

675 3.2.1 Fundamentals of simulating a proton collision

676 The procedure of to generate $\text{pp} \rightarrow \text{X}$ events can be subdivided into sequential steps [99–101],
 677 as shown in Figure 3.3.

678 The interaction of two incoming protons is often soft and elastic leading to events that are not
 679 interesting in the framework of this thesis. More intriguing are the hard interaction between two
 680 partons from the incoming protons. The matrix elements of a hard scattering process of interest
 681 is the starting point of the generation of events. Monte Carlo techniques are used to sample the
 682 corresponding cross section integral and the resulting sample of events reflect the probability
 683 distribution of a process over its final state phase space. After obtaining the sample of events of
 684 the hard interaction, a parton shower (PS) program is used to simulate the hadronisation of
 685 final state partons into hadrons which then decay further. Additionally, radiation of soft gluons
 686 or quarks from initial or final state partons is simulated. These are respectively referred to as
 687 initial state radiation (ISR) or final state radiation (FSR). Contributions from soft secondary
 688 interactions, the so-called underlying event (UE), and colour reconnection effects are also taken
 689 into account. A brief overview of the employed programs used for the event generation of the
 690 signal and main background processes used in the search presented in the thesis are given in
 691 Section 3.2.2.

NOTE: Should I add more details?

692 3.2.2 Programs for event generation

693 The FEYNRULES package [102] allows the calculation of the Feynman rules in momentum space
 694 for any quantum field theory model. By use of a Lagrangian, the set of Feynman rules associated
 695 with this Lagrangian are calculated. Via the Universal FeynRules Output (UFO) [103] the
 696 results are then passed to matrix element generators.

697 The MadGraph program [104] is used to interpret the physics model and calculate the cor-
 698 responding Feynman diagrams and matrix elements. After this, MadEvent [105] is used to




Figure 3.3: Sketch of a hadron collision as simulated by a Monte-Carlo event generator. The red blob in the centre represents the hard collision, surrounded by a tree-like structure representing Bremsstrahlung as simulated by parton showers. The purple blob indicates a secondary hard scattering event. Parton-to-hadron transitions are represented by light green blobs, dark green blobs indicate hadron decays, while yellow lines signal soft photon radiation. Figure taken from [101].

699 calculate the corresponding partons. These generated parton configurations are then merged
 700 with Pythia [106–108] parton showers using the MLM merging scheme [109].

701 The MadGraph5_aMC@NLO program [110] combines the LO MadGraph [104] and the aMC@NLO
 702 program into a common framework. This combination supports the generation of samples at
 703 LO or next to NLO together with a dedicated matching to parton showers using the MLM [109]
 704 or FXFX [111] schemes respectively. The FXFX scheme produces a certain fraction of events
 705 with negative weights originating from the subtraction of amplitudes that contain additional
 706 emissions from the NLO matrix element to prevent double-counting.

707 The POWHEG box (versions 1,2) [112–117] contains predefined implementations of various
 708 processes at NLO. It applies the POWHEG method for ME- to PS- matching, where the hardest
 709 radiation generated from the ME has priority over subsequent PS emission to remove the overlap
 710 with the PS simulation.

711 The JHU generator (version 7.02) [118–121] is used to generate the parton level information
 712 including full spin and polarization correlations. It is commonly used for studying the spin and
 713 parity properties of new resonances such as $ab \rightarrow X \rightarrow VV$, where $V = Z, W, \gamma$.

714 The generation of events from processes involving the production and decay of resonances
 715 creates a computational heavy load, especially at NLO. The narrow width approximation the

resonant particle is assumed to be on-shell. This makes the production and decay amplitude factorize, allowing to perform the simulation of the production and decay of heavy resonances like top quarks or Higgs bosons to be performed in separate steps. The MadSpin program [122] extends this approach and accounts for off-shell effects through a partial reweighting of the events. Additionally, spin correlation effects between production and decay products are taken into account.

The Pythia program (versions 6,8) [106–108] generates events of various processes at LO. Usually in the analysis, it is however only used for its PS simulation and it is interfaced with other LO and NLO event generators to perform subsequent parton showering, hadronisation, and simulation of the underlying event. In this thesis the underlying event tunes [123] are the CUETP8M2T4, CUETP8M1 and CUETP8M2.

The detector response is simulated via the Geant4 [98] program. This program tracks the particles through the detector material via a detailed description of the detector and generates several hits throughout several sensitive layers. In addition, the response of the detector electronics to these hits are simulated.

3.2.3 Generating FCNC top-Z interactions

The FCNC processes are generated by interfacing the Lagrangian in Equation 1.25 with MadGraph5_aMC@NLO by means of the FeynRules package and its Universal FeynRules Output format. The complex chiral parameters are arbitrary chosen to be $f_{Xq}^L = 0$ and $f_{Xq}^R = 1$. The signal rates are estimated by use of the MadGraph5_aMC@NLO program for estimating the partial widths. The anomalous couplings are left free to float for this estimation, and only one coupling allowed to be non-vanishing at a time. The results are presented in Table 3.1. The

NOTE: Why
LH and not
RH?

Table 3.1: Leading order partial widths related to the anomalous decay modes of the top quark, where the new physics scale Λ is given in GeV.

Anomalous coupling	vertex	Partial decay width (GeV)	
κ_{gqt}/Λ	t g u	$3.665220 \cdot 10^5$	$(\kappa_{tg u}/\Lambda)^2$
	t g c	$3.664620 \cdot 10^5$	$(\kappa_{tg c}/\Lambda)^2$
$\kappa_{t\gamma q}/\Lambda$	t γ u	$1.989066 \cdot 10^4$	$(\kappa_{t\gamma u}/\Lambda)^2$
	t γ c	$1.988904 \cdot 10^4$	$(\kappa_{t\gamma c}/\Lambda)^2$
κ_{tZq}/Λ	t Z u	$1.637005 \cdot 10^4$	$(\kappa_{tZ u}/\Lambda)^2$
	t Z c	$1.636554 \cdot 10^4$	$(\kappa_{tZ c}/\Lambda)^2$
ζ_{tZq}	t Z u	$1.685134 \cdot 10^{-1}$	$(\zeta_{tZ u})^2$
	t Z c	$1.684904 \cdot 10^{-1}$	$(\zeta_{tZ c})^2$
η_{tHq}	t H u	$1.904399 \cdot 10^{-1}$	$(\eta_{tH u})^2$
	t H c	$1.904065 \cdot 10^{-1}$	$(\eta_{tH c})^2$

738 anomalous single top cross sections are calculated by convolution of the hard scattering matrix
 739 elements with the LO order set of CTEQ6 partons densities [124]. The NLO effects are modelled
 740 by multiplying each LO cross section by a global k -factor. The LO single top production cross
 741 section and the global k -factors for the top-Z production are shown in Table 3.2. The hard
 742 scattering events are then matched to parton showers to Pythia to account for the simulation
 of the QCD environment relevant for hadronic collisions.

Table 3.2: Leading order single top production cross section for $pp \rightarrow tZ$ or $\bar{t}Z$, where the new physics scale is given in GeV. The NLO k -factors [125] are given in the last column.

Anomalous coupling	Cross section (pb)	NLO k -factor
κ_{tg_u}/Λ	$3.272 \cdot 10^7$	$(\kappa_{tg_u}/\Lambda)^2$
κ_{tg_c}/Λ	$3.021 \cdot 10^6$	$(\kappa_{tg_c}/\Lambda)^2$
$\kappa_{t\gamma_u}/\Lambda$	$2.260 \cdot 10^5$	$(\kappa_{t\gamma_u}/\Lambda)^2$
$\kappa_{t\gamma_c}/\Lambda$	$2.654 \cdot 10^4$	$(\kappa_{t\gamma_c}/\Lambda)^2$
κ_{tZ_u}/Λ	$1.728 \cdot 10^6$	$(\kappa_{tZ_u}/\Lambda)^2$
κ_{tZ_c}/Λ	$2.040 \cdot 10^5$	$(\kappa_{tZ_c}/\Lambda)^2$
ζ_{tZ_u}	7.484	$(\zeta_{tZ_u})^2$
ζ_{tZ_c}	1.038	$(\zeta_{tZ_c})^2$

743

The top pair cross sections are derived from the SM $t\bar{t}$ cross section, calculated with MadGraph5_aMC@NLO at NLO ($\sigma_{t\bar{t}} = 6.741 \cdot 10^2$ pb), and considering the decay $t\bar{t} \rightarrow (bW^\pm)(X_{qt})$. The branching ratio $\mathcal{B}(t \rightarrow bW^\pm)$ is assumed to be equal to one and the FCNC branching ratio is calculated as

$$\mathcal{B}(t \rightarrow qX) = \frac{\Gamma_{t \rightarrow qX}}{\Gamma_t^{\text{SM}} + \Gamma_t^{\text{FCNC}}} \approx \frac{\Gamma_{t \rightarrow qX}}{\Gamma_t^{\text{SM}}}, \quad (3.5)$$

744 where $\Gamma_{t \rightarrow qX}$ is given in Table 3.1, and the assumption $\Gamma_t^{\text{FCNC}} \ll \Gamma_t^{\text{SM}}$ is made. In Table 3.3 the
 745 resulting NLO cross sections for the top-Z FCNC interactions are given.

746 3.2.4 Generating SM background events

747 The SM tZ events were generated using the MadGraph5_aMC@NLO generator, interfaced with
 748 Pythia version 8.2 [108] for parton showering and hadronisation. The $WZ + \text{jets}$, $t\bar{t}Z$, tZq ,
 749 and $t\bar{t}W$ samples are produced using the MadGraph5_aMC@NLO(version 5.222) [110], which
 750 includes up to one hadronic jet at next to leading order (NLO) QCD accuracy. Other minor
 751 background (e.g. WW , ZZ , tWZ and $t\bar{t}H$) are simulated using different generators such as
 752 MadGraph [104], MadSpin [122] and JHU [118–121]. All events are interfaced to Pythia for
 753 parton shower and hadronisation.

NOTE:
these partial widths
are at LO,
how does
this relate
to NLO that
is used? Or
is there no
difference?

The complete list of SM samples is given in Table 3.4, along with their cross sections. The
 cross sections without a reference are coming from the generator with which the sample has

NOTE: Add
source

Table 3.3: Next to leading order top pair cross section for the top-Z FCNC interactions with with a full leptonic decay.

Anomalous coupling	Process	Cross section (pb)
κ_{tZu}/Λ	$t\bar{t} \rightarrow (b\ell^+\nu)(\bar{u}\ell^+\ell^-)$	$2.727008 \cdot 10^5 \left(\kappa_{tZu}/\Lambda\right)^2$
	$t\bar{t} \rightarrow (\bar{b}\ell^-\bar{\nu})(u\ell^+\ell^-)$	$2.727008 \cdot 10^5 \left(\kappa_{tZu}/\Lambda\right)^2$
κ_{tZc}/Λ	$t\bar{t} \rightarrow (b\ell^+\nu)(\bar{c}\ell^+\ell^-)$	$2.72625710^5 \left(\kappa_{tZc}/\Lambda\right)^2$
	$t\bar{t} \rightarrow (\bar{b}\ell^-\bar{\nu})(c\ell^+\ell^-)$	$2.726257 \cdot 10^5 \left(\kappa_{tZc}/\Lambda\right)^2$
ζ_{tZu}	$t\bar{t} \rightarrow (b\ell^+\nu)(\bar{u}\ell^+\ell^-)$	$2.827184 \left(\zeta_{tZu}\right)^2$
	$t\bar{t} \rightarrow (\bar{b}\ell^-\bar{\nu})(u\ell^+\ell^-)$	$2.827184 \left(\zeta_{tZu}\right)^2$
ζ_{tZc}	$t\bar{t} \rightarrow (b\ell^+\nu)(\bar{c}\ell^+\ell^-)$	$2.806801 \left(\zeta_{tZc}\right)^2$
	$t\bar{t} \rightarrow (\bar{b}\ell^-\bar{\nu})(c\ell^+\ell^-)$	$2.806801 \left(\zeta_{tZc}\right)^2$

been made, for some of them the uncertainties are provided by the Generator Group. For each MC sample, the integrated luminosity that the sample represents is estimated as the number of simulated events divided by the cross section of the generated process. For processes generated with MadGraph5_aMC@NLO, the effective number of simulated events is used, taking into account positive and negative event weights. The correction factor for those events is defined as

$$C = \frac{\text{Nb. of pos. weights} + \text{Nb. of neg. weights}}{\text{Nb. of pos. weights} - \text{Nb. of neg. weights}} \times \text{mc baseweight} \quad (3.6)$$

Table 3.4: SM MC samples used in this analysis with their corresponding cross section and MadGraph5_aMC@NLO correction C when applicable. The generators used for each sample are indicated.

Process	Generator	Cross section (pb)	C
$WZ \rightarrow 3\ell\nu$	MadGraph5_aMC@NLO+Pythia	5.26	1.61
tZq with $Z \rightarrow \ell^+\ell^-$	MadGraph5_aMC@NLO+Pythia	0.0758	3.77
tqH with $H \rightarrow ZZ \rightarrow \ell^+\ell^-\ell^+\ell^-$	JHU+Pythia	$8.80 \cdot 10^{-6}$	-
$t\bar{t}W + \text{jets}$ with $W \rightarrow \ell\nu$	MadGraph5_aMC@NLO+MadSpin+Pythia	0.2043 ± 0.0020	1.94
$t\bar{t}Z \rightarrow 2\ell + 2\nu + \text{other}$, with $m_{\ell\ell} > 10$ GeV	MadGraph5_aMC@NLO+Pythia	0.2529 ± 0.0004	2.15
$t\bar{t}H, \text{no } b\bar{b} \text{ decays}$	POWHEG+Pythia	0.2151	-
$t\bar{t}H, b\bar{b} \text{ decays}$	POWHEG+Pythia	0.2934	-
$WW \rightarrow 2\ell 2\nu$	POWHEG+Pythia	12.178	-
$ZZ \rightarrow 4\ell$	POWHEG+Pythia	0.3366	-
WZZ	MadGraph5_aMC@NLO+Pythia	0.05565	1.14
ZZZ	MadGraph5_aMC@NLO+Pythia	0.01398	1.17
single top tWZ , with $Z_\mu \rightarrow \ell^+\ell^-$	MadGraph+Pythia	0.001123	-
single top t-channel \bar{t}	POWHEG+MadSpin+Pythia	$44.33^{+1.76}_{-1.49}$	-
single top t-channel t	POWHEG+MadSpin+Pythia	$26.38^{+1.32}_{-1.18}$	-
single top $\bar{t}W$	POWHEG+Pythia	$35.85 \pm 0.90 \text{ (scale)} \pm 1.70 \text{ (PDF)}$	-
single top tW	POWHEG+Pythia	$35.85 \pm 0.90 \text{ (scale)} \pm 1.70 \text{ (PDF)}$	-
$t\bar{t}$	POWHEG+Pythia	$831.76^{+19.77+35.06}_{-29.20-35.06}$	-
$Z/\gamma^* + \text{jets}$, with $m_{\ell\ell} > 50$ GeV	MadGraph5_aMC@NLO+Pythia	$3 \times (1921.8 \pm 0.6 \pm 33.2)$	1.49
$Z/\gamma^* + \text{jets}$, with $10 \text{ GeV} < m_{\ell\ell} < 50 \text{ GeV}$	MadGraph+Pythia	18610	-

754 3.3 Multivariate analysis techniques: Boosted Decision Trees

755 The need of processing large quantities of data and discriminating between events with largely
 756 similar experimental signatures makes multivariate statistical analysis (MVA) a largely used
 757 method in the physics community. Multivariate classification methods based on machine
 758 learning techniques are a fundamental ingredient to most analyses. The advantage of using
 759 a MVA classifier is that it can achieve a better discrimination power with respect to a simple
 760 cut and count analysis with a poorly discriminating variables. These variables are referred to
 761 as weak variables and have similar distributions for signal and background samples. A risk of
 762 using MVA classifiers is overtraining. This happens when there are too many model parameters
 763 of an algorithm adjusted to too few data points. This leads to an increase in the classification
 764 performance over the objectively achievable one.

765 There are many software tools that exist for MVA. In this thesis the Tool for Multivariate
 766 Analysis (TMVA) [126] is used. This software is an open source project included into
 767 ROOT [127]. All multivariate techniques in TMVA belong to supervised learning algorithms. By
 768 training on events for which the outcome is known, a mapping function is determined that
 769 describes a classification or an approximation of the underlying behaviour defining the target
 770 value (regression).

771 In this thesis boosted decision trees (BDT) are employed for the classification of events as
 772 implemented in the TMVA framework [126]. This multivariate technique is based on a set of
 773 decision trees where each yields a binary output depending on the fact that an event is signal- or
 774 background-like. The advantage of such a multivariate technique is that several discriminating
 775 variables can be combined into a powerful one-dimensional discriminant D.

In Figure 3.4 a schematic view of a decision tree is shown. The starting point is the root node. Then a consecutive set of a total of i questions (nodes) regarding discriminating variables x_i are asked with only two possible answers per question (binary splits). The decision tree is constructed by training on a dataset for which the outcome is already provided, such as simulation dataset with signal and background processes (supervised learning). For each node a criterion $x_i > C_i$ is found by maximizing the separation gain between nodes

$$\text{separationgain} \approx \text{gain}(\text{parent}) - \text{gain}(\text{daughter, Signal}) - \text{gain}(\text{daughter, Background}), \quad (3.7)$$

with the gain computed using the Gini index

$$\text{gain}(\text{cell}) \approx p(1-p), \quad (3.8)$$

776 where p denotes the purity of a selection $x > C$. This is repeated until the maximum of nodes is
 777 reached and at the end of the sequence, the leaf nodes are labelled either signal S or background
 778 B, depending on the majority of events that end up on those nodes.

Different trees can be combined into a forest where the final output is determined by the majority vote of all trees, forming the sum of so-called weak learners into one strong learner. From one training collection, trees are derived by reweighting events, and combined into a single classifier as the weighted average of each individual decision tree. A method for making such forests is boosting a tree. In this method, misclassified events are weighted higher so




Figure 3.4: Schematic view of a decision tree. Figure taken from [126].

that future learner concentrate on these events. This has as advantage that the response of the decision trees are stabilised against fluctuations in the training sample which enhances the performance. Additionally, the trees can be kept very shallow, in this thesis $i = 3$, which improves the robustness against overtraining. Examples of such boosting algorithms are Adaptive Boosting (AdaBoost) and Gradient Boosting [128]. In AdaBoost, each weight of the misclassified events are enhanced while reducing the weight of correctly classified events after each training such that future events learn those better

$$\alpha_{n+1} = \left(\frac{1 - \epsilon_n}{\epsilon_n} \right)^\beta, \quad (3.9)$$

where ϵ_n denotes the misclassification error of the current tree n and β is a learning rate. The weight w_i at node i is then equal to $w_i = \ln \alpha_i$. The final weight is the sum of all classifiers weighted by their errors. The learning rate is typically chosen to be $\beta \leq 0.5$ to allow more boosting steps. Gradient boosting has a similar approach and combines a gradient descent with boosting. Instead of fitting the base-learner to the reweighted data as in AdaBoost, it is fitted to the negative gradient vector of the loss function evaluated at the previous node. Misclassified events will result in a majority vote with large gradients of the loss function. Also for the Gradient boost, the learning rate is typically slow, this also known as shrinkage. In this thesis Gradient boost is used with a shrinkage of 0.2-0.3.

In this thesis, the Gradient boost is used in combination with bagging, so-called stochastic gradient boosting. Bagging is a resampling technique draws a subset of events from the training data where the same event is allowed to be randomly picked several times from the parent sample. The tree is then trained on this subset and this is repeated many times. It is based on the assumption that sampling from a dataset that follows a distribution is the same as sampling from the distribution itself [129]. If one draws an event out of the parent sample, it is more likely to draw an event out of the phase space that has a high probability density, as the original dataset will have more events in the regions. Since the selected event is kept in the original sample, the parent sample stays unchanged so that randomly extracted samples have

797 the same parent distribution, albeit statistically fluctuated. Bagging smears over the statistical
 798 fluctuations in the training data, making it suitable for stabilising the response of the classifier
 799 and increasing the performance by eliminating overtraining. In stochastic gradient boosting the
 800 bagging resampling procedure uses random sub-samples of the training events for growing the
 801 trees.

802 The discriminating power of a BDT is assessed by analysing the receiver operating statistics
 803 (ROC) curve. This curves show the background rejection over the signal efficiency of the
 804 remaining sample. By looking at the area under the curve with respect to random guessing
 805 (AUC), the best classifier can be identified. This follows the Neyman-Pearson lemma that
 806 the best ROC curve is given by the likelihood ratio $\mathcal{L}(x|Signal)/\mathcal{L}(x|Background)$ [129]. No
 807 discrimination power will result in an AUC of 0%, while 50% means fully separated event
 classes. In [Figure 3.5](#) an example of ROC curve is shown.




Figure 3.5: Example of ROC curves. In this example, the green method is better than the red one, which is better than the blue one. The dashed line represents a case where there is no separation. Figure taken from [130].

808

809 3.4 Statistical methodology

810 The search performed in the framework of this thesis requires the simultaneous analysis of data
 811 from different decay channels. The statistical methodology used for this search is developed
 812 by the ATLAS and CMS collaborations in the context of the LHC Higgs Combination group.
 813 The description of the methodology can be found in Refs. [131–134]. The Higgs Combined
 814 Tool [135] is a RooStats [136] framework which runs different statistical methods. In this
 815 section, only the statistical tools necessary for the performed search are described. The results
 816 presented in this thesis are obtained using the asymptotic formulae [137].

817 In general the event yields of signal and background processes are denoted as s and b
 818 respectively. These represent event counts in multiple bins or for unbinned probability density

functions . By use of simulation, predictions on both signal and background yields are made. These predictions are subject to multiple uncertainties that are accounted for by introducing nuisance parameters θ such that $s = s(\theta)$ and $b = b(\theta)$. In the following, the actual observed events are denoted as data or observation.

3.4.1 The absence of signal: limits

The absence of a signal is characterised in high energy physics by the Bayesian and modified classical frequentist statistical approaches. They allow to quantify the level of incompatibility of data with a signal hypothesis in terms of confidence levels (CL). The convention is to require a 95% CL for excluding a signal.

An analysis targeting a certain signal production mechanism can either set approximate model-independent limits on signal cross sections times branching ratio ($\sigma \times \mathcal{B}$) or on the signal cross section times branching ratio times detector acceptance ($\sigma \times \mathcal{B} \times \mathcal{A}$). In order to test various theories, the latter is not useful unless the acceptance \mathcal{A} is provided. However, many analysis are not able to present result in a form of limits on $\sigma \times \mathcal{B} (\times \mathcal{A})$, therefore an alternative is adopted to set limits in the signal strength modifier μ . The signal strength modifier is defined to equally change all the cross sections of all production mechanisms of the signal by the same scale. sections.

In this thesis, the modified frequentist approach confidence levels are used [138, 139]. The classical frequentist uses a test statistic q_μ based on the profile likelihood ratio to determine how signal- or background-like the data is. However, it does not allow nuisance parameters and is modified to incorporate these. First a likelihood $\mathcal{L}(\text{data} | \mu, \theta)$ is constructed as

$$\mathcal{L}(\text{data} | \mu, \theta) = \text{Poisson}(\text{data} | \mu s(\theta) + b(\theta)) p(\tilde{\theta} | \theta). \quad (3.10)$$

The probability density function (pdf) $p(\tilde{\theta} | \theta)$ describes all sources of uncertainty and is described in Section 3.4.2. The data in Equation 3.10 represents either the actual observation or pseudo-data to construct sampling distributions. For a binned likelihood, the Poisson probabilities to observe n_i events in bin i is given as

$$\text{Poisson}(\text{data} | \mu s(\theta) + b(\theta)) = \prod_i \frac{(\mu s_i(\theta) + b_i(\theta))^{n_i}}{n_i!} e^{-\mu s_i(\theta) - b_i(\theta)}. \quad (3.11)$$

At the LHC, the test statistic is defined as

$$q_\mu = -2 \ln \frac{\mathcal{L}(\text{data} | \mu, \hat{\theta}_\mu)}{\mathcal{L}(\text{data} | \hat{\mu}, \hat{\theta}_\mu)}, \quad (3.12)$$

where the likelihood is maximised in the numerator (maximum likelihood estimator, MLE) for a given μ and (pseudo) data at $\hat{\theta}_\mu$, while $\hat{\mu}$ combined with $\hat{\theta}$ defines the point for which the likelihood reaches its global maximum. The estimated signal strength modifier $\hat{\mu}$ can not become negative since a signal rate is positive defined by physics. Furthermore, an upper constraint on the MLE $\hat{\mu} \leq \mu$ is imposed to guarantee a one sided confidence interval. This has

as consequence that upward fluctuations of the data ($\hat{\mu} > \mu$) are not considered against the signal hypothesis of data with a signal with strength μ .

The criterion for excluding the signal at $1 - \alpha$ confidence level is the ratio of the probabilities to observe a value of the test statistic at least as large as the one observed in data q_μ^{obs} , under the signal plus background ($s + b$) and background only (b) hypothesis is defined as

$$\text{CL} = \frac{P(q_\mu \geq q_\mu^{\text{obs}} | \mu s + b)}{P(q_\mu \geq q_\mu^{\text{obs}} | b)} \leq \alpha. \quad (3.13)$$

These probabilities are defined as

$$\begin{aligned} p_\mu &= P(q_\mu \geq q_\mu^{\text{obs}} | \mu s + b) = \int_{q_\mu^{\text{obs}}}^{\infty} f(q_\mu | \mu, \hat{\theta}_\mu^{\text{obs}}) dq_\mu, \\ 1 - p_b &= P(q_\mu \geq q_\mu^{\text{obs}} | b) = \int_{q_{\mu=0}^{\text{obs}}}^{\infty} f(q_\mu | \mu = 0, \hat{\theta}_{\mu=0}^{\text{obs}}) dq_\mu, \end{aligned} \quad (3.14)$$

where p_μ and p_b are called the p-values associated to the two hypothesis, and $f(q_\mu | \mu, \hat{\theta}_\mu^{\text{obs}})$ and $f(q_\mu | \mu = 0, \hat{\theta}_{\mu=0}^{\text{obs}})$ are the pdfs of the signal plus background and background only hypothesis constructed from toy Monte Carlo pseudo data. These pdfs are shown in Figure 3.6 and are generated with nuisance parameters fixed to $\hat{\theta}_{\mu=0}^{\text{obs}}$ and $\hat{\theta}_\mu^{\text{obs}}$. These values of the nuisance parameters for the background only $\hat{\theta}_{\mu=0}^{\text{obs}}$ and signal plus background $\hat{\theta}_\mu^{\text{obs}}$ hypothesis that best describe the data are found by maximising the likelihood from Equation 3.10. The 95% CL level upper limit on μ is achieved by adjusting μ until $\text{CL} = 0.05$




Figure 3.6: Test statistic distributions for pseudo data generated for the signal plus background ($\mu = 1$) and background only ($\mu = 0$) hypothesis. Figure taken from [134].

849

850 The expected median upper limit and the $\pm 1\sigma$ and $\pm 2\sigma$ bands for a hypothesis is generated
851 by a large set of pseudo data and calculate the CLs and the value of μ at 95% CL for each of
852 them. A cumulative probability distribution can be build by starting the integration from the

853 side corresponding to low event yields. The median expected value is where the cumulative
 854 distribution function crosses the 50% quantile. The $\pm 1\sigma$ (68%) and $\pm 2\sigma$ (95%) bands are
 855 defined by the crossings of the 16% and 84%, and 2.5% and 97.5% quantiles.

856 3.4.2 Adding sources of uncertainty

857 In this thesis, all sources of uncertainties are assumed to be either 100% correlated or uncor-
 858 related. Partially correlated uncertainties are broken down to subcomponents that fit those
 859 requirements, allowing to include all constraints in the likelihoods in a clean factorised form.

A systematic uncertainty pdf $p(\theta|\tilde{\theta})$ for the nuisance θ with nominal value $\tilde{\theta}$ is used. It reflects the degree of belief of what the true value of the θ is. In this thesis, the approach from the Higgs Combined Tool is used where the pdfs $p(\theta|\tilde{\theta})$ are re-interpret as posteriors of real or imaginary measurements $\tilde{\theta}$

$$p(\theta|\tilde{\theta}) \sim p(\theta|\tilde{\theta}) \pi_\theta(\theta), \quad (3.15)$$

860 where $\pi_\theta(\theta)$ is the hyper prior for the (imaginary) measurements. For the pdfs used by the
 861 Higgs Combine Tool (normal, log normal, gamma distribution), hyper priors can remain flat.
 862 This allows to use the pdf $p(\theta|\tilde{\theta})$ to constrain the likelihood of the main measurement in a
 863 frequentist calculation. Additionally this allows to build a sampling distribution of the test
 864 statistic [134].

The statistical uncertainties on the Monte Carlo prediction in each bin are obtained following the Barlow-Beeston-light approach [140]. In this approach a single Gaussian constrained nuisance parameter is assigned to scale the sum of the process yields in each bin, constrained by the total uncertainty. This method has as advantage that it minimises the number of parameters required in the maximum likelihood fit. Considering n_{tot} events in a bin with background process i in the bin

$$n_{\text{tot}} = \sum_{i \in \text{bkg}} n_i, \quad (3.16)$$

the total uncertainty e_{tot} is given by

$$e_{\text{tot}} = \sqrt{\sum_{i \in \text{bkg}} e_i^2}, \quad (3.17)$$

865 with e_i the uncertainty on background i and is given by the sum of squares of weights used to
 866 fill the bins. The Gaussian constrained parameter x has then a nominal value of zero and scales
 867 the yield as $n_{\text{tot}} + x e_{\text{tot}}$.

NOTE: Hoe
is de stat
unc op data
behandeld?

868 Choices of systematic uncertainty density functions

For uncertainties that are unconstrained by a priori measurements that do not involve the data going into the statistical analysis, flat priors are used. When there are a priori measurements available such as those from control regions, one can use either a Gaussian pdf, a log-normal pdf, or a gamma distribution. The Gaussian pdf is suited for describing uncertainties on parameters

with both positive and negative values. This prior is however not suitable for positively defined observables such as cross sections, cut efficiencies, luminosity, etc and is not used in this thesis. An alternative option is the log normal pdf which is used in the rest of this thesis

$$\rho(\theta) = \frac{1}{\sqrt{2\pi} \ln(\kappa)} \exp\left(-\frac{(\ln(\theta/\tilde{\theta}))^2}{2(\ln \kappa)^2}\right) \frac{1}{\theta}. \quad (3.18)$$

The parameter κ characterises the width of the log normal pdf. For example $\kappa = 1.10$ implies that the observable can be larger or smaller by a factor 1.10, both deviation having a chance of 16%. The gamma distribution is used for describing statistical uncertainties associated with a number of Monte Carlo events in simulation or a number of observed events in a data control sample. In this thesis, the gamma distribution is only used for the latter. The event rate in the signal region n is related to the number of events in the control region N as $n = \alpha N$. Ignoring the uncertainties on α , the predicted rate follows

$$\rho(n) = \frac{1}{\alpha} \frac{n/\alpha)^N}{N!} \exp(-n/\alpha). \quad (3.19)$$

869 The mapping between the posteriors $\rho(\theta|\tilde{\theta})$ and the auxiliary measurement pdfs $p(\tilde{\theta}|\theta)$ are
870 given in [134].

871 3.4.3 Asymptotic approximation of the CL method

872 In order to significantly reduce computing time, the Asymptotic CL method is used. This method
873 avoids an ensemble of toy Monte Carlo samples and instead replaces it by one representative
874 dataset, called Asimov dataset. This dataset is constructed such that all observed quantities are
875 set equal to their MLE values ($\hat{\theta}_{\text{Asimov}} = \theta_0$). More information about this procedure can be
found in Refs. [132].

NOTE: A [§176](#)
explanation
at Denys

877 3.4.4 Extracting the signal model parameters

From a scan of the profile likelihood ratio,

$$q(a) = -2 \ln \frac{\mathcal{L}(\text{obs} | s(a) + b, \hat{\theta}_a)}{\mathcal{L}(\text{obs} | s(\hat{a}) + b, \hat{\theta})}, \quad (3.20)$$

the signal model parameters are evaluated. The likelihood is maximised by the parameters \hat{a} and $\hat{\theta}$. The likelihood

$$\mathcal{L}_{\max} = \mathcal{L}(\text{obs} | s(\hat{a}) + b, \hat{\theta}) \quad (3.21)$$

878 is called the best-fit set.

879 The 68% and 95% CL on a given parameter of interest a_i is then evaluated from $q(a_i) = 1$ or
880 $q(a_i) = 3.84$ respectively, where all other unconstrained model parameters are treated in the
same way as the nuisance parameters [133].

NOTE: A [§181](#)
explanation
at Denys

Event reconstruction and selection

4

883 After the detector simulation described in [Section 3.2](#), the simulated data has the exact same
 884 format as the real collision data recorded at the CMS experiment. Therefore the same software
 885 can be used for the reconstruction of both simulation and real data. In [Section 4.1](#), the event
 886 reconstruction for physics analysis is shown. After reconstructing events, a basic event selection
 887 is made for selecting signal like events. The necessary event requirement are discussed in
 888 [Section 4.2](#).

889 The analysis uses signal and background regions to constrain the huge SM background
 890 compared to the expected signal. [Section 4.3](#) discusses each region that is entering the analysis.
 891 On top of the use of background estimation from control regions, backgrounds that have prompt
 892 leptons contaminated by real leptons either from decays of tau leptons or from hadronized
 893 mesons or baryons (collectively commonly referred as “non-prompt leptons”) as well as by
 894 hadrons or jets misidentified as leptons¹ are evaluated with a data-driven method discussed in
 895 [Section 4.4](#).

896 4.1 Event reconstruction

897 Muon reconstruction

898 The muon reconstruction [\[141\]](#) has three subdivisions: local reconstruction, regional reconstruction
 899 and global reconstruction. The local reconstruction is performed on individual detector
 900 elements such as strip and pixel hits in the inner tracking system, and muon hits and/or seg-
 901 ments in the muon chambers. Independent tracks are reconstructed in the inner tracker -
 902 called tracker tracks - and in the muon system, called standalone tracks. Based on these tracks,
 903 two reconstructions are considered. The outside-in approach is referred to as Global Muon
 904 reconstruction. For each standalone track, a tracker track is found by comparing the parameters
 905 of the two tracks propagated onto a common surface. Combining the hits from the tracker track
 906 and the standalone track, gives a fit via the Kalman filter technique [\[142, 143\]](#) for a global
 907 muon track. The second approach is an inside-out reconstruction, creating tracker muons. All

¹These two classes of contamination will be referred to as not prompt-lepton (NPL) samples.

candidate tracker tracks are extrapolated to the muon system taking into account the magnetic field, the average expected energy losses, and multiple Coulomb scattering in the detector material. When at least one muon segment - DT or CSC hits - matches the extrapolated track, the corresponding tracker track is indicated as a tracker muon.

For low transverse momenta ($p_T \lesssim 5$ GeV), the tracker muon reconstruction is more efficient than the global muon approach. This is due to the fact that tracker muons only require a single muon segment in muon system, while the global muon approach requires typically segments in at least two muon stations. The global muon approach typically improves the tracker reconstruction for $p_T \gtrsim 200$ GeV.

917 Track reconstruction

An iterative tracking algorithm is responsible for the reconstruction of the tracks made by charged particles in the inner tracking system. Each iteration consists of four steps[64]: the track-seed generation, the pattern recognition algorithm, removal of track-hit ambiguities and a final track fit.

The seed generation is the first step. It consists of finding reconstructed hits that are usable for seeding the subsequent track-finding algorithm. They are identified from a group of at least three reconstructed hits in the tracker, or from a pair of hits while requiring the origin of the track segment to be compatible with the nominal beam-collision point. Since the pixel has a higher granularity compared to the strip tracker, its seed generation efficiency is higher. The overall efficiency exceeds 99%. The second step of each iteration, the pattern recognition algorithm, uses the seeds as a starting point for a Kalman filter method [142, 143]. This algorithm extrapolates the seed trajectory towards the next tracker layer taking into account the magnetic field and multiple scattering effects. The track parameters are updated when a compatible hit in the next layer is found. This procedure continues until the outermost layer is reached. Since the Kalman filter method can result in multiple tracks associated to the same seed, or different tracks sharing the same hits, a removal of ambiguities is necessary. This ambiguity resolving is done by removing tracks that are sharing too many hits from the list of track candidates. The tracks with highest number of hits or with the lowest χ^2 if the track fit is kept. The updated track parameters are then refitted using the Kalman filter method, where all hits found in the pattern recognition step are taken into account. The fit is done twice - once outwards from the beam line towards the calorimeters, and inwards from the outermost track hit to the beam line -, improving the estimation of the track parameters.

All hits that are unambiguously associated to the final track are removed from the list of available hits. In order to associate the remaining hits, the procedure is repeated with looser track reconstruction criteria. The use of the iterative track reconstruction procedure has a high track finding efficiency, where the fake track reconstruction rate is negligible. For muons, this results in a global track reconstruction efficiency exceeding 98%, and 75-98% for charged hadrons.

946 Primary vertex reconstruction

The primary vertex reconstruction should be able to measure the location of all proton interaction vertices in each event: the signal vertex and all vertices from pile up events. It consists of a vertex

finding and a vertex fitting algorithm and happens in three steps. Tracks are selected to be consistent with being produced promptly in the primary interaction by imposing requirements on the track parameters[76] By grouping reconstructed tracks according to the z coordinate of their closest approach to the beam line, vertices for all interaction in the same beam crossing are found, at CMS this is done by a deterministic annealing algorithm [144] . On top of this, a vertex fitting algorithm like the Adaptive Vertex fitter [145], is performed. This creates the three-dimensional primary-vertex position. With this fit, the contribution from long-lived hadron decays is reduced by down weighting the tracks with a larger distance to the vertex. The primary vertex corresponding to the highest sum of squared track transverse momenta is noted as the point of the main interaction. The resolution on the primary vertex is about 14 μm in $r\phi$ and about 19 μm in the z direction for primary vertices with the sum of the track $p_T > 100$ GeV for 2016 data taking.

4.2 Event selection

4.3 Regions and channels

4.4 Data driven background simulation

The search for FCNC involving a top quark and a Z boson

5

964 **5.1 Construction of template distributions**

966 **5.2 Systematic uncertainties**

967 **5.3 Limit setting procedure**

968 **5.4 Result and discussion**

6

969

Conclusion and outlook




Figure 6.1:

Bibliography

- 971 [1] MICHAEL E PESKIN and DANIEL V SCHROEDER: **An introduction to quantum field**
 972 **theory; 1995 ed.** Includes exercises. Boulder, CO: Westview, 1995. URL: <https://cds.cern.ch/record/257493> (see pp. 1, 5).
- 973
- 974 [2] C. P. BURGESS: **Introduction to Effective Field Theory**. In: *Ann. Rev. Nucl. Part.*
 975 *Sci.*, **57**: (2007), pp. 329–362. DOI: [10.1146/annurev.nucl.56.080805.140508](https://doi.org/10.1146/annurev.nucl.56.080805.140508). arXiv:
 976 [hep-th/0701053 \[hep-th\]](https://arxiv.org/abs/hep-th/0701053) (see p. 1).
- 977 [3] NADIA ROBOTTI: **The discovery of the electron: I**. In: *European Journal of Physics*,
 978 **18**:3 (1997), p. 133. URL: <http://stacks.iop.org/0143-0807/18/i=3/a=002> (see p. 2).
- 979 [4] C. PATRIGNANI et al.: **Review of Particle Physics**. In: *Chin. Phys.*, **C40**:10 (2016),
 980 p. 100001. DOI: [10.1088/1674-1137/40/10/100001](https://doi.org/10.1088/1674-1137/40/10/100001) (see pp. 2–3, 6–7, 9, 34–35).
- 981 [5] S. ABACHI et al.: **Observation of the top quark**. In: *Phys. Rev. Lett.*, **74**: (1995),
 982 pp. 2632–2637. DOI: [10.1103/PhysRevLett.74.2632](https://doi.org/10.1103/PhysRevLett.74.2632). arXiv: [hep-ex/9503003 \[hep-ex\]](https://arxiv.org/abs/hep-ex/9503003)
 983 (see p. 2).
- 984 [6] F. ABE et al.: **Observation of top quark production in $\bar{p}p$ collisions**. In: *Phys. Rev.*
 985 *Lett.*, **74**: (1995), pp. 2626–2631. DOI: [10.1103/PhysRevLett.74.2626](https://doi.org/10.1103/PhysRevLett.74.2626). arXiv: [hep-ex/9503002 \[hep-ex\]](https://arxiv.org/abs/hep-ex/9503002) (see p. 2).
- 986
- 987 [7] SERGUEI CHATRHYAN et al.: **Observation of a new boson at a mass of 125 GeV**
 988 **with the CMS experiment at the LHC**. In: *Phys. Lett.*, **B716**: (2012), pp. 30–61. DOI:
 989 [10.1016/j.physletb.2012.08.021](https://doi.org/10.1016/j.physletb.2012.08.021). arXiv: [1207.7235 \[hep-ex\]](https://arxiv.org/abs/1207.7235) (see pp. 2, 17).
- 990 [8] GEORGES AAD et al.: **Observation of a new particle in the search for the Standard**
 991 **Model Higgs boson with the ATLAS detector at the LHC**. In: *Phys. Lett.*, **B716**:
 992 (2012), pp. 1–29. DOI: [10.1016/j.physletb.2012.08.020](https://doi.org/10.1016/j.physletb.2012.08.020). arXiv: [1207.7214 \[hep-ex\]](https://arxiv.org/abs/1207.7214)
 993 (see pp. 2, 17).
- 994 [9] NICOLA CABIBBO: **Unitary Symmetry and Leptonic Decays**. In: *Phys. Rev. Lett.*, **10**:
 995 (12 June 1963), pp. 531–533. DOI: [10.1103/PhysRevLett.10.531](https://doi.org/10.1103/PhysRevLett.10.531) (see p. 5).
- 996 [10] S. L. GLASHOW, J. ILIOPoulos, and L. MAIANI: **Weak Interactions with Lepton-Hadron**
 997 **Symmetry**. In: *Phys. Rev. D*, **2**: (7 Oct. 1970), pp. 1285–1292. DOI: [10.1103/PhysRevD.2.1285](https://doi.org/10.1103/PhysRevD.2.1285) (see p. 5).
- 998
- 999 [11] B.J. BJØRKEN and S.L. GLASHOW: **Elementary particles and SU(4)**. In: *Physics Letters*,
 1000 **11**:3 (1964), pp. 255–257. DOI: [https://doi.org/10.1016/0031-9163\(64\)90433-0](https://doi.org/10.1016/0031-9163(64)90433-0) (see
 1001 p. 5).

- 1002 [12] LUCIANO MAIANI: **The GIM Mechanism: origin, predictions and recent uses.** In:
 1003 *Proceedings, 48th Rencontres de Moriond on Electroweak Interactions and Unified Theories:*
 1004 *La Thuile, Italy, March 2-9, 2013.* 2013, pp. 3–16. arXiv: 1303.6154 [hep-ph]. URL:
 1005 <https://inspirehep.net/record/1225307/files/arXiv:1303.6154.pdf> (see p. 5).
- 1006 [13] PATRICK KOPPENBURG and SEBASTIEN DESCOTES-GENON: **The CKM Parameters.** In:
 1007 (2017). arXiv: 1702.08834 [hep-ex] (see p. 6).
- 1008 [14] J. A. AGUILAR-SAAVEDRA: **Top flavor-changing neutral interactions: Theoretical
 1009 expectations and experimental detection.** In: *Acta Phys. Polon.*, B35: (2004),
 1010 pp. 2695–2710. arXiv: hep-ph/0409342 [hep-ph] (see pp. 6, 13–14).
- 1011 [15] ANDREA GIAMMANCO and JEANNINE WAGNER-KUHR: **Measurement of the t-channel
 1012 single Top-quark production rates in pp collisions at 7 TeV.** 2011. URL: <http://cms.web.cern.ch/news/measurement-t-channel-single-top-quark-production-rates-pp-collisions-7-tev> (see p. 8).
- 1013 [16] LHCTOP WORKING GROUP: **ATLAS-CMS recommended predictions for single top
 1014 cross sections using the Hathor v2.1 program.** 2017. URL: https://twiki.cern.ch/twiki/bin/view/LHCPhysics/SingleTopRefXsec#Predictions_at_7_8_13_and_14_TeV
 (see p. 9).
- 1015 [17] CMS COLLABORATION: **Summaries of CMS cross section measurements.** 2017. URL:
 1016 <https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsCombined> (see
 1017 p. 9).
- 1018 [18] LHCTOP WORKING GROUP: **LHCTopWG Summary plots.** 2017. URL: <https://twiki.cern.ch/twiki/bin/view/LHCPhysics/LHCTopWGSummaryPlots> (see pp. 10–11).
- 1019 [19] B. GRZADKOWSKI, M. ISKRZYNSKI, M. MISIAK, and J. ROSIEK: **Dimension-Six Terms
 1020 in the Standard Model Lagrangian.** In: *JHEP*, 10: (2010), p. 085. DOI: 10.1007/
 1021 JHEP10(2010)085. arXiv: 1008.4884 [hep-ph] (see p. 11).
- 1022 [20] ANDY BUCKLEY, CHRISTOPH ENGLERT, JAMES FERRANDO, et al.: **Constraining top
 1023 quark effective theory in the LHC Run II era.** In: *JHEP*, 04: (2016), p. 015. DOI:
 10.1007/JHEP04(2016)015. arXiv: 1512.03360 [hep-ph] (see p. 11).
- 1024 [21] YORIKIYO NAGASHIMA: **Beyond the standard model of elementary particle physics.**
 1025 Weinheim, Germany: Wiley-VCH Verlag, 2014. URL: <http://www-spires.fnal.gov/spires/find/books/www?cl=QC793.2.N34::2014> (see p. 12).
- 1026 [22] Y. FUKUDA et al.: **Evidence for oscillation of atmospheric neutrinos.** In: *Phys. Rev.
 1027 Lett.*, 81: (1998), pp. 1562–1567. DOI: 10.1103/PhysRevLett.81.1562. arXiv: hep-
 1028 ex/9807003 [hep-ex] (see p. 12).
- 1029 [23] Y. ABE et al.: **Indication of Reactor $\bar{\nu}_e$ Disappearance in the Double Chooz Experiment.** In: *Phys. Rev. Lett.*, 108: (13 Mar. 2012), p. 131801. DOI: 10.1103/PhysRevLett.108.131801 (see p. 12).
- 1030 [24] P. A. R. ADE et al.: **Planck 2015 results. XIII. Cosmological parameters.** In: *Astron.
 1031 Astrophys.*, 594: (2016), A13. DOI: 10.1051/0004-6361/201525830. arXiv: 1502.01589
 1032 [astro-ph.CO] (see p. 12).

- 1042 [25] P. J. E. PEEBLES and BHARAT RATRA: **The Cosmological constant and dark energy.**
 1043 In: *Rev. Mod. Phys.*, **75**: (2003), pp. 559–606. doi: [10.1103/RevModPhys.75.559](https://doi.org/10.1103/RevModPhys.75.559). arXiv:
 1044 [astro-ph/0207347 \[astro-ph\]](https://arxiv.org/abs/astro-ph/0207347) (see p. 12).
- 1045 [26] A. D. SAKHAROV: **Violation of CP Invariance, c Asymmetry, and Baryon Asym-**
 1046 **metry of the Universe.** In: *Pisma Zh. Eksp. Teor. Fiz.*, **5**: (1967). [*Usp. Fiz.*
 1047 *Nauk*161,61(1991)], pp. 32–35. doi: [10.1070/PU1991v034n05ABEH002497](https://doi.org/10.1070/PU1991v034n05ABEH002497) (see p. 12).
- 1048 [27] GUSTAVO BURDMAN: **New solutions to the hierarchy problem.** In: *Braz. J. Phys.*, **37**:
 1049 (2007), pp. 506–513. doi: [10.1590/S0103-97332007000400006](https://doi.org/10.1590/S0103-97332007000400006). arXiv: [hep-ph/0703194](https://arxiv.org/abs/hep-ph/0703194)
 1050 [hep-ph] (see p. 12).
- 1051 [28] T. ET AL. AALTONEN: **Search for the Flavor-Changing Neutral-Current Decay $t \rightarrow Zq$**
 1052 **in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV.** In: *Phys. Rev. Lett.*, **101**: (19 Nov. 2008),
 1053 p. 192002. doi: [10.1103/PhysRevLett.101.192002](https://doi.org/10.1103/PhysRevLett.101.192002) (see p. 13).
- 1054 [29] VICTOR MUKHAMEDOVICH ABAZOV et al.: **Search for flavour changing neutral cur-**
 1055 **rents via quark-gluon couplings in single top quark production using 2.3 fb^{-1} of**
 1056 **$p\bar{p}$ collisions.** In: *Phys. Lett.*, **B693**: (2010), pp. 81–87. doi: [10.1016/j.physletb.2010.08.011](https://doi.org/10.1016/j.physletb.2010.08.011). arXiv: [1006.3575 \[hep-ex\]](https://arxiv.org/abs/1006.3575) (see p. 13).
- 1058 [30] GEORGES AAD et al.: **Search for flavour-changing neutral current top-quark decays**
 1059 **to qZ in pp collision data collected with the ATLAS detector at $\sqrt{s} = 8$ TeV.** In: *Eur. Phys. J.*, **C76**:1 (2016), p. 12. doi: [10.1140/epjc/s10052-015-3851-5](https://doi.org/10.1140/epjc/s10052-015-3851-5). arXiv:
 1061 [1508.05796 \[hep-ex\]](https://arxiv.org/abs/1508.05796) (see p. 13).
- 1062 [31] GEORGES AAD et al.: **Search for single top-quark production via flavour-changing**
 1063 **neutral currents at 8 TeV with the ATLAS detector.** In: *Eur. Phys. J.*, **C76**:2 (2016),
 1064 p. 55. doi: [10.1140/epjc/s10052-016-3876-4](https://doi.org/10.1140/epjc/s10052-016-3876-4). arXiv: [1509.00294 \[hep-ex\]](https://arxiv.org/abs/1509.00294) (see pp. 13,
 1065 16).
- 1066 [32] GEORGES AAD et al.: **Search for flavour-changing neutral current top quark decays**
 1067 **$t \rightarrow Hq$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector.** In: *JHEP*, **12**:
 1068 (2015), p. 061. doi: [10.1007/JHEP12\(2015\)061](https://doi.org/10.1007/JHEP12(2015)061). arXiv: [1509.06047 \[hep-ex\]](https://arxiv.org/abs/1509.06047) (see
 1069 p. 13).
- 1070 [33] MORAD AABOUD et al.: **Search for top quark decays $t \rightarrow qH$, with $H \rightarrow \gamma\gamma$, in**
 1071 **$\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector.** In: (2017). arXiv: [1707.01404](https://arxiv.org/abs/1707.01404)
 1072 [hep-ex] (see pp. 13, 16).
- 1073 [34] ALBERT M SIRUNYAN et al.: **Search for associated production of a Z boson with a**
 1074 **single top quark and for tZ flavour-changing interactions in pp collisions at \sqrt{s}**
 1075 **= 8 TeV.** In: (2017). arXiv: [1702.01404 \[hep-ex\]](https://arxiv.org/abs/1702.01404) (see pp. 13, 16).
- 1076 [35] SERGUEI CHATRCHYAN et al.: **Search for Flavor-Changing Neutral Currents in Top-**
 1077 **Quark Decays $t \rightarrow Zq$ in pp Collisions at $\sqrt{s} = 8$ TeV.** In: *Phys. Rev. Lett.*, **112**:17
 1078 (2014), p. 171802. doi: [10.1103/PhysRevLett.112.171802](https://doi.org/10.1103/PhysRevLett.112.171802). arXiv: [1312.4194 \[hep-ex\]](https://arxiv.org/abs/1312.4194)
 1079 (see p. 13).
- 1080 [36] VARDAN KHACHATRYAN et al.: **Search for anomalous single top quark production in**
 1081 **association with a photon in pp collisions at $\sqrt{s} = 8$ TeV.** In: *JHEP*, **04**: (2016),
 1082 p. 035. doi: [10.1007/JHEP04\(2016\)035](https://doi.org/10.1007/JHEP04(2016)035). arXiv: [1511.03951 \[hep-ex\]](https://arxiv.org/abs/1511.03951) (see pp. 13, 16).

- 1083 [37] VARDAN KHACHATRYAN et al.: **Search for top quark decays via Higgs-boson-mediated**
 1084 **flavor-changing neutral currents in pp collisions at $\sqrt{s} = 8$ TeV.** In: *JHEP*, **02**:
 1085 (2017), p. 079. DOI: [10.1007/JHEP02\(2017\)079](https://doi.org/10.1007/JHEP02(2017)079). arXiv: [1610.04857 \[hep-ex\]](https://arxiv.org/abs/1610.04857) (see
 1086 p. 13).
- 1087 [38] **Search for the flavor-changing interactions of the top quark with the Higgs boson**
 1088 **in $H \rightarrow b\bar{b}$ channel at $\sqrt{s} = 13$ TeV.** Tech. rep. CMS-PAS-TOP-17-003. Geneva:
 1089 CERN, 2017. URL: <https://cds.cern.ch/record/2284743> (see p. 13).
- 1090 [39] J. A. AGUILAR-SAAVEDRA: **A Minimal set of top anomalous couplings.** In: *Nucl. Phys.*,
 1091 **B812:** (2009), pp. 181–204. DOI: [10.1016/j.nuclphysb.2008.12.012](https://doi.org/10.1016/j.nuclphysb.2008.12.012). arXiv: [0811.3842](https://arxiv.org/abs/0811.3842)
 1092 [hep-ph] (see p. 13).
- 1093 [40] J. A. AGUILAR-SAAVEDRA: **A Minimal set of top-Higgs anomalous couplings.** In: *Nucl.*
 1094 *Phys.*, **B821:** (2009), pp. 215–227. DOI: [10.1016/j.nuclphysb.2009.06.022](https://doi.org/10.1016/j.nuclphysb.2009.06.022). arXiv:
 1095 [0904.2387 \[hep-ph\]](https://arxiv.org/abs/0904.2387) (see p. 13).
- 1096 [41] M. BENEKE et al.: **Top quark physics.** In: *1999 CERN Workshop on standard model*
 1097 *physics (and more) at the LHC, CERN, Geneva, Switzerland, 25-26 May: Proceedings.*
 1098 2000, pp. 419–529. arXiv: [hep-ph/0003033 \[hep-ph\]](https://arxiv.org/abs/hep-ph/0003033). URL: <http://weblib.cern.ch/abstract?CERN-TH-2000-100> (see p. 14).
- 1100 [42] JUN GAO, CHONG SHENG LI, and HUA XING ZHU: **Top Quark Decay at Next-to-Next-**
 1101 **to Leading Order in QCD.** In: *Phys. Rev. Lett.*, **110**:4 (2013), p. 042001. DOI:
 1102 [10.1103/PhysRevLett.110.042001](https://doi.org/10.1103/PhysRevLett.110.042001). arXiv: [1210.2808 \[hep-ph\]](https://arxiv.org/abs/1210.2808) (see p. 15).
- 1103 [43] STEPHEN MYERS: **The LEP Collider, from design to approval and commissioning.**
 1104 John Adams' Lecture. Delivered at CERN, 26 Nov 1990. Geneva: CERN, 1991. URL:
 1105 [http://cds.cern.ch/record/226776](https://cds.cern.ch/record/226776) (see p. 17).
- 1106 [44] STEPHEN HOLMES, RONALD S MOORE, and VLADIMIR SHILTSEV: **Overview of the**
 1107 **Tevatron collider complex: goals, operations and performance.** In: *Journal of In-*
 1108 *strumentation*, **6**:08 (2011), T08001. URL: <http://stacks.iop.org/1748-0221/6/i=08/a=T08001> (see p. 17).
- 1110 [45] R. BARATE et al.: **Search for the standard model Higgs boson at LEP.** In: *Phys.*
 1111 *Lett.*, **B565:** (2003), pp. 61–75. DOI: [10.1016/S0370-2693\(03\)00614-2](https://doi.org/10.1016/S0370-2693(03)00614-2). arXiv: [hep-ex/0306033 \[hep-ex\]](https://arxiv.org/abs/hep-ex/0306033) (see p. 17).
- 1113 [46] KENNETH HERNER: **Higgs Boson Studies at the Tevatron.** In: *Nucl. Part. Phys. Proc.*,
 1114 **273-275:** (2016), pp. 852–856. DOI: [10.1016/j.nuclphysbps.2015.09.131](https://doi.org/10.1016/j.nuclphysbps.2015.09.131) (see p. 17).
- 1115 [47] ABDELHAK DJOUADI: **The Anatomy of electro-weak symmetry breaking. I: The**
 1116 **Higgs boson in the standard model.** In: *Phys. Rept.*, **457:** (2008), pp. 1–216. DOI:
 1117 [10.1016/j.physrep.2007.10.004](https://doi.org/10.1016/j.physrep.2007.10.004). arXiv: [hep-ph/0503172 \[hep-ph\]](https://arxiv.org/abs/hep-ph/0503172) (see p. 17).
- 1118 [48] LYNDON EVANS and PHILIP BRYANT: **LHC Machine.** In: *Journal of Instrumentation*,
 1119 **3**:08 (2008), S08001. URL: <http://stacks.iop.org/1748-0221/3/i=08/a=S08001> (see
 1120 p. 17).
- 1121 [49] THOMAS SVEN PETTERSSON and P LEFÈVRE: **The Large Hadron Collider: conceptual**
 1122 **design.** Tech. rep. CERN-AC-95-05-LHC. Oct. 1995, p. 20 and 22. URL: <https://cds.cern.ch/record/291782> (see p. 17).

- 1124 [50] JORG WENNINGER and EZIO TODESCO: **Large Hadron Collider momentum calibration**
 1125 **and accuracy.** Tech. rep. CERN-ACC-2017-0007. Geneva: CERN, Feb. 2017. URL:
 1126 <https://cds.cern.ch/record/2254678> (see p. 18).
- 1127 [51] CINZIA DE MELIS: **The CERN accelerator complex. Complex des accélérateurs du**
 1128 **CERN.** In: (July 2016). General Photo. URL: <https://cds.cern.ch/record/2197559>
 1129 (see p. 18).
- 1130 [52] OLIVER SIM BRÜNING, PAUL COLLIER, P LEBRUN, et al.: **LHC Design Report.** CERN
 1131 Yellow Reports: Monographs. Geneva: CERN, 2004. URL: <https://cds.cern.ch/record/782076> (see p. 18).
- 1133 [53] G. AAD et al.: **The ATLAS Experiment at the CERN Large Hadron Collider.** In: *JINST*,
 1134 3: (2008), S08003. DOI: [10.1088/1748-0221/3/08/S08003](https://doi.org/10.1088/1748-0221/3/08/S08003) (see p. 19).
- 1135 [54] S. CHATRCHYAN et al.: **The CMS Experiment at the CERN LHC.** In: *JINST*, 3: (2008),
 1136 S08004. DOI: [10.1088/1748-0221/3/08/S08004](https://doi.org/10.1088/1748-0221/3/08/S08004) (see pp. 19, 26–28).
- 1137 [55] K. AAMODT et al.: **The ALICE experiment at the CERN LHC.** In: *JINST*, 3: (2008),
 1138 S08002. DOI: [10.1088/1748-0221/3/08/S08002](https://doi.org/10.1088/1748-0221/3/08/S08002) (see p. 19).
- 1139 [56] A. AUGUSTO ALVES JR. et al.: **The LHCb Detector at the LHC.** In: *JINST*, 3: (2008),
 1140 S08005. DOI: [10.1088/1748-0221/3/08/S08005](https://doi.org/10.1088/1748-0221/3/08/S08005) (see p. 19).
- 1141 [57] M. BONGI et al.: **Astroparticle physics at LHC: The LHCf experiment ready for data**
 1142 **taking.** In: *Nucl. Instrum. Meth.*, A612: (2010), pp. 451–454. DOI: [10.1016/j.nima.2009.08.039](https://doi.org/10.1016/j.nima.2009.08.039) (see p. 19).
- 1144 [58] G. ANELLI et al.: **The TOTEM experiment at the CERN Large Hadron Collider.** In: *JINST*, 3: (2008), S08007. DOI: [10.1088/1748-0221/3/08/S08007](https://doi.org/10.1088/1748-0221/3/08/S08007) (see p. 19).
- 1146 [59] B. ACHARYA et al.: **The Physics Programme Of The MoEDAL Experiment At The LHC.**
 1147 In: *Int. J. Mod. Phys.*, A29: (2014), p. 1430050. DOI: [10.1142/S0217751X14300506](https://doi.org/10.1142/S0217751X14300506).
 1148 arXiv: [1405.7662 \[hep-ph\]](https://arxiv.org/abs/1405.7662) (see p. 19).
- 1149 [60] BY JAMES GILLIES: **Luminosity? Why don't we just say collision rate?** In: (Mar.
 1150 2011). URL: [http://cds.cern.ch/record/1997001](https://cds.cern.ch/record/1997001) (see p. 19).
- 1151 [61] BY HARRIET JARLETT and HARRIET KIM JARLETT: **LHC pushes limits of performance.**
 1152 In: (Aug. 2016). URL: [http://cds.cern.ch/record/2212301](https://cds.cern.ch/record/2212301) (see p. 20).
- 1153 [62] **Technical proposal.** LHC Tech. Proposal. Cover title : CMS, the Compact Muon
 1154 Solenoid : technical proposal. Geneva: CERN, 1994. URL: <https://cds.cern.ch/record/290969> (see p. 20).
- 1156 [63] G. L. BAYATIAN et al.: **CMS physics: Technical design report.** In: (2006) (see pp. 20,
 1157 29).
- 1158 [64] G L BAYATIAN, S CHATRCHYAN, G HMAYAKYAN, et al.: **CMS Physics: Technical Design**
 1159 **Report Volume 1: Detector Performance and Software.** Technical Design Report
 1160 CMS. There is an error on cover due to a technical problem for some items. Geneva:
 1161 CERN, 2006. URL: <https://cds.cern.ch/record/922757> (see pp. 20, 25, 50).
- 1162 [65] CMS COLLABORATION: **Detector Drawings.** CMS Collection. Mar. 2012. URL: <https://cds.cern.ch/record/1433717> (see p. 21).

- 1164 [66] S CHATRCHYAN, V KHACHATRYAN, A M SIRUNYAN, et al.: **Performance of the CMS Drift**
 1165 **Tube Chambers with Cosmic Rays**. In: *J. Instrum.*, 5:arXiv:0911.4855. CMS-CFT-
 1166 09-012 (Nov. 2009), T03015 . 47 p. URL: <http://cds.cern.ch/record/1223944> (see
 1167 pp. 23–24).
- 1168 [67] TOM DODINGTON: **News from the CMS experimental site: 22 November 2013**. 2013.
 1169 URL: <http://cms.web.cern.ch/news/news-point-5-22-november-2013> (see p. 24).
- 1170 [68] INSTITUTE OF RESEARCH INTO THE FUNDAMENTAL LAWS THE UNIVERSE: **The CMS**
 1171 **detector superconducting solenoid**. 2006. URL: http://irfu.cea.fr/en/Phocea/Vie_des_labos/Ast/ast_visu.php?id_ast=839 (see p. 26).
- 1173 [69] G BAIATIAN, SIRUNYAN, et al.: **Design, Performance, and Calibration of CMS Hadron-
 1174 Barrel Calorimeter Wedges**. Tech. rep. CMS-NOTE-2006-138. Geneva: CERN, May
 1175 2007. URL: <https://cds.cern.ch/record/1049915>.
- 1176 [70] FERGUS WILSON: **Experimental Particle Physics**. 2012. URL: <http://slideplayer.com/slide/794631/> (see p. 26).
- 1178 [71] EMRAH TIRAS, BURAK BILKI, and YASAR ONEL: **Commissioning of CMS Forward**
 1179 **Hadron Calorimeters with Upgraded Multi-anode PMTs and μTCA Readout**. In:
 1180 (2016). arXiv: [1611.05232 \[physics.ins-det\]](1611.05232) (see p. 27).
- 1181 [72] LUCAS TAYLOR: **Experimental Particle Physics**. 2011. URL: <http://cms.web.cern.ch/news/electromagnetic-calorimeter> (see p. 27).
- 1183 [73] **Proceedings, 34th International Conference on High Energy Physics (ICHEP 2008)**■
 1184 URL: <http://www.slac.stanford.edu/econf/C080730> (see p. 27).
- 1185 [74] L. BRIANZA: **Precision crystal calorimetry in LHC Run II with the CMS ECAL**. In:
 1186 *Journal of Instrumentation*, 12:01 (2017), p. C01069. URL: <http://stacks.iop.org/1748-0221/12/i=01/a=C01069> (see pp. 28, 32).
- 1188 [75] SERGUEI CHATRCHYAN et al.: **Energy Calibration and Resolution of the CMS Elec-
 1189 **tromagnetic Calorimeter in pp Collisions at $\sqrt{s} = 7$ TeV**. In: *JINST*, 8: (2013).
 1190 [JINST8,9009(2013)], P09009. DOI: <10.1088/1748-0221/8/09/P09009>. arXiv: [1306.2016 \[hep-ex\]](1306.2016) (see p. 32).**
- 1192 [76] SERGUEI CHATRCHYAN, VARDAN KHACHATRYAN, ALBERT M SIRUNYAN, et al.: **Description**
 1193 **and performance of track and primary-vertex reconstruction with the CMS**
 1194 **tracker**. In: *J. Instrum.*, 9:arXiv:1405.6569. CERN-PH-EP-2014-070. CMS-TRK-11-001
 1195 (May 2014). Comments: Replaced with published version. Added journal reference
 1196 and DOI, P10009. 80 p. URL: <http://cds.cern.ch/record/1704291> (see pp. 28, 51).
- 1197 [77] BY CHRISTINE SUTTON: **Chronicles of CMS: the saga of LS1**. In: (May 2015). URL:
 1198 <http://cds.cern.ch/record/2024986> (see p. 29).
- 1199 [78] **A beautiful barrel for CMS**. In: (Mar. 2014). URL: <http://cds.cern.ch/record/1998635> (see p. 29).
- 1201 [79] **Cool running for CMS tracker**. In: (Mar. 2014). URL: <http://cds.cern.ch/record/1998606> (see p. 30).
- 1203 [80] BY CORINNE PRALAVORIO and CORINNE PRALAVORIO: **Major work to ready the LHC**
 1204 **experiments for Run 2**. In: (May 2015). URL: <http://cds.cern.ch/record/2024977>
 1205 (see pp. 30–31).

- 1206 [81] L. CADAMURO: **The CMS Level-1 trigger system for LHC Run II.** In: *Journal of*
 1207 *Instrumentation*, **12**:03 (2017), p. C03021. URL: <http://stacks.iop.org/1748-0221/12/i=03/a=C03021> (see p. 31).
- 1209 [82] VARDAN KHACHATRYAN et al.: **The CMS trigger system.** In: *JINST*, **12**:01 (2017),
 1210 P01020. DOI: [10.1088/1748-0221/12/01/P01020](https://doi.org/10.1088/1748-0221/12/01/P01020). arXiv: [1609.02366 \[physics.ins-det\]](https://arxiv.org/abs/1609.02366) (see p. 31).
- 1212 [83] LUIGI GUIDUCCI: **CMS muon system towards LHC Run 2 and beyond.** Tech. rep.
 1213 CMS-CR-2014-333. Geneva: CERN, Oct. 2014. URL: <https://cds.cern.ch/record/1966038> (see p. 32).
- 1215 [84] CARLO BATTILANA: **The CMS muon system status and upgrades for LHC run-2 and**
 1216 **performance of muon reconstruction with 13 TeV data.** Tech. rep. CMS-CR-2016-
 1217 437. Geneva: CERN, Dec. 2016. URL: <https://cds.cern.ch/record/2239185> (see
 1218 p. 32).
- 1219 [85] CLAUDIO GRANDI, DAVID STICKLAND, LUCAS TAYLOR, ACHILLE PETRILLI, and ALAIN
 1220 HERVÉ: **CMS Computing Model: The "CMS Computing Model RTAG".** Tech. rep.
 1221 CMS-NOTE-2004-031. CERN-LHCC-2004-035. LHCC-G-083. Geneva: CERN, Dec.
 1222 2004. URL: [http://cds.cern.ch/record/814248](https://cds.cern.ch/record/814248) (see p. 32).
- 1223 [86] CHRISTOPH ECK, J KNOBLOCH, LESLIE ROBERTSON, et al.: **LHC computing Grid: Technical**
 1224 **Design Report. Version 1.06 (20 Jun 2005).** Technical Design Report LCG.
 1225 Geneva: CERN, 2005. URL: <https://cds.cern.ch/record/840543> (see p. 32).
- 1226 [87] JOHN C. COLLINS, DAVISON E. SOPER, and GEORGE F. STERMAN: **Factorization of Hard**
 1227 **Processes in QCD.** In: *Adv. Ser. Direct. High Energy Phys.*, **5**: (1989), pp. 1–91. DOI:
 1228 [10.1142/9789814503266_0001](https://doi.org/10.1142/9789814503266_0001). arXiv: [hep-ph/0409313 \[hep-ph\]](https://arxiv.org/abs/hep-ph/0409313) (see p. 33).
- 1229 [88] RINGAILE PLACAKYTE: **Parton Distribution Functions.** In: *Proceedings, 31st International Conference on Physics in collisions (PIC 2011): Vancouver, Canada, August 28-September 1, 2011.* 2011. arXiv: [1111.5452 \[hep-ph\]](https://arxiv.org/abs/1111.5452). URL: <https://inspirehep.net/record/954990/files/arXiv:1111.5452.pdf> (see p. 33).
- 1233 [89] RICHARD D. BALL, VALERIO BERTONE, STEFANO CARRAZZA, et al.: **Parton distributions**
 1234 **for the LHC run II.** In: *Journal of High Energy Physics*, **2015**:4 (2015), p. 40. DOI:
 1235 [10.1007/JHEP04\(2015\)040](https://doi.org/10.1007/JHEP04(2015)040) (see p. 33).
- 1236 [90] JON BUTTERWORTH et al.: **PDF4LHC recommendations for LHC Run II.** In: *J. Phys.*,
 1237 **G43:** (2016), p. 023001. DOI: [10.1088/0954-3899/43/2/023001](https://doi.org/10.1088/0954-3899/43/2/023001). arXiv: [1510.03865 \[hep-ph\]](https://arxiv.org/abs/1510.03865) (see pp. 33–34).
- 1239 [91] H. ABRAMOWICZ and A. CALDWELL: **HERA collider physics.** In: *Rev. Mod. Phys.*, **71**:
 1240 (1999), pp. 1275–1410. DOI: [10.1103/RevModPhys.71.1275](https://doi.org/10.1103/RevModPhys.71.1275). arXiv: [hep-ex/9903037 \[hep-ex\]](https://arxiv.org/abs/hep-ex/9903037) (see p. 33).
- 1242 [92] STEPHEN HOLMES, RONALD S. MOORE, and VLADIMIR SHILTSEV: **Overview of the Tevatron Collider Complex: Goals, Operations and Performance.** In: *JINST*, **6**: (2011),
 1243 T08001. DOI: [10.1088/1748-0221/6/08/T08001](https://doi.org/10.1088/1748-0221/6/08/T08001). arXiv: [1106.0909 \[physics.acc-ph\]](https://arxiv.org/abs/1106.0909) (see p. 34).
- 1246 [93] JUAN ROJO et al.: **The PDF4LHC report on PDFs and LHC data: Results from Run I**
 1247 **and preparation for Run II.** In: *J. Phys.*, **G42**: (2015), p. 103103. DOI: [10.1088/0954-3899/42/10/103103](https://doi.org/10.1088/0954-3899/42/10/103103). arXiv: [1507.00556 \[hep-ph\]](https://arxiv.org/abs/1507.00556) (see p. 34).

- 1249 [94] J. PUMPLIN, D. STUMP, R. BROCK, et al.: **Uncertainties of predictions from parton dis-**
 1250 **distribution functions. 2. The Hessian method.** In: *Phys. Rev.*, **D65**: (2001), p. 014013.
 1251 DOI: [10.1103/PhysRevD.65.014013](https://doi.org/10.1103/PhysRevD.65.014013). arXiv: [hep-ph/0101032](https://arxiv.org/abs/hep-ph/0101032) [hep-ph] (see p. 34).
- 1252 [95] FRANZ MANDL and GRAHAM G SHAW: **Quantum field theory; 2nd ed.** New York, NY:
 1253 Wiley, 2010. URL: <https://cds.cern.ch/record/1236742> (see p. 35).
- 1254 [96] S BANERJEE: **CMS Simulation Software.** In: *Journal of Physics: Conference Series*,
 1255 **396**:2 (2012), p. 022003. URL: <http://stacks.iop.org/1742-6596/396/i=2/a=022003>
 1256 (see p. 36).
- 1257 [97] M HILDRETH, V N IVANCHENKO, D J LANGE, and M J KORTELAINEN: **CMS Full Simu-**
 1258 **lation for Run-2.** In: *Journal of Physics: Conference Series*, **664**:7 (2015), p. 072022.
 1259 URL: <http://stacks.iop.org/1742-6596/664/i=7/a=072022> (see p. 36).
- 1260 [98] S. AGOSTINELLI, J. ALLISON, K. AMAKO, et al.: **Geant4-a simulation toolkit.** In: *Nuclear*
 1261 *Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers,*
 1262 *Detectors and Associated Equipment*, **506**:3 (2003), pp. 250–303. DOI: [https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8) (see pp. 36, 38).
- 1264 [99] MICHAEL H. SEYMOUR and MARILYN MARX: **Monte Carlo Event Generators.** In: *Pro-*
 1265 *ceedings, 69th Scottish Universities Summer School in Physics : LHC Phenomenology*
 1266 *(SUSSP69): St.Andrews, Scotland, August 19-September 1, 2012.* 2013, pp. 287–319.
 1267 DOI: [10.1007/978-3-319-05362-2_8](https://doi.org/10.1007/978-3-319-05362-2_8). arXiv: [1304.6677](https://arxiv.org/abs/1304.6677) [hep-ph] (see p. 36).
- 1268 [100] TORBJORN SJÖSTRAND: **Monte Carlo Tools.** In: *Proceedings, 65th Scottish Universities*
 1269 *Summer School in Physics: LHC Physics (SUSSP65): St. Andrews, UK, August 16-29,*
 1270 *2009.* 2009, pp. 309–339. DOI: [10.1201/b11865-14](https://doi.org/10.1201/b11865-14). arXiv: [0911.5286](https://arxiv.org/abs/0911.5286) [hep-ph] (see
 1271 p. 36).
- 1272 [101] STEFAN HÖCHE: **Introduction to parton-shower event generators.** In: *Proceedings,*
 1273 *Theoretical Advanced Study Institute in Elementary Particle Physics: Journeys Through*
 1274 *the Precision Frontier: Amplitudes for Colliders (TASI 2014): Boulder, Colorado, June*
 1275 *2-27, 2014.* 2015, pp. 235–295. DOI: [10.1142/9789814678766_0005](https://doi.org/10.1142/9789814678766_0005). arXiv: [1411.4085](https://arxiv.org/abs/1411.4085)
 1276 [hep-ph] (see pp. 36–37).
- 1277 [102] ADAM ALLOUL, NEIL D. CHRISTENSEN, CELINE DEGRANDE, CLAUDE DUHR, and BEN-
 1278 JAMIN FUKS: **FeynRules 2.0 - A complete toolbox for tree-level phenomenology.** In:
 1279 *Comput. Phys. Commun.*, **185**: (2014), pp. 2250–2300. DOI: [10.1016/j.cpc.2014.04.012](https://doi.org/10.1016/j.cpc.2014.04.012). arXiv: [1310.1921](https://arxiv.org/abs/1310.1921) [hep-ph] (see p. 36).
- 1281 [103] CELINE DEGRANDE, CLAUDE DUHR, BENJAMIN FUKS, et al.: **UFO - The Universal**
 1282 **FeynRules Output.** In: *Comput. Phys. Commun.*, **183**: (2012), pp. 1201–1214. DOI:
 1283 [10.1016/j.cpc.2012.01.022](https://doi.org/10.1016/j.cpc.2012.01.022). arXiv: [1108.2040](https://arxiv.org/abs/1108.2040) [hep-ph] (see p. 36).
- 1284 [104] JOHAN ALWALL, MICHEL HERQUET, FABIO MALTONI, OLIVIER MATTELAER, and TIM
 1285 STELZER: **MadGraph 5 : Going Beyond.** In: *JHEP*, **06**: (2011), p. 128. DOI: [10.1007/JHEP06\(2011\)128](https://doi.org/10.1007/JHEP06(2011)128). arXiv: [1106.0522](https://arxiv.org/abs/1106.0522) [hep-ph] (see pp. 36–37, 39).
- 1287 [105] MICHELANGELO L. MANGANO, MAURO MORETTI, FULVIO PICCININI, and MICHELE TREC-
 1288 **CANI: Matching matrix elements and shower evolution for top-quark production**
 1289 **in hadronic collisions.** In: *JHEP*, **01**: (2007), p. 013. DOI: [10.1088/1126-6708/2007/01/013](https://doi.org/10.1088/1126-6708/2007/01/013). arXiv: [hep-ph/0611129](https://arxiv.org/abs/hep-ph/0611129) [hep-ph] (see p. 36).

- 1291 [106] TORBJORN SJOSTRAND, STEFAN ASK, JESPER R. CHRISTIANSEN, et al.: **An introduction**
 1292 to **{PYTHIA} 8.2**. In: *Computer Physics Communications*, **191**: (2015), pp. 159–177.
 1293 DOI: <http://dx.doi.org/10.1016/j.cpc.2015.01.024> (see pp. 37–38).
- 1294 [107] TORBJORN SJOSTRAND, STEPHEN MRENNA, and PETER Z. SKANDS: **PYTHIA 6.4 Physics**
 1295 and **Manual**. In: *JHEP*, **0605**: (2006), p. 026. DOI: <10.1088/1126-6708/2006/05/026>.
 1296 arXiv: [hep-ph/0603175 \[hep-ph\]](hep-ph/0603175) (see pp. 37–38).
- 1297 [108] TORBJORN SJOSTRAND, STEFAN ASK, JESPER R. CHRISTIANSEN, et al.: **An Introduction**
 1298 to **PYTHIA 8.2**. In: *Comput. Phys. Commun.*, **191**: (2015), pp. 159–177. DOI:
 1299 <10.1016/j.cpc.2015.01.024>. arXiv: [1410.3012 \[hep-ph\]](1410.3012) (see pp. 37–39).
- 1300 [109] JOHAN ALWALL et al.: **Comparative study of various algorithms for the merging**
 1301 **of parton showers and matrix elements in hadronic collisions**. In: *Eur. Phys. J.*,
 1302 **C53**: (2008), pp. 473–500. DOI: <10.1140/epjc/s10052-007-0490-5>. arXiv: <0706.2569>
 1303 [hep-ph] (see p. 37).
- 1304 [110] J. ALWALL, R. FREDERIX, S. FRIXIONE, et al.: **The automated computation of tree-level**
 1305 **and next-to-leading order differential cross sections, and their matching to parton**
 1306 **shower simulations**. In: *JHEP*, **07**: (2014), p. 079. DOI: [10.1007/JHEP07\(2014\)079](10.1007/JHEP07(2014)079).
 1307 arXiv: [1405.0301 \[hep-ph\]](1405.0301) (see pp. 37, 39).
- 1308 [111] RIKKERT FREDERIX and STEFANO FRIXIONE: **Merging meets matching in MC@NLO**.
 1309 In: *JHEP*, **12**: (2012), p. 061. DOI: [10.1007/JHEP12\(2012\)061](10.1007/JHEP12(2012)061). arXiv: <1209.6215>
 1310 [hep-ph] (see p. 37).
- 1311 [112] SIMONE ALIOLI, PAOLO NASON, CARLO OLEARI, and EMANUELE RE: **A general frame-**
 1312 **work for implementing NLO calculations in shower Monte Carlo programs: the**
 1313 **POWHEG BOX**. In: *Journal of High Energy Physics*, **2010**:6 (2010), p. 43. DOI: [10.1007/JHEP06\(2010\)043](10.1007/JHEP06(2010)043) (see p. 37).
- 1315 [113] SIMONE ALIOLI, PAOLO NASON, CARLO OLEARI, and EMANUELE RE: **NLO single-top**
 1316 **production matched with shower in POWHEG: s - and t -channel contributions**.
 1317 In: *Journal of High Energy Physics*, **2009**:09 (2009), p. 111. URL: <http://stacks.iop.org/1126-6708/2009/i=09/a=111> (see p. 37).
- 1319 [114] STEFANO FRIXIONE, PAOLO NASON, and CARLO OLEARI: **Matching NLO QCD com-**
 1320 **putations with parton shower simulations: the POWHEG method**. In: *Journal of*
 1321 *High Energy Physics*, **2007**:11 (2007), p. 070. URL: <http://stacks.iop.org/1126-6708/2007/i=11/a=070> (see p. 37).
- 1323 [115] SIMONE ALIOLI, PAOLO NASON, CARLO OLEARI, and EMANUELE RE: **A general frame-**
 1324 **work for implementing NLO calculations in shower Monte Carlo programs: the**
 1325 **POWHEG BOX**. In: *JHEP*, **06**: (2010), p. 043. DOI: [10.1007/JHEP06\(2010\)043](10.1007/JHEP06(2010)043). arXiv:
 1326 [1002.2581 \[hep-ph\]](1002.2581) (see p. 37).
- 1327 [116] STEFANO FRIXIONE, PAOLO NASON, and CARLO OLEARI: **Matching NLO QCD com-**
 1328 **putations with Parton Shower simulations: the POWHEG method**. In: *JHEP*, **11**:
 1329 (2007), p. 070. DOI: <10.1088/1126-6708/2007/11/070>. arXiv: [0709.2092 \[hep-ph\]](0709.2092) (see
 1330 p. 37).
- 1331 [117] PAOLO NASON: **A New method for combining NLO QCD with shower Monte Carlo**
 1332 **algorithms**. In: *JHEP*, **11**: (2004), p. 040. DOI: <10.1088/1126-6708/2004/11/040>.
 1333 arXiv: [hep-ph/0409146 \[hep-ph\]](hep-ph/0409146) (see p. 37).

- 1334 [118] ANDREI V. GRITSAN, RAOUL RÖNTSCH, MARKUS SCHULZE, and MENG XIAO: **Constraining**
 1335 **anomalous Higgs boson couplings to the heavy flavor fermions using matrix**
 1336 **element techniques.** In: *Phys. Rev.*, D94:5 (2016), p. 055023. doi: [10.1103/PhysRevD.94.055023](https://doi.org/10.1103/PhysRevD.94.055023). arXiv: [1606.03107 \[hep-ph\]](https://arxiv.org/abs/1606.03107) (see pp. 37, 39).
- 1338 [119] IAN ANDERSON et al.: **Constraining anomalous HVV interactions at proton and**
 1339 **lepton colliders.** In: *Phys. Rev.*, D89:3 (2014), p. 035007. doi: [10.1103/PhysRevD.89.035007](https://doi.org/10.1103/PhysRevD.89.035007). arXiv: [1309.4819 \[hep-ph\]](https://arxiv.org/abs/1309.4819) (see pp. 37, 39).
- 1341 [120] SARA BOLOGNESI, YANYAN GAO, ANDREI V. GRITSAN, et al.: **On the spin and parity of**
 1342 **a single-produced resonance at the LHC.** In: *Phys. Rev.*, D86: (2012), p. 095031.
 1343 doi: [10.1103/PhysRevD.86.095031](https://doi.org/10.1103/PhysRevD.86.095031). arXiv: [1208.4018 \[hep-ph\]](https://arxiv.org/abs/1208.4018) (see pp. 37, 39).
- 1344 [121] YANYAN GAO, ANDREI V. GRITSAN, ZIJIN GUO, et al.: **Spin determination of single-**
 1345 **produced resonances at hadron colliders.** In: *Phys. Rev.*, D81: (2010), p. 075022.
 1346 doi: [10.1103/PhysRevD.81.075022](https://doi.org/10.1103/PhysRevD.81.075022). arXiv: [1001.3396 \[hep-ph\]](https://arxiv.org/abs/1001.3396) (see pp. 37, 39).
- 1347 [122] PIERRE ARTOISENET, RIKKERT FREDERIX, OLIVIER MATTELAER, and ROBBERT RIETKERK:
 1348 **Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations.** In: *JHEP*, 03: (2013), p. 015. doi: [10.1007/JHEP03\(2013\)015](https://doi.org/10.1007/JHEP03(2013)015). arXiv: [1212.3460 \[hep-ph\]](https://arxiv.org/abs/1212.3460) (see pp. 38–39).
- 1351 [123] V. KHACHATRYAN and ETAL: **Event generator tunes obtained from underlying event**
 1352 **and multiparton scattering measurements.** In: *The European Physical Journal C*,
 1353 76:3 (Mar. 17, 2016), p. 155. doi: [10.1140/epjc/s10052-016-3988-x](https://doi.org/10.1140/epjc/s10052-016-3988-x) (see p. 38).
- 1354 [124] J. PUMPLIN, D. R. STUMP, J. HUSTON, et al.: **New generation of parton distributions**
 1355 **with uncertainties from global QCD analysis.** In: *JHEP*, 07: (2002), p. 012. doi:
 1356 [10.1088/1126-6708/2002/07/012](https://doi.org/10.1088/1126-6708/2002/07/012). arXiv: [hep-ph/0201195 \[hep-ph\]](https://arxiv.org/abs/hep-ph/0201195) (see p. 39).
- 1357 [125] YUE ZHANG, Bo HUA LI, CHONG SHENG LI, JUN GAO, and HUA XING ZHU: **Next-to-**
 1358 **leading order QCD corrections to the top quark associated with γ production via**
 1359 **model-independent flavor-changing neutral-current couplings at hadron collid-**
 1360 **ers.** In: *Phys. Rev.*, D83: (2011), p. 094003. doi: [10.1103/PhysRevD.83.094003](https://doi.org/10.1103/PhysRevD.83.094003). arXiv:
 1361 [1101.5346 \[hep-ph\]](https://arxiv.org/abs/1101.5346) (see p. 39).
- 1362 [126] A. HOECKER, P. SPECKMAYER, J. STELZER, et al.: **TMVA - Toolkit for Multivariate**
 1363 **Data Analysis.** In: *ArXiv Physics e-prints*, (Mar. 2007). eprint: [physics/0703039](https://arxiv.org/abs/physics/0703039) (see
 1364 pp. 42–43).
- 1365 [127] R. BRUN and F. RADEMAKERS: **ROOT: An object oriented data analysis framework.**
 1366 In: *Nucl. Instrum. Meth.*, A389: (1997), pp. 81–86. doi: [10.1016/S0168-9002\(97\)00048-X](https://doi.org/10.1016/S0168-9002(97)00048-X) (see p. 42).
- 1368 [128] A. MAYR, H. BINDER, O. GEFELLER, and M. SCHMID: **The Evolution of Boosting**
 1369 **Algorithms - From Machine Learning to Statistical Modelling.** In: *ArXiv e-prints*,
 1370 (Mar. 2014). arXiv: [1403.1452 \[stat.ME\]](https://arxiv.org/abs/1403.1452) (see p. 43).
- 1371 [129] OLAF BEHNKE, KEVIN KRONINGER, GREGORY SCHOTT, and THOMAS SCHORNER-SADENIUS: **Data**
 1372 **Analysis in High Energy Physics: A Practical Guide to Statistical Methods.** 1st.
 1373 Wiley-VCH, 2013 (see pp. 43–44).

- 1374 [130] CHRISTIAN BÖSER, SIMON FINK, and STEFFEN RÖCKER: **Introduction to Boosted**
 1375 **Decision Trees: A multivariate approach to classification problems.** Presented at
 1376 the 'KSETA Doktoranden' workshop, Lauterbad. July 2014. URL: <https://indico.scc.kit.edu/indico/event/48/session/4/contribution/35/material/slides/0.pdf> (see
 1377 p. 44).
- 1379 [131] SERGUEI CHATRCHYAN et al.: **Combined results of searches for the standard model**
 1380 **Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV.** In: *Phys. Lett.*, **B710**: (2012), pp. 26–
 1381 48. DOI: [10.1016/j.physletb.2012.02.064](https://doi.org/10.1016/j.physletb.2012.02.064). arXiv: [1202.1488 \[hep-ex\]](https://arxiv.org/abs/1202.1488) (see p. 44).
- 1382 [132] GLEN COWAN, KYLE CRANMER, EILAM GROSS, and OFER VITELLS: **Asymptotic formulae**
 1383 **for likelihood-based tests of new physics.** In: *Eur. Phys. J.*, **C71**: (2011). [Erratum:
 1384 Eur. Phys. J.C73,2501(2013)], p. 1554. DOI: [10.1140/epjc/s10052-011-1554-0](https://doi.org/10.1140/epjc/s10052-011-1554-0), [10.1140/epjc/s10052-013-2501-z](https://doi.org/10.1140/epjc/s10052-013-2501-z). arXiv: [1007.1727 \[physics.data-an\]](https://arxiv.org/abs/1007.1727) (see
 1385 pp. 44, 48).
- 1387 [133] **Observation of a new boson with a mass near 125 GeV.** Tech. rep. CMS-PAS-HIG-
 1388 12-020. Geneva: CERN, 2012. URL: <https://cds.cern.ch/record/1460438> (see pp. 44,
 1389 48).
- 1390 [134] **Procedure for the LHC Higgs boson search combination in Summer 2011.** Tech.
 1391 rep. CMS-NOTE-2011-005. ATL-PHYS-PUB-2011-11. Geneva: CERN, Aug. 2011. URL:
 1392 [http://cds.cern.ch/record/1379837](https://cds.cern.ch/record/1379837) (see pp. 44, 46–48).
- 1393 [135] HIGGS WORKING GROUP: **Documentation of the RooStats based statistics tools**
 1394 **for Higgs PAG.** 2017. URL: <https://twiki.cern.ch/twiki/bin/viewauth/CMS/SWGuideHiggsAnalysisCombinedLimit> (see p. 44).
- 1396 [136] LORENZO MONETA, KEVIN BELASCO, KYLE S. CRANMER, et al.: **The RooStats Project.**
 1397 In: *PoS, ACAT2010*: (2010), p. 057. arXiv: [1009.1003 \[physics.data-an\]](https://arxiv.org/abs/1009.1003) (see p. 44).
- 1398 [137] GLEN COWAN, KYLE CRANMER, EILAM GROSS, and OFER VITELLS: **Asymptotic formulae**
 1399 **for likelihood-based tests of new physics.** In: *The European Physical Journal C*, **71**:2
 1400 (Feb. 2011). DOI: [10.1140/epjc/s10052-011-1554-0](https://doi.org/10.1140/epjc/s10052-011-1554-0). arXiv: [1007.1727 \[hep-ph\]](https://arxiv.org/abs/1007.1727) (see
 1401 p. 44).
- 1402 [138] THOMAS JUNK: **Confidence level computation for combining searches with small**
 1403 **statistics.** In: *Nuclear Instruments and Methods in Physics Research Section A: Accelerators,*
 1404 *Spectrometers, Detectors and Associated Equipment*, **434**:2 (1999), pp. 435–443.
 1405 DOI: [https://doi.org/10.1016/S0168-9002\(99\)00498-2](https://doi.org/10.1016/S0168-9002(99)00498-2) (see p. 45).
- 1406 [139] A L READ: **Presentation of search results: the CL s technique.** In: *Journal of Physics G: Nuclear and Particle Physics*, **28**:10 (2002), p. 2693. URL: <http://stacks.iop.org/0954-3899/28/i=10/a=313> (see p. 45).
- 1409 [140] J. S. CONWAY: **Incorporating Nuisance Parameters in Likelihoods for Multisource Spectra.** In: *Proceedings, PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding, CERN, Geneva, Switzerland 17-20 January 2011.* 2011, pp. 115–120. DOI: [10.5170/CERN-2011-006.115](https://doi.org/10.5170/CERN-2011-006.115). arXiv: [1103.0354 \[physics.data-an\]](https://arxiv.org/abs/1103.0354) (see p. 47).
- 1414 [141] SERGUEI CHATRCHYAN et al.: **Performance of CMS muon reconstruction in pp collision events at $\sqrt{s} = 7$ TeV.** In: *JINST*, **7**: (2012), P10002. DOI: [10.1088/1748-0221/7/10/P10002](https://doi.org/10.1088/1748-0221/7/10/P10002). arXiv: [1206.4071 \[physics.ins-det\]](https://arxiv.org/abs/1206.4071) (see p. 49).

- 1417 [142] R. FRÜHWIRTH: **Application of Kalman filtering to track and vertex fitting**. In:
1418 *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrom-
1419 eters, Detectors and Associated Equipment*, **262**:2 (1987), pp. 444–450. doi: [http://dx.doi.org/10.1016/0168-9002\(87\)90887-4](http://dx.doi.org/10.1016/0168-9002(87)90887-4) (see pp. 49–50).
- 1421 [143] PIERRE BILLOIR: **Progressive track recognition with a Kalman like fitting proce-
1422 dure**. In: *Comput. Phys. Commun.*, **57**: (1989), pp. 390–394. doi: [10.1016/0010-4655\(89\)90249-X](https://doi.org/10.1016/0010-4655(89)90249-X) (see pp. 49–50).
- 1424 [144] K. ROSE: **Deterministic annealing for clustering, compression, classification, re-
1425 gression, and related optimization problems**. In: *Proceedings of the IEEE*, **86**:11
1426 (Nov. 1998), pp. 2210–2239. doi: [10.1109/5.726788](https://doi.org/10.1109/5.726788) (see p. 51).
- 1427 [145] WOLFGANG WALTENBERGER: **Adaptive Vertex Reconstruction**. Tech. rep. CMS-NOTE-
1428 2008-033. Geneva: CERN, July 2008. url: <https://cds.cern.ch/record/1166320> (see
1429 p. 51).