

Quantum Theory of Toroidal Solitons

Non-Perturbative Quantization, Internal Spectrum, and Soliton Statistics

Ivan Salines

Independent Researcher

December 2025

Abstract

This thirteenth document formulates the full quantum description of the toroidal solitons introduced in Documents 1–12. While previous works established classical stability, nonlinear dynamics, and EM/GR phenomenology, the present document constructs the non-perturbative quantum mechanics of solitons. We develop collective-coordinate quantization, analyze the internal vibrational spectrum, construct soliton creation operators, define a Hilbert space of multi-torus states, and derive the statistical and thermodynamic behavior of a gas of toroidal quanta. This provides the final layer needed to interpret the toroidal soliton as a genuine quantum object.

1 Introduction

Quantum solitons behave neither like point particles nor like classical extended objects. The toroidal soliton possesses:

- an internal phase degree of freedom,
- topologically quantized winding n ,
- discrete vibrational modes,
- a conserved charge Q ,
- a stable geometric profile.

These features yield a non-perturbative quantum structure analogous to:

- instantons (topology),
- Skyrmions (collective coordinates),

- Q-balls (charge quantization),
- flux tubes (vibrational spectra).

But the combination is unique to the unified motion theory.

2 Collective-Coordinate Quantization

The toroidal soliton solution depends on four key zero modes:

1. three translations,
2. one global phase rotation (U(1) symmetry).

Introduce collective coordinates:

$$\mathbf{X}(t), \quad \Theta(t),$$

representing the soliton's center and phase.

Substitute into the action and expand:

$$\psi(\mathbf{x}, t) = \rho_0(\mathbf{x} - \mathbf{X}(t)) e^{i(\omega t + \Theta(t) + \delta\theta)}.$$

The effective Lagrangian becomes:

$$L_{\text{eff}} = \frac{1}{2} M_{\text{eff}} \dot{\mathbf{X}}^2 + Q \dot{\Theta} - E(Q, n),$$

with

$$M_{\text{eff}} = \int d^3x T_{00}.$$

Canonical momenta:

$$\mathbf{P} = M_{\text{eff}} \dot{\mathbf{X}}, \quad P_{\Theta} = Q.$$

Quantization:

$$[\hat{X}_i, \hat{P}_j] = i\delta_{ij}, \quad [\hat{\Theta}, \hat{Q}] = i.$$

Thus Q is the generator of phase rotations, consistent with Noether charge quantization.

3 Quantum Internal Spectrum

From the nonlinear stability analysis (Document 10), internal modes satisfy:

$$-\Omega_k^2 M \zeta_k = K \zeta_k.$$

Quantization gives creation/annihilation operators:

$$\hat{a}_k, \hat{a}_k^\dagger, \quad [\hat{a}_k, \hat{a}_{k'}^\dagger] = \delta_{kk'}.$$

Energy levels:

$$E_k = \Omega_k \left(n_k + \frac{1}{2} \right).$$

Key quantized modes:

- **Breathing mode:** $\Omega_{\text{br}},$
- **Torsional mode:** $\Omega_{\text{tors}} \propto n/R_c,$
- **Kelvin-like bending modes:** $\Omega_m \propto m/R_c,$
- **Azimuthal phonons:** $\Omega_k \propto k/R_c.$

This predicts a discrete internal spectrum observable in GW/EM signals.

4 Soliton Creation and Annihilation Operators

Define a non-perturbative creation operator:

$$\hat{\mathcal{O}}^\dagger(Q, n) = \exp \left(\int d^3x \Phi_0(x; Q, n) \hat{\psi}^\dagger(x) \right).$$

Acting on the vacuum:

$$|\mathcal{T}_{Q,n}\rangle = \hat{\mathcal{O}}^\dagger(Q, n) |0\rangle.$$

This state is:

- not perturbative,
- not Gaussian,
- a coherent configuration of infinitely many quanta of $\hat{\psi},$
- stabilized by nonlinearity and topology.

Multi-soliton states follow naturally:

$$|\mathcal{T}_{Q_1, n_1} \mathcal{T}_{Q_2, n_2} \cdots \rangle = \prod_i \hat{\mathcal{O}}^\dagger(Q_i, n_i) |0\rangle.$$

5 Quantum Interactions and Scattering

Soliton scattering is governed by:

$$S = \exp \left(i \int d^4x \mathcal{L}_{\text{int}}(\psi, \rho_0, \theta_0) \right).$$

Collective-coordinate quantization yields:

$$\langle \mathcal{T}_f | S | \mathcal{T}_i \rangle = \int \mathcal{D}X(t) \mathcal{D}\Theta(t) e^{iS_{\text{eff}}[X, \Theta]}.$$

Predictions:

- quantized scattering angles,
- resonant enhancement when Ω_k matches impact frequency,
- tunneling between different n -sectors is forbidden,
- fusion creates quantum superpositions:

$$|\mathcal{T}_{Q_1} \mathcal{T}_{Q_2}\rangle \rightarrow \sum_{\alpha} c_{\alpha} |\mathcal{T}_{Q_1+Q_2}, \alpha\rangle.$$

6 Quantum Statistics of Toroidal Solitons

The soliton carries:

- conserved charge Q ,
- winding number n ,
- internal phase Θ ,
- geometric radius R_c ,
- internal excitations.

Because it is an extended object with topological data:

- it is not a boson or fermion,
- it behaves as a **geometric anyon-like excitation**,
- exchange induces a Berry-phase-like rotation:

$$\phi_{\text{exch}} = n_1 n_2 \Delta\theta.$$

Thus toroidal solitons obey **generalized quantum statistics**.

7 Thermodynamics of Soliton Gases

A dilute gas of toroidal quanta has partition function:

$$Z = \sum_{\{N_k\}} \exp \left[-\beta \left(\sum_k N_k E_k + \sum_{i < j} V_{\text{int}}(i, j) \right) \right].$$

Phenomena:

- condensation into lowest torsional mode,
- ring-lattice formation,
- soliton crystals for large Q ,
- topological phase transitions.

These phases have no analogue in standard scalar QFT.

8 Conclusion

The toroidal soliton is a fully consistent quantum object with:

- quantized collective coordinates,
- discrete internal spectrum,
- creation and annihilation operators,
- generalized anyonic statistics,
- well-defined thermodynamics,
- non-perturbative scattering amplitudes.

This completes the quantum layer of the unified motion theory and positions the toroidal soliton as a true *elementary quantum of geometry and motion*.