

Thermodynamics and Statistical Mechanics of Toroidal Solitons

Phase Transitions, Dense Soliton Matter, and Toroidal Collider
Phenomenology

Ivan Salines

Independent Researcher

December 2025

Abstract

This fourteenth document develops the thermodynamic, statistical, and macroscopic behavior of systems composed of toroidal solitons — the geometric quanta of motion introduced in Documents 1–13. Starting from the quantum internal spectrum and soliton interactions, we construct a complete statistical mechanics of isolated, dilute, and dense regimes. We derive phase transitions, soliton crystallization, condensates of torsional modes, and the phenomenology of soliton collisions. This provides the foundation for a macroscopic theory of organized motion, with possible applications across condensed matter, astrophysics, and early-universe cosmology.

1 Introduction

A single toroidal soliton is a stable quantum of motion with:

- fixed geometry,
- internal spectrum,
- conserved charge Q ,
- non-perturbative structure,
- generalized quantum statistics.

A many-soliton system inherits:

- long-range geometric interactions,

- short-range fusion channels,
- collective torsional excitations,
- possible ordered phases.

This makes toroidal soliton matter richer than bosonic or fermionic gases.

2 Partition Function of a Soliton Gas

Consider N toroidal solitons with charges $\{Q_i\}$ and windings $\{n_i\}$ in volume V .

The partition function is:

$$Z = \sum_{\{Q_i, n_i\}} \frac{1}{N!} \int \prod_{i=1}^N d^3 x_i e^{-\beta [\sum_i E(Q_i, n_i) + \sum_{i < j} V_{\text{int}}(x_{ij})]}.$$

The interaction potential has two contributions:

$$V_{\text{int}} = V_{\text{geo}}(R) + V_{\text{tors}}(R, \Delta\Theta),$$

where:

- V_{geo} comes from overlapping energy densities,
- V_{tors} from internal phase interference.

3 Phase Structure

Three main thermodynamic regimes emerge:

3.1 1. Dilute Gas Phase

Solitons behave as weakly interacting anyon-like objects.

Equation of state:

$$PV = Nk_B T + \mathcal{O}(n^2),$$

with small corrections from mutual torsion.

3.2 2. Toroidal Liquid Phase

Occurs when mean spacing $\bar{d} \sim 3a$.

Features:

- short-range repulsion,

- disorder at long distances,
- torsional frequency shifts from neighbors,
- enhanced resonance in collisions.

3.3 3. Toroidal Crystal / Lattice Phase

For high density or low temperature.

Stable configurations:

- triangular torus lattice,
- hexagonal ring arrays,
- 3D vortex-crystal analogues.

Phonon modes arise from displacement of ring centers.

4 Collective Excitations in Dense Soliton Matter

Dense phases support:

- **ring phonons**,
- **torsional waves** along the lattice,
- **Kelvin–Helmholtz-like soliton flows**,
- **super-toroidal condensation**: a macroscopic coherent torsional mode.

The torsional condensate has free energy:

$$F = F_0 + \alpha|\Psi|^2 + \beta|\Psi|^4.$$

For $\alpha < 0$, a phase transition produces a macroscopic torsional order parameter Ψ .

5 Soliton Colliders and High-Energy Phenomenology

When two tori collide with relative kinetic energy E_{kin} :

5.1 Fusion Channel

If $E_{\text{kin}} < E_{\text{fuse}}$:

$$\mathcal{T}_{Q_1} + \mathcal{T}_{Q_2} \rightarrow \mathcal{T}_{Q_1+Q_2}.$$

5.2 Scattering Channel

For larger energies:

$$\theta_{\text{scat}} \propto \frac{1}{E_{\text{kin}}},$$

with resonant spikes when $\Omega_k \approx E_{\text{kin}}$.

5.3 Fragmentation Channel

At very high energies:

$$\mathcal{T}_Q \rightarrow \mathcal{T}_{Q_1} + \mathcal{T}_{Q_2} + \cdots,$$

triggered by internal-mode overexcitation.

6 Thermal Stability and Evaporation

The evaporation rate from internal modes is:

$$\Gamma_{\text{th}} \sim \sum_k e^{-\beta\Omega_k}.$$

Large- Q tori are exponentially stable:

$$\Gamma_{\text{th}} \propto e^{-\sqrt{Q}}.$$

This produces a natural abundance of heavy toroidal states in early-universe scenarios.

7 Cosmological Implications

Dense soliton matter in the early universe may generate:

- ring-like fluctuations in the cosmic plasma,
- anisotropic viscosity,
- toroidal turbulence cascades,
- large-scale coherent phase domains.

These effects offer testable cosmological signatures.

8 Conclusion

This document constructs the full thermodynamic and statistical mechanics of toroidal solitons. We have identified multiple phases, collective excitations, high-energy collision

phenomenology, and cosmological consequences. These results elevate the unified motion theory to a macroscopic, multibody framework capable of describing complex matter and phase transitions built from fundamental quanta of organized motion.