

Brazil-Malvinas Confluence

Ivenis Pita

Final Project


Large Scale Ocean Circulation

Content

- Brazil-Malvinas Confluence (BMC)
- Drifter Dataset
- Surface Velocity
- Location of the BMC
- Takeaway notes


Brazil-Malvinas Confluence (BMC)

- The mean position of the BMC is located near 38 °S
 - Influenced by seasonality (Olson et al. 1988; Garzoli and Garaffo 1989).
 - The driving processes for this seasonal variability remain unclear (Julion et al. 2010).
 - Variability of the BMC position is partially influenced by variations in the MC transport (MODEL)
 - linked to fluctuations in the ACC transport and in the wind stress over the Southern Ocean (Matano 1993; Wainer et al. 2000; Fetter and Matano 2008).
 - Variability of the position of the BMC is forced primarily by the local wind field (Garzoli and Giulivi 1994) (OBS).





Brazil-Malvinas Confluence (BMC)

- The circulation of the BMC is greatly influenced by the presence of the Zapiola Rise
 - a sediment mount that extends zonally along 45 °S between 36° and 51 °W.
 - Strong (~ 100 Sv) anticyclonic circulation (Saunders and King 1995).
 - Region of low eddy kinetic energy (EKE) surrounded by high EKE.
 - Saraceno et al. (2004) argued that a closed planetary vorticity contour around the rise acts as a boundary isolating the center of the circulation cell from the highly variable BMC.




Drifter Dataset

- 6 hourly- interpolated data
 - From 2000 to 2019
 - -50 to -20° S
 - -60 to -30° W





AOML receives position data computed from Doppler measurements from Argos-tracked drifters or from GPS installed in all Iridium and some Argos drifters. Historically, these data are irregularly distributed in time and they also contain erroneous positions and/or bad Sea Surface Temperature values, therefore not suited for many kinds of analysis or displays. The Drifter Data Assembly Center (DAC) at AOML applies quality control procedures to edit these data (position and temperature) and interpolates them to 6-hour intervals using an optimum interpolation procedure called kriging, which is commonly used for two and three-dimensional analyses. Some drifters have other sensors attached such as barometric pressure, salinity, wind speed and direction, but these data are stored in their raw form and kept in the archives with no further editing done at the present time. See Hansen, D.V. and Pouliquen



Drifter Dataset




Surface Velocity




Surface Velocity




Surface Velocity (Seasons)




Location of the BMC





Location of the BMC



Location of the BMC



Eddy Kinetic Energy



Takeaway notes

- Zapiola Rise limits the BMC location
- Local and remote winds (Southern Ocean) can influence BMC variability
- Location of BMC is better inferred by drifter database during summer and fall months
- BMC has greater EKE - > possible instability growth
- Lower EKE region close to Zapiola Rise

Thanks!