

“Intra-Seasonal Waning” as Methodological Artifact

Ivo M. Foppa

10/10/2018

Lipsitch editorial (CID, 2018):

Lipsitch editorial (CID, 2018):

“Even if the vaccine effect does not wane over time, then such waning will nonetheless appear to occur in most studies [. . .]” (“leaky” vaccines)

Lipsitch editorial (CID, 2018):

“Even if the vaccine effect does not wane over time, then such waning will nonetheless appear to occur in most studies [. . .]” (“leaky” vaccines)

This will happen if:

1. “there is heterogeneous risk of becoming infected within those who are vaccinated”
2. “some trial participants during the course of the trial become infected but are not counted as cases”

Lipsitch editorial (CID, 2018):

“Even if the vaccine effect does not wane over time, then such waning will nonetheless appear to occur in most studies [. . .]” (“leaky” vaccines)

This will happen if:

1. “there is heterogeneous risk of becoming infected within those who are vaccinated”
2. “some trial participants during the course of the trial become infected but are not counted as cases” ← **Not relevant to TND studies! But:**

Lipsitch editorial (CID, 2018):

“Even if the vaccine effect does not wane over time, then such waning will nonetheless appear to occur in most studies [...]” (“leaky” vaccines)

This will happen if:

1. “there is heterogeneous risk of becoming infected within those who are vaccinated”
2. “some trial participants during the course of the trial become infected but are not counted as cases” ← **Not relevant to TND studies! But:** Faster depletion of unvaccinated than vaccinated ($e^{-\Lambda_t} < e^{-\Lambda_t\theta}$), shifting the odds of case odds in vaccinated.

Lipsitch editorial (CID, 2018):

“Even if the vaccine effect does not wane over time, then such waning will nonetheless appear to occur in most studies [...]” (“leaky” vaccines)

This will happen if:

1. “there is heterogeneous risk of becoming infected within those who are vaccinated”
2. “some trial participants during the course of the trial become infected but are not counted as cases” ← **Not relevant to TND studies! But:** Faster depletion of unvaccinated than vaccinated ($e^{-\Lambda_t} < e^{-\Lambda_t\theta}$), shifting the odds of case odds in vaccinated.
3. What if the vaccine is not leaky?

Vaccine models

- ▶ “Leaky” model: Those susceptible before vaccination have a risk of $\lambda_1 = \lambda_0 \theta$ of becoming infected during a contact if an unvaccinated susceptible has risk λ_0 .

$$VE = 1 - \frac{\lambda_1}{\lambda_0} = 1 - \theta$$

Vaccine models

- ▶ “Leaky” model: Those susceptible before vaccination have a risk of $\lambda_1 = \lambda_0 \theta$ of becoming infected during a contact if an unvaccinated susceptible has risk λ_0 .

$$VE = 1 - \frac{\lambda_1}{\lambda_0} = 1 - \theta$$

- ▶ “All-or-none” model: A proportion φ of those vaccinated becomes fully immunized against “influenza virus”, while the rest remains fully susceptible. $VE = \varphi$

Vaccine models

- ▶ “Leaky” model: Those susceptible before vaccination have a risk of $\lambda_1 = \lambda_0\theta$ of becoming infected during a contact if an unvaccinated susceptible has risk λ_0 .

$$VE = 1 - \frac{\lambda_1}{\lambda_0} = 1 - \theta$$

- ▶ “All-or-none” model: A proportion φ of those vaccinated becomes fully immunized against “influenza virus”, while the rest remains fully susceptible. $VE = \varphi$
- ▶ Hybrid models:
 1. A **proportion** φ of vaccinated develop **“leaky” immunity** (Lewnard, 2018) $VE = \varphi(1 - \theta)$
 2. “Leaky” model with time-variant θ
 3. ...

Vaccine models

- ▶ “Leaky” model: Those susceptible before vaccination have a risk of $\lambda_1 = \lambda_0 \theta$ of becoming infected during a contact if an unvaccinated susceptible has risk λ_0 .

$$VE = 1 - \frac{\lambda_1}{\lambda_0} = 1 - \theta$$

- ▶ “All-or-none” model: A proportion φ of those vaccinated becomes fully immunized against “influenza virus”, while the rest remains fully susceptible. $VE = \varphi$
- ▶ Hybrid models:
 1. A **proportion** φ of vaccinated develop “**leaky**” **immunity** (Lewnard, 2018) $VE = \varphi(1 - \theta)$
 2. “Leaky” model with time-variant $\theta \leftarrow$
 3. ...

Waning

└ Methodological sources of "waning effect"

└ Approach

Approach

Approach


1.
 - ▶ Simulation of seasonal influenza epidemics using simple SIR ODE models
 - ▶ Implement two scenarios (“leaky”, “all-or-none” with two viruses)
 - ▶ Use numerical solutions to ODEs to generate TND data
 - ▶ Calculate VE “estimates” and true VE

Approach


1.
 - ▶ Simulation of seasonal influenza epidemics using simple SIR ODE models
 - ▶ Implement two scenarios (“leaky”, “all-or-none” with two viruses)
 - ▶ Use numerical solutions to ODEs to generate TND data
 - ▶ Calculate VE “estimates” and true VE
2.
 - ▶ Simulation of seasonal influenza epidemics using a stochastic SIR model, keeping track of time since vaccination
 - ▶ Implement “all-or-none” with two viruses: A certain proportion of the population, if vaccinated, remains susceptible to virus 1, 2, 1 and 2 or neither
 - ▶ Use Ray’s analytic approach (only vaccinees, conditional logistic regression)

First scenario: "Leaky" vaccine

Vaccination coverage: 0.47, constant; $R_0 = 1.6$; $\delta = 0.25$




Rel. Bias




Second scenario: 2 viruses, “all-or-none”

Vaccination coverage: 0.3; $R_0 = 1.25, 1.75$; $\delta = 0.33$



Time-dependent θ



“All-or-none”, 2 viruses, stochastic simulation, time-since-vaccination

- ▶ Stochastic simulation with time-since vaccination (TSV)
- ▶ Normally-dist. vaccination uptake (cumm. 0.47), starting 100 days before transmission, continuing until 200 days after seeding;
 $R_0 = 1.8, 1.7$; Proportion of population susceptible (after vacc.) to virus 1, 2, neither or both: 0.2, 0.4, 0.3, 0.1
- ▶ All infections used as cases, control-case ratio 1:3
- ▶ Conditional logistic regression, with time since vaccination: < 60 , $61 - 120, 121 - 180, > 181$ days; conditioning on day of enrollment

“All-or-none”, 2 viruses, stochastic simulation, time-since-vaccination

- ▶ Stochastic simulation with time-since vaccination (TSV)
- ▶ Normally-dist. vaccination uptake (cumm. 0.47), starting 100 days before transmission, continuing until 200 days after seeding;
 $R_0 = 1.8, 1.7$; Proportion of population susceptible (after vacc.) to virus 1, 2, neither or both: 0.2, 0.4, 0.3, 0.1
- ▶ All infections used as cases, control-case ratio 1:3
- ▶ Conditional logistic regression, with time since vaccination: < 60 , $61 - 120$, $121 - 180$, > 181 days; conditioning on day of enrollment
- ▶ One simulation:

“All-or-none”, 2 viruses, stochastic simulation, time-since-vaccination

- ▶ Stochastic simulation with time-since vaccination (TSV)
- ▶ Normally-dist. vaccination uptake (cumm. 0.47), starting 100 days before transmission, continuing until 200 days after seeding;
 $R_0 = 1.8, 1.7$; Proportion of population susceptible (after vacc.) to virus 1, 2, neither or both: 0.2, 0.4, 0.3, 0.1
- ▶ All infections used as cases, control-case ratio 1:3
- ▶ Conditional logistic regression, with time since vaccination: < 60 , $61 - 120, 121 - 180, > 181$ days; conditioning on day of enrollment
- ▶ One simulation:

| Variable | OR |
|----------|------|
| TSV 1 | Ref. |
| TSV 2 | 3.62 |
| TSV 3 | 2.13 |
| TSV 4 | 5.92 |