Formulas where Subject Appears Once

Exam code: 0607

Download PDF

Written by: Mark Curtis
Reviewed by: Dan Finlay
Updated on 27 January 2026

Simple rearranging

What are formulas?

- A **formula** is a rule, definition or relationship between different quantities, written in shorthand using **letters** (**variables**)
  - Formulas include an **equals** sign
- Some examples you should be **familiar** with are:
  - The equation of a straight line
    - $y = mx + c$
  - The area of a trapezium
    - $\text{Area} = \frac{(a + b)h}{2}$
  - Pythagoras' theorem
    - $a^2 + b^2 = c^2$

How do I rearrange formulas?

- The letter (**variable**) that is on its **own** on one side is called the **subject**
  - y is the subject of $y = mx + c$
- To make a different letter the subject, we need to **rearrange the formula**
  - This is also called **changing the subject**
- The method is as follows:
  - First, **remove** any **fractions**
    - Multiply both sides by the lowest common denominator
  - Then use **inverse** (opposite) **operations** to get the variable on its own
    - This is similar to solving equations
- For example, make x the subject of $\frac{5x + 6}{2} = y$
  - First remove fractions
    - Multiply both sides by 2
      $5x + 6 = 2y$
  - Then get x on its own

Stuck on something?
Ask questions and get help with exactly what you're revising.
Got it

$$x = \frac{2y - 6}{5}$$

- There may be more than one correct way to write an answer
  - The following are acceptable **alternative** forms

$$x = \frac{2y}{5} - \frac{6}{5}$$
$$x = \frac{2(y - 3)}{5}$$
$$x = 0.4(y - 3)$$
$$x = 0.4y - 1.2$$

Should I expand brackets?

- If the variable is inside the brackets, you can either:
  - **expand** the brackets
  - or **divide** both by the coefficient if it is not zero
- For example, make x the subject of $3(1 + x) = y$
  - You can expand and then rearrange
    - $3 + 3x = y$
    - $3x = y - 3$
    - $x = \frac{y - 3}{3}$
  - You can divide and then rearrange
    - $1 + x = \frac{y}{3}$
    - $x = \frac{y}{3} - 1$
  - Both answers are equivalent
    - $\frac{y - 3}{3}$ is the same as $\frac{y}{3} - 1$

Examiner Tips and Tricks

Stop and think when you see a bracket. If the variable you are trying to make the subject is not inside the bracket, then you do not need to expand.

For example, to make x the subject of $(1 + k)x = y$, you can simply divide both sides by $(1 + k)$ to get $x = \frac{y}{1 + k}$.

What if I get fractions in fractions?

- Some rearrangements can lead to **fractions in fractions**
  - $x = \frac{\frac{3}{t}}{2}$
- Either rewrite with a **divide sign**, $\div$, then use the method of **dividing two fractions**

$$x = \frac{3}{t} \times \frac{1}{2}$$

$$x = \frac{3}{2t}$$

- Or **multiply top and bottom** by the **lowest common denominator** of the two fractions and **cancel**
  - $x = \dfrac{\frac{5}{y}}{\frac{t}{8}}$ becomes $x = \dfrac{\frac{5}{y} \times 8y}{\frac{t}{8} \times 8y} = \dfrac{40}{ty}$

What if I end up dividing by a negative?

- Remember that $\dfrac{a}{-b}$ (minus **below**) is the same as $\dfrac{-a}{b}$ (minus **above**) and the same as $-\dfrac{a}{b}$ (minus **outside**)
  - Though be careful, as $\dfrac{-a}{-b}$ is $\dfrac{a}{b}$
- $-2x = y - 3$ becomes $x = \dfrac{y-3}{-2}$ (minus **below**)
  - This is the same as $x = \dfrac{-(y-3)}{2}$ (minus **above**) or $-\dfrac{y-3}{2}$ (minus **outside**)
    - **brackets** are **required** for minus **above**
    - **brackets** are **assumed** for minus **outside**
  - You can also **expand** the brackets $\dfrac{-(y-3)}{2} = \dfrac{-y+3}{2} = \dfrac{3-y}{2}$

Examiner Tips and Tricks

Mark schemes will accept different forms of the same answer, as long as they are correct and fully simplified.

Worked Example

Make x the subject of the following.

(a) $4m + 5x = 3$

Answer:

Get $5x$ on its own by subtracting $4m$ from both sides

$$5x = 3 - 4m$$

Get x on its own by dividing both sides by 5

(b) $3t = \frac{2}{x}$

Answer:

Remove fractions by multiplying both sides by the denominator, x

$$3tx = 2$$

Get x on its own by dividing both sides by $3t$

$$x = \frac{2}{3t}$$

(c) $A = \frac{9(1-4x)}{2g}$

Answer:

Remove fractions by multiplying both sides by the denominator, $2g$

$$2gA = 9(1-4x)$$

x is inside the brackets
Expand the brackets to release the x term

$$2gA = 9 - 36x$$

One way to get x on its own is by subtracting 9 then dividing by -36
Or you can first add $36x$ to both sides, to create positive $36x$ on the left

$$2gA + 36x = 9$$

Now get x on its own by subtracting $2gA$ then dividing by 36

$$36x = 9 - 2gA$$
$$x = \frac{9-2gA}{36}$$

$$x = \frac{9-2gA}{36}$$

Other accepted forms of the answer are

$$\frac{2gA-9}{-36}, \quad \frac{-(2gA-9)}{36}, \quad -\frac{2gA-9}{36}, \quad \frac{1}{4} - \frac{gA}{18}$$

Teacher only

Test your students on this in just a few clicks

- Create exam-aligned tests
- Tailor format & difficulty
- Quizzes or homework in minutes

Test yourself

Was this revision note helpful? Yes No

Previous Next

Course Study tools All topics

Resources

Learning Hub
Scholarship
Join
Past Papers
Solution Banks
Sitemap

Dashboard

Overview
Brand ambassador
Profile
Subscription
Settings
Log out

Company

About us
Exam Specificity
Content Quality
Promotions
Jobs
Terms
Privacy
Cookie Policy
Help and Support

Subjects

Biology
Chemistry
Physics
Maths
Geography
English Literature
Psychology
All Subjects

© Copyright 2015–2026 Save My Exams Ltd. All Rights Reserved.