

CHAPTER 9

9.1

$$\begin{bmatrix} 2 & 0 & 5 \\ 0 & 1 & 1 \\ 8 & 3 & 0 \end{bmatrix} \begin{Bmatrix} x_1 \\ x_2 \\ x_3 \end{Bmatrix} = \begin{Bmatrix} 40 \\ 10 \\ 20 \end{Bmatrix} \quad [A]^T = \begin{bmatrix} 2 & 0 & 8 \\ 0 & 1 & 3 \\ 5 & 1 & 0 \end{bmatrix}$$

9.2 (a) Possible multiplications:

$$[A][B] = \begin{bmatrix} 23.5 & -2 \\ 52 & 16 \\ -18 & 8 \end{bmatrix} \quad [A][C] = \begin{bmatrix} 15 & -13 \\ 0 & -16 \\ -22 & 14 \end{bmatrix} \quad [B][C] = \begin{bmatrix} -7 & 1 \\ -5 & 1 \end{bmatrix} \quad [C][B] = \begin{bmatrix} 7 & -4 \\ -11.5 & -2 \end{bmatrix}$$

Note: Some students might recognize that we can also compute $[B][B]$ and $[C][C]$:

$$[B][B] = \begin{bmatrix} 16 & 0 \\ 3 & 4 \end{bmatrix} \quad [C][C] = \begin{bmatrix} 10 & -6 \\ -9 & 7 \end{bmatrix}$$

(b) $[B][A]$ and $[C][A]$ are impossible because the inner dimensions do not match:

$$(2 \times 2) * (3 \times 2)$$

(c) According to **(a)**, $[B][C] \neq [C][B]$

$$\begin{array}{llll} \mathbf{9.3 (a)} & [A] = 3 \times 2 & [B] = 3 \times 3 & [C] = 3 \times 1 & [D] = 2 \times 4 \\ & [E] = 3 \times 3 & [F] = 2 \times 3 & [G] = 1 \times 3 & \end{array}$$

(b) Square: $[B]$ and $[E]$ Column: $[C]$ Row: $[G]$

$$\begin{array}{lll} \mathbf{(c)} & a_{12} = 5 & b_{23} = 6 & d_{32} = \text{does not exist} \\ & e_{22} = 2 & f_{12} = 0 & g_{12} = 5 \end{array}$$

(d)

$$(1) [E] + [B] = \begin{bmatrix} 5 & 8 & 16 \\ 8 & 4 & 9 \\ 6 & 0 & 10 \end{bmatrix} \quad (2) [A] + [F] = \text{not possible}$$

$$(3) [B] - [E] = \begin{bmatrix} 3 & -2 & -2 \\ -6 & 0 & 3 \\ -2 & 0 & -2 \end{bmatrix} \quad (4) 7[B] = \begin{bmatrix} 28 & 21 & 49 \\ 7 & 14 & 42 \\ 14 & 0 & 28 \end{bmatrix}$$

$$(5) [A] \times [B] = \text{not possible} \quad (6) \{C\}^T = \begin{bmatrix} 3 & 5 & 1 \end{bmatrix}$$

$$(7) [B] \times [A] = \begin{bmatrix} 54 & 68 \\ 36 & 45 \\ 28 & 34 \end{bmatrix} \quad (8) \{D\}^T = \begin{bmatrix} 9 & 2 \\ 4 & -1 \\ 3 & 6 \\ -6 & 5 \end{bmatrix}$$


$$(9) [A] \times [C] = \text{not possible} \quad (10) [I] \times [B] = [B]$$

$$(11) [E]^T [E] = \begin{bmatrix} 66 & 19 & 54 \\ 19 & 29 & 51 \\ 54 & 51 & 126 \end{bmatrix} \quad (12) [C]^T [C] = 35$$

9.4 The equations can be rearranged into a format for plotting x_2 versus x_1 :

$$x_2 = 3 - 0.3333x_1$$

$$x_2 = 5 - \frac{1}{8}x_1$$



Therefore, the solution is $x_1 = 9.6$, $x_2 = 6.2$. The results can be checked by substituting them back into the original equations:


$$2(9.6) - 6(6.2) = -18$$

$$-9.6 + 8(6.2) = 40$$


9.5 (a) The equations can be rearranged into a format for plotting x_2 versus x_1 :

$$x_2 = 14.25 - 0.77x_1$$

$$x_2 = 11.76471 - \frac{1.2}{1.7}x_1$$



If you zoom in, it appears that there is a root at about $(38.7, -15.6)$.



The results can be checked by substituting them back into the original equations:

$$0.77(38.7) - 15.6 = 14.2 \cong 14.25$$

$$1.2(38.7) + 1.7(-15.6) = 19.92 \cong 20$$

(b) The plot suggests that the system may be ill-conditioned because the slopes are so similar.

(c) The determinant can be computed as

$$D = 0.77(1.7) - 1(1.2) = 0.11$$

which is relatively small. Note that if the system is normalized first by dividing each equation by the largest coefficient,

$$0.77x_1 + x_2 = 14.25$$

$$0.705882x_1 + x_2 = 11.76471$$

the determinant is even smaller

$$D = 0.77(1) - 1(0.705882) = 0.064$$

(d) Using Eqs. (9.10) and (9.11) yields

$$x_1 = \frac{14.25(1.7) - 1(20)}{0.109} = 38.761$$

$$x_2 = \frac{0.77(20) - 14.25(1.2)}{0.109} = -15.596$$

9.6 (a) The determinant can be computed as:

$$A_1 = \begin{vmatrix} 1 & 1 \\ 1 & 0 \end{vmatrix} = 1(0) - 1(1) = -1$$

$$A_2 = \begin{vmatrix} 2 & 1 \\ 3 & 0 \end{vmatrix} = 2(0) - 1(3) = -3$$

$$A_3 = \begin{vmatrix} 2 & 1 \\ 3 & 1 \end{vmatrix} = 2(1) - 1(3) = -1$$

$$D = 0(-1) - 2(-3) + 5(-1) = 1$$

(b) Cramer's rule

$$x_1 = \frac{\begin{vmatrix} 1 & 2 & 5 \\ 1 & 1 & 1 \\ 2 & 1 & 0 \end{vmatrix}}{D} = \frac{-2}{1} = -2$$

$$x_2 = \frac{\begin{vmatrix} 0 & 1 & 5 \\ 2 & 1 & 1 \\ 3 & 2 & 0 \end{vmatrix}}{D} = \frac{8}{1} = 8$$

$$x_3 = \frac{\begin{vmatrix} 0 & 2 & 1 \\ 2 & 1 & 1 \\ 3 & 1 & 2 \end{vmatrix}}{D} = \frac{-3}{1} = -3$$

(c) The results can be checked by substituting them back into the original equations:

$$2(8) + 5(-3) = 1$$


$$2(-2) + 8 - 3 = 1$$

$$3(-2) + 8 = 2$$

9.7 (a) The equations can be rearranged into a format for plotting x_2 versus x_1 :

$$x_2 = 9.5 + 0.5x_1$$

$$x_2 = 9.4 + 0.51x_1$$



The solution is $x_1 = 10$, $x_2 = 14.5$. Notice that the lines have very similar slopes.

(b) The determinant can be computed as

$$D = 0.5(-2) - (-1)1.02 = 0.02$$

(c) The plot and the low value of the determinant both suggest that the system is ill-conditioned.

(d) Using Eqs. (9.10) and (9.11) yields

$$x_1 = \frac{-9.5(-2) - (-1)(-18.8)}{0.02} = 10$$

$$x_2 = \frac{0.5(-18.8) - (-9.5)1.02}{0.02} = 14.5$$

(e) Using Eqs. (9.10) and (9.11) yields

$$x_1 = \frac{-9.5(-2) - (-1)(-18.8)}{-0.02} = -10$$

$$x_2 = \frac{0.52(-18.8) - (-9.5)1.02}{-0.02} = 4.3$$

The ill-conditioned nature of the system is illustrated by the fact that a small change in one of the coefficients results in a huge change in the results.

9.8 (a) The system is first expressed as an augmented matrix:

$$\begin{bmatrix} 10 & 2 & -1 & 27 \\ -3 & -6 & 2 & -61.5 \\ 1 & 1 & 5 & -21.5 \end{bmatrix}$$

Forward elimination:

a_{21} is eliminated by multiplying row 1 by $-3/10$ and subtracting the result from row 2. a_{31} is eliminated by multiplying row 1 by $1/10$ and subtracting the result from row 3.

$$\begin{bmatrix} 10 & 2 & -1 & 27 \\ 0 & -5.4 & 1.7 & -53.4 \\ 0 & 0.8 & 5.1 & -24.2 \end{bmatrix}$$

a_{32} is eliminated by multiplying row 2 by $0.8/(-5.4)$ and subtracting the result from row 3.

$$\begin{bmatrix} 10 & 2 & -1 & 27 \\ 0 & -5.4 & 1.7 & -53.4 \\ 0 & 0 & 5.351852 & -32.1111 \end{bmatrix}$$

Back substitution:

$$x_3 = \frac{-32.1111}{5.351852} = -6$$

$$x_2 = \frac{-53.4 - 1.7(-6)}{-5.4} = 8$$

$$x_1 = \frac{27 - (-1)(-6) - 2(8)}{10} = 0.5$$

(b) Check:

$$10(0.5) + 2(8) - (-6) = 27$$

$$-3(0.5) - 6(8) + 2(-6) = -61.5$$

$$0.5 + 8 + 5(-6) = -21.5$$

9.9 (a) The system is first expressed as an augmented matrix:

PROPRIETARY MATERIAL. © The McGraw-Hill Companies, Inc. All rights reserved. No part of this Manual may be displayed, reproduced or distributed in any form or by any means, without the prior written permission of the publisher, or used beyond the limited distribution to teachers and educators permitted by McGraw-Hill for their individual course preparation. If you are a student using this Manual, you are using it without permission.

$$\begin{bmatrix} 4 & 1 & -1 & -2 \\ 5 & 1 & 2 & 4 \\ 6 & 1 & 1 & 6 \end{bmatrix}$$

Forward elimination: First, we pivot by switching rows 1 and 3:

$$\begin{bmatrix} 6 & 1 & 1 & 6 \\ 5 & 1 & 2 & 4 \\ 4 & 1 & -1 & -2 \end{bmatrix}$$

Multiply row 1 by 5/6 and subtract from row 2 to eliminate a_{21} . Multiply row 1 by 4/6 and subtract from row 3 to eliminate a_{31} .

$$\begin{bmatrix} 6 & 1 & 1 & 6 \\ 0 & 0.16667 & 1.16667 & -1 \\ 0 & 0.33333 & -1.6667 & -6 \end{bmatrix}$$

Pivot:

$$\begin{bmatrix} 6 & 1 & 1 & 6 \\ 0 & 0.33333 & 1.6667 & -6 \\ 0 & 0.16667 & -1.1667 & -1 \end{bmatrix}$$

Multiply row 2 by $0.16667/0.33333 = 0.5$ and subtract from row 3 to eliminate a_{32} .

$$\begin{bmatrix} 6 & 1 & 1 & 6 \\ 0 & 0.33333 & 1.6667 & -6 \\ 0 & 0 & 2 & 2 \end{bmatrix}$$

Back substitution:

$$x_3 = \frac{2}{2} = 1$$

$$x_2 = \frac{-6 - (-1.6667)(1)}{0.33333} = -13$$

$$x_1 = \frac{6 - 1 - (-13)}{6} = 3$$

Check:

$$4(3) - 13 - 3 = -2$$

$$5(3) - 13 + 2(1) = 4$$

$$6(3) - 13 + 1 = 6$$

9.10 (a) The system is first expressed as an augmented matrix:

$$\begin{bmatrix} 2 & -6 & -1 & -38 \\ -3 & -1 & 7 & -34 \\ -8 & 1 & -2 & -20 \end{bmatrix}$$

Forward elimination: First, we pivot by switching rows 1 and 3:

$$\begin{bmatrix} -8 & 1 & -2 & -20 \\ -3 & -1 & 7 & -34 \\ 2 & -6 & -1 & -38 \end{bmatrix}$$

Multiply row 1 by $-3/(-8) = 0.375$ and subtract from row 2 to eliminate a_{21} . Multiply row 1 by $2/(-8) = -0.25$ and subtract from row 3 to eliminate a_{31} .

$$\begin{bmatrix} -8 & 1 & -2 & -20 \\ 0 & -1.375 & 7.75 & -26.5 \\ 0 & -5.75 & -1.5 & -43 \end{bmatrix}$$

Pivot:

$$\begin{bmatrix} -8 & 1 & -2 & -20 \\ 0 & -5.75 & -1.5 & -43 \\ 0 & -1.375 & 7.75 & -26.5 \end{bmatrix}$$

Multiply row 2 by $-1.375/-5.75 = 0.23913$ and subtract from row 3 to eliminate a_{32} .

$$\begin{bmatrix} -8 & 1 & -2 & -20 \\ 0 & -5.75 & -1.5 & -43 \\ 0 & 0 & 8.108696 & -16.2174 \end{bmatrix}$$

Back substitution:

$$x_3 = \frac{-16.2174}{8.108696} = -2$$

$$x_2 = \frac{-43 - (-1.5)(-2)}{-5.75} = 8$$

$$x_1 = \frac{-20 + 2(-2) - 1(8)}{-8} = 4$$

(b) Check:

$$2(4) - 6(8) - (-2) = -38$$

$$-3(4) - (8) + 7(-2) = -34$$

$$-8(4) + (8) - 2(-2) = -20$$

9.11 (a) The determinant can be computed as:

$$A_1 = \begin{vmatrix} 2 & -1 \\ -2 & 0 \end{vmatrix} = 2(0) - (-1)(-2) = -2$$

$$A_2 = \begin{vmatrix} 1 & -1 \\ 5 & 0 \end{vmatrix} = 1(0) - (-1)(5) = 5$$

$$A_3 = \begin{vmatrix} 1 & 2 \\ 5 & -2 \end{vmatrix} = 1(-2) - 2(5) = -12$$

$$D = 0(-2) - (-3)5 + 7(-12) = -69$$

(b) Cramer's rule

$$x_1 = \frac{\begin{vmatrix} 2 & -3 & 7 \\ 3 & 2 & -1 \\ 2 & -2 & 0 \end{vmatrix}}{D} = \frac{-68}{-69} = 0.985507$$

$$x_2 = \frac{\begin{vmatrix} 0 & 2 & 7 \\ 1 & 3 & -1 \\ 5 & 2 & 0 \end{vmatrix}}{D} = \frac{-101}{-69} = 1.463768$$

$$x_3 = \frac{\begin{vmatrix} 0 & -3 & 2 \\ 1 & 2 & 3 \\ 5 & -2 & 2 \end{vmatrix}}{D} = \frac{-63}{-69} = 0.913043$$

(c) The system is first expressed as an augmented matrix:

$$\left[\begin{array}{cccc} 0 & -3 & 7 & 2 \\ 1 & 2 & -1 & 3 \\ 5 & -2 & 0 & 2 \end{array} \right]$$

Forward elimination: First, we pivot by switching rows 1 and 3:

$$\left[\begin{array}{cccc} 5 & -2 & 0 & 2 \\ 1 & 2 & -1 & 3 \\ 0 & -3 & 7 & 2 \end{array} \right]$$

Multiply row 1 by $1/5 = 0.2$ and subtract from row 2 to eliminate a_{21} . Because a_{31} already equals zero, it does not have to be eliminated.

$$\left[\begin{array}{cccc} 5 & -2 & 0 & 2 \\ 0 & 2.4 & -1 & 2.6 \\ 0 & -3 & 7 & 2 \end{array} \right]$$

Pivot:

$$\left[\begin{array}{cccc} 5 & -2 & 0 & 2 \\ 0 & -3 & 7 & 2 \\ 0 & 2.4 & -1 & 2.6 \end{array} \right]$$

Multiply row 2 by $2.4/(-3) = -0.8$ and subtract from row 3 to eliminate a_{32} .

$$\left[\begin{array}{cccc} 5 & -2 & 0 & 2 \\ 0 & -3 & 7 & 2 \\ 0 & 0 & 4.6 & 4.2 \end{array} \right]$$

Back substitution:

$$x_3 = \frac{4.2}{4.6} = 0.913043$$

$$x_2 = \frac{2 - 7(0.913043)}{-3} = 1.463768$$

$$x_1 = \frac{2 - 0(0.913043) - (-2)(1.463768)}{5} = 0.985507$$

(d) Check:

$$-3(1.463768) + 7(0.913043) = 2$$

$$(0.985507) + 2(1.463768) - (0.913043) = 3$$

$$5(0.985507) - 2(1.463768) = 2$$

9.12 The system is first expressed as an augmented matrix:

$$\left[\begin{array}{cccc} 2 & 1 & -1 & 1 \\ 5 & 2 & 2 & -4 \\ 3 & 1 & 1 & 5 \end{array} \right]$$

Normalize the first row and then eliminate a_{21} and a_{31} ,

$$\left[\begin{array}{cccc} 1 & 0.5 & -0.5 & 0.5 \\ 0 & -0.5 & 4.5 & -6.5 \\ 0 & -0.5 & 2.5 & 3.5 \end{array} \right]$$

Normalize the second row and eliminate a_{12} and a_{32} ,

$$\left[\begin{array}{cccc} 1 & 0 & 4 & -6 \\ 0 & 1 & -9 & 13 \\ 0 & 0 & -2 & 10 \end{array} \right]$$

Normalize the third row and eliminate a_{13} and a_{23} ,

$$\left[\begin{array}{cccc} 1 & 0 & 0 & 14 \\ 0 & 1 & 0 & -32 \\ 0 & 0 & 1 & -5 \end{array} \right]$$

Thus, the answers are $x_1 = 14$, $x_2 = -32$, and $x_3 = -5$. Check:

$$2(14) + (-32) - (-5) = 1$$

$$5(14) + 2(-32) + 2(-5) = -4$$

$$3(14) + (-32) + (-5) = 5$$

9.13 (a) The system is first expressed as an augmented matrix:

$$\left[\begin{array}{cccc} 1 & 1 & -1 & -3 \\ 6 & 2 & 2 & 2 \\ -3 & 4 & 1 & 1 \end{array} \right]$$

Forward elimination: a_{21} is eliminated by multiplying row 1 by $6/1 = 6$ and subtracting the result from row 2. a_{31} is eliminated by multiplying row 1 by $-3/1 = -3$ and subtracting the result from row 3.

$$\begin{bmatrix} 1 & 1 & -1 & -3 \\ 0 & -4 & 8 & 20 \\ 0 & 7 & -2 & -8 \end{bmatrix}$$

a_{32} is eliminated by multiplying row 2 by $7/(-4) = -1.75$ and subtracting the result from row 3.

$$\begin{bmatrix} 1 & 1 & -1 & -3 \\ 0 & -4 & 8 & 20 \\ 0 & 0 & 12 & 27 \end{bmatrix}$$

Back substitution:

$$x_3 = \frac{27}{12} = 2.25$$

$$x_2 = \frac{20 - 8(2.25)}{-4} = -0.5$$

$$x_1 = \frac{-3 - (-1)(2.25) - 1(-0.5)}{1} = -0.25$$

(b) The system is first expressed as an augmented matrix:

$$\left[\begin{array}{cccc} 1 & 1 & -1 & -3 \\ 6 & 2 & 2 & 2 \\ -3 & 4 & 1 & 1 \end{array} \right]$$

Forward elimination: First, we pivot by switching rows 1 and 2:

$$\left[\begin{array}{cccc} 6 & 2 & 2 & 2 \\ 1 & 1 & -1 & -3 \\ -3 & 4 & 1 & 1 \end{array} \right]$$

Multiply row 1 by $1/6 = 0.16667$ and subtract from row 2 to eliminate a_{21} . Multiply row 1 by $-3/6 = -0.5$ and subtract from row 3 to eliminate a_{31} .

$$\left[\begin{array}{cccc} 6 & 2 & 2 & 2 \\ 0 & 0.66667 & -1.33333 & -3.33333 \\ 0 & 5 & 2 & 2 \end{array} \right]$$

Pivot:

$$\left[\begin{array}{cccc} 6 & 2 & 2 & 2 \\ 0 & 5 & 2 & 2 \\ 0 & 0.66667 & -1.33333 & -3.33333 \end{array} \right]$$

Multiply row 2 by $0.66667/5 = 0.133333$ and subtract from row 3 to eliminate a_{32} .

$$\left[\begin{array}{cccc} 6 & 2 & 2 & 2 \\ 0 & 5 & 2 & 2 \\ 0 & 0 & -1.6 & -3.6 \end{array} \right]$$

Back substitution:

$$x_3 = \frac{-3.6}{1.6} = 2.25$$

$$x_2 = \frac{2 - 2(2.25)}{5} = -0.5$$

$$x_1 = \frac{2 - 2(2.25) - 2(-0.5)}{6} = -0.25$$

(c) The system is first expressed as an augmented matrix:

$$\left[\begin{array}{cccc} 1 & 1 & -1 & -3 \\ 6 & 2 & 2 & 2 \\ -3 & 4 & 1 & 1 \end{array} \right]$$

Normalize the first row, and then eliminate a_{21} and a_{31} ,

$$\left[\begin{array}{cccc} 1 & 1 & -1 & -3 \\ 0 & -4 & 8 & 20 \\ 0 & 7 & -2 & -8 \end{array} \right]$$

Normalize the second row and eliminate a_{12} and a_{32} ,

$$\left[\begin{array}{cccc} 1 & 0 & 1 & 2 \\ 0 & 1 & -2 & -5 \\ 0 & 0 & 12 & 27 \end{array} \right]$$

Normalize the third row and eliminate a_{13} and a_{23} ,

$$\left[\begin{array}{cccc} 1 & 0 & 0 & -0.25 \\ 0 & 1 & 0 & -0.5 \\ 0 & 0 & 1 & 2.25 \end{array} \right]$$

9.14 In a fashion similar to Example 9.11, vertical force balances can be written to give the following system of equations,

$$\begin{aligned} m_1 g - T_{12} - c_1 v &= m_1 a \\ m_2 g + T_{12} - c_2 v - T_{23} &= m_2 a \\ m_3 g - c_3 v + T_{23} - T_{34} &= m_3 a \\ m_4 g - c_4 v + T_{34} - T_{45} &= m_4 a \\ m_5 g - c_5 v + T_{45} &= m_5 a \end{aligned}$$

After substituting the known values, the equations can be expressed in matrix form ($g = 9.8$),

$$\left[\begin{array}{ccccc} 55 & 1 & 0 & 0 & 0 \\ 75 & -1 & 1 & 0 & 0 \\ 60 & 0 & -1 & 1 & 0 \\ 75 & 0 & 0 & -1 & 1 \\ 90 & 0 & 0 & 0 & -1 \end{array} \right] \left\{ \begin{array}{c} a \\ T_{12} \\ T_{23} \\ T_{34} \\ T_{45} \end{array} \right\} = \left\{ \begin{array}{c} 498 \\ 627 \\ 453 \\ 591 \\ 792 \end{array} \right\}$$

The system can be solved for

$$\begin{array}{lll} a = 8.225 & T_{12} = 4.5 & T_{23} = 14.625 \\ T_{34} = -25.875 & T_{12} = -51.75 & \end{array}$$

9.15 Using the format of Eq. 9.27,

$$\begin{array}{ll} [A] = \begin{bmatrix} 3 & 4 \\ 0 & 1 \end{bmatrix} & [B] = \begin{bmatrix} 2 & 0 \\ -1 & 0 \end{bmatrix} \\ \{U\} = \begin{Bmatrix} 3 \\ 3 \end{Bmatrix} & \{V\} = \begin{Bmatrix} 1 \\ 0 \end{Bmatrix} \end{array}$$

The set of equations to be solved are

$$\begin{array}{rcl} 3x_1 + 4x_2 - 2y_1 & & = 3 \\ & x_2 + y_1 & = 3 \\ 2x_1 & + 3y_1 + 4y_2 & = 1 \\ -x_1 & & + y_2 = 0 \end{array}$$

These can be solved for $x_1 = -0.46667$, $x_2 = 1.73333$, $y_1 = 1.26667$, and $y_2 = -0.46667$. Therefore, the solution is $z_1 = -0.46667 + 1.73333 i$ and $z_2 = 1.26667 - 0.46667 i$.

9.16 Here is a VBA program to implement matrix multiplication and solve Prob. 9.2 for the case of $[A] \times [B]$.

```
Option Explicit

Sub Mult()
    Dim i As Integer, j As Integer
    Dim l As Integer, m As Integer, n As Integer
    Dim a(10, 10) As Double, b(10, 10) As Double
    Dim c(10, 10) As Double
    l = 2
    m = 2
    n = 3
    a(1, 1) = 6: a(1, 2) = -1
    a(2, 1) = 12: a(2, 2) = 8
    a(3, 1) = -5: a(3, 2) = 4
    b(1, 1) = 4: b(1, 2) = 0
    b(2, 1) = 0.5: b(2, 2) = 2
    Call Mmult(a, b, c, m, n, l)
    For i = 1 To n
        For j = 1 To l
            MsgBox c(i, j)
        Next j
    Next i
End Sub

Sub Mmult(a, b, c, m, n, l)
    Dim i As Integer, j As Integer, k As Integer
    Dim sum As Double
    For i = 1 To n
        For j = 1 To l
            sum = 0
            For k = 1 To m
```

```

        sum = sum + a(i, k) * b(k, j)
    Next k
    c(i, j) = sum
Next j
Next i
End Sub

```

9.17 Here is a VBA program to implement the matrix transpose and solve Prob. 9.2 for the case of $[A]^T$.

```

Option Explicit
Sub TransTest()
Dim i As Integer, j As Integer
Dim m As Integer, n As Integer
Dim a(10, 10) As Double, aT(10, 10) As Double
n = 3
m = 2
a(1, 1) = 6: a(1, 2) = -1
a(2, 1) = 12: a(2, 2) = 8
a(3, 1) = -5: a(3, 2) = 4
Call Transpose(a, aT, n, m)
For i = 1 To m
    For j = 1 To n
        MsgBox aT(i, j)
    Next j
Next i
End Sub

Sub Transpose(a, b, n, m)
Dim i As Integer, j As Integer
For i = 1 To m
    For j = 1 To n
        b(i, j) = a(j, i)
    Next j
Next i
End Sub

```

9.18 Here is a VBA program to implement the Gauss elimination algorithm and solve the test case in the problem statement.

```

Option Explicit
Sub GaussElim()
Dim n As Integer, er As Integer, i As Integer
Dim a(10, 10) As Double, b(10) As Double, x(10) As Double
n = 3
a(1, 1) = 1: a(1, 2) = 2: a(1, 3) = -1
a(2, 1) = 5: a(2, 2) = 2: a(2, 3) = 2
a(3, 1) = -3: a(3, 2) = 5: a(3, 3) = -1
b(1) = 2: b(2) = 9: b(3) = 1
Call Gauss(a, b, n, x, er)
If er = 0 Then
    For i = 1 To n
        MsgBox "x(" & i & ") = " & x(i)
    Next i
Else
    MsgBox "ill-conditioned system"
End If
End Sub

Sub Gauss(a, b, n, x, er)
Dim i As Integer, j As Integer
Dim s(10) As Double

```

```

Const tol As Double = 0.000001
er = 0
For i = 1 To n
    s(i) = Abs(a(i, 1))
    For j = 2 To n
        If Abs(a(i, j)) > s(i) Then s(i) = Abs(a(i, j))
    Next j
Next i
Call Eliminate(a, s, n, b, tol, er)
If er <> -1 Then
    Call Substitute(a, n, b, x)
End If
End Sub

```

```

Sub Pivot(a, b, s, n, k)
Dim p As Integer, ii As Integer, jj As Integer
Dim factor As Double, big As Double, dummy As Double
p = k
big = Abs(a(k, k) / s(k))
For ii = k + 1 To n
    dummy = Abs(a(ii, k) / s(ii))
    If dummy > big Then
        big = dummy
        p = ii
    End If
Next ii
If p <> k Then
    For jj = k To n
        dummy = a(p, jj)
        a(p, jj) = a(k, jj)
        a(k, jj) = dummy
    Next jj
    dummy = b(p)
    b(p) = b(k)
    b(k) = dummy
    dummy = s(p)
    s(p) = s(k)
    s(k) = dummy
End If
End Sub

```

```

Sub Substitute(a, n, b, x)
Dim i As Integer, j As Integer
Dim sum As Double
x(n) = b(n) / a(n, n)
For i = n - 1 To 1 Step -1
    sum = 0
    For j = i + 1 To n
        sum = sum + a(i, j) * x(j)
    Next j
    x(i) = (b(i) - sum) / a(i, i)
Next i
End Sub

```

```

Sub Eliminate(a, s, n, b, tol, er)
Dim i As Integer, j As Integer, k As Integer
Dim factor As Double
For k = 1 To n - 1
    Call Pivot(a, b, s, n, k)
    If Abs(a(k, k) / s(k)) < tol Then
        er = -1
    Exit For

```

```

End If
For i = k + 1 To n
    factor = a(i, k) / a(k, k)
    For j = k + 1 To n
        a(i, j) = a(i, j) - factor * a(k, j)
    Next j
    b(i) = b(i) - factor * b(k)
Next i
Next k
If Abs(a(k, k) / s(k)) < tol Then er = -1
End Sub

```

Its application yields a solution of (1, 1, 1).

9.19 The position of the three masses can be modeled by the following steady-state force balances

$$0 = k(x_2 - x_1) + m_1 g - kx_1$$

$$0 = k(x_3 - x_2) + m_2 g - k(x_2 - x_1)$$

$$0 = m_3 g - k(x_3 - x_2)$$

Terms can be combined to yield

$$2kx_1 - kx_2 = m_1 g$$

$$-kx_1 + 2kx_2 - kx_3 = m_2 g$$

$$-kx_2 + kx_3 = m_3 g$$

Substituting the parameter values

$$\begin{bmatrix} 20 & -10 & 0 \\ -10 & 20 & -10 \\ 0 & -10 & 10 \end{bmatrix} \begin{Bmatrix} x_1 \\ x_2 \\ x_3 \end{Bmatrix} = \begin{Bmatrix} 19.62 \\ 29.43 \\ 24.525 \end{Bmatrix}$$

The system can be solved for 7.3575, 12.7530 and 15.2055.

9.20 Here is MATLAB M-file function to implement the Newton-Raphson method to solve simultaneous nonlinear systems:

```

function [x,f,ea,iter]=newtmult(func,x0,es,maxit,varargin)
% newtmult: Newton-Raphson root zeroes nonlinear systems
% [x,f,ea,iter]=newtmult(func,x0,es,maxit,p1,p2,...):
% uses the Newton-Raphson method to find the roots of
% a system of nonlinear equations
% input:
% func = name of function that returns f and J
% x0 = initial guess
% es = desired percent relative error (default = 0.0001%)
% maxit = maximum allowable iterations (default = 50)
% p1,p2,... = additional parameters used by function
% output:
% x = vector of roots
% f = vector of functions evaluated at roots
% ea = approximate percent relative error (%)
% iter = number of iterations
if nargin<2,error('at least 2 input arguments required'),end
if nargin<3||isempty(es),es=0.0001;end
if nargin<4||isempty(maxit),maxit=50;end

```

```

iter = 0;x=x0;
while (1)
    [J,f]=func(x,varargin{:});
    dx=J\f;
    x=x-dx;
    iter = iter + 1;
    ea=100*max(abs(dx./x));
    if iter>=maxit|ea<=es, break, end
end

```

9.21 MATLAB M-files can be used to compute both the function values and the Jacobian as

```

function [J,f]=jfrain(x,varargin)
del=0.00001;
df1dx1=(f1(x(1)+del*x(1),x(2),x(3),x(4),x(5),varargin{:})-...
    f1(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df1dx2=(f1(x(1),x(2)+del*x(2),x(3),x(4),x(5),varargin{:})-...
    f1(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df1dx3=(f1(x(1),x(2),x(3)+del*x(3),x(4),x(5),varargin{:})-...
    f1(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df1dx4=(f1(x(1),x(2),x(3),x(4)+del*x(4),x(5),varargin{:})-...
    f1(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df1dx5=(f1(x(1),x(2),x(3),x(4),x(5)+del*x(5),varargin{:})-...
    f1(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df2dx1=(f2(x(1)+del*x(1),x(2),x(3),x(4),x(5),varargin{:})-...
    f2(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df2dx2=(f2(x(1),x(2)+del*x(2),x(3),x(4),x(5),varargin{:})-...
    f2(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df2dx3=(f2(x(1),x(2),x(3)+del*x(3),x(4),x(5),varargin{:})-...
    f2(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df2dx4=(f2(x(1),x(2),x(3),x(4)+del*x(4),x(5),varargin{:})-...
    f2(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df2dx5=(f2(x(1),x(2),x(3),x(4),x(5)+del*x(5),varargin{:})-...
    f2(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df3dx1=(f3(x(1)+del*x(1),x(2),x(3),x(4),x(5),varargin{:})-...
    f3(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df3dx2=(f3(x(1),x(2)+del*x(2),x(3),x(4),x(5),varargin{:})-...
    f3(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df3dx3=(f3(x(1),x(2),x(3)+del*x(3),x(4),x(5),varargin{:})-...
    f3(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df3dx4=(f3(x(1),x(2),x(3),x(4)+del*x(4),x(5),varargin{:})-...
    f3(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df3dx5=(f3(x(1),x(2),x(3),x(4),x(5)+del*x(5),varargin{:})-...
    f3(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df4dx1=(f4(x(1)+del*x(1),x(2),x(3),x(4),x(5),varargin{:})-...
    f4(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df4dx2=(f4(x(1),x(2)+del*x(2),x(3),x(4),x(5),varargin{:})-...
    f4(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df4dx3=(f4(x(1),x(2),x(3)+del*x(3),x(4),x(5),varargin{:})-...
    f4(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df4dx4=(f4(x(1),x(2),x(3),x(4)+del*x(4),x(5),varargin{:})-...
    f4(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df4dx5=(f4(x(1),x(2),x(3),x(4),x(5)+del*x(5),varargin{:})-...
    f4(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df5dx1=(f5(x(1)+del*x(1),x(2),x(3),x(4),x(5),varargin{:})-...
    f5(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df5dx2=(f5(x(1),x(2)+del*x(2),x(3),x(4),x(5),varargin{:})-...
    f5(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df5dx3=(f5(x(1),x(2),x(3)+del*x(3),x(4),x(5),varargin{:})-...
    f5(x(1),x(2),x(3),x(4),x(5),varargin{:}))/del;
df5dx4=(f5(x(1),x(2),x(3),x(4)+del*x(4),x(5),varargin{:})-...

```



```

    f5(x(1),x(2) ,x(3),x(4),x(5),varargin{:}))/ (del*x(4));
df5dx5=(f5(x(1),x(2),x(3),x(4),x(5)+del*x(5),varargin{:}))-...
    f5(x(1),x(2) ,x(3),x(4),x(5),varargin{:}))/ (del*x(5));
J=[df1dx1 df1dx2 df1dx3 df1dx4 df1dx5
df2dx1 df2dx2 df2dx3 df2dx4 df2dx5
df3dx1 df3dx2 df3dx3 df3dx4 df3dx5
df4dx1 df4dx2 df4dx3 df4dx4 df4dx5
df5dx1 df5dx2 df5dx3 df5dx4 df5dx5];
f11=f1(x(1),x(2),x(3),x(4),x(5),varargin{:});
f22=f2(x(1),x(2),x(3),x(4),x(5),varargin{:});
f33=f3(x(1),x(2),x(3),x(4),x(5),varargin{:});
f44=f4(x(1),x(2),x(3),x(4),x(5),varargin{:});
f55=f5(x(1),x(2),x(3),x(4),x(5),varargin{:});
f=[f11;f22;f33;f44;f55];
end

function f=f1(H,OH,cT,HCO3,CO3,KH,K1,K2,Kw,pco2)
f = 1e6*H*HCO3/KH/pco2-K1;
end
function f=f2(H,OH,cT,HCO3,CO3,KH,K1,K2,Kw,pco2)
f = H*CO3/HCO3-K2;
end
function f=f3(H,OH,cT,HCO3,CO3,KH,K1,K2,Kw,pco2)
f = H*OH-Kw;
end
function f=f4(H,OH,cT,HCO3,CO3,KH,K1,K2,Kw,pco2)
f = KH*pco2/1e6+HCO3+CO3-cT;
end
function f=f5(H,OH,cT,HCO3,CO3,KH,K1,K2,Kw,pco2)
f = HCO3+2*CO3+OH-H;
end

```

A script can then be written to invoke the function developed in Prob. 9.20,

```

format short g
x0=[10^-7;10^-7;1e-5;2e-6;5e-11];
KH=10^-1.46;K1=10^-6.3;K2=10^-10.3;Kw=10^-14;pco2=315;
[x,f,ea,iter]=newtmult(@jfrain,x0,[],[],KH,K1,K2,Kw,pco2);
roots=x'
fprintf('Maximum relative error = %8.4g percent\n',ea)
fprintf('Number of iterations = %5d',iter)
pH=-log10(x(1))

```

This script, which was saved as Prob0921NM6.m can then be run to determine the roots and display the pH

```

>> Prob0921NM6

roots =
    2.3419e-006    4.2701e-009    1.326e-005    2.3375e-006    5.0025e-011
Maximum relative error = 1.265e-012 percent
Number of iterations =      6
pH =
    5.6304

```