

Extremal no-three-in-line subsets of a modular hyperbola in the Hall–Jackson–Sudbery–Wild window

Alex Komang*

August 2026 — draft v0.6

Abstract

Let p be an odd prime, $c \in \mathbb{F}_p^*$, and let $H(c, p) = \{(x, y) \in \mathbb{Z}^2 : xy \equiv c \pmod{p}\}$ be the modular hyperbola. Hall, Jackson, Sudbery and Wild (1975) observed that the $2p \times 2p$ window $G(p) = [-\frac{p-1}{2}, \frac{3p-1}{2}] \times [0, 2p-1]$ contains exactly four points of each of the $p-1$ residue classes of $H(c, p)$, at the vertices of a $p \times p$ square, and that keeping three of the four (the twelve outer blocks T_2 of the 4×4 block subdivision of $G(p)$) yields $3(p-1)$ points with no three collinear. Kovács, Nagy and Szabó (2025/26) state that this deletion is optimal, i.e. that $H(c, p) \cap T_2$ is a largest no-three-in-line subset of $H(c, p) \cap G(p)$; the value $3(p-1)$ is therefore not new. We give what is, to our knowledge, the first complete proof of this statement, for every c , and we determine the complete extremal structure: we classify all lines carrying three or more points of $H(c, p) \cap G(p)$ (they have slope ± 1 , and their number is $\frac{3}{2}(p-1) - s$), we exhibit an explicit linear-programming certificate for the bound $3(p-1)$ (so even the fractional relaxation has value $3(p-1)$), and we show that the number of maximum no-three-in-line subsets is exactly 9^s , where $s \in \{0, 1, 2\}$ is the number of quadratic residues among c and $-c$. In particular the HJSW set is the unique maximum if and only if $p \equiv 1 \pmod{4}$ and c is a quadratic non-residue. The bound does not depend on the position of the window: for every $2p \times 2p$ box $[x_0, x_0 + 2p) \times [y_0, y_0 + 2p)$ the modular hyperbola contains at most $3(p-1)$ points in general position, and we determine the exact maximum and the number of maximum sets for every box (an “orbit lemma”: a box separates 0 or 2 of the four partner pairs of each generic orbit of the Klein four-group generated by $(a, b) \mapsto (-b, -a)$ and $(a, b) \mapsto (b, a)$, and the maximum is $12n_2 + 10n_1 + 8n_0 + 6s$ in terms of the numbers n_2, n_1, n_0 of generic orbits with 2, 1, 0 split pairs of each kind); the HJSW box is among the boxes where $3(p-1)$ is attained. The classification also gives the no-four-in-line analogue: the largest no-four-in-line subset of $H(c, p) \cap G(p)$ has exactly $\frac{7}{2}(p-1) + s$ points, so the set S_3 of Kovács–Nagy–Szabó is optimal as well. The proofs are elementary and self-contained; independent programs check the statements about the HJSW box for all $p \leq 101$ (all c for $p \leq 31$), the bound $3(p-1)$ for all boxes with $p \leq 61$, and the box formula and orbit lemma for all boxes with $p \leq 13$ and $p \leq 31$ respectively.

1 Introduction

The no-three-in-line problem asks for the largest subset of the $n \times n$ grid with no three collinear points; at most $2n$ points are possible, and the best general lower bound, $\frac{3}{2}n - o(n)$, is due to Hall, Jackson, Sudbery and Wild [1]: for $n = 2p$, p prime, they place $3(p-1)$ points of a modular hyperbola $H(c, p)$ in the window $G(p)$. Their construction rests on two facts: every line meets $H(c, p)$ in at most two residue classes, and $G(p)$ contains exactly four points of each class, at the vertices of a $p \times p$ square; keeping the three copies outside the middle block of the 4×4 block subdivision of $G(p)$ leaves no three collinear.

Kovács, Nagy and Szabó [2] recently used the same window and its refinements for the no- $(k+1)$ -in-line problem. In their setting the HJSW set is the case $|W| = 1$, $k = 2$ of a general

*Independent researcher, Bali, Indonesia. See the AI/tool-use disclosure at the end.

randomised construction (all four copies of every class are taken, then a minimum set R of points is deleted so that every line of slope ± 1 retains at most k points), and they state that for $|W| = 1$, $k = 2$ “Subsection 2.2 essentially details an optimal choice for the points R to be removed to gain the largest no-3-in-line set $S_2(c, p)$ inside $S(W)$ ” [2, App. A]. The general estimate of [2, Rem. 3.4], $|R| \geq \frac{1}{2}X$ with $X = \sum_{\ell}(|\ell \cap S(W)| - k)^+$, gives for $|W| = 1$, $k = 2$ the bound $|R| \geq p - 1 - s$ (see Remark 9 below), i.e. $|S| \leq 3(p - 1) + s$ for every no-three-in-line $S \subseteq H(c, p) \cap G(p)$: exact when neither c nor $-c$ is a square modulo p , but one or two short of the truth otherwise. We are not aware of a proof of the exact statement in [2] or elsewhere; the purpose of this note is to give one, and to determine the whole extremal structure of the problem. To be explicit about what is new: the value $3(p - 1)$ and the qualitative optimality of S_2 and S_3 are asserted in [2]; the proof, the classification of the rich lines, the LP certificate, the count 9^s with its uniqueness criterion, the exact value and count for every $2p \times 2p$ box, and the exact no-four-in-line value and count are, as far as we could determine, new.

Context. Modular hyperbolas $xy \equiv c \pmod{m}$ have a rich literature (see Shparlinski’s survey [3]; for combinatorial-geometric questions about them, e.g. [4]); the general problem of selecting a largest subset in general position from a given point set is NP-hard and APX-hard [5], with general bounds in [6], whereas here the structure of the point set makes the problem solvable exactly. Variants of the no-three-in-line problem with exact structural answers include the extensible version [7] and the torus version [8]. For $k \geq 3$ points per line the asymptotic answer kn was recently established [9], while for $k = 2$ the HJSW construction is still the best known general lower bound (see Green’s Problem 72 [10]); the state of the art for exact $2n$ -point configurations is documented in Flammenkamp’s database [11] and in [12].

Results. Throughout, p is an odd prime, $h = (p - 1)/2$, $c \in \mathbb{F}_p^*$, and

$$H(c, p) = \{(x, y) \in \mathbb{Z}^2 : xy \equiv c \pmod{p}\}, \quad G(p) = \{-h, \dots, 3h + 1\} \times \{0, \dots, 2p - 1\},$$

and $\mathcal{P} := H(c, p) \cap G(p)$. A set is *lawful* (no-three-in-line, in general position) if it contains no three collinear points. A line is *rich* if it contains at least three points of $\mathcal{P}$; $\mathcal{L}$ denotes the set of rich lines. Let $s = [c \text{ is a QR}] + [-c \text{ is a QR}] \in \{0, 1, 2\}$ (QR = quadratic residue modulo p).

Theorem 1. (i) Every rich line has slope $+1$ or -1 ; there are exactly $\frac{3}{2}(p - 1) - s$ rich lines, $p - 1$ of them carrying three points of $\mathcal{P}$ and $\frac{1}{2}(p - 1) - s$ carrying four (Proposition 8 lists them); exactly $2s$ points of $\mathcal{P}$ lie on no rich line.

(ii) Every lawful subset of $\mathcal{P}$ has at most $3(p - 1)$ points; the bound is attained. Even the fractional relaxation (maximise $\sum_q x_q$ subject to $\sum_{q \in \ell} x_q \leq 2$ for $\ell \in \mathcal{L}$, $0 \leq x_q \leq 1$) has optimum $3(p - 1)$.

(iii) The number of lawful subsets of size $3(p - 1)$ is exactly 9^s . Every maximum set coincides with the HJSW set $H(c, p) \cap T_2$ outside the (at most four) classes $(a, \pm a)$ with $a^2 = \pm c$.

Corollary 2. If $p \equiv 3 \pmod{4}$ then $s = 1$ for every c : there are exactly 9 maximum sets. If $p \equiv 1 \pmod{4}$ then $s = 2$ for c a QR (81 maximum sets) and $s = 0$ for c a non-residue: the HJSW set is the unique maximum. Thus $H(c, p) \cap T_2$ is the unique largest lawful subset of $H(c, p) \cap G(p)$ if and only if $p \equiv 1 \pmod{4}$ and c is a quadratic non-residue.

Theorem 3. Let $x_0, y_0 \in \mathbb{Z}$ and $W = [x_0, x_0 + 2p) \times [y_0, y_0 + 2p)$. Every lawful subset of $H(c, p) \cap W$ has at most $3(p - 1)$ points. More precisely, the maximum equals $12n_2 + 10n_1 + 8n_0 + 6s$, where n_2, n_1, n_0 are the numbers of generic orbits of the Klein four-group whose partner pairs are split by the box in the patterns $\{(0, 2), (2, 0)\}$, $(1, 1)$, $(0, 0)$ respectively ($n_2 + n_1 + n_0 = \frac{1}{4}(p - 1 - 2s)$; no other pattern occurs), and the number of maximum sets is $144^{n_1} 1296^{n_0} 9^{s_1} 6^{s_2}$, where s_1 (resp. s_2) is the number of exceptional orbits whose pair is split (resp. shared), $s_1 + s_2 = s$. In particular $3(p - 1)$ is attained iff $n_1 = n_0 = 0$; the HJSW box has $n_2 = \frac{1}{4}(p - 1 - 2s)$, $s_1 = s$. (Maximum sets

are counted as subsets of $\mathbb{Z}^2$, not up to symmetry. The pattern of an orbit and the split/shared label of an exceptional pair depend only on the orbit and the box, not on the class chosen to name the orbit.)

The proof of Theorem 1 is a double count over the rich lines: for a lawful S , $|S| \leq Z + 2|\mathcal{L}|$ where Z is the number of points on no rich line, and the identity $Z + 2|\mathcal{L}| = 3(p-1)$ follows from the classification of the rich lines. The count is exact precisely because the $2s$ points that lie on no rich line are accounted separately; this is the correction to the averaging bound of [2] mentioned above. All rich lines lie inside orbits of a Klein four-group acting on the classes, and the equality analysis reduces to two small “gadgets” (Figure 1), which gives 9^s . For a general box the rich lines still lie inside the orbits, so the maximum is a sum of orbit maxima; the orbit lemma (Section 5) says that a box splits 0 or 2 of the four partner pairs of a generic orbit, and the four possible patterns have maxima 8, 10, 12, 12 — whence the exact formula of Theorem 3. For boxes other than the HJSW box the maximum can be smaller than $3(p-1)$ (Table 1). The four-point rich lines are pairwise disjoint, which settles the no-four-in-line analogue as well (Corollary 14: the maximum is $\frac{7}{2}(p-1) + s$ in the HJSW box, attained by the set S_3 of [2]).

2 Notation

Two points are *congruent* if their coordinates agree modulo p ; a *class* is a congruence class, identified with an element of $\mathbb{F}_p \times \mathbb{F}_p$. Points of $H(c, p)$ have both coordinates prime to p , so the classes of $H(c, p)$ are the $p-1$ pairs $\kappa = (a, b) \in \mathbb{F}_p^* \times \mathbb{F}_p^*$ with $ab = c$. Since each coordinate range of $G(p)$ is an interval of $2p$ consecutive integers, it contains exactly two representatives of every residue class; hence every class κ of $H(c, p)$ has exactly four points in $G(p)$ and $|\mathcal{P}| = 4(p-1)$. We write (x_κ, y_κ) for the *base copy* of κ , its unique point in $\{-h, \dots, h\} \times \{1, \dots, p-1\}$ (necessarily $x_\kappa \neq 0$), and

$$\kappa(r, s) := (x_\kappa + rp, y_\kappa + sp), \quad r, s \in \{0, 1\},$$

for its four points in $G(p)$. Following [1, 2] we sort the classes into four *types* according to the position of the base copy:

$$A : x_\kappa > 0, y_\kappa > h; \quad B : x_\kappa < 0, y_\kappa > h; \quad C : x_\kappa > 0, y_\kappa \leq h; \quad D : x_\kappa < 0, y_\kappa \leq h.$$

(In the notation of [2, Def. 2.3]: $\kappa(0, 0) \in A_{0,0}, B_{-1,0}, C_{0,0}, D_{-1,0}$ respectively; the HJSW set T_2 omits, for each type, the copy $A_{0,0}, B_{0,0}, C_{0,1}, D_{0,1}$, i.e. $\kappa(0, 0)$ for A , $\kappa(1, 0)$ for B , $\kappa(0, 1)$ for C , $\kappa(1, 1)$ for D ; the union of these omitted blocks is the middle block M of [2, §2.3]. We call this copy the *middle copy* of κ .) We write n_A, n_B, n_C, n_D for the numbers of classes of each type, so $n_A + n_B + n_C + n_D = p-1$. Finally, for a class κ put

$$d_\kappa := x_\kappa - y_\kappa, \quad e_\kappa := x_\kappa + y_\kappa, \quad D_\kappa := \{x - y = d_\kappa\}, \quad E_\kappa := \{x + y = e_\kappa + p\};$$

D_κ (the *diagonal* of κ) contains $\kappa(0, 0), \kappa(1, 1)$ and E_κ (the *antidiagonal*) contains $\kappa(0, 1), \kappa(1, 0)$. For a real number x let $x^+ = \max\{x, 0\}$.

Three involutions act on the classes of $H(c, p)$:

$$\sigma(a, b) = (-b, -a), \quad \tau(a, b) = (b, a), \quad \nu(a, b) = (-a, -b) = \sigma\tau(a, b) = \tau\sigma(a, b),$$

so $V = \{1, \sigma, \tau, \nu\}$ is a Klein four-group acting on the $p-1$ classes. σ has fixed classes iff $-c$ is a QR (then exactly two: $(a, -a)$ with $a^2 = -c$), τ has fixed classes iff c is a QR (two: (a, a) with $a^2 = c$), and ν has none. Put $f_\sigma := \#\{\kappa : \sigma\kappa = \kappa\} \in \{0, 2\}$, $f_\tau := \#\{\kappa : \tau\kappa = \kappa\} \in \{0, 2\}$; thus $f_\sigma + f_\tau = 2s$. The V -orbits are of size 4 (*generic*: four distinct classes) or of size 2 (*exceptional*: $\{\kappa, \nu\kappa\}$ with κ and $\nu\kappa$ both fixed by σ , or both fixed by τ); there are exactly s exceptional orbits.

3 Rich lines

Lemma 4. *Let ℓ be a line containing at least two points of $H(c, p)$. Then the points of $H(c, p) \cap \ell$ lie in at most two classes. More precisely, if ℓ contains a point of class κ , then the classes met by ℓ are: only κ if ℓ has slope 0 or ∞ ; among $\{\kappa, \sigma\kappa\}$ if ℓ has slope +1; among $\{\kappa, \tau\kappa\}$ if ℓ has slope -1.*

Proof. Let $(u, v) \in \mathbb{Z}^2$ be a primitive direction vector of ℓ and $(x_0, y_0) \in H(c, p) \cap \ell$. The lattice points of ℓ are $(x_0 + tu, y_0 + tv)$, $t \in \mathbb{Z}$, and such a point lies on $H(c, p)$ iff $uv t^2 + (x_0 v + y_0 u) t \equiv 0 \pmod{p}$ (we used $x_0 y_0 \equiv c$). The class of the point depends only on $t \pmod{p}$. If $uv \not\equiv 0$, the congruence has at most two roots modulo p ; if exactly one of u, v is divisible by p (both cannot be, as $\gcd(u, v) = 1$), it reduces to $x_0 v t \equiv 0$ or $y_0 u t \equiv 0$ with a unit coefficient, so $t \equiv 0$: one class. For $(u, v) = (1, 0), (0, 1)$ this gives the claim for slopes 0, ∞ . For $(u, v) = (1, 1)$ the roots are $t \equiv 0$ and $t \equiv -(x_0 + y_0)$, the latter giving the point $(-y_0, -x_0)$ modulo p , of class $\sigma\kappa$. For $(u, v) = (1, -1)$ the roots are $t \equiv 0$ and $t \equiv y_0 - x_0$, giving (y_0, x_0) , of class $\tau\kappa$. $\square$

Lemma 5. *Two distinct points of one class in $G(p)$ differ by $\pm(p, 0)$, $\pm(0, p)$, $\pm(p, p)$ or $\pm(p, -p)$; the line through them has slope 0, ∞ , +1 or -1 and contains no third point of that class. The point $\kappa(r, s)$ lies on the slope-(+1) line $x - y = d_\kappa + (r - s)p$ and on the slope-(-1) line $x + y = e_\kappa + (r + s)p$. In particular D_κ and E_κ are the only lines of slope ± 1 carrying two points of κ .*

Proof. The four points of a class are the vertices of an axis-parallel square of side p ; no three are collinear. The rest is a direct computation. $\square$

Corollary 6. *A rich line has slope +1 or -1. Consequently a set $S \subseteq \mathcal{P}$ is lawful iff $|S \cap \ell| \leq 2$ for every $\ell \in \mathcal{L}$.*

Proof. By Lemma 4 three points of $\mathcal{P}$ on a line lie in at most two classes, so two of them are congruent, and by Lemma 5 the line has slope 0, ∞ , ± 1 . A line of slope 0 or ∞ meets a single class (Lemma 4), which has at most two points on it (Lemma 5). $\square$

Lemma 7 (position of the partner classes). *For every class κ the types of $\sigma\kappa$, $\tau\kappa$ and the shifts $d_{\sigma\kappa} - d_\kappa$, $e_{\tau\kappa} - e_\kappa$ are given by the following table:*

type of κ	type of $\sigma\kappa$	$d_{\sigma\kappa} - d_\kappa$	type of $\tau\kappa$	$e_{\tau\kappa} - e_\kappa$
A	A	0	D	$-p$
B	C	$+p$	B	0
C	B	$-p$	C	0
D	D	0	A	$+p$

In particular σ preserves the types A, D and interchanges B, C ; τ preserves B, C and interchanges A, D ; ν interchanges $A \leftrightarrow D$ and $B \leftrightarrow C$; the fixed classes of σ lie in $A \cup D$, those of τ in $B \cup C$; and $n_A = n_D$, $n_B = n_C$.

Proof. Write $\kappa = (a, b)$ with base copy (x_κ, y_κ) , so $x_\kappa \equiv a$, $y_\kappa \equiv b$. The base copy of $\sigma\kappa = (-b, -a)$ has first coordinate $-y_\kappa$ if $y_\kappa \leq h$ and $p - y_\kappa$ if $y_\kappa > h$, and second coordinate $p - x_\kappa$ if $x_\kappa > 0$ and $-x_\kappa$ if $x_\kappa < 0$ (these are the representatives of $-b$ in $[-h, h]$ and of $-a$ in $[1, p-1]$). Going through the four types: $\kappa \in A$: $(p - y_\kappa, p - x_\kappa)$, again of type A , difference $x_\kappa - y_\kappa$; $\kappa \in D$: $(-y_\kappa, -x_\kappa)$, type D , difference $x_\kappa - y_\kappa$; $\kappa \in C$: $(-y_\kappa, p - x_\kappa)$, type B , difference $x_\kappa - y_\kappa - p$; $\kappa \in B$: $(p - y_\kappa, -x_\kappa)$, type C , difference $x_\kappa - y_\kappa + p$. Similarly, the base copy of $\tau\kappa = (b, a)$ has first coordinate y_κ if $y_\kappa \leq h$ and $y_\kappa - p$ otherwise, and second coordinate x_κ if $x_\kappa > 0$ and $x_\kappa + p$ otherwise: $\kappa \in B$: $(y_\kappa - p, x_\kappa + p)$, type B , sum e_κ ; $\kappa \in C$: (y_κ, x_κ) , type C , sum e_κ ; $\kappa \in A$: $(y_\kappa - p, x_\kappa)$, type D , sum $e_\kappa - p$; $\kappa \in D$: $(y_\kappa, x_\kappa + p)$, type A , sum $e_\kappa + p$. The remaining assertions follow. $\square$

Proposition 8. $\mathcal{L}$ consists of the following lines, and no others; for each we list its points in $\mathcal{P}$.

- (a) For $\kappa \in A \cup D$ with $\sigma\kappa \neq \kappa$: the common diagonal $D_\kappa = D_{\sigma\kappa}$, with the four points $\kappa(0,0), \kappa(1,1), \sigma\kappa(0,0), \sigma\kappa(1,1)$.
- (b) For $\kappa \in B \cup C$ with $\tau\kappa \neq \kappa$: the common antidiagonal $E_\kappa = E_{\tau\kappa}$, with the four points $\kappa(0,1), \kappa(1,0), \tau\kappa(0,1), \tau\kappa(1,0)$.
- (c) For every $\kappa \in C$: D_κ with the three points $\kappa(0,0), \kappa(1,1), \sigma\kappa(1,0)$ ($\sigma\kappa \in B$); for every $\kappa \in B$: D_κ with the three points $\kappa(0,0), \kappa(1,1), \sigma\kappa(0,1)$ ($\sigma\kappa \in C$).
- (d) For every $\kappa \in A$: E_κ with the three points $\kappa(0,1), \kappa(1,0), \tau\kappa(1,1)$ ($\tau\kappa \in D$); for every $\kappa \in D$: E_κ with the three points $\kappa(0,1), \kappa(1,0), \tau\kappa(0,0)$ ($\tau\kappa \in A$).

The lines listed in (a)–(d) are pairwise distinct (apart from the identifications $D_\kappa = D_{\sigma\kappa}$ in (a) and $E_\kappa = E_{\tau\kappa}$ in (b) that are built into the list), and each of them lies inside a single V -orbit of classes. Consequently

$$|\mathcal{L}| = \frac{n_A + n_D - f_\sigma}{2} + \frac{n_B + n_C - f_\tau}{2} + (n_A + n_D) + (n_B + n_C) = \frac{3}{2}(p-1) - s, \quad (1)$$

with $p-1$ three-point lines and $\frac{1}{2}(p-1) - s$ four-point lines. Moreover, the points of $\mathcal{P}$ lying on no rich line are exactly $\kappa(1,1)$ for σ -fixed $\kappa \in A$, $\kappa(0,0)$ for σ -fixed $\kappa \in D$, $\kappa(0,1)$ for τ -fixed $\kappa \in B$ and $\kappa(1,0)$ for τ -fixed $\kappa \in C$; their number is

$$Z := \#\{q \in \mathcal{P} : q \text{ lies on no rich line}\} = f_\sigma + f_\tau = 2s. \quad (2)$$

The points lying on two rich lines are exactly the middle copies of the classes κ whose partner ($\sigma\kappa$ for $\kappa \in A \cup D$, $\tau\kappa$ for $\kappa \in B \cup C$) is different from κ ; no point lies on three rich lines.

Proof. Let $\ell \in \mathcal{L}$. By Corollary 6 its slope is ± 1 ; say $+1$ (the case -1 is symmetric, with τ, e_κ, E_κ in place of $\sigma, d_\kappa, D_\kappa$). By Lemma 4 the points of $\ell \cap \mathcal{P}$ lie in two classes $\kappa, \sigma\kappa$ (a single class carries at most two points on a line), and one of the classes, say κ , has two points on ℓ ; by Lemma 5 these are $\kappa(0,0), \kappa(1,1)$ and $\ell = D_\kappa$, and $\sigma\kappa \neq \kappa$. The points of $\sigma\kappa$ on D_κ are the $\sigma\kappa(r,s)$ with $d_{\sigma\kappa} + (r-s)p = d_\kappa$. By Lemma 7: for $\kappa \in A \cup D$, $r=s$, giving (a); for $\kappa \in C$, $r-s=1$, giving $\sigma\kappa(1,0)$; for $\kappa \in B$, $r-s=-1$, giving $\sigma\kappa(0,1)$; this is (c). Conversely each listed line has at least three points.

Distinctness: if $D_\kappa = D_\lambda$ then this line of slope $+1$ meets both classes κ, λ , so $\lambda \in \{\kappa, \sigma\kappa\}$ by Lemma 4; and $D_\kappa = D_{\sigma\kappa}$ holds for $\kappa \in A \cup D$ (shift 0) but not for $\kappa \in B \cup C$ (shift $\pm p$). Likewise $E_\kappa = E_\lambda$ forces $\lambda \in \{\kappa, \tau\kappa\}$, and $E_\kappa = E_{\tau\kappa}$ holds exactly for $\kappa \in B \cup C$. A rich line meets only the classes $\kappa, \sigma\kappa$ (or $\kappa, \tau\kappa$), which lie in one V -orbit. Counting: (a) gives $(n_A + n_D - f_\sigma)/2$ lines, (b) $(n_B + n_C - f_\tau)/2$, (c) $n_B + n_C$, (d) $n_A + n_D$, which is (1).

For the incidences of a single point, take $\kappa \in A$. Its points $\kappa(0,1), \kappa(1,0)$ lie on $E_\kappa \in \mathcal{L}$ (d), and on no line of type (a)/(c) (their slope- $(+1)$ lines $x-y = d_\kappa \pm p$ carry no second point of κ and, by the table, no point of $\sigma\kappa$). $\kappa(1,1)$ lies on D_κ , which is rich iff $\sigma\kappa \neq \kappa$, and on $x+y = e_\kappa + 2p = e_{\tau\kappa} + 3p$, which carries no point of $\tau\kappa$: so $\kappa(1,1)$ is on one rich line if $\sigma\kappa \neq \kappa$ and on none otherwise. $\kappa(0,0)$ lies on D_κ and on $x+y = e_\kappa = e_{\tau\kappa} + p$, i.e. on $E_{\tau\kappa} \in \mathcal{L}$: two rich lines if $\sigma\kappa \neq \kappa$, one otherwise. The types D, B, C are analogous (for B : $\kappa(1,0) \in D_{\sigma\kappa}$ since $d_\kappa + p = d_{\sigma\kappa}$, while $\kappa(0,1)$ lies on $x-y = d_\kappa - p = d_{\sigma\kappa} - 2p$, which carries no point of $\sigma\kappa$; for C the roles of $(1,0), (0,1)$ are exchanged; for D : $\kappa(1,1) \in E_{\tau\kappa}$ since $e_\kappa + 2p = e_{\tau\kappa} + p$, and $\kappa(0,0)$ lies on $x+y = e_\kappa = e_{\tau\kappa} - p$). This gives (2) and the last two assertions. $\square$




Figure 1: The two gadgets of the equality analysis, drawn to scale for $p = 11, c = 1$ (dotted squares: the four copies of a class; grey: rich lines). Generic orbit: the circled points are excluded and the other 12 are forced ($= T_2$). Exceptional orbit: the circled points are forced and each rich line keeps two of its three points, $3 \cdot 3 = 9$ choices.

4 Proof of Theorem 1 and Corollary 2

Part (i) is Corollary 6 and Proposition 8.

Upper bound. Let $S \subseteq \mathcal{P}$ be lawful. For $q \in \mathcal{P}$ let $m(q)$ be the number of rich lines through q . Every rich line carries at most two points of S (Corollary 6), hence

$$\begin{aligned}
 |S| &= \#\{q \in S : m(q) = 0\} + \#\{q \in S : m(q) \geq 1\} \leq Z + \sum_{q \in S} m(q) - \sum_{q \in S} (m(q) - 1)^+ \\
 &= Z + \sum_{\ell \in \mathcal{L}} |S \cap \ell| - \sum_{q \in S} (m(q) - 1)^+ \leq Z + 2|\mathcal{L}|,
 \end{aligned} \tag{3}$$

and by (1), (2), $Z + 2|\mathcal{L}| = 2s + 3(p - 1) - 2s = 3(p - 1)$. The HJSW set attains the bound [1]; the equality analysis below exhibits all sets attaining it, so attainment does not depend on [1].

LP certificate. The same count is valid for fractional $x \in [0, 1]^{\mathcal{P}}$ with $\sum_{q \in \ell} x_q \leq 2$ ($\ell \in \mathcal{L}$): $\sum_q x_q \leq \sum_{q: m(q)=0} x_q + \sum_{\ell \in \mathcal{L}} \sum_{q \in \ell} x_q \leq Z + 2|\mathcal{L}|$. In LP-duality terms the dual solution assigns weight 1 to every rich line and weight 1 to the bound $x_q \leq 1$ of each of the Z points on no rich line. This proves (ii). (The statement concerns the optimum value; we do not claim that the polytope of the relaxation is integral.)

Equality. $|S| = 3(p - 1)$ holds iff all inequalities in (3) are equalities, i.e. iff (I) S contains all Z points lying on no rich line; (II) $|S \cap \ell| = 2$ for every $\ell \in \mathcal{L}$; (III) no point of S lies on two rich lines. Since every rich line lies inside one V -orbit of classes (Proposition 8), conditions (I)–(III) split into independent conditions on the orbits, and the number of maximum sets is the product of the numbers of admissible choices in the orbits. In a generic orbit $\{\kappa, \sigma\kappa, \tau\kappa, \nu\kappa\}$ all four classes have distinct partners; the six rich lines meeting the orbit (two of type (a) or (b), four of type (c) or (d)) meet no other orbit, its four middle copies have $m = 2$ and are excluded by (III), and each of the six lines then retains exactly two points, both with $m = 1$, which are forced into S by (II): on a generic orbit S coincides with $H(c, p) \cap T_2$, and there is exactly one admissible choice. An exceptional orbit $\{\kappa, \nu\kappa\}$ (κ and $\nu\kappa$ fixed by σ , or both fixed by τ) meets

exactly two rich lines, each with three points of $m = 1$, and contains two points with $m = 0$: conditions (I)–(III) allow $\binom{3}{2}\binom{3}{2} = 9$ choices, and all of them are lawful by Corollary 6. There are s exceptional orbits, whence 9^s maximum sets, all agreeing with the HJSW set outside the exceptional orbits, i.e. outside the classes $(a, \pm a)$ with $a^2 = \pm c$. This proves (iii).

Corollary 2. -1 is a QR modulo p iff $p \equiv 1 \pmod{4}$. If $p \equiv 3 \pmod{4}$, exactly one of $c, -c$ is a QR, so $s = 1$; if $p \equiv 1 \pmod{4}$, c and $-c$ have the same character, so $s \in \{0, 2\}$ according as c is a non-residue or a residue. $\square$

Remark 9. (a) In the language of [2, Constr. 3.2, Rem. 3.4] with $W = \{c\}$, $k = 2$ (so $S(W) = \mathcal{P}$ and L' is the set of lines of slope ± 1): $X_{2,p}(\{c\}) = \sum_{\ell} (|\ell \cap \mathcal{P}| - 2)^+ = (p-1) \cdot 1 + (\frac{1}{2}(p-1) - s) \cdot 2 = 2(p-1) - 2s$ by Theorem 1(i), so the general bound $|R| \geq \frac{1}{2}X$ gives $|R| \geq p-1-s$ and hence, for every lawful $S \subseteq \mathcal{P}$, $|S| \leq |\mathcal{P}| - |R| \leq 3(p-1) + s$ (a lawful S is a trimming, by Corollary 6). The averaging loses exactly one point per exceptional orbit, because each such orbit contains two points on no rich line and no point on two rich lines; the count (3) keeps track of these points and is exact.

(b) The theorem says that no lawful set can use the middle block M profitably: adding fourth copies to the HJSW set is impossible without removing at least as many points. Any improvement of the HJSW construction must therefore use points outside $H(c, p) \cap G(p)$ (another hyperbola, a larger window — by Theorem 3 not merely a shifted $2p \times 2p$ box — or non-hyperbola points).

(c) The proof uses only: at most two classes per line (Lemma 4), the square arrangement of the four copies, and the position of the partner classes (Lemma 7). It does not use $p \geq 7$; $p = 3$ and $p = 5$ are covered as well.

5 All $2p \times 2p$ boxes

Let $W = [x_0, x_0 + 2p) \times [y_0, y_0 + 2p)$ with $x_0, y_0 \in \mathbb{Z}$ (only x_0, y_0 modulo p matter), $\mathcal{P}_W = H(c, p) \cap W$, and let $\rho_x(u) \in [x_0, x_0 + p)$, $\rho_y(u) \in [y_0, y_0 + p)$ denote the representatives of a residue u . Every class $\kappa = (a, b)$ of $H(c, p)$ again has exactly four points $\kappa(r, s) = (\rho_x(a) + rp, \rho_y(b) + sp)$, $r, s \in \{0, 1\}$, in W ; put $d_\kappa = \rho_x(a) - \rho_y(b)$, $e_\kappa = \rho_x(a) + \rho_y(b)$, $D_\kappa = \{x - y = d_\kappa\}$, $E_\kappa = \{x + y = e_\kappa + p\}$. Lemmas 4, 5 and Corollary 6 hold verbatim (they do not use the position of the box), so a rich line is D_κ for a class with $\sigma\kappa \neq \kappa$ or E_κ for a class with $\tau\kappa \neq \kappa$, and its points are the two copies $\kappa(0, 0), \kappa(1, 1)$ (resp. $\kappa(0, 1), \kappa(1, 0)$) together with the copies of the partner class on the same line. Since $d_\kappa \equiv d_{\sigma\kappa} \pmod{p}$ and both lie in an interval of length $2p - 2$, $d_{\sigma\kappa} - d_\kappa \in \{-p, 0, p\}$; we call the σ -pair $\{\kappa, \sigma\kappa\}$ *shared* if $d_{\sigma\kappa} = d_\kappa$ (then $D_\kappa = D_{\sigma\kappa}$ carries the four points $\kappa(0, 0), \kappa(1, 1), \sigma\kappa(0, 0), \sigma\kappa(1, 1)$) and *split* otherwise (then D_κ carries $\kappa(0, 0), \kappa(1, 1)$ and one copy of $\sigma\kappa$, and $D_{\sigma\kappa}$ carries $\sigma\kappa(0, 0), \sigma\kappa(1, 1)$ and one copy of κ). Likewise a τ -pair is shared ($E_\kappa = E_{\tau\kappa}$, four points) or split (two lines with three points). In the HJSW box the σ -pairs in $A \cup D$ and the τ -pairs in $B \cup C$ are shared and the others are split (Lemma 7); in the box $[0, 2p)^2$ all pairs are shared.

Lemma 10 (orbit lemma). *Let $O = \{\kappa, \sigma\kappa, \tau\kappa, \nu\kappa\}$ be a generic orbit, and let e_O (resp. f_O) be the number of split σ -pairs (resp. τ -pairs) among the two σ -pairs and two τ -pairs of O . Then $e_O + f_O \in \{0, 2\}$; in particular $e_O + f_O \leq 2$.*

Proof. Write $\kappa = (a, b)$, $X_u = \rho_x(u)$, $X'_u = \rho_x(-u)$, $Y_u = \rho_y(u)$, $Y'_u = \rho_y(-u)$ for $u \in \{a, b\}$. The base copies are $P = (X_a, Y_b)$ of κ , $Q = (X'_b, Y'_a)$ of $\sigma\kappa$, $R = (X_b, Y_a)$ of $\tau\kappa$ and $T = (X'_a, Y'_b)$ of $\nu\kappa$; the σ -pairs are $\{P, Q\}$, $\{R, T\}$ and the τ -pairs $\{P, R\}$, $\{Q, T\}$. With $d(x, y) = x - y$, $e(x, y) = x + y$, the four split conditions are $E_1 : d_P \neq d_Q$, $E_2 : d_T \neq d_R$, $F_1 : e_P \neq e_R$, $F_2 : e_Q \neq e_T$; each of the four differences is $\equiv 0 \pmod{p}$ and lies in $[-p, p]$, hence is one of $-p, 0, p$.

Let m_x be the least multiple of p exceeding $2x_0$ and $\mu_x = m_x - 2x_0 \in [1, p]$; similarly m_y, μ_y . Since $X_u + X'_u \equiv 0 \pmod{p}$ and $X_u + X'_u \in [2x_0 + 1, 2x_0 + 2p - 3]$, we have $X_u + X'_u = m_x + \varepsilon_x(u)p$ with $\varepsilon_x(u) \in \{0, 1\}$, and explicitly $\varepsilon_x(u) = [X_u - x_0 > \mu_x]$; likewise $Y_u + Y'_u = m_y + \varepsilon_y(u)p$,

$\varepsilon_y(u) = [Y_u - y_0 > \mu_y]$. Put $\Delta = \varepsilon_x(a) - \varepsilon_x(b)$, $\Gamma = \varepsilon_y(a) - \varepsilon_y(b)$, $A = \varepsilon_x(b) - \varepsilon_y(a)$, $B = \varepsilon_x(a) - \varepsilon_y(b)$, and define the integers θ, η by

$$(X_a + X_b) - (Y_a + Y_b) - (m_x - m_y) = \theta p, \quad (X_b - X_a) - (Y_b - Y_a) = \eta p.$$

Substituting $X'_u = m_x + \varepsilon_x(u)p - X_u$, $Y'_u = m_y + \varepsilon_y(u)p - Y_u$ one finds

$$d_P - d_Q = (\theta - A)p, \quad d_T - d_R = (B - \theta)p, \quad e_R - e_P = \eta p, \quad e_Q - e_T = (\Gamma - \Delta - \eta)p,$$

so that $E_1 \Leftrightarrow \theta \neq A$, $E_2 \Leftrightarrow \theta \neq B$, $F_1 \Leftrightarrow \eta \neq 0$, $F_2 \Leftrightarrow \eta \neq \Gamma - \Delta$, and $\theta - A$, $\theta - B$, η , $\eta - (\Gamma - \Delta) \in \{-1, 0, 1\}$.

Parity. For $x \in \{-1, 0, 1\}$, $[x \neq 0] \equiv x \pmod{2}$. Hence $E_1 + E_2 + F_1 + F_2 \equiv (\theta - A) + (\theta - B) + \eta + (\eta - \Gamma + \Delta) \equiv A + B + \Gamma - \Delta = 2\varepsilon_x(b) - 2\varepsilon_y(b) \equiv 0 \pmod{2}$: the number of split pairs in O is even.

If F_1 holds, then E_1 and E_2 do not both hold. Put $s = X_a - x_0$, $t = X_b - x_0 \in [0, p-1]$ and $v = (x_0 - y_0) \bmod p$; since $Y_u \equiv X_u$, we have $Y_a - y_0 = s + v - c_a p$ and $Y_b - y_0 = t + v - c_b p$ with $c_a = [s + v \geq p]$, $c_b = [t + v \geq p]$, and therefore $\eta = c_b - c_a$. So F_1 means $c_a \neq c_b$; the substitution $a \leftrightarrow b$ exchanges $E_1 \leftrightarrow E_2$ and fixes F_1 , so we may assume $c_a = 0$, $c_b = 1$, i.e. $s \leq p-1-v < t$. Since $\mu_x - \mu_y \equiv -2v \pmod{p}$ and $|\mu_x - \mu_y| \leq p-1$, we have $\mu_x - \mu_y = -2v + kp$ with $k \in \{0, 1, 2\}$, and $\theta p = (2x_0 + s + t) - (2y_0 + s + t + 2v - p) - (\mu_x + 2x_0 - \mu_y - 2y_0) = p - 2v - (\mu_x - \mu_y) = (1-k)p$, i.e. $\theta = 1 - k$. With $\varepsilon_x(a) = [s > \mu_x]$, $\varepsilon_x(b) = [t > \mu_x]$, $\varepsilon_y(a) = [s + v > \mu_y]$, $\varepsilon_y(b) = [t + v - p > \mu_y]$ and $\mu_y = \mu_x + 2v - kp$:

$k = 1$: $E_1 \Leftrightarrow [t > \mu_x] \neq [s > \mu_x + v - p]$ and $E_2 \Leftrightarrow [s > \mu_x] \neq [t > \mu_x + v]$. If $s > \mu_x$ then $t > \mu_x$ and $s > \mu_x + v - p$, so E_1 fails. If $s \leq \mu_x$, E_2 forces $t > \mu_x + v$, so $\mu_x + v \leq p-2$ and $[t > \mu_x] = 1$, and E_1 forces $s \leq \mu_x + v - p < 0$, impossible.

$k = 0$: here $\mu_y = \mu_x + 2v \leq p$, so $\varepsilon_y(b) = 0$ and $E_2 \Leftrightarrow s \leq \mu_x$; then $\varepsilon_y(a) = [s > \mu_x + v] = 0$ and $E_1 \Leftrightarrow t \leq \mu_x$, but $t \geq p - v > p - 2v \geq \mu_x$.

$k = 2$: here $\mu_y = \mu_x + 2v - 2p \geq 1$, so $\mu_x \geq 2p + 1 - 2v$ and $\varepsilon_y(a) = 1$; $E_1 \Leftrightarrow t > \mu_x$ and $E_2 \Leftrightarrow (s > \mu_x) \vee (t \leq \mu_x + v - p)$. Now $s > \mu_x$ would give $2p + 2 - 2v \leq s \leq p - 1 - v$, i.e. $v \geq p + 3$; so $s \leq \mu_x$ and E_2 needs $t \leq \mu_x + v - p < \mu_x$, contradicting E_1 .

Consequently, if F_1 holds then $E_1 + E_2 + F_1 + F_2 \leq 3$, and if F_1 fails then again $E_1 + E_2 + F_1 + F_2 \leq 3$; by parity the sum is 0 or 2. $\square$

Lemma 11 (gadget maxima). *Let O be an orbit and $\mathcal{P}_O \subseteq \mathcal{P}_W$ its 16 (or 8) points. Every rich line meeting $\mathcal{P}_O$ lies in $\mathcal{P}_O$, and the largest lawful subset of $\mathcal{P}_O$ has size and multiplicity as follows: exceptional orbit: 6 points, 9 sets if the pair is split and 6 if it is shared; generic orbit with $(e_O, f_O) = (0, 0)$: 8 points, $6^4 = 1296$ sets; $(1, 1)$: 10 points, 144 sets; $(0, 2)$ or $(2, 0)$: 12 points, one set.*

Proof. A rich line meets only a class and its σ - or τ -partner (Lemma 4), hence lies in one orbit. *Exceptional orbit* $\{\kappa, \nu\kappa\}$, say $\sigma\kappa = \kappa$: the rich lines are $E_\kappa = E_{\nu\kappa}$ (four points) if the τ -pair is shared, resp. $E_\kappa, E_{\nu\kappa}$ (three points each) if it is split; the remaining four (resp. two) points are free: $2 + 4 = 6$ points in $\binom{4}{2} = 6$ ways, resp. $2 + 2 + 2 = 6$ points in $3 \cdot 3 = 9$ ways. *Pattern* $(0, 0)$: the two shared diagonals and the two shared antidiagonals are four disjoint four-point lines covering $\mathcal{P}_O$: $4 \cdot 2 = 8$ points in 6^4 ways. *Pattern* $(2, 0)$ (and symmetrically $(0, 2)$): the four diagonals D_λ ($\lambda \in O$) are rich with three points each and the two antidiagonals $E_\kappa = E_{\tau\kappa}$, $E_{\sigma\kappa} = E_{\nu\kappa}$ have four points; every class λ has exactly one point of multiplicity $m = 2$, namely its off-diagonal copy lying on $D_{\sigma\lambda}$ (which is also on the antidiagonal of λ). Each D_λ carries $\lambda(0, 0)$, $\lambda(1, 1)$ ($m = 1$) and one $m = 2$ point, each antidiagonal carries two $m = 1$ and two $m = 2$ points. By (3) restricted to O (with $Z = 0$), $|S \cap \mathcal{P}_O| \leq 12$ with equality iff S avoids the four $m = 2$ points and takes both remaining points of each of the six lines — which is lawful; so

the maximum is 12 and it is unique. (For the HJSW box this is the generic orbit of Section 4.) *Pattern* (1, 1): the split σ -pair and the split τ -pair share exactly one class κ (any σ -edge and any τ -edge of the four-cycle $\kappa - \tau\kappa - \nu\kappa - \sigma\kappa - \kappa$ meet in one vertex). The six rich lines are

$$\begin{aligned} A &:= D_\kappa, & B &:= E_\kappa, & A' &:= D_{\sigma\kappa}, & B' &:= E_{\tau\kappa} & \text{(three points each),} \\ A'' &:= D_{\tau\kappa} = D_{\nu\kappa}, & B'' &:= E_{\sigma\kappa} = E_{\nu\kappa} & & & & & \text{(four points each).} \end{aligned}$$

Write κ_A, κ'_A for the two copies of κ on A (these are $\kappa(0,0), \kappa(1,1)$) and κ_B, κ'_B for the two on B ; the third point of A is a copy $\bar{\sigma}$ of $\sigma\kappa$ off its own diagonal, hence on B'' ; the third point of B is a copy $\bar{\tau}$ of $\tau\kappa$ off its own antidiagonal, hence on A'' ; the third point of A' is a copy of κ off A , i.e. one of κ_B, κ'_B — name it κ_B ; the third point of B' is one of κ_A, κ'_A — name it κ_A . No other incidences occur (a rich line of slope +1 through a copy of κ is A or A' , of slope -1 it is B or B' , and $A \cap B = \emptyset$ because a common lattice point would have $x = x_\kappa + p/2$). Hence $m = 2$ exactly for the four points $\kappa_A \in A \cap B'$, $\kappa_B \in B \cap A'$, $\bar{\sigma} \in A \cap B''$, $\bar{\tau} \in B \cap A''$, and $m = 1$ for the other twelve: A and B consist of one $m = 1$ point and two $m = 2$ points, A' and B' of two $m = 1$ points and one $m = 2$ point, A'' and B'' of three $m = 1$ points and one $m = 2$ point. For lawful $S \subseteq \mathcal{P}_O$ let t be the number of $m = 2$ points in S ; since every point of $\mathcal{P}_O$ lies on a rich line, $|S| = \sum_\ell |S \cap \ell| - t$ over the six lines, each term being ≤ 2 . If $t = 0$ then $|S \cap A|, |S \cap B| \leq 1$ and $|S| \leq 1 + 1 + 2 + 2 + 2 + 2 = 10$; if $t = 1$, the $m = 2$ point lies on exactly one of A, B , the other of the two carries at most one point of S , and $|S| \leq 11 - 1 = 10$; if $t \geq 2$ then $|S| \leq 12 - t \leq 10$. So the maximum is at most 10, and a set of size 10 makes every inequality used tight. Counting these sets by t (every set listed is lawful, since each rich line receives at most two of its points):

$t = 0$: S contains κ'_A, κ'_B , both $m = 1$ points of A' and both of B' (forced), and two of the three $m = 1$ points of A'' and of B'' : $3 \cdot 3 = 9$ sets.

$t = 1$: the $m = 2$ point q is one of $\kappa_A, \bar{\sigma} \in A$ or $\kappa_B, \bar{\tau} \in B$; say $q \in A$ (the case $q \in B$ is symmetric). Then $S \cap A = \{q, \kappa'_A\}$ and $S \cap B = \{\kappa'_B\}$; the second line through q receives q and one of its $m = 1$ points, and each of the three remaining lines receives two of its $m = 1$ points. For $q = \kappa_A$: B' in 2 ways, A' forced, A'' and B'' in $3 \cdot 3$ ways — 18 sets; for $q = \bar{\sigma}$: B'' in 3 ways, A' and B' forced, A'' in 3 ways — 9 sets. With the symmetric case $q \in B$: $2 \cdot (18 + 9) = 54$ sets.

$t = 2$: the two $m = 2$ points cannot both lie on A (then $|S \cap B| \leq 1$ and $|S| \leq 11 - 2 = 9$), nor both on B ; so one is on A and one on B , every line receives exactly two points, and $S \cap A = \{q, \kappa'_A\}$, $S \cap B = \{q', \kappa'_B\}$. For $\{q, q'\} = \{\kappa_A, \kappa_B\}$: B' and A' each receive their $m = 2$ point and one of two $m = 1$ points ($2 \cdot 2$), A'' , B'' two of three ($3 \cdot 3$): 36 sets; for $\{\kappa_A, \bar{\tau}\}$: B' 2 ways, A'' ($\ni \bar{\tau}$) 3 ways, A' forced, B'' 3 ways: 18; for $\{\bar{\sigma}, \kappa_B\}$: symmetrically 18; for $\{\bar{\sigma}, \bar{\tau}\}$: B'' 3, A'' 3, A' , B' forced: 9. Total $36 + 18 + 18 + 9 = 81$.

$t \geq 3$: $|S| \leq 12 - t \leq 9$: no maximum set.

Altogether $9 + 54 + 81 = 144$ maximum sets of size 10. (An independent brute-force enumeration of the 16-point gadgets, `slack/gadget11_check.py`, confirms the line structure and the counts 9, 54, 81 for every (1, 1)-orbit encountered in all boxes for $p \leq 13$ and in samples of boxes for $p \leq 31$.) $\square$

Proof of Theorem 3. By Lemma 11 the rich lines of $\mathcal{P}_W$ lie inside the orbits, so a set is lawful iff its intersection with every orbit is lawful, and the maximum and the number of maximum sets are the sum, resp. the product, of the orbit values. By Lemma 10 every generic orbit has pattern (0, 0), (1, 1), (0, 2) or (2, 0), and Lemma 11 gives the formula. Since $n_2 + n_1 + n_0 = \frac{1}{4}(p - 1 - 2s)$, the maximum is $12n_2 + 10n_1 + 8n_0 + 6s \leq 12(n_2 + n_1 + n_0) + 6s = 3(p - 1)$, with equality iff $n_1 = n_0 = 0$. For the HJSW box, Lemma 7 gives $(e, f) = (0, 2)$ for the A/D -orbits and (2, 0) for the B/C -orbits, and the exceptional pairs are split; this recovers Theorem 1(ii),(iii). $\square$

Remark 12 (algorithmic consequence). Selecting a largest subset in general position from an arbitrary point set is NP-hard [5]; for $H(c, p) \cap W$ Theorems 1 and 3 reduce it to bookkeeping: after listing the $p - 1$ classes and their V -orbits, the maximum, the number of maximum sets and an explicit maximum set are obtained in $O(p)$ operations (the pattern of an orbit is read off from the split/shared status of its four pairs).

Remark 13. The double count of Section 4 gives, for every box, $|S| \leq Z + 2|\mathcal{L}| = 2(p - 1) + 2s + 2\sum_O(e_O + f_O) = 12n_2 + 12n_1 + 8n_0 + 6s$ (one checks $Z + 2|\mathcal{L}| = 2(p - 1) + 2s + \#\{m = 2\}$ and $\#\{m = 2\} = 2\sum_O(e_O + f_O)$ exactly as in Section 4); it is off by exactly 2 on every $(1, 1)$ -orbit, where two rich three-point lines consist of one point of multiplicity one and two of multiplicity two.

Corollary 14 (no-four-in-line). *In every box W the four-point rich lines are pairwise disjoint; hence the largest subset of $\mathcal{P}_W$ with no four collinear points has exactly $4(p - 1) - N_{sh}(W)$ points, where $N_{sh}(W)$ denotes the number of shared pairs (σ -pairs and τ -pairs together; equivalently, the number of four-point rich lines of $\mathcal{P}_W$), and there are $4^{N_{sh}(W)}$ such subsets. By the orbit lemma $N_{sh}(W) \geq 2 \cdot \#\{\text{generic orbits}\} = \frac{1}{2}(p - 1) - s$, so the maximum is at most $\frac{7}{2}(p - 1) + s$; in the HJSW box $N_{sh} = \frac{1}{2}(p - 1) - s$ and the set $S_3 = H(c, p) \cap T_3$ of [2, Constr. 2.10] (with $|S_3| = \frac{7}{2}(p - 1) + s$) is a largest no-four-in-line subset of $H(c, p) \cap G(p)$, as asserted in [2, App. A]. (No line carries five points of $\mathcal{P}_W$, so the no-five-in-line maximum is $|\mathcal{P}_W| = 4(p - 1)$.)*

Proof. A four-point rich line is D_κ for a shared σ -pair (points $\kappa(0, 0), \kappa(1, 1), \sigma\kappa(0, 0), \sigma\kappa(1, 1)$) or E_κ for a shared τ -pair (points with $r + s = 1$). The slope- (-1) line through a point $\kappa(r, r)$ is not of the second kind: if it were E_λ then $\kappa \in \{\lambda, \tau\lambda\}$ by Lemma 4, so $E_\lambda = E_\kappa$, which contains only the copies $\kappa(0, 1), \kappa(1, 0)$ of κ ; likewise the slope- $(+1)$ line through a point $\kappa(r, 1 - r)$ is not of the first kind. Hence the four-point lines are pairwise disjoint, a no-four-in-line subset must miss at least one point of each of them, and deleting exactly one point from each is sufficient (all other lines carry at most three points). For the HJSW box, $N_{sh} = \frac{1}{2}(p - 1) - s$ by Lemma 7, and $|S_3| = |S_2| + |M \cap E \cap H(c, p)| = 3(p - 1) + \frac{1}{2}(p - 1) + s$ because the classes fixed by σ or τ lie in E and the remaining classes split evenly between E and F [2, Obs. 2.9]. $\square$

For the box $[0, 2p)^2$ all pairs are shared ($n_2 = n_1 = 0, s_2 = s$): the maximum is $8n_0 + 6s = 2(p - 1) + 2s$ with $1296^{n_0}6^s$ maximum sets. Table 1 shows the exact maxima for all boxes at $p = 11, c = 1$ (two generic orbits, $s = 1$): the values 22, 24, 26, 28, 30 correspond to $(b, c) = (0, 2), (1, 1), (2, 0)$ or $(0, 1), (1, 0), (0, 0)$. The number of boxes attaining $3(p - 1)$ depends on c (for $p = 13$: 28 of the 169 boxes for $c = 1$, 4 for $c = 2$); it always includes the HJSW box $x_0 = -h, y_0 \in \{0, 1\}$.

Remark 15 (larger boxes). In boxes larger than $2p \times 2p$ the classes have more copies; the maximum is then at least $3(p - 1)$ (equal to it for $N = 2p, 2p + 1$ in the cases computed and larger from $N = 2p + 2$ on), and the ratio maximum/ N does not exceed the HJSW value $3(p - 1)/2p$ in the range computed (it is not monotone in N ; for $p = 7$ the $3p \times 3p$ box ties, $27/21 = 18/14$). Exact values (MIP with zero gap, $c = 1$, box $[-h, -h + N) \times [0, N)$): for $p = 11$ and $N = 22, 23, \dots, 33$ the maxima are 30, 30, 31, 31, 33, 36, 36, 38, 39, 40, 42, 43; for $p = 13, N = 26, \dots, 39$: 36, 36, 38, 38, 38, 38, 40, 40, 43, 45, 47, 49, 51, 52; for $p = 17, N = 34, \dots, 51$: 48, 48, 49, 49, 50, 51, 52, 53, 56, 56, 58, 61, 62, 63, 65, 66, 67, 69. In the $3p \times 3p$ box the maximum is 27, 43, 52, 69, 77, 93 for $p = 7, 11, 13, 17, 19, 23$: about $4.3(p - 1)$, i.e. ratio 1.29–1.35 to the side $3p$, against $3(p - 1)/2p = 1.29, 1.36, 1.38, 1.41, 1.42, 1.43$ for the HJSW box at the same primes (the value depends on c : 51, 52, 54 for $p = 13, c = 2, 1, 3$). In the $2p \times 4p$ box the trivial bound $4(p - 1)$ (two of the four collinear copies in each column) is attained for $p = 7, 11, 13$. Data: slack/verification/windows_2p_to_3p.txt.

$x_0 + h \setminus y_0$	0	1	2	3	4	5	6	7	8	9	10
0	30	30	30	28	26	24	22	24	26	28	30
1	28	28	28	30	28	26	24	26	28	30	28
2	26	26	26	28	30	28	26	28	30	28	26
3	24	24	24	26	28	30	28	30	28	26	24
4	22	22	22	24	26	28	30	28	26	24	22
5	22	22	22	24	26	28	30	28	26	24	22
6	22	22	22	24	26	28	30	28	26	24	22
7	22	22	22	24	26	28	30	28	26	24	22
8	24	24	24	26	28	30	28	30	28	26	24
9	26	26	26	28	30	28	26	28	30	28	26
10	28	28	28	30	28	26	24	26	28	30	28

Table 1: $p = 11$, $c = 1$: exact maximum lawful subset of $H(1, 11)$ in the box $[x_0, x_0 + 22) \times [y_0, y_0 + 22)$, indexed by $x_0 + h \bmod p$ (rows; 0 is the HJSW position $x_0 = -h$) and $y_0 \bmod p$ (columns). Values (HiGHS MIP, witnesses through the law; they agree with the formula of Theorem 3) range from $2(p-1) + 2s = 22$ to $3(p-1) = 30$; the double-count bound $Z + 2|\mathcal{L}|$ equals 30 for 76 boxes but is attained for 20 (the others have a $(1, 1)$ -orbit).

6 Computational observations beyond one hyperbola

Nothing in this section is a theorem: it records finite computations and the picture they suggest. Theorems 1 and 3 concern a single hyperbola. Unions of two hyperbolae in the same box do exceed $3(p-1)$, but — as far as we can compute — only by a few points. For $\mathcal{P}_k = (H(1, p) \cup H(k, p)) \cap G(p)$ ($8(p-1)$ candidate points) maximum lawful subsets were computed by SAT solving (kissat, CaDiCaL; two cardinality encodings), mixed-integer programming (HiGHS) or constraint programming (OR-tools CP-SAT), as indicated row by row; a value is called exact when a solver proved optimality (SAT: unsatisfiability at the next size; MIP/CP-SAT: zero gap); every witness was verified by an independent checker of all triples:

p	$3(p-1)$	k	max	gain	LP relax.	method
11	30	3	35	+5	36.00	MIP; all k : 32–35 ($k = 2$: 33, also SAT twice)
13	36	2	41	+5	47.54	SAT (two encodings), MIP; all k : 38–41
17	48	−1	54	+6	60.15	MIP; all k : 49–54 ($k = 2$: 53)
19	54	−1	59	+5	63.64	MIP (2 min); local search 59
23	66	−1	70–75	$\geq +4$	—	CP-SAT (8 workers, 4 h): 70, dual bound 78; MIP: 70, dual bound 75.4 (running); $k = 2$: 68–80, $k = 3$: 69–78

Three observations. (i) The double count of Section 4 does not extend: rich lines of general slope appear (a line meets each hyperbola in at most two classes, and copies of four different classes can be collinear), their number grows quickly with p , and the LP relaxation (“at most two per rich line”) is far from the truth (column “LP”), whereas for one hyperbola it is exact (Theorem 1(ii)). (ii) The gain over $3(p-1)$, maximised over k (+5, +5, +6, +5 for $p = 11, 13, 17, 19$; all k computed for $p \leq 17$, $k \in \{-1, 3\}$ for $p = 19$; at $p = 23$ the exact value is not settled, $70 \leq \max \leq 75$ for $k = -1$), shows no growth. A parabola $y = x^2 + k$ together with $H(1)$ in the same box behaves alike: exact maxima 34, 39, 51, 58 for $p = 11, 13, 17, 19$ (+4, +3, +3, +4; MIP). Three hyperbolae $H(1) \cup H(2) \cup H(3)$ ($12(p-1)$ candidates) give 37 at $p = 11$ (exact, MIP; +7) and at least 42 at $p = 13$ (MIP, dual bound 47; +6 so far) — again a few points more than two hyperbolae, not a new order of growth. (iii) A heuristic explanation: fix the HJSW set S_2 of $H(1, p)$ and call a point q of $H(k, p) \cap G(p)$ *blocked* if some line through q carries two points of S_2 . The average number of blocking pairs per point grows like $\log p$ (about 3.1 at $p = 11$ and 7 at $p = 101$ for $k \in \{2, 3, -1\}$), and the number of unblocked points is 0–2 for every $p \leq 101$ tested; freeing a point of the second hyperbola costs deletions in the first. This suggests that unions of hyperbolae in the $2p \times 2p$ box gain $O(1)$ (or $o(p)$) over the HJSW value, i.e. that their

asymptotic ratio is still $3/2$; we have no proof. The same picture appears for hyperbolae with different moduli in one box: for $H(1, p) \cup H(1, q)$ in the $2p \times 2p$ box (each hyperbola with its own HJSW shift) the exact maxima (MIP) exceed $3(p-1)$ by 1, 2 or 3 in all eleven pairs computed, $7 \leq p < q \leq 29$ — e.g. 32 vs 30 for (11, 13), 38 vs 36 for (13, 17), 50 vs 48 for (17, 19), 68 vs 66 for (23, 29) — and the number of collinear triples with points on both hyperbolae grows, in the range computed ($N \leq 586$), like $N \log N$ — the ratio (number of such triples)/($N \ln N$) stabilises near 11 — as for random sets of the same density. (Definitions and data: `slack/crosstripples.py`, `docs/research/slack_experiments.md`.)

7 Independent computer verification

The theorem has a short self-contained proof; the following check is an independent sanity test of the bookkeeping (the classification of the rich lines, the exceptional classes and the count of the maximum sets), performed by a program that trusts no part of the proof. The script `slack/hjsw_window_check.py` of the repository recomputes, from the point set alone: all lines with at least three points of $\mathcal{P}$ (found by brute force over all pairs of points; all have slope ± 1 , three or four points, and lie inside one V -orbit); the identity $2|\mathcal{L}| + Z = 3(p-1)$; the exact maximum by exhaustive search inside each orbit (at most 2^{16} subsets), assembling a maximum witness which is passed through the independent verifier `saturation.the.law` in grid coordinates; and the number of maximum sets, compared with 9^s . All checks pass for every prime $3 \leq p \leq 101$ (all $c \in \mathbb{F}_p^*$ for $p \leq 31$; $c \in \{1, 2, 3, -1, -2, (p+1)/2\}$ for larger p); log: `slack/verification/hjsw_window_check_all.log`. For Theorem 3 the script `slack/hjsw_boxes_check.py` (log `slack/verification/hjsw_boxes_check.log`) checks, for every box $(x_0, y_0 \bmod p)$ and every c : the formula $Z + 2|\mathcal{L}| = 2(p-1) + 2s + 2\sum_O(e_O + f_O)$ against brute-force line enumeration (random boxes, $p \leq 19$); $Z + 2|\mathcal{L}| \leq 3(p-1)$ for all primes $p \leq 61$ (with equality for some boxes); the orbit lemma orbit by orbit ($(e_O, f_O) \in \{(0, 0), (0, 2), (1, 1), (2, 0)\}$ only) for all $p \leq 31$; every identity used in the proof of Lemma 10 ($\theta = c_a + c_b - k$, $\eta = c_b - c_a$, the four split conditions, parity, and “ $F_1 \Rightarrow \neg(E_1 \wedge E_2)$ ”) on random instances up to $p = 199$; the maxima and multiplicities of Lemma 11 orbit by orbit (exhaustive search over the 2^{16} subsets) for all $p \leq 13$, all c and all boxes; and the formula $12n_2 + 10n_1 + 8n_0 + 6s$ against MIP optima on random boxes with $p \leq 31$. The exact maxima of Table 1 are MIP optima (HiGHS) with witnesses through the law. The values agree with the exact maxima obtained earlier in the project by SAT solving (kissat/CaDiCaL, two encodings, witnesses through the law): 18, 30, 36, 48, 54, 66 for $p = 7, 11, 13, 17, 19, 23$. Two external LLM-assisted audits (ChatGPT deep-research reports commissioned by the author of record on v0.1 and on v0.5 of this note; kept in `docs/reviews/`) re-derived the point sets and the rich lines with clean-room implementations and confirmed $3(p-1)$, 9^s , the LP value and the no-four values for all c with $p \leq 31$ (samples for larger p), all box patterns and gadget counts through $p = 13$, and the shifted-box table for $p = 11$; the second audit also identified a gap in the earlier presentation of Lemma 11 (the $(1, 1)$ count), which is now proved above.

Repository (public): <https://github.com/iwasborninbali/saturation>.

This version of the note is the immutable tag `hjsw-note-v0.6`: file `paper/hjsw_window.tex`; scripts `slack/hjsw_window_check.py`, `slack/hjsw_boxes_check.py`, `slack/gadget11_check.py`; logs in `slack/verification/`; checksums and the list of tags in `docs/RELEASES.md`.

AI/tool-use disclosure and authorship statement. The theorems, their proofs, the verification programs and the text of this note were produced by autonomous large-language-model agents (Anthropic Claude, model *Claude Fable 5*, August 2026) working in a shared repository: one agent found and proved the theorems and wrote the note; two others had established the computational setup and observations (Section 6) and revised the note after the external audits. The author of record posed the question, directed and organised the work, commissioned the

p	c	$ \mathcal{P} $	$ \mathcal{L} $	Z	max	#max sets
7	1	24	8	2	18	9
11	1	40	14	2	30	9
13	1	48	16	4	36	81
13	2	48	18	0	36	1
17	1	64	22	4	48	81
19	1	72	26	2	54	9
23	1	88	32	2	66	9
101	1	400	148	4	300	81

Table 2: Sample values from the verification $\log(|\mathcal{L}| = \frac{3}{2}(p-1) - s, Z = 2s)$.

external audits, decided what to claim and publish, and takes responsibility for the content; the extent of his independent verification of the proofs is recorded in `docs/HUMAN_AUDIT.md` of the repository and will be stated precisely in any submitted version. The AI systems are not authors. All code, logs and the correspondence of the agents are public in the repository.

References

- [1] R. R. Hall, T. H. Jackson, A. Sudbery, K. Wild, Some advances in the no-three-in-line problem, *J. Combin. Theory Ser. A* 18 (1975) 336–341.
- [2] B. Kovács, Z. L. Nagy, D. R. Szabó, Randomised algebraic constructions for the no- $(k+1)$ -in-line problem, arXiv:2508.07632 (v1 11 Aug 2025, v2 21 Jul 2026); *Advances in Combinatorics* 2026:7, doi:10.19086/aic.2026.7 (to appear).
- [3] I. E. Shparlinski, Modular hyperbolas, *Jpn. J. Math.* 7 (2012) 235–294; arXiv:1103.2879.
- [4] M. R. Khan, R. Magner, S. Senger, A. Winterhof, Two combinatorial geometric problems involving modular hyperbolas, arXiv:1304.6943.
- [5] V. Froese, I. Kanj, A. Nichterlein, R. Niedermeier, Finding points in general position, *Internat. J. Comput. Geom. Appl.* 27 (2017) 277–296; arXiv:1508.01097.
- [6] M. S. Payne, D. R. Wood, On the general position subset selection problem, *SIAM J. Discrete Math.* 27 (2013) 1727–1733.
- [7] D. T. Nagy, Z. L. Nagy, R. Woodroffe, The extensible no-three-in-line problem, *European J. Combin.* 114 (2023) 103796.
- [8] C. Y. Ku, K. B. Wong, On no-three-in-line problem on m -dimensional torus, *Graphs Combin.* 34 (2018) 355–364.
- [9] A. Ghosal, R. Goenka, A. Grebennikov, P. Keevash, M. Kwan, H. T. Pham, No- $(k+1)$ -in-line problem for $k \geq 3$, arXiv:2607.05255 (2026).
- [10] B. Green, 100 open problems, manuscript (version of December 2025), Problem 72.
- [11] A. Flammenkamp, The no-three-in-line problem, database and tables, <https://www.homes.uni-bielefeld.de/achim/no3in/readme.html>.
- [12] T. Prellberg, Constraint satisfaction programming for the no-three-in-line problem, arXiv:2602.07751 (2026).