

Directed equality with dinaturality

Andrea Laretto, Fosco Loregian, Niccolò Veltri

Tallinn University of Technology

5th Itaca Workshop, Padova

December 20th, 2024

Type theory and symmetric equality

- A very interesting aspect of Martin-Löf type theory/logic: *equality*.
- For simplicity, consider first order logic:

$$\frac{A \text{ type}}{[a : A, b : A] \ a = b \text{ prop}}$$

Type theory and symmetric equality

- A very interesting aspect of Martin-Löf type theory/logic: *equality*.
- For simplicity, consider first order logic:

$$\frac{A \text{ type}}{[a : A, b : A] \ a = b \text{ prop}}$$

$$\frac{}{[x : A] \ F \vdash x = x} \text{ (refl)}$$

Type theory and symmetric equality

- A very interesting aspect of Martin-Löf type theory/logic: *equality*.
- For simplicity, consider first order logic:

$$\frac{}{[x : A] \Gamma \vdash x = x} \text{ (refl)} \quad \frac{\frac{A \text{ type}}{[a : A, b : A] a = b \text{ prop}} \quad \frac{[z : A, x : C] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : A, b : A, x : C] a = b, \Gamma(a, b, x) \vdash P(a, b, x)} \text{ (J)}}{[a : A, b : A, x : C] a = b, \Gamma(a, b, x) \vdash P(a, b, x)}$$

Type theory and symmetric equality

- A very interesting aspect of Martin-Löf type theory/logic: *equality*.
- For simplicity, consider first order logic:

$$\frac{\frac{}{[x : A] \Gamma \vdash x = x} \text{ (refl)} \quad \frac{\frac{A \text{ type}}{[a : A, b : A] a = b \text{ prop}} \quad \frac{[z : A, x : C] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : A, b : A, x : C] a = b, \Gamma(a, b, x) \vdash P(a, b, x)} \text{ (J)}}{[a : A, b : A, x : C] a = b, \Gamma(a, b, x) \vdash P(a, b, x)} \text{ (J)}$$

- Transitivity of equality:

$$\frac{\frac{[z : A, c : A] \quad z = c \vdash z = c}{[a : A, b : A, c : A] a = b, b = c \vdash a = c} \text{ (J)}}{[a : A, b : A, c : A] a = b, b = c \vdash a = c} \text{ (id)}$$

Type theory and symmetric equality

- A very interesting aspect of Martin-Löf type theory/logic: *equality*.
- For simplicity, consider first order logic:

$$\frac{\frac{\frac{}{A \text{ type}}}{[a : A, b : A] \ a = b \text{ prop}}}{[x : A] \ \Gamma \vdash x = x} \text{ (refl)} \quad \frac{[z : A, x : C] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : A, b : A, x : C] \ a = b, \ \Gamma(a, b, x) \vdash P(a, b, x)} \text{ (J)}$$

- Transitivity of equality:

$$\frac{\frac{[z : A, c : A] \quad z = c \vdash z = c}{[a : A, b : A, c : A] \ a = b, \ b = c \vdash a = c} \text{ (J)}}{} \text{ (id)}$$

- Equality in MLTT is inherently symmetric:

$$\frac{\frac{[z : A] \quad \top \vdash z = z}{[a : A, b : A] \ a = b \vdash b = a} \text{ (refl)}}{} \text{ (J)}$$

What can types be?

Martin-Löf type theory admits a sound interpretation in:

- Types as sets. [Martin-Löf 1971]
- Types as groupoids. [Hoffmann-Streicher 1999]
- Types as ∞ -groupoids. [Voevodsky 2013], [van der Berg-Garner 2010]
- Types as **sSets**, **Top**, cubical sets, ... [Awodey-Warren 2007], [CCHM 2007]
- Types as reflexive graphs for parametricity. [Atkey-Ghani-Johann 2007]

What can types be?

Martin-Löf type theory admits a sound interpretation in:

- Types as sets. [Martin-Löf 1971]
- Types as groupoids. [Hoffmann-Streicher 1999]
- Types as ∞ -groupoids. [Voevodsky 2013], [van der Berg-Garner 2010]
- Types as **sSets**, **Top**, cubical sets, ... [Awodey-Warren 2007], [CCHM 2007]
- Types as reflexive graphs for parametricity. [Atkey-Ghani-Johann 2007]

Intuition: types are groupoids, equalities are always-invertible morphisms.

What can types be?

Martin-Löf type theory admits a sound interpretation in:

- Types as sets. [Martin-Löf 1971]
- Types as groupoids. [Hoffmann-Streicher 1999]
- Types as ∞ -groupoids. [Voevodsky 2013], [van der Berg-Garner 2010]
- Types as **sSets**, **Top**, cubical sets, ... [Awodey-Warren 2007], [CCHM 2007]
- Types as reflexive graphs for parametricity. [Atkey-Ghani-Johann 2007]

Intuition: types are groupoids, equalities are always-invertible morphisms.

Some familiar structures are missing...

What can types be?

Martin-Löf type theory admits a sound interpretation in:

- Types as sets. [Martin-Löf 1971]
- Types as groupoids. [Hoffmann-Streicher 1999]
- Types as ∞ -groupoids. [Voevodsky 2013], [van der Berg-Garner 2010]
- Types as **sSets**, **Top**, cubical sets, ... [Awodey-Warren 2007], [CCHM 2007]
- Types as reflexive graphs for parametricity. [Atkey-Ghani-Johann 2007]

Intuition: types are groupoids, equalities are always-invertible morphisms.

Some familiar structures are missing... *what about categories and posets?!*

Motivation 1: Directed type theory

Martin-Löf type theory with refl/J is intrinsically about symmetric equality.

Directed type theory is the generalization to “directed equality”.

Motivation 1: Directed type theory

Martin-Löf type theory with refl/J is intrinsically about symmetric equality.

Directed type theory is the generalization to “directed equality”.

The interpretation of directed type theory with *(1-)categories*:

Types \rightsquigarrow Categories

Terms \rightsquigarrow Functors

Relations \rightsquigarrow Profunctors

Points of a type \rightsquigarrow Objects of a category

Equalities $e : a = b \rightsquigarrow$ Morphisms $e : \text{hom}(a, b)$

Equality types $=_A : A \times A \rightarrow \text{Type} \rightsquigarrow$ Hom types $\text{hom}_{\mathbb{C}} : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$

\rightarrow Now types have a *polarity*, \mathbb{C} and \mathbb{C}^{op} , i.e., the opposite category.

Motivation 1: Directed type theory

Martin-Löf type theory with refl/J is intrinsically about symmetric equality.

Directed type theory is the generalization to “directed equality”.

The interpretation of directed type theory with $(1-)$ categories:

Types \rightsquigarrow Categories

Terms \rightsquigarrow Functors

Relations \rightsquigarrow Profunctors

Points of a type \rightsquigarrow Objects of a category

Equalities $e : a = b \rightsquigarrow$ Morphisms $e : \text{hom}(a, b)$

Equality types $=_A : A \times A \rightarrow \text{Type} \rightsquigarrow$ Hom types $\text{hom}_{\mathbb{C}} : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$

→ Now types have a *polarity*, \mathbb{C} and \mathbb{C}^{op} , i.e., the opposite category.

→ Now equalities $e : \text{hom}(a, b)$ have *directionality*: rewrites, trans., processes.

Motivation 1: Directed type theory

Martin-Löf type theory with refl/J is intrinsically about symmetric equality.

Directed type theory is the generalization to “directed equality”.

The interpretation of directed type theory with $(1-)$ categories:

Types \rightsquigarrow Categories

Terms \rightsquigarrow Functors

Relations \rightsquigarrow Profunctors

Points of a type \rightsquigarrow Objects of a category

Equalities $e : a = b \rightsquigarrow$ Morphisms $e : \text{hom}(a, b)$

Equality types $=_A : A \times A \rightarrow \text{Type} \rightsquigarrow$ Hom types $\text{hom}_{\mathbb{C}} : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$

→ Now types have a *polarity*, \mathbb{C} and \mathbb{C}^{op} , i.e., the opposite category.

→ Now equalities $e : \text{hom}(a, b)$ have *directionality*: rewrites, trans., processes.

We want to find which syntactic restriction of MLTT
allow for types can be interpreted as categories.

Current approaches to directed type theory

- Semantically, refl should be $\text{id}_c \in \text{hom}_{\mathbb{C}}(c, c)$ for $c : \mathbb{C}$.

Current approaches to directed type theory

- Semantically, refl should be $\text{id}_c \in \text{hom}_{\mathbb{C}}(c, c)$ for $c : \mathbb{C}$.
- Transitivity of directed equality \rightsquigarrow composition of morphisms in \mathbb{C} .

Current approaches to directed type theory

- Semantically, refl should be $\text{id}_c \in \text{hom}_{\mathbb{C}}(c, c)$ for $c : \mathbb{C}$.
- Transitivity of directed equality \rightsquigarrow composition of morphisms in \mathbb{C} .

$$\frac{\frac{[z : \mathbb{C}^{\text{op}}, c : \mathbb{C}] \quad \text{hom}(z, c) \vdash \text{hom}(z, c)}{\text{(id)}}}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}, c : \mathbb{C}] \text{ hom}(a, b), \text{ hom}(b, c) \vdash \text{hom}(a, c)} \text{(J)}$$

Current approaches to directed type theory

- Semantically, refl should be $\text{id}_c \in \text{hom}_{\mathbb{C}}(c, c)$ for $c : \mathbb{C}$.
- Transitivity of directed equality \rightsquigarrow composition of morphisms in \mathbb{C} .

$$\frac{[z : \mathbb{C}^{\text{op}}, c : \mathbb{C}] \quad \text{hom}(z, c) \vdash \text{hom}(z, c)}{\text{refl}_c : \text{hom}_{\mathbb{C}}(c, c)} \quad (\text{id})$$
$$\frac{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}, c : \mathbb{C}] \quad \text{hom}(a, b), \text{hom}(b, c) \vdash \text{hom}(a, c)}{\text{refl}_a \circ \text{refl}_b : \text{hom}_{\mathbb{C}}(a, c)} \quad (J)$$

- However, directed type theory is not so straightforward:

$$\frac{a : \mathbb{C}}{\text{refl}_a : \text{hom}_{\mathbb{C}}(a, a)}$$

- *Problem:* rule is not functorial w.r.t. variance of $\text{hom}_{\mathbb{C}} : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, since $a : \mathbb{C}$ appears both contravariantly and covariantly.

Current approaches to directed type theory

- Semantically, refl should be $\text{id}_c \in \text{hom}_{\mathbb{C}}(c, c)$ for $c : \mathbb{C}$.
- Transitivity of directed equality \rightsquigarrow composition of morphisms in \mathbb{C} .

$$\frac{\frac{[z : \mathbb{C}^{\text{op}}, c : \mathbb{C}]}{\text{hom}(z, c) \vdash \text{hom}(z, c)} \quad (\text{id})}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}, c : \mathbb{C}] \text{hom}(a, b), \text{hom}(\bar{b}, c) \vdash \text{hom}(a, c)} \quad (J)$$

- However, directed type theory is not so straightforward:

$$\frac{a : \mathbb{C}}{\text{refl}_a \dots ? : \text{hom}_{\mathbb{C}}(a, a)} \rightsquigarrow \frac{a : \mathbb{C}^{\text{core}}}{\text{refl}_a : \text{hom}(\text{i}^{\text{op}}(a), \text{i}(a))} \quad [\text{North 2018}]$$

- *Problem:* rule is not functorial w.r.t. variance of $\text{hom}_{\mathbb{C}} : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, since $a : \mathbb{C}$ appears both contravariantly and covariantly.
- Current approach to DTT in **Cat**: [North 2018], [Neumann 2024]
 \rightarrow Use the maximal subgroupoid \mathbb{C}^{core} to collapse the two variances,

Current approaches to directed type theory

- Semantically, refl should be $\text{id}_c \in \text{hom}_{\mathbb{C}}(c, c)$ for $c : \mathbb{C}$.
- Transitivity of directed equality \rightsquigarrow composition of morphisms in \mathbb{C} .

$$\frac{\frac{[z : \mathbb{C}^{\text{op}}, c : \mathbb{C}]}{\text{hom}(z, c) \vdash \text{hom}(z, c)} \quad (\text{id})}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}, c : \mathbb{C}] \text{hom}(a, b), \text{hom}(\bar{b}, c) \vdash \text{hom}(a, c)} \quad (J)$$

- However, directed type theory is not so straightforward:

$$\frac{a : \mathbb{C}}{\text{refl}_a \dots ? : \text{hom}_{\mathbb{C}}(a, a)} \rightsquigarrow \frac{a : \mathbb{C}^{\text{core}}}{\text{refl}_a : \text{hom}(i^{\text{op}}(a), i(a))} \quad [\text{North 2018}]$$

- Problem:* rule is not functorial w.r.t. variance of $\text{hom}_{\mathbb{C}} : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, since $a : \mathbb{C}$ appears both contravariantly and covariantly.
- Current approach to DTT in **Cat**: [North 2018], [Neumann 2024]
 \rightarrow Use the maximal subgroupoid \mathbb{C}^{core} to collapse the two variances, since $(\mathbb{C}^{\text{core}})^{\text{op}} \cong \mathbb{C}^{\text{core}}$, then use $i : \mathbb{C}^{\text{core}} \rightarrow \mathbb{C}$ and $i^{\text{op}} : \mathbb{C}^{\text{core}} \rightarrow \mathbb{C}^{\text{op}}$.

Current approaches to directed type theory

- Semantically, refl should be $\text{id}_c \in \text{hom}_{\mathbb{C}}(c, c)$ for $c : \mathbb{C}$.
- Transitivity of directed equality \rightsquigarrow composition of morphisms in \mathbb{C} .

$$\frac{\frac{[z : \mathbb{C}^{\text{op}}, c : \mathbb{C}]}{\text{hom}(z, c) \vdash \text{hom}(z, c)} \quad (\text{id})}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}, c : \mathbb{C}] \text{hom}(a, b), \text{hom}(\bar{b}, c) \vdash \text{hom}(a, c)} \quad (J)$$

- However, directed type theory is not so straightforward:

$$\frac{a : \mathbb{C}}{\text{refl}_a \dots ? : \text{hom}_{\mathbb{C}}(a, a)} \rightsquigarrow \frac{a : \mathbb{C}^{\text{core}}}{\text{refl}_a : \text{hom}(i^{\text{op}}(a), i(a))} \quad [\text{North 2018}]$$

- Problem:* rule is not functorial w.r.t. variance of $\text{hom}_{\mathbb{C}} : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, since $a : \mathbb{C}$ appears both contravariantly and covariantly.
- Current approach to DTT in **Cat**: [North 2018], [Neumann 2024]
 \rightarrow Use the maximal subgroupoid \mathbb{C}^{core} to collapse the two variances, since $(\mathbb{C}^{\text{core}})^{\text{op}} \cong \mathbb{C}^{\text{core}}$, then use $i : \mathbb{C}^{\text{core}} \rightarrow \mathbb{C}$ and $i^{\text{op}} : \mathbb{C}^{\text{core}} \rightarrow \mathbb{C}^{\text{op}}$.
- Then a J -like rule is validated, but *again using groupoidal structure*.

Contribution 1 – Towards dinatural directed type theory

Is there a way to interpret a directed type theory in 1-categories without having to collapse categories back to the "undirected" groupoidal case?

Contribution 1 – Towards dinatural directed type theory

Is there a way to interpret a directed type theory in 1-categories without having to collapse categories back to the "undirected" groupoidal case?

Our approach: yes, by validating rules with dinatural transformations.

Contribution 1 – Towards dinatural directed type theory

Is there a way to interpret a directed type theory in 1-categories without having to collapse categories back to the "undirected" groupoidal case?

Our approach: yes, by validating rules with dinatural transformations.

- *Intuition:* dinatural transformations allow for the same variable to appear both covariantly and contravariantly \rightarrow solving the issue with refl.

Contribution 1 – Towards dinatural directed type theory

Is there a way to interpret a directed type theory in 1-categories without having to collapse categories back to the "undirected" groupoidal case?

Our approach: yes, by validating rules with dinatural transformations.

- *Intuition:* dinatural transformations allow for the same variable to appear both covariantly and contravariantly \rightarrow solving the issue with refl.
- We will see how dinaturality validates both
 \rightarrow a refl rule (directed equality introduction),

Contribution 1 – Towards dinatural directed type theory

Is there a way to interpret a directed type theory in 1-categories without having to collapse categories back to the "undirected" groupoidal case?

Our approach: yes, by validating rules with dinatural transformations.

- *Intuition:* dinatural transformations allow for the same variable to appear both covariantly and contravariantly \rightarrow solving the issue with refl.
- We will see how dinaturality validates both
 - \rightarrow a refl rule (directed equality introduction),
 - \rightarrow a J rule (directed equality elimination).

Contribution 1 – Towards dinatural directed type theory

Is there a way to interpret a directed type theory in 1-categories without having to collapse categories back to the "undirected" groupoidal case?

Our approach: yes, by validating rules with dinatural transformations.

- *Intuition:* dinatural transformations allow for the same variable to appear both covariantly and contravariantly \rightarrow solving the issue with refl.
- We will see how dinaturality validates both
 - \rightarrow a refl rule (directed equality introduction),
 - \rightarrow a J rule (directed equality elimination).
- The directed J rule is extremely similar to the standard MLTT J rule,

Contribution 1 – Towards dinatural directed type theory

Is there a way to interpret a directed type theory in 1-categories without having to collapse categories back to the "undirected" groupoidal case?

Our approach: yes, by validating rules with dinatural transformations.

- *Intuition:* dinatural transformations allow for the same variable to appear both covariantly and contravariantly \rightarrow solving the issue with refl.
- We will see how dinaturality validates both
 - \rightarrow a refl rule (directed equality introduction),
 - \rightarrow a J rule (directed equality elimination).
- The directed J rule is extremely similar to the standard MLTT J rule, but with a syntactic restriction which does **not** allow for symmetry.
- This comes at the cost of compositionality:
we have a cut rule only if variables appear *naturally* in an entailment.

Contribution 1 – Directed type theory

Comparison between symmetric equality and our directed equality rule:

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] \quad a = b, \Gamma(a, b, x) \vdash P(a, b, x)}}{(\text{eq})}$$

Contribution 1 – Directed type theory

Comparison between symmetric equality and our directed equality rule:

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] \quad a = b, \Gamma(a, b, x) \vdash P(a, b, x)}}{(\text{eq})}$$

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{z}, z, \bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad \text{hom}(a, b), \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash P(a, b, \bar{x}, x)}}{(\text{hom})}$$

Contribution 1 – Directed type theory

Comparison between symmetric equality and our directed equality rule:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] \quad a = b, \Gamma(a, b, x) \vdash P(a, b, x)} \text{ (eq)}$$

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{z}, z, \bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad \text{hom}(a, b), \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash P(a, b, \bar{x}, x)} \text{ (hom)}$$

- **Idea:** now variables have variance, *positive* ($a : \mathbb{A}$) or *negative* ($\bar{a} : \mathbb{A}^{\text{op}}$).
- **Restriction:** a, b must appear only *positively* in P , and *negatively* in Γ .

Contribution 1 – Directed type theory

Comparison between symmetric equality and our directed equality rule:

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] \quad a = b, \Gamma(a, b, x) \vdash P(a, b, x)}}{(\text{eq})}$$

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{z}, z, \bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad \text{hom}(a, b), \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash P(a, b, \bar{x}, x)}}{(\text{hom})}$$

- **Idea:** now variables have variance, *positive* ($a : \mathbb{A}$) or *negative* ($\bar{a} : \mathbb{A}^{\text{op}}$).
- **Restriction:** a, b must appear only *positively* in P , and *negatively* in Γ .
- (Again, dinaturality allows z to appear twice in $P(z, z, x)$.)

Contribution 1 – Directed type theory

Comparison between symmetric equality and our directed equality rule:

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] \quad a = b, \Gamma(a, b, x) \vdash P(a, b, x)}}{(\text{eq})}$$

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{z}, z, \bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad \text{hom}(a, b), \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash P(a, b, \bar{x}, x)}}{(\text{hom})}$$

- **Idea:** now variables have variance, *positive* ($a : \mathbb{A}$) or *negative* ($\bar{a} : \mathbb{A}^{\text{op}}$).
 - **Restriction:** a, b must appear only *positively* in P , and *negatively* in Γ .
 - (Again, dinaturality allows z to appear twice in $P(z, z, x)$.)
- Same terms provable in MLTT about equality:
transitivity, transport, congruence, *but not symmetry!*

Contribution 1 – Directed type theory

Comparison between symmetric equality and our directed equality rule:

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] \quad a = b, \Gamma(a, b, x) \vdash P(a, b, x)}}{(\text{eq})}$$

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{z}, z, \bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad \text{hom}(a, b), \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash P(a, b, \bar{x}, x)}}{(\text{hom})}$$

- **Idea:** now variables have variance, *positive* ($a : \mathbb{A}$) or *negative* ($\bar{a} : \mathbb{A}^{\text{op}}$).
- **Restriction:** a, b must appear only *positively* in P , and *negatively* in Γ .
- (Again, dinaturality allows z to appear twice in $P(z, z, x)$.)
- ▶ Same terms provable in MLTT about equality:
transitivity, transport, congruence, *but not symmetry!*
- ▶ Proving properties about them follows precisely the steps in MLTT.
(e.g., we have the same computation rules.)

Contribution 2 – (Co)end calculus

Another important open question:

Another important open question:

What should the quantifiers of directed type theory be?

Another important open question:

What should the quantifiers of directed type theory be?

- Dinaturality again gives us a hint: *ends and coends*.

Contribution 2 – (Co)end calculus

Another important open question:

What should the quantifiers of directed type theory be?

- Dinaturality again gives us a hint: *ends and coends*.
- We present a series of “logical rules”, inspired by the doctrinal approach, to view (co)ends as the *directed quantifiers* of DTT.

Contribution 2 – (Co)end calculus

Another important open question:

What should the quantifiers of directed type theory be?

- Dinaturality again gives us a hint: *ends and coends*.
- We present a series of “logical rules”, inspired by the doctrinal approach, to view (co)ends as the *directed quantifiers* of DTT.
- ▶ We then use the rules to give *simple proofs* of theorems in category theory, using dinaturality and viewing \hom as directed equality.

Contribution 2 – (Co)end calculus

Another important open question:

What should the quantifiers of directed type theory be?

- Dinaturality again gives us a hint: *ends and coends*.
- We present a series of “logical rules”, inspired by the doctrinal approach, to view (co)ends as the *directed quantifiers* of DTT.
- ▶ We then use the rules to give *simple proofs* of theorems in category theory, using dinaturality and viewing \hom as directed equality.
- **Claim:** *(co)end calculus is the first-order instantiation of a directed type theory with quantifiers, with semantics in 1-categories.*

Motivation 2: Profunctors

Another story, now in CT:

Rel \rightsquigarrow **Prof**

Sets \rightsquigarrow Categories

Relations $A \times B \rightarrow \{\top, \perp\} \rightsquigarrow$ Profunctors $\mathbb{C}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set}$

Existential quantifiers $\exists x \rightsquigarrow$ Coends \int^x

Composition: $\exists b \in B. R(a, b) \wedge Q(b, c) \rightsquigarrow \int^{b:\mathbb{B}} R(a, b) \times Q(b, c)$

Conjunction of truth values \rightsquigarrow Cartesian products of **Sets**

Identity relation: $=_{\mathbb{C}} \rightsquigarrow$ Identity profunctor: $\text{hom}_{\mathbb{C}}$

Motivation 2: Profunctors

Another story, now in CT:

Rel \rightsquigarrow **Prof**

Sets \rightsquigarrow Categories

Relations $A \times B \rightarrow \{\top, \perp\} \rightsquigarrow$ Profunctors $\mathbb{C}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set}$

Existential quantifiers $\exists x \rightsquigarrow$ Coends \int^x

Composition: $\exists b \in B. R(a, b) \wedge Q(b, c) \rightsquigarrow \int^{b:\mathbb{B}} R(a, b) \times Q(b, c)$

Conjunction of truth values \rightsquigarrow Cartesian products of **Sets**

Identity relation: $=_{\mathbb{C}} \rightsquigarrow$ Identity profunctor: $\text{hom}_{\mathbb{C}}$

Why do unitality and associativity hold?

Because of rules for equality in FOL. \rightsquigarrow Because of the coYoneda lemma.

Because of rules for \exists and \wedge in FOL. \rightsquigarrow Because of properties of coends.

Motivation 2: Profunctors

Another story, now in CT:

Rel \rightsquigarrow **Prof**

Sets \rightsquigarrow Categories

Relations $A \times B \rightarrow \{\top, \perp\} \rightsquigarrow$ Profunctors $\mathbb{C}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set}$

Existential quantifiers $\exists x \rightsquigarrow$ Coends \int^x

Composition: $\exists b \in B. R(a, b) \wedge Q(b, c) \rightsquigarrow \int^{b:\mathbb{B}} R(a, b) \times Q(b, c)$

Conjunction of truth values \rightsquigarrow Cartesian products of **Sets**

Identity relation: $=_{\mathbb{C}} \rightsquigarrow$ Identity profunctor: $\text{hom}_{\mathbb{C}}$

Why do unitality and associativity hold?

Because of rules for equality in FOL. \rightsquigarrow Because of the coYoneda lemma.

Because of rules for \exists and \wedge in FOL. \rightsquigarrow Because of properties of coends.

- We use FOL in **Rel**: there is no *directed logic* for **Prof**.

Motivation 2: Profunctors

Another story, now in CT:

	Rel \rightsquigarrow Prof
	Sets \rightsquigarrow Categories
Relations $A \times B \rightarrow \{\top, \perp\}$	Profunctors $\mathbb{C}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set}$
Existential quantifiers $\exists x$	Coends \int^x
Composition: $\exists b \in B. R(a, b) \wedge Q(b, c)$	$\int^{b:\mathbb{B}} R(a, b) \times Q(b, c)$
Conjunction of truth values	Cartesian products of Sets
Identity relation: $=_{\mathbb{C}}$	Identity profunctor: $\text{hom}_{\mathbb{C}}$

Why do unitality and associativity hold?

Because of rules for equality in FOL. \rightsquigarrow Because of the coYoneda lemma.

Because of rules for \exists and \wedge in FOL. \rightsquigarrow Because of properties of coends.

- We use FOL in **Rel**: there is no *directed logic* for **Prof**.
- A proper treatment of coends *as quantifiers*.

Motivation 2: Profunctors

Another story, now in CT:

Rel	\rightsquigarrow	Prof
Sets	\rightsquigarrow	Categories
Relations $A \times B \rightarrow \{\top, \perp\}$	\rightsquigarrow	Profunctors $\mathbb{C}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set}$
Existential quantifiers $\exists x$	\rightsquigarrow	Coends \int^x
Composition: $\exists b \in B. R(a, b) \wedge Q(b, c)$	\rightsquigarrow	$\int^{b:\mathbb{B}} R(a, b) \times Q(b, c)$
Conjunction of truth values	\rightsquigarrow	Cartesian products of Sets
Identity relation: $=_{\mathbb{C}}$	\rightsquigarrow	Identity profunctor: $\text{hom}_{\mathbb{C}}$

Why do unitality and associativity hold?

Because of rules for equality in FOL. \rightsquigarrow Because of the coYoneda lemma.

Because of rules for \exists and \wedge in FOL. \rightsquigarrow Because of properties of coends.

- We use FOL in **Rel**: there is no *directed logic* for **Prof**.
- A proper treatment of coends *as quantifiers*.
- A syntactic treatment of directed logic (with directed equality and its rules, and quantifiers), e.g., using a doctrinal approach.

Motivation 3: Lawvere equality for categories

*Can we interpret logic in **Cat**, viewed as a category of contexts?*

Motivation 3: Lawvere equality for categories

*Can we interpret logic in **Cat**, viewed as a category of contexts?*

From type theory: why **Cat** is *not* a model for Martin-Löf type theory:

- **Cat** is *not* locally cartesian closed,
- **Cat** is *not* even regular (but it is 2-regular [Street CBS 1982])

Motivation 3: Lawvere equality for categories

*Can we interpret logic in **Cat**, viewed as a category of contexts?*

From type theory: why **Cat** is *not* a model for Martin-Löf type theory:

- **Cat** is *not* locally cartesian closed,
- **Cat** is *not* even regular (but it is 2-regular [Street CBS 1982])

From the doctrinal perspective: **Set** plays the role of *generalized truth values* in **Prof**, and presheaves are “generalized logical predicates”, but:

Motivation 3: Lawvere equality for categories

*Can we interpret logic in **Cat**, viewed as a category of contexts?*

From type theory: why **Cat** is *not* a model for Martin-Löf type theory:

- **Cat** is *not* locally cartesian closed,
- **Cat** is *not* even regular (but it is 2-regular [Street CBS 1982])

From the doctrinal perspective: **Set** plays the role of *generalized truth values* in **Prof**, and presheaves are “generalized logical predicates”, but:

- Equality in the presheaf hyperdoctrine

$$\mathcal{P}\text{sh}(\mathbb{C}) := [\mathbb{C}^{\text{op}}, \mathbf{Set}] : \mathbf{Cat}^{\text{op}} \rightarrow \mathbf{CAT}$$

does **not** validate Frobenius and Beck-Chevalley: [Melliès-Zeilberger 2016]

Motivation 3: Lawvere equality for categories

*Can we interpret logic in **Cat**, viewed as a category of contexts?*

From type theory: why **Cat** is *not* a model for Martin-Löf type theory:

- **Cat** is *not* locally cartesian closed,
- **Cat** is *not* even regular (but it is 2-regular [Street CBS 1982])

From the doctrinal perspective: **Set** plays the role of *generalized truth values* in **Prof**, and presheaves are “generalized logical predicates”, but:

- Equality in the presheaf hyperdoctrine

$$\mathcal{Psh}(\mathbb{C}) := [\mathbb{C}^{\text{op}}, \mathbf{Set}] : \mathbf{Cat}^{\text{op}} \rightarrow \mathbf{CAT}$$

does **not** validate Frobenius and Beck-Chevalley: [Melliès-Zeilberger 2016]

$$\text{Eq}_{\mathbb{C}} := \exists_{\Delta_{\mathbb{C}}}(\top_{\mathbb{C}}) \in \mathcal{Psh}(\mathbb{C} \times \mathbb{C})$$

$$\text{Eq}_{\mathbb{C}} = (a, b) \mapsto \int^{x \in \mathbb{C}} \text{hom}_{\mathbb{C}}(a, x) \times \text{hom}_{\mathbb{C}}(b, x)$$

i.e., any two objects forming a cospan for some x should be equated.

Note: These conditions also fail in groupoids.

Motivation 3: Lawvere on hom as equality

Lawvere commenting on the failure of Frobenius/Beck-Chevalley for $\mathcal{P}sh$:

[Lawvere 1970, Equality in Hyperdoctrines]

Motivation 3: Lawvere on \mathbf{hom} as equality

Lawvere commenting on the failure of Frobenius/Beck-Chevalley for $\mathcal{P}sh$:

- [...] *This should not be taken as indicative of a lack of vitality of $\mathcal{P}sh$ as hyperdoctrine, or even of a lack of a satisfactory theory of equality for it.*

[Lawvere 1970, Equality in Hyperdoctrines]

Motivation 3: Lawvere on \mathbf{hom} as equality

Lawvere commenting on the failure of Frobenius/Beck-Chevalley for $\mathcal{P}sh$:

- [...] *This should not be taken as indicative of a lack of vitality of $\mathcal{P}sh$ as hyperdoctrine, or even of a lack of a satisfactory theory of equality for it.*
- *Rather, it indicates that we have probably been too naive in defining equality in a manner too closely suggested by the classical conception.*

[Lawvere 1970, Equality in Hyperdoctrines]

Motivation 3: Lawvere on \mathbf{hom} as equality

Lawvere commenting on the failure of Frobenius/Beck-Chevalley for $\mathcal{P}sh$:

- [...] *This should not be taken as indicative of a lack of vitality of $\mathcal{P}sh$ as hyperdoctrine, or even of a lack of a satisfactory theory of equality for it.*
- *Rather, it indicates that we have probably been too naive in defining equality in a manner too closely suggested by the classical conception.*
- *Equality should be the “graph” of the identity term. But present categorical conceptions indicate that the graph of a functor $F : \mathbb{C} \rightarrow \mathbb{D}$ should be [...] a binary attribute of mixed variance $\mathcal{P}sh(\mathbb{C}^{\text{op}} \times \mathbb{D})$.*
- *Thus in particular “equality” should be the functor $\mathbf{hom}_{\mathbb{C}}$ [...].*

[Lawvere 1970, Equality in Hyperdoctrines]

Motivation 3: Lawvere on hom as equality

Lawvere commenting on the failure of Frobenius/Beck-Chevalley for $\mathcal{P}sh$:

- [...] *This should not be taken as indicative of a lack of vitality of $\mathcal{P}sh$ as hyperdoctrine, or even of a lack of a satisfactory theory of equality for it.*
- *Rather, it indicates that we have probably been too naive in defining equality in a manner too closely suggested by the classical conception.*
- *Equality should be the “graph” of the identity term. But present categorical conceptions indicate that the graph of a functor $F : \mathbb{C} \rightarrow \mathbb{D}$ should be [...] a binary attribute of mixed variance $\mathcal{P}sh(\mathbb{C}^{\text{op}} \times \mathbb{D})$.*
- *Thus in particular “equality” should be the functor $\text{hom}_{\mathbb{C}}$ [...].*
- *The term which would take the place of $\Delta_{\mathbb{C}}$ in such a more enlightened theory of equality would be the forgetful functor $\text{Tw}(\mathbb{C}) \rightarrow \mathbb{C}^{\text{op}} \times \mathbb{C}$. [...]*
- *Of course to abstract from this example would require at least the addition of a functor $T \xrightarrow{\text{op}} T$ to the structure of a [doctrine].*

[Lawvere 1970, Equality in Hyperdoctrines]

Background on dinaturality and (co)ends

Background: Dinatural transformations

- A **difunctor** from \mathbb{C} to \mathbb{D} is a functor $F, G : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$.

Background: Dinatural transformations

- A **difunctor** from \mathbb{C} to \mathbb{D} is a functor $F, G : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$.
- A **dipresheaf** on \mathbb{C} is an endoprofunctor $P : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$.

Background: Dinatural transformations

- A **difunctor** from \mathbb{C} to \mathbb{D} is a functor $F, G : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$.
- A **dipresheaf** on \mathbb{C} is an endoprofunctor $P : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$.
- Given difunctors $F, G : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$, a **dinatural transformation** $\alpha : F \rightrightarrows G$ is a family of morphisms of \mathbb{D} for each $x : \mathbb{C}$,

$$\alpha_x : F(x, x) \longrightarrow G(x, x)$$

Background: Dinatural transformations

- A **difunctor** from \mathbb{C} to \mathbb{D} is a functor $F, G : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$.
- A **dipresheaf** on \mathbb{C} is an endoprofunctor $P : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$.
- Given difunctors $F, G : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$, a **dinatural transformation** $\alpha : F \rightrightarrows G$ is a family of morphisms of \mathbb{D} for each $x : \mathbb{C}$,

$$\alpha_x : F(x, x) \longrightarrow G(x, x)$$

such that for every $a, b : \mathbb{C}$ and $f : a \rightarrow b$ this hexagon commutes:

$$\begin{array}{ccccc} & & F(b, b) & \xrightarrow{\alpha_b} & G(b, b) \\ & \nearrow^{F(\text{id}, f)} & & & \searrow^{G(f, \text{id})} \\ F(b, a) & & & & G(a, b) \\ & \searrow_{F(f, \text{id})} & & & \nearrow_{G(\text{id}, f)} \\ & & F(a, a) & \xrightarrow{\alpha_a} & G(a, a) \end{array}$$

Background: Dinatural transformations

- A **difunctor** from \mathbb{C} to \mathbb{D} is a functor $F, G : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$.
- A **dipresheaf** on \mathbb{C} is an endoprofunctor $P : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$.
- Given difunctors $F, G : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$, a **dinatural transformation** $\alpha : F \rightrightarrows G$ is a family of morphisms of \mathbb{D} for each $x : \mathbb{C}$,

$$\alpha_x : F(x, x) \longrightarrow G(x, x)$$

such that for every $a, b : \mathbb{C}$ and $f : a \rightarrow b$ this hexagon commutes:


$$\begin{array}{ccccc} & & F(b, b) & \xrightarrow{\alpha_b} & G(b, b) \\ & \nearrow^{F(\text{id}, f)} & & & \searrow^{G(f, \text{id})} \\ F(b, a) & & & & G(a, b) \\ & \searrow_{F(f, \text{id})} & & & \nearrow_{G(\text{id}, f)} \\ & & F(a, a) & \xrightarrow{\alpha_a} & G(a, a) \end{array}$$

Theorem (Naturals are dinaturals [Dubuc and Street 1969])

A dinatural between functors which do not depend on \mathbb{C}^{op} is just a natural.


Background: Dinaturals don't always compose

- Dinaturals don't always compose: $\alpha : F \rightrightarrows G$, $\beta : G \rightrightarrows H$,




Background: Dinaturals don't always compose

- Dinaturals don't always compose: $\alpha : F \rightrightarrows G$, $\beta : G \rightrightarrows H$,



- They always compose with naturals: $\alpha : F \rightrightarrows G$, $\beta : G \rightarrow H$,



(Co)end calculus in one slide

- (Co)ends give elegant characterizations of theorems of category theory.

(Co)end calculus in one slide

- (Co)ends give elegant characterizations of theorems of category theory.
- Natural transformations as ends:

$$\mathrm{Nat}(F, G) \cong \int_{x:\mathbb{C}} \mathrm{hom}_{\mathbb{D}}(F(\overline{x}), G(x))$$

(Co)end calculus in one slide

- (Co)ends give elegant characterizations of theorems of category theory.
- Natural transformations as ends:

$$\mathrm{Nat}(F, G) \cong \int_{x:\mathbb{C}} \mathrm{hom}_{\mathbb{D}}(F(\overline{x}), G(x))$$

- Quantifier exchange

$$(Q : (\mathbb{C}^{\mathrm{op}} \times \mathbb{C}) \\ \times (\mathbb{D}^{\mathrm{op}} \times \mathbb{D}) \rightarrow \mathbf{Set}):$$

$$\begin{aligned} & \int_{x:\mathbb{C}} \int_{y:\mathbb{D}} Q(\overline{x}, x, \overline{y}, y) \\ & \cong \int_{y:\mathbb{D}} \int_{x:\mathbb{C}} Q(\overline{x}, x, \overline{y}, y) \\ & \cong \int_{(x,y):\mathbb{C} \times \mathbb{D}} Q(\overline{x}, x, \overline{y}, y) \end{aligned}$$

(Co)end calculus in one slide

- (Co)ends give elegant characterizations of theorems of category theory.
- Natural transformations as ends:

$$\mathrm{Nat}(F, G) \cong \int_{x:\mathbb{C}} \mathrm{hom}_{\mathbb{D}}(F(\bar{x}), G(x))$$

- Quantifier exchange

$$(Q : (\mathbb{C}^{\mathrm{op}} \times \mathbb{C}) \\ \times (\mathbb{D}^{\mathrm{op}} \times \mathbb{D}) \rightarrow \mathbf{Set}):$$

$$\begin{aligned} & \int_{x:\mathbb{C}} \int_{y:\mathbb{D}} Q(\bar{x}, x, \bar{y}, y) \\ & \cong \int_{y:\mathbb{D}} \int_{x:\mathbb{C}} Q(\bar{x}, x, \bar{y}, y) \\ & \cong \int_{(x,y):\mathbb{C} \times \mathbb{D}} Q(\bar{x}, x, \bar{y}, y) \end{aligned}$$

- (Co)Yoneda lemma: $(P : \mathbb{C} \rightarrow \mathbf{Set})$

$$P(a) \cong \int_{x:\mathbb{C}} \mathrm{hom}_{\mathbb{C}}(a, \bar{x}) \Rightarrow P(x)$$

$$P(a) \cong \int^{x:\mathbb{C}} \mathrm{hom}_{\mathbb{C}}(\bar{x}, a) \times P(x)$$

(Co)end calculus in one slide

- (Co)ends give elegant characterizations of theorems of category theory.
- Natural transformations as ends:

$$\mathrm{Nat}(F, G) \cong \int_{x:\mathbb{C}} \mathrm{hom}_{\mathbb{D}}(F(\bar{x}), G(x))$$

- Quantifier exchange

$$(Q : (\mathbb{C}^{\mathrm{op}} \times \mathbb{C}) \times (\mathbb{D}^{\mathrm{op}} \times \mathbb{D}) \rightarrow \mathbf{Set}):$$

$$\int_{x:\mathbb{C}} \int_{y:\mathbb{D}} Q(\bar{x}, x, \bar{y}, y)$$

$$\cong \int_{y:\mathbb{D}} \int_{x:\mathbb{C}} Q(\bar{x}, x, \bar{y}, y)$$

$$\cong \int_{(x,y):\mathbb{C} \times \mathbb{D}} Q(\bar{x}, x, \bar{y}, y)$$

- (Co)Yoneda lemma: $(P : \mathbb{C} \rightarrow \mathbf{Set})$

$$P(a) \cong \int_{x:\mathbb{C}} \mathrm{hom}_{\mathbb{C}}(a, \bar{x}) \Rightarrow P(x)$$

$$P(a) \cong \int^{x:\mathbb{C}} \mathrm{hom}_{\mathbb{C}}(\bar{x}, a) \times P(x)$$

- Kan extensions: $(F : \mathbb{C} \rightarrow \mathbb{D})$

$$(\mathrm{Ran}_F P)(a) \cong \int_{x:\mathbb{C}} \mathrm{hom}_{\mathbb{D}}(a, F(\bar{x})) \Rightarrow P(x)$$

$$(\mathrm{Lan}_F P)(a) \cong \int^{x:\mathbb{C}} \mathrm{hom}_{\mathbb{D}}(F(\bar{x}), a) \times P(x)$$

Logical interpretation of (co)end calculus

- (Co)Yoneda lemma:

$$\begin{aligned} P(a) &\cong \int_{x:\mathbb{C}} \text{hom}_{\mathbb{C}}(a, \bar{x}) \Rightarrow P(x) \\ P(a) &\Leftrightarrow \forall (x:\mathbb{C}). \quad a =_{\mathbb{C}} x \Rightarrow P(x) \end{aligned}$$

$$\begin{aligned} P(a) &\cong \int^{x:\mathbb{C}} \text{hom}_{\mathbb{C}}(\bar{x}, a) \times P(x) \\ P(a) &\Leftrightarrow \exists (x:\mathbb{C}). \quad x =_{\mathbb{C}} a \wedge P(x) \end{aligned}$$

Logical interpretation of (co)end calculus

- (Co)Yoneda lemma:

$$\begin{aligned} P(a) &\cong \int_{x:\mathbb{C}} \text{hom}_{\mathbb{C}}(a, \bar{x}) \Rightarrow P(x) \\ P(a) &\Leftrightarrow \forall (x : \mathbb{C}). \quad a =_{\mathbb{C}} x \Rightarrow P(x) \end{aligned}$$

$$\begin{aligned} P(a) &\cong \int^{x:\mathbb{C}} \text{hom}_{\mathbb{C}}(\bar{x}, a) \times P(x) \\ P(a) &\Leftrightarrow \exists (x : \mathbb{C}). \quad x =_{\mathbb{C}} a \wedge P(x) \end{aligned}$$

- (Pointwise) right/left Kan extensions using ends/coends:

$$\begin{aligned} (\text{Ran}_F P)(a) &\cong \int_{x:\mathbb{C}} \text{hom}_{\mathbb{D}}(a, F(x)) \Rightarrow P(x) \\ (\forall_f(P))(y) &\Leftrightarrow \forall (x : C). \quad y =_D f(x) \Rightarrow P(x) \end{aligned}$$

$$\begin{aligned} (\text{Lan}_F P)(a) &\cong \int^{x:\mathbb{C}} \text{hom}_{\mathbb{D}}(F(x), a) \times P(x) \\ (\exists_f(P))(y) &\Leftrightarrow \exists (x : C). \quad f(x) =_D y \wedge P(x) \end{aligned}$$

Motivation 4: Computing adjoints to reindexings

- Consider a doctrine $\mathcal{P} : \mathbb{C}^{\text{op}} \rightarrow \mathbf{Pos}$ with $\wedge, \Rightarrow, \forall_{A[R]}, \exists_{A[R]}, \text{Eq}_A$.

Motivation 4: Computing adjoints to reindexings

- Consider a doctrine $\mathcal{P} : \mathbb{C}^{\text{op}} \rightarrow \mathbf{Pos}$ with $\wedge, \Rightarrow, \forall_{A[R]}, \exists_{A[R]}, \text{Eq}_A$.
- With just these assumptions, we can compute the adjoints $\forall_f, \exists_f : \mathcal{P}(C) \rightarrow \mathcal{P}(D)$ along any $f : C \rightarrow D$, not just projections.

Motivation 4: Computing adjoints to reindexings

- Consider a doctrine $\mathcal{P} : \mathbb{C}^{\text{op}} \rightarrow \mathbf{Pos}$ with $\wedge, \Rightarrow, \forall_{A[R]}, \exists_{A[R]}, \text{Eq}_A$.
- With just these assumptions, we can compute the adjoints $\forall_f, \exists_f : \mathcal{P}(C) \rightarrow \mathcal{P}(D)$ along any $f : C \rightarrow D$, not just projections.
- In the syntactic model, for $P : \mathcal{P}(C)$, $f : C \rightarrow D$:

$$(\forall_f(P))(y) := \forall x. (y =_D f(x) \Rightarrow P(x))$$

$$(\exists_f(P))(y) := \exists x. (f(x) =_D y \wedge P(x))$$

Motivation 4: Computing adjoints to reindexings

- Consider a doctrine $\mathcal{P} : \mathbb{C}^{\text{op}} \rightarrow \mathbf{Pos}$ with $\wedge, \Rightarrow, \forall_{A[I]}, \exists_{A[I]}, \text{Eq}_A$.
- With just these assumptions, we can compute the adjoints $\forall_f, \exists_f : \mathcal{P}(C) \rightarrow \mathcal{P}(D)$ along any $f : C \rightarrow D$, not just projections.
- In the syntactic model, for $P : \mathcal{P}(C)$, $f : C \rightarrow D$:

$$(\forall_f(P))(y) := \forall x. (y =_D f(x) \Rightarrow P(x))$$

$$(\exists_f(P))(y) := \exists x. (f(x) =_D y \wedge P(x))$$

- Compare these to the formula to compute Kan extensions via (co)ends:

$$(\text{Ran}_F P)(y) \cong \int_{x:\mathbb{C}} \text{hom}_{\mathbb{D}}(y, F(\overline{x})) \Rightarrow P(x)$$

$$(\text{Lan}_F P)(y) \cong \int^{x:\mathbb{C}} \text{hom}_{\mathbb{D}}(F(\overline{x}), y) \times P(x)$$

Motivation 4: Computing adjoints to reindexings

- Consider a doctrine $\mathcal{P} : \mathbb{C}^{\text{op}} \rightarrow \mathbf{Pos}$ with $\wedge, \Rightarrow, \forall_{A[R]}, \exists_{A[R]}, \text{Eq}_A$.
- With just these assumptions, we can compute the adjoints $\forall_f, \exists_f : \mathcal{P}(C) \rightarrow \mathcal{P}(D)$ along any $f : C \rightarrow D$, not just projections.
- In the syntactic model, for $P : \mathcal{P}(C)$, $f : C \rightarrow D$:

$$(\forall_f(P))(y) := \forall x. (y =_D f(x) \Rightarrow P(x))$$

$$(\exists_f(P))(y) := \exists x. (f(x) =_D y \wedge P(x))$$

- Compare these to the formula to compute Kan extensions via (co)ends:

$$(\text{Ran}_F P)(y) \cong \int_{x:\mathbb{C}} \text{hom}_{\mathbb{D}}(y, F(\overline{x})) \Rightarrow P(x)$$

$$(\text{Lan}_F P)(y) \cong \int^{x:\mathbb{C}} \text{hom}_{\mathbb{D}}(F(\overline{x}), y) \times P(x)$$

- there is yet no decomposition of these properties for doctrines, e.g., coends as quantifiers/adjoints, or "having directed equality".

Motivation 4: Computing Kan extensions with (co)ends

- A logical proof that $\forall_f(P)$ is right adjoint to precomposition with f :

$$\frac{\frac{[y : D] \quad \Gamma(y) \vdash (\forall_f P)(y) \quad := \forall x. (y = f(x) \Rightarrow P(x))}{[x : C, y : D] \quad \Gamma(y) \vdash y = f(x) \Rightarrow P(x)}}{[x : C, y : D] \quad y = f(x), \Gamma(y) \vdash P(x)} \\ \frac{}{[x : C] \quad \Gamma(f(x)) \vdash P(x)}$$

Motivation 4: Computing Kan extensions with (co)ends

- A logical proof that $\forall_f(P)$ is right adjoint to precomposition with f :

$$\frac{\frac{[y : D] \quad \Gamma(y) \vdash (\forall_f P)(y) \quad := \forall x. (y = f(x) \Rightarrow P(x))}{[x : C, y : D] \quad \Gamma(y) \vdash y = f(x) \Rightarrow P(x)}}{[x : C, y : D] \quad y = f(x), \Gamma(y) \vdash P(x)} \\ \frac{}{[x : C] \quad \Gamma(f(x)) \vdash P(x)}$$

- There is yet no formal system to do this for the directed case;

Motivation 4: Computing Kan extensions with (co)ends

- A logical proof that $\forall_f(P)$ is right adjoint to precomposition with f :

$$\frac{\frac{[y : D] \quad \Gamma(y) \vdash (\forall_f P)(y) \quad := \forall x.(y = f(x) \Rightarrow P(x))}{[x : C, y : D] \quad \Gamma(y) \vdash y = f(x) \Rightarrow P(x)}}{[x : C, y : D] \quad y = f(x), \Gamma(y) \vdash P(x)} \\ \frac{}{[x : C] \quad \Gamma(f(x)) \vdash P(x)}$$

- ▶ There is yet no formal system to do this for the directed case;
- ▶ We follow exactly this proof for Lan/Ran, using our rules for dinaturality.

We present the semantics for a *first-order* non-dependent directed type theory using dinaturality, where types are interpreted by categories, directed equality by hom-functors, quantifiers by (co)ends.

Setting

We present the semantics for a *first-order* non-dependent directed type theory using dinaturality, where types are interpreted by categories, directed equality by hom-functors, quantifiers by (co)ends.

We consider the following interpretation:

Types \rightsquigarrow Categories (possibly with $-^{\text{op}}$)

Contexts \rightsquigarrow Lists of categories

Terms \rightsquigarrow Functors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$

Propositions \rightsquigarrow Endoprofunctors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$

Entailments \rightsquigarrow Dinatural transformations (not required to compose)

Directed equality \rightsquigarrow hom-functors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$.

We present the semantics for a *first-order* non-dependent directed type theory using dinaturality, where types are interpreted by categories, directed equality by hom-functors, quantifiers by (co)ends.

We consider the following interpretation:

Types \rightsquigarrow Categories (possibly with $-^{\text{op}}$)

Contexts \rightsquigarrow Lists of categories

Terms \rightsquigarrow Functors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$

Propositions \rightsquigarrow Endoprofunctors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$

Entailments \rightsquigarrow Dinatural transformations (not required to compose)

Directed equality \rightsquigarrow hom-functors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$.

- **Warning:** we do not give a doctrinal presentation of this logic, precisely because dinaturals do not compose in general.

We present the semantics for a *first-order* non-dependent directed type theory using dinaturality, where types are interpreted by categories, directed equality by hom-functors, quantifiers by (co)ends.

We consider the following interpretation:

Types \rightsquigarrow Categories (possibly with $-^{\text{op}}$)

Contexts \rightsquigarrow Lists of categories

Terms \rightsquigarrow Functors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$

Propositions \rightsquigarrow Endoprofunctors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$

Entailments \rightsquigarrow Dinatural transformations (not required to compose)

Directed equality \rightsquigarrow hom-functors $\mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$.

- **Warning:** we do not give a doctrinal presentation of this logic, precisely because dinaturals do not compose in general.
- Guiding intuition: we study the “doctrine” (in paracategories)

Dinats : $\mathbf{Cat}^{\text{op}} \rightarrow \mathbf{PARACAT}$

Dinats(\mathbb{C}) := $[\mathbb{C}^{\text{op}} \times \mathbb{C}, \mathbf{Set}]_{\text{dinaturals}}$

Directed type theory: notation

- Example of an entailment:

$$[x : \mathbb{C}, y : \mathbb{D}] P(\bar{x}, \bar{y}, x, y) \vdash \alpha : Q(\bar{x}, \bar{y}, x, y)$$

Semantics: " α is a dinatural from P to $Q : (\mathbb{C} \times \mathbb{D})^{\text{op}} \times (\mathbb{C} \times \mathbb{D}) \rightarrow \mathbf{Set}$."

- **Crucial:** variables now can appear both as $x : \mathbb{C}$ or $\bar{x} : \mathbb{C}^{\text{op}}$.
- We give names to assumptions p, q :

$$[x : \mathbb{C}] p : P(\bar{x}, x), q : Q(\bar{x}, x) \vdash h[p, q] : R(\bar{x}, x)$$

prop. context \rightarrow interpreted as pointwise product of presheaves in **Set**.

Directed type theory: notation

- Example of an entailment:

$$[x : \mathbb{C}, y : \mathbb{D}] P(\bar{x}, \bar{y}, x, y) \vdash \alpha : Q(\bar{x}, \bar{y}, x, y)$$

Semantics: " α is a dinatural from P to $Q : (\mathbb{C} \times \mathbb{D})^{\text{op}} \times (\mathbb{C} \times \mathbb{D}) \rightarrow \mathbf{Set}$."

- **Crucial:** variables now can appear both as $x : \mathbb{C}$ or $\bar{x} : \mathbb{C}^{\text{op}}$.
- We give names to assumptions p, q :

$$[x : \mathbb{C}] p : P(\bar{x}, x), q : Q(\bar{x}, x) \vdash h[p, q] : R(\bar{x}, x)$$

prop. context \rightarrow interpreted as pointwise product of presheaves in **Set**.

- The following dinaturals are the same:

$$\begin{aligned} [x : \mathbb{C}] F(\bar{x}) \vdash \alpha : G(\bar{x}, x) \\ [x : \mathbb{C}^{\text{op}}] F(x) \vdash \alpha' : G(x, \bar{x}) \end{aligned}$$

Directed type theory: propositional rules – Products

- Dinaturals support propositional conjunction, using products in **Set**.

Directed type theory: propositional rules – Products

- Dinaturals support propositional conjunction, using products in **Set**.

Theorem (Product of dipresheaves)

There is an isomorphism of sets natural in $\Gamma, P, Q : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$:

$$\frac{[x : \mathbb{C}] \Gamma(\bar{x}, x) \vdash P(\bar{x}, x) \times Q(\bar{x}, x)}{[x : \mathbb{C}] \Gamma(\bar{x}, x) \vdash P(\bar{x}, x), \quad [x : \mathbb{C}] \Gamma(\bar{x}, x) \vdash Q(\bar{x}, x)}.$$

Bottom side: product of sets of dinaturals.

Similarly, $\top_{\mathbb{C}} : \mathbb{C}^{\diamond} \rightarrow \mathbf{Set} := (c, c') \mapsto \{\}$ has a unique dinat $\Gamma \multimap \top_{\mathbb{C}}$.*

Directed type theory: propositional rules – Exponentials

- Contrary to naturals, “exponentials” of dipresheaves are *pointwise*:

Directed type theory: propositional rules – Exponentials

- Contrary to naturals, “exponentials” of dipresheaves are *pointwise*:

Theorem (“Exponential” of dipresheaves) ()

There is an isomorphism of sets natural in $F, G, H : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$:

$$\frac{[x : \mathbb{C}] \quad F(\bar{x}, x) \times G(\bar{x}, x) \vdash H(\bar{x}, x)}{[x : \mathbb{C}] \quad G(\bar{x}, x) \vdash F^{\text{op}}(x, \bar{x}) \Rightarrow H(\bar{x}, x)} \quad (\text{exp})$$

Proof. *Obvious by currying the families of morphisms.*

Directed type theory: propositional rules – Exponentials

- Contrary to naturals, “exponentials” of dipresheaves are *pointwise*:

Theorem (“Exponential” of dipresheaves) ()

There is an isomorphism of sets natural in $F, G, H : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$:

$$\frac{[x : \mathbb{C}] \quad F(\bar{x}, x) \times G(\bar{x}, x) \vdash H(\bar{x}, x)}{[x : \mathbb{C}] \quad G(\bar{x}, x) \vdash F^{\text{op}}(x, \bar{x}) \Rightarrow H(\bar{x}, x)} \quad (\text{exp})$$

Proof. *Obvious by currying the families of morphisms.*

- Why are exponentials of presheaves with naturals not pointwise?*

$$\frac{[x : \mathbb{C}] \quad F(x) \times G(x) \vdash H(x)}{[x : \mathbb{C}] \quad G(x) \vdash F^{\text{op}}(\bar{x}) \Rightarrow H(x)} \quad (\text{exp})$$

Directed type theory: propositional rules – Exponentials

- Contrary to naturals, “exponentials” of dipresheaves are *pointwise*:

Theorem (“Exponential” of dipresheaves)

There is an isomorphism of sets natural in $F, G, H : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$:

$$\frac{[x : \mathbb{C}] \quad F(\bar{x}, x) \times G(\bar{x}, x) \vdash H(\bar{x}, x)}{[x : \mathbb{C}] \quad G(\bar{x}, x) \vdash F^{\text{op}}(x, \bar{x}) \Rightarrow H(\bar{x}, x)} \quad (\text{exp})$$

Proof. Obvious by currying the families of morphisms.

- Why are exponentials of presheaves with naturals not pointwise?

$$\frac{[x : \mathbb{C}] \quad F(x) \times G(x) \vdash H(x)}{[x : \mathbb{C}] \quad G(x) \vdash F^{\text{op}}(\bar{x}) \Rightarrow H(x)} \quad (\text{exp})$$

but, the bottom family would be *dinatural* in x .

Directed type theory: propositional rules – Exponentials

- Contrary to naturals, “exponentials” of dipresheaves are *pointwise*:

Theorem (“Exponential” of dipresheaves) (↪)

There is an isomorphism of sets natural in $F, G, H : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$:

$$\frac{[x : \mathbb{C}] \quad F(\bar{x}, x) \times G(\bar{x}, x) \vdash H(\bar{x}, x)}{[x : \mathbb{C}] \quad G(\bar{x}, x) \vdash F^{\text{op}}(x, \bar{x}) \Rightarrow H(\bar{x}, x)} \quad (\text{exp})$$

Proof. Obvious by currying the families of morphisms.

- Why are exponentials of presheaves with naturals not pointwise?

$$\frac{[x : \mathbb{C}] \quad F(x) \times G(x) \vdash H(x)}{[x : \mathbb{C}] \quad G(x) \vdash F^{\text{op}}(\bar{x}) \Rightarrow H(x)} \quad (\text{exp})$$

but, the bottom family would be *dinatural* in x .

- We will show how to use our rules to justify the usual exponential.

Directed type theory: term rules – Reindexing

- Entailments/dinaturals can be reindexed by terms/difunctors:

Theorem (Reindexing with difunctors) (🌀)

Take a difunctor $F : \mathbb{C}^\diamond \rightarrow \mathbb{D}$ and a dinat $\alpha : P \rightrightarrows Q$ for $P, Q : \mathbb{D}^\diamond \rightarrow \mathbb{E}$.

$$\frac{[x : \mathbb{D}] \quad P(\bar{x}, x) \vdash \alpha : Q(\bar{x}, x)}{[x : \mathbb{C}] \quad P(F^{\text{op}}(x, \bar{x}), F(\bar{x}, x)) \vdash F^*(\alpha) : Q(F^{\text{op}}(x, \bar{x}), F(\bar{x}, x))} \text{ (reindex)}$$

defined by $F^*(\alpha)_x := \alpha_{F(x, x)}$.

- Dinaturality allows us to solve the variance issue of refl: [North 2018]

- Dinaturality allows us to solve the variance issue of refl: [North 2018]

Theorem (Directed equality introduction) ()

There is a dinatural transformation $\text{refl}_{\mathbb{C}} : \top \dashrightarrow \text{hom}$,

$$\frac{}{[x : \mathbb{C}] \top \vdash \text{refl}_{\mathbb{C}} : \text{hom}_{\mathbb{C}}(\bar{x}, x)} \quad (\text{refl})$$

where \top denotes the dipresheaf constant in \top_{Set} .

Proof. $\alpha_x(*) := \text{id}_x$. *Dinaturality:* for any $f : a \rightarrow b$, $f ; \text{id}_b = \text{id}_a ; f$.

- Dinaturality allows us to solve the variance issue of refl: [North 2018]

Theorem (Directed equality introduction) ()

There is a dinatural transformation $\text{refl}_{\mathbb{C}} : \top \xrightarrow{\bullet\bullet} \text{hom}$,

$$\frac{}{[x : \mathbb{C}] \top \vdash \text{refl}_{\mathbb{C}} : \text{hom}_{\mathbb{C}}(\overline{x}, x)} \text{ (refl)}$$

where \top denotes the dipresheaf constant in \top_{Set} .

Proof. $\alpha_x(*) := \text{id}_x$. *Dinaturality:* for any $f : a \rightarrow b$, $f ; \text{id}_b = \text{id}_a ; f$.

- This is reflexivity of directed equality, via identities.

Directed type theory with dinaturality – Intermezzo

- Before introducing the directed J rule, we show its fundamental idea: the connection between naturality and dinaturality.

Directed type theory with dinaturality – Intermezzo

- Before introducing the directed J rule, we show its fundamental idea: the connection between naturality and dinaturality.

Theorem (Characterization of dinaturals via naturality)

There is an isomorphism, natural in $P, Q : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, between the set of dinaturals $P \dashrightarrow Q$ and certain natural transformations:

$$\frac{[x : \mathbb{C}] \ P(\bar{x}, x) \dashrightarrow Q(\bar{x}, x)}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}] \ \text{hom}(a, b) \longrightarrow P^{\text{op}}(b, a) \Rightarrow Q(a, b)}$$

Directed type theory with dinaturality – Intermezzo

- Before introducing the directed J rule, we show its fundamental idea: the connection between naturality and dinaturality.

Theorem (Characterization of dinaturals via naturality)

There is an isomorphism, natural in $P, Q : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, between the set of dinaturals $P \rightrightarrows Q$ and certain natural transformations:

$$\frac{[x : \mathbb{C}] \ P(\bar{x}, x) \rightrightarrows Q(\bar{x}, x)}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}] \ \text{hom}(a, b) \longrightarrow P^{\text{op}}(b, a) \Rightarrow Q(a, b)}$$

Proof.

(\Downarrow) Given $\alpha : P \rightrightarrows Q$ and $f : \text{hom}(a, b)$, the map $P(b, a) \rightarrow Q(a, b)$ is exactly the side of the hexagon in the definition of dinaturality. This is obtained via the functorial action of P, Q .

Directed type theory with dinaturality – Intermezzo

- Before introducing the directed J rule, we show its fundamental idea: the connection between naturality and dinaturality.

Theorem (Characterization of dinaturals via naturality)

There is an isomorphism, natural in $P, Q : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, between the set of dinaturals $P \rightrightarrows Q$ and certain natural transformations:

$$\frac{[x : \mathbb{C}] \ P(\bar{x}, x) \rightrightarrows Q(\bar{x}, x)}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}] \ \text{hom}(a, b) \longrightarrow P^{\text{op}}(b, a) \Rightarrow Q(a, b)}$$

Proof.

- (\Downarrow) Given $\alpha : P \rightrightarrows Q$ and $f : \text{hom}(a, b)$, the map $P(b, a) \rightarrow Q(a, b)$ is exactly the side of the hexagon in the definition of dinaturality. This is obtained via the functorial action of P, Q .
- (\Uparrow) Take $a = b$ and precompose with $\text{id}_a \in \text{hom}(a, a)$.

Directed type theory with dinaturality – Intermezzo

- Before introducing the directed J rule, we show its fundamental idea: the connection between naturality and dinaturality.

Theorem (Characterization of dinaturals via naturality)

There is an isomorphism, natural in $P, Q : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, between the set of dinaturals $P \rightrightarrows Q$ and certain natural transformations:

$$\frac{[x : \mathbb{C}] \ P(\bar{x}, x) \rightrightarrows Q(\bar{x}, x)}{\frac{}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}] \ \text{hom}(a, b) \longrightarrow P^{\text{op}}(b, a) \Rightarrow Q(a, b)}}$$

Proof.

(\Downarrow) Given $\alpha : P \rightrightarrows Q$ and $f : \text{hom}(a, b)$, the map $P(b, a) \rightarrow Q(a, b)$ is exactly the side of the hexagon in the definition of dinaturality.

This is obtained via the functorial action of P, Q .

(\Uparrow) Take $a = b$ and precompose with $\text{id}_a \in \text{hom}(a, a)$.

The isomorphism follows from (di)naturality of both maps.

Directed type theory with dinaturality – Directed J

- Directed equality elimination is just this result, uncurried.

Theorem (Directed equality elimination) ()

Take $\Gamma, P : (\mathbb{A}^{\text{op}}) \times (\mathbb{A}) \times (\mathbb{C}^{\text{op}} \times \mathbb{C}) \rightarrow \mathbf{Set}$.

Given a dinatural $h : \Gamma \rightrightarrows P$, there is a dinatural $J(h)$ as follows:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash h[k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \ e : \text{hom}(a, b), k : \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash J(h)[e, k] : P(a, b, \bar{x}, x)} \quad (J)$$

Directed type theory with dinaturality – Directed J

- Directed equality elimination is just this result, uncurried.

Theorem (Directed equality elimination) ()

Take $\Gamma, P : (\mathbb{A}^{\text{op}}) \times (\mathbb{A}) \times (\mathbb{C}^{\text{op}} \times \mathbb{C}) \rightarrow \mathbf{Set}$.

Given a dinatural $h : \Gamma \dashrightarrow P$, there is a dinatural $J(h)$ as follows:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash h[k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \ e : \text{hom}(a, b), k : \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash J(h)[e, k] : P(a, b, \bar{x}, x)} \quad (J)$$

The dinatural $J(h)$ satisfies the following “computation rule”,

$$J(h)_{zzx}[\text{refl}_{\mathbb{A}_z}, k] = h_{zx}[k]$$

for any object $z : \mathbb{A}, x : \mathbb{C}$ and $k \in \Gamma(z, z, x, x)$.

Directed type theory with dinaturality – Directed J

- Directed equality elimination is just this result, uncurried.

Theorem (Directed equality elimination) ()

Take $\Gamma, P : (\mathbb{A}^{\text{op}}) \times (\mathbb{A}) \times (\mathbb{C}^{\text{op}} \times \mathbb{C}) \rightarrow \mathbf{Set}$.

Given a dinatural $h : \Gamma \dashrightarrow P$, there is a dinatural $J(h)$ as follows:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash h[k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \ e : \text{hom}(a, b), k : \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash J(h)[e, k] : P(a, b, \bar{x}, x)} \quad (J)$$

The dinatural $J(h)$ satisfies the following “computation rule”,

$$J(h)_{zzx}[\text{refl}_{\mathbb{A}_z}, k] = h_{zx}[k]$$

for any object $z : \mathbb{A}, x : \mathbb{C}$ and $k \in \Gamma(z, z, x, x)$.

Proof. Explicitly, the dinatural $J(h)$ is given by

$$J(h)_{abx}[e, k] := (\Gamma(\text{id}_b, e, \text{id}_x, \text{id}_x) ; h_{bx} ; P(e, \text{id}_b, \text{id}_x, \text{id}_x))[k].$$

Computation clearly holds for $e = \text{id}_z$, without dinaturality.

Intution for the directed J rule

- ▶ Whenever there are two positions $a : \mathbb{A}^{\text{op}}, b : \mathbb{A}$ in the conclusion P ,

Intution for the directed J rule

- ▶ Whenever there are two positions $a : \mathbb{A}^{\text{op}}, b : \mathbb{A}$ in the conclusion P ,
- ▶ and I have a directed equality in context $\text{hom}_{\mathbb{A}}(a, b)$,

Intution for the directed J rule

- ▶ Whenever there are two positions $a : \mathbb{A}^{\text{op}}, b : \mathbb{A}$ in the conclusion P ,
 - ▶ and I have a directed equality in context $\text{hom}_{\mathbb{A}}(a, b)$,
-
- ▶ it is enough to prove that P holds “on the diagonal”,
where a, b are identified with the same dinatural variable $z : \mathbb{A}$.

Intuition for the directed J rule

- ▶ Whenever there are two positions $a : \mathbb{A}^{\text{op}}, b : \mathbb{A}$ in the conclusion P ,
 - ▶ and I have a directed equality in context $\text{hom}_{\mathbb{A}}(a, b)$,
-
- ▶ it is enough to prove that P holds “on the diagonal”, where a, b are identified with the same dinatural variable $z : \mathbb{A}$.
 - ▶ Moreover, a, b can be identified in context *only if they appear negatively*.
-
- With the directed J rule we can define maps+properties directed equality.
 - Examples: *transitivity, congruence, transport* for directed equality.

Example (Composition in a category)

Transitivity of directed equality \rightsquigarrow categories \mathbb{C} have *composition maps*.

Example (Composition in a category)

Transitivity of directed equality \rightsquigarrow categories \mathbb{C} have *composition maps*.
Composition is natural in $a : \mathbb{C}^{\text{op}}, c : \mathbb{C}$ and dinatural in $b : \mathbb{C}$:

$$\frac{\frac{[z : \mathbb{C}, c : \mathbb{C}]}{\text{hom}(\bar{z}, c) \vdash \text{id} : \text{hom}(\bar{z}, c)} \text{ (id)}}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}, c : \mathbb{C}] \text{ hom}(a, b), \text{hom}(\bar{b}, c) \vdash J(\text{id}) : \text{hom}(a, c)} \text{ (J)}$$

Example (Composition in a category)

Transitivity of directed equality \rightsquigarrow categories \mathbb{C} have *composition maps*.
Composition is natural in $a : \mathbb{C}^{\text{op}}, c : \mathbb{C}$ and dinatural in $b : \mathbb{C}$:

$$\frac{\frac{}{[z : \mathbb{C}, c : \mathbb{C}]}}{\frac{}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}, c : \mathbb{C}]}} \quad \frac{}{\text{hom}(\bar{z}, c) \vdash \text{id} : \text{hom}(\bar{z}, c)} \text{(id)} \quad \frac{}{\text{hom}(a, b), \text{hom}(\bar{b}, c) \vdash J(\text{id}) : \text{hom}(a, c)} \text{(J)}$$

We contract $\text{hom}(a, b)$. Rule (J) can be applied: a, b appear only negatively in ctx (a does not) and positively in conclusion (\bar{b} does not).

Example (Composition in a category)

Transitivity of directed equality \rightsquigarrow categories \mathbb{C} have *composition maps*. Composition is natural in $a : \mathbb{C}^{\text{op}}, c : \mathbb{C}$ and dinatural in $b : \mathbb{C}$:

$$\frac{\frac{}{[z : \mathbb{C}, c : \mathbb{C}]}}{\text{hom}(\bar{z}, c) \vdash \text{id} : \text{hom}(\bar{z}, c)} \text{(id)} \quad \frac{}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}, c : \mathbb{C}] \text{ hom}(a, b), \text{hom}(\bar{b}, c) \vdash J(\text{id}) : \text{hom}(a, c)} \text{(J)}$$

We contract $\text{hom}(a, b)$. Rule (J) can be applied: a, b appear only negatively in ctx (a does not) and positively in conclusion (\bar{b} does not).

The computation rule for $f ; g := J(\text{id})[f, g]$ states unitality on the left.

Failure of symmetry for directed equality

These restrictions do *not* allow us to obtain directed equality is symmetric:

$$[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}] \ e : \text{hom}(a, b) \not\vdash \text{sym} : \text{hom}(\bar{b}, \bar{a})$$

$\text{hom}(a, b)$ cannot be contracted: a, b must appear *positively* in conclusion.

- As in the classical case, the rule for directed equality is an isomorphism.

Theorem (J as isomorphism)

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash h[k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad e : \text{hom}(a, b), k : \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash J(h)[e, k] : P(a, b, \bar{x}, x)} \quad (\text{hom})$$

- As in the classical case, the rule for directed equality is an isomorphism.

Theorem (J as isomorphism)

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash h[k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad e : \text{hom}(a, b), k : \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash J(h)[e, k] : P(a, b, \bar{x}, x)} \quad (\text{hom})$$

Proof. The inverse is $J^{-1}(\alpha)_{zx}[k] := \alpha_{zzx}[\text{refl}_{\mathbb{A}z}, k]$.

- As in the classical case, the rule for directed equality is an isomorphism.

Theorem (J as isomorphism)

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash h[k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad e : \text{hom}(a, b), k : \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash J(h)[e, k] : P(a, b, \bar{x}, x)} \quad (\text{hom})$$

Proof. The inverse is $J^{-1}(\alpha)_{zx}[k] := \alpha_{zzx}[\text{refl}_{\mathbb{A}_z}, k]$.

The computation rule for hom-elimination is precisely $J ; J^{-1} = \text{id}$.

On the other hand, $J^{-1} ; J = \text{id}$ follows using (di)naturality.

Directed type theory – dependent directed J

- Back to the transitivity example.
- *How do we prove associativity of transitivity and unitality on the right?*

- Back to the transitivity example.
- *How do we prove associativity of transitivity and unitality on the right?*
- Using the same method of MLTT: by *dependent* hom induction.

Directed type theory – dependent directed J

- Back to the transitivity example.
- *How do we prove associativity of transitivity and unitality on the right?*
- Using the same method of MLTT: by *dependent* hom induction.
- We work with *proof-relevant entailments*: we need a judgm. for $\alpha = \beta$.

Directed type theory – dependent directed J

- Back to the transitivity example.
- *How do we prove associativity of transitivity and unitality on the right?*
- Using the same method of MLTT: by *dependent* hom induction.
- We work with *proof-relevant entailments*: we need a judgm. for $\alpha = \beta$.

Theorem (Dependent J rule for judgemental equality) (👉)

Take $\Gamma, P : (\mathbb{A}^{\text{op}}) \times (\mathbb{A}) \times (\mathbb{C}^{\text{op}} \times \mathbb{C}) \rightarrow \mathbf{Set}$. Given dinats α, β where J can be applied, then the equality above implies the one below:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \ k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash \alpha[\text{refl}_z, k] = \beta[\text{refl}_z, k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \ e : \text{hom}(a, b), k : \Gamma(\bar{a}, \bar{b}, \bar{x}, x) \vdash \alpha[e, k] = \beta[e, k] : P(a, b, \bar{x}, x)} \quad (J\text{-eq})$$

Directed type theory – dependent directed J

- Back to the transitivity example.
- *How do we prove associativity of transitivity and unitality on the right?*
- Using the same method of MLTT: by *dependent* hom induction.
- We work with *proof-relevant entailments*: we need a judgm. for $\alpha = \beta$.

Theorem (Dependent J rule for judgemental equality) (👉)

Take $\Gamma, P : (\mathbb{A}^{\text{op}}) \times (\mathbb{A}) \times (\mathbb{C}^{\text{op}} \times \mathbb{C}) \rightarrow \mathbf{Set}$. Given dinats α, β where J can be applied, then the equality above implies the one below:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \ k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash \alpha[\text{refl}_z, k] = \beta[\text{refl}_z, k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \ e : \text{hom}(a, b), k : \Gamma(\bar{a}, \bar{b}, \bar{x}, x) \vdash \alpha[e, k] = \beta[e, k] : P(a, b, \bar{x}, x)} \quad (J\text{-eq})$$

i.e., $\forall k : P(z, z, x, x). \alpha_{zzx}[\text{refl}_z, k] = \beta_{zzx}[\text{refl}_z, k]$ implies $\alpha = \beta$.

Directed type theory – dependent directed J

- Back to the transitivity example.
- *How do we prove associativity of transitivity and unitality on the right?*
- Using the same method of MLTT: by *dependent* hom induction.
- We work with *proof-relevant entailments*: we need a judgm. for $\alpha = \beta$.

Theorem (Dependent J rule for judgemental equality) (👉)

Take $\Gamma, P : (\mathbb{A}^{\text{op}}) \times (\mathbb{A}) \times (\mathbb{C}^{\text{op}} \times \mathbb{C}) \rightarrow \mathbf{Set}$. Given dinats α, β where J can be applied, then the equality above implies the one below:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \ k : \Gamma(\bar{z}, z, \bar{x}, x) \vdash \alpha[\text{refl}_z, k] = \beta[\text{refl}_z, k] : P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \ e : \text{hom}(a, b), k : \Gamma(\bar{a}, \bar{b}, \bar{x}, x) \vdash \alpha[e, k] = \beta[e, k] : P(a, b, \bar{x}, x)} \quad (J\text{-eq})$$

i.e., $\forall k : P(z, z, x, x). \ \alpha_{zzx}[\text{refl}_z, k] = \beta_{zzx}[\text{refl}_z, k]$ implies $\alpha = \beta$.

Proof. By hypothesis, $J^{-1}(\alpha) = J^{-1}(\beta)$, simply apply J .

Example (Properties of composition)

Back to composition: we want to prove unitality and associativity.

Example (Properties of composition)

Back to composition: we want to prove unitality and associativity.

$$\frac{[z : \mathbb{C}, c : \mathbb{C}] \quad g : \text{hom}(\bar{z}, c) \vdash \text{refl}_z ; g = g : \text{hom}(\bar{z}, c)}{(J\text{-comp})}$$

Example (Properties of composition)

Back to composition: we want to prove unitality and associativity.

$$\frac{}{[z : \mathbb{C}, c : \mathbb{C}] \ g : \text{hom}(\bar{z}, c) \vdash \text{refl}_z ; g = g : \text{hom}(\bar{z}, c)} \quad (J\text{-comp})$$

Unitality on the right is shown by dependent hom induction:

$$\frac{}{[w : \mathbb{C}] \top \vdash \text{refl}_w ; \text{refl}_w = \text{refl}_w : \text{hom}(\bar{w}, w)} \quad (J\text{-comp})$$

$$\frac{}{[a : \mathbb{C}^{\text{op}}, z : \mathbb{C}] \ f : \text{hom}(a, z) \vdash f ; \text{refl}_z = f : \text{hom}(a, z)} \quad (J\text{-eq})$$

Example (Properties of composition)

Back to composition: we want to prove unitality and associativity.

$$\frac{}{[z : \mathbb{C}, c : \mathbb{C}] \ g : \text{hom}(\bar{z}, c) \vdash \text{refl}_z ; g = g : \text{hom}(\bar{z}, c)} \quad (J\text{-comp})$$

Unitality on the right is shown by dependent hom induction:

$$\frac{}{[w : \mathbb{C}] \ \top \vdash \text{refl}_w ; \text{refl}_w = \text{refl}_w : \text{hom}(\bar{w}, w)} \quad (J\text{-comp})$$

$$\frac{[a : \mathbb{C}^{\text{op}}, z : \mathbb{C}] \ f : \text{hom}(a, z) \vdash f ; \text{refl}_z = f : \text{hom}(a, z)}{} \quad (J\text{-eq})$$

To prove associativity, simply contract $f : \text{hom}(a, b)$:

$$\frac{[z : \mathbb{C}, c : \mathbb{C}, d : \mathbb{C}] \quad g : \text{hom}(\bar{z}, c), h : \text{hom}(\bar{c}, d) \vdash \text{refl}_z ; (g ; h) = (\text{refl}_z ; g) ; h : \text{hom}(\bar{z}, d)}{[a : \mathbb{C}, b : \mathbb{C}, c : \mathbb{C}, d : \mathbb{C}] \ f : \text{hom}(\bar{a}, b), g : \text{hom}(\bar{b}, c), h : \text{hom}(\bar{c}, d) \vdash f ; (g ; h) = (f ; g) ; h : \text{hom}(\bar{a}, d)} \quad (J\text{-eq})$$

where the top sequent $= g ; h$ by computation rules for $\text{comp} := J(\text{id})$.

Example (Properties of composition)

Back to composition: we want to prove unitality and associativity.

$$\frac{}{[z : \mathbb{C}, c : \mathbb{C}] \ g : \text{hom}(\bar{z}, c) \vdash \text{refl}_z ; g = g : \text{hom}(\bar{z}, c)} \quad (J\text{-comp})$$

Unitality on the right is shown by dependent hom induction:

$$\frac{}{[w : \mathbb{C}] \ \top \vdash \text{refl}_w ; \text{refl}_w = \text{refl}_w : \text{hom}(\bar{w}, w)} \quad (J\text{-comp})$$

$$\frac{[a : \mathbb{C}^{\text{op}}, z : \mathbb{C}] \ f : \text{hom}(a, z) \vdash f ; \text{refl}_z = f : \text{hom}(a, z)}{} \quad (J\text{-eq})$$

To prove associativity, simply contract $f : \text{hom}(a, b)$:

$$\frac{\frac{[z : \mathbb{C}, c : \mathbb{C}, d : \mathbb{C}] \quad g : \text{hom}(\bar{z}, c), h : \text{hom}(\bar{c}, d) \vdash \text{refl}_z ; (g ; h) = (\text{refl}_z ; g) ; h : \text{hom}(\bar{z}, d)}{} \quad (J\text{-comp})}{[a : \mathbb{C}, b : \mathbb{C}, c : \mathbb{C}, d : \mathbb{C}] \ f : \text{hom}(\bar{a}, b), g : \text{hom}(\bar{b}, c), h : \text{hom}(\bar{c}, d) \vdash f ; (g ; h) = (f ; g) ; h : \text{hom}(\bar{a}, d)} \quad (J\text{-eq})$$

where the top sequent $= g ; h$ by computation rules for $\text{comp} := J(\text{id})$.

- *Note!* These are exactly the steps in MLTT for transitivity of paths.

Directed type theory – congr for directed equality

- Every term respects directed equality, i.e., it is a “congruence”:
semantically, this is just the functorial action of terms F on morphisms.

Directed type theory – congr for directed equality

- Every term respects directed equality, i.e., it is a “congruence”:
semantically, this is just the functorial action of terms F on morphisms.

Example (Directed equality is a congruence)

Let $F : \mathbb{C} \rightarrow \mathbb{D}$ be a functor.

$$\frac{\frac{[z : \mathbb{C}] \top \vdash \text{hom}_{\mathbb{D}}(F^{\text{op}}(\bar{z}), F(z))}{[x : \mathbb{C}, y : \mathbb{C}] \text{hom}_{\mathbb{C}}(\bar{x}, y) \vdash \text{hom}_{\mathbb{D}}(F^{\text{op}}(\bar{x}), F(y))} \quad (J)}{\quad} \quad (\text{reindex}) + (\text{refl})$$

Directed type theory – congr for directed equality

- Every term respects directed equality, i.e., it is a “congruence”:
semantically, this is just the functorial action of terms F on morphisms.

Example (Directed equality is a congruence)

Let $F : \mathbb{C} \rightarrow \mathbb{D}$ be a functor.

$$\frac{\frac{[z : \mathbb{C}] \top \vdash \text{hom}_{\mathbb{D}}(F^{\text{op}}(\bar{z}), F(z))}{[x : \mathbb{C}, y : \mathbb{C}] \text{hom}_{\mathbb{C}}(\bar{x}, y) \vdash \text{hom}_{\mathbb{D}}(F^{\text{op}}(\bar{x}), F(y))} \text{ (reindex)+(refl)} \quad (J)$$

Take $\text{map}_F[f] := J(F^*(\text{refl}_{\mathbb{C}}))$. Computation rule: F maps refl to refl :

$$\frac{[z : \mathbb{C}] \top \vdash \text{map}_F[\text{refl}_z] = F^*(\text{refl}_z) : \text{hom}_{\mathbb{D}}(F^{\text{op}}(\bar{x}), F(x))}{\text{ (J-comp)}}$$

Directed type theory – congr for directed equality

- Every term respects directed equality, i.e., it is a “congruence”:
semantically, this is just the functorial action of terms F on morphisms.

Example (Directed equality is a congruence)

Let $F : \mathbb{C} \rightarrow \mathbb{D}$ be a functor.

$$\frac{\frac{}{[z : \mathbb{C}] \top \vdash \text{hom}_{\mathbb{D}}(F^{\text{op}}(\bar{z}), F(z))} \text{ (reindex)+(refl)}}{[x : \mathbb{C}, y : \mathbb{C}] \text{ hom}_{\mathbb{C}}(\bar{x}, y) \vdash \text{hom}_{\mathbb{D}}(F^{\text{op}}(\bar{x}), F(y))} (J)$$

Take $\text{map}_F[f] := J(F^*(\text{refl}_{\mathbb{C}}))$. Computation rule: F maps refl to refl :

$$\frac{}{[z : \mathbb{C}] \top \vdash \text{map}_F[\text{refl}_z] = F^*(\text{refl}_z) : \text{hom}_{\mathbb{D}}(F^{\text{op}}(\bar{x}), F(x))} (J\text{-comp})$$

Functoriality holds, since both top sides = $\text{map}_F[g]$ via computation rules:

$$\frac{[z : \mathbb{C}, c : \mathbb{C}] \quad g : \text{hom}(\bar{z}, c) \vdash \text{map}_F[\text{refl}_z ; g] = \text{refl}_{F(z)} ; \text{map}_F[g] : \text{hom}(\bar{z}, d)}{(J\text{-comp})} \quad \frac{}{(J\text{-eq})}$$

$$[a : \mathbb{C}, b : \mathbb{C}, c : \mathbb{C}] \quad f : \text{hom}(\bar{a}, b), g : \text{hom}(\bar{b}, c) \vdash \text{map}_F[f ; g] = \text{map}_F[f] ; \text{map}_F[g] : \text{hom}(\bar{a}, d)$$

Directed type theory – transport along directed equalities

- Transporting points of predicates (i.e., presheaves) along directed equalities is the functorial action of $P : \mathbb{C} \rightarrow \mathbf{Set}$:

Directed type theory – transport along directed equalities

- Transporting points of predicates (i.e., presheaves) along directed equalities is the functorial action of $P : \mathbb{C} \rightarrow \mathbf{Set}$:

Example (Transporting along directed equality)

For any $P : \mathbb{C} \rightarrow \mathbf{Set}$:

$$\frac{\overline{[z : \mathbb{C}] \ P(z) \vdash P(z)}}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}] \ \text{hom}(a, b), P(\bar{a}) \vdash P(b)} \begin{array}{l} (\text{id}) \\ (J) \end{array}$$

Directed type theory – transport along directed equalities

- Transporting points of predicates (i.e., presheaves) along directed equalities is the functorial action of $P : \mathbb{C} \rightarrow \mathbf{Set}$:

Example (Transporting along directed equality)

For any $P : \mathbb{C} \rightarrow \mathbf{Set}$:

$$\frac{\overline{[z : \mathbb{C}] P(z) \vdash P(z)}}{[a : \mathbb{C}^{\text{op}}, b : \mathbb{C}] \text{hom}(a, b), P(\bar{a}) \vdash P(b)} \begin{array}{l} (\text{id}) \\ (J) \end{array}$$

Computation rule for $\text{subst}[f, k] := J(\text{id})$:

“transporting a point of $P(a)$ along the path refl_z is the identity”:

$$\overline{[z : \mathbb{C}] k : P(z) \vdash \text{subst}[\text{refl}_z, k] = k : P(z)} \quad (J\text{-comp})$$

Directed type theory – Groupoidal case

- When $\mathbb{A} \cong \mathbb{A}^{\text{op}}$ is a groupoid, hom is the characterization of symmetric equality as left adjoint to reindexing on diagonals. (with Frobenius)

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{z}, z, \bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{\text{hom}(a, b), \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash P(a, b, \bar{x}, x)}} \quad (\text{hom})$$

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] \quad a \cong b, \Gamma(b, a, x) \vdash P(a, b, x)}} \quad (\text{eq})$$

Directed type theory – Groupoidal case

- When $\mathbb{A} \cong \mathbb{A}^{\text{op}}$ is a groupoid, hom is the characterization of symmetric equality as left adjoint to reindexing on diagonals. (with Frobenius)

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{z}, z, \bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{\quad}}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \text{ hom}(a, b), \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash P(a, b, \bar{x}, x)} \text{ (hom)}$$

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{\quad}}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] a \cong b, \Gamma(b, a, x) \vdash P(a, b, x)} \text{ (eq)}$$

- The *proof-relevant directness* of **Cat** seems to be the fundamental obstacle to a fully compositional theory of dinaturals:

Theorem (Dinaturals in groupoids compose)

Given a groupoid $\mathbb{C} \cong \mathbb{C}^{\text{op}}$ and any \mathbb{D} for $F, G, H : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$, all dinaturals $\alpha : F \rightrightarrows G$, $\beta : G \rightrightarrows H$ compose.

Directed type theory – Groupoidal case

- When $\mathbb{A} \cong \mathbb{A}^{\text{op}}$ is a groupoid, hom is the characterization of symmetric equality as left adjoint to reindexing on diagonals. (with Frobenius)

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{z}, z, \bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{\text{hom}(a, b), \Gamma(\bar{b}, \bar{a}, \bar{x}, x) \vdash P(a, b, \bar{x}, x)}} \quad (\text{hom})$$

$$\frac{\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(z, z, x) \vdash P(z, z, x)}{[a : \mathbb{A}, b : \mathbb{A}, x : \mathbb{C}] \quad a \cong b, \Gamma(b, a, x) \vdash P(a, b, x)}} \quad (\text{eq})$$

- The *proof-relevant directness* of **Cat** seems to be the fundamental obstacle to a fully compositional theory of dinaturals:

Theorem (Dinaturals in groupoids compose)

Given a groupoid $\mathbb{C} \cong \mathbb{C}^{\text{op}}$ and any \mathbb{D} for $F, G, H : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbb{D}$, all dinaturals $\alpha : F \rightrightarrows G$, $\beta : G \rightrightarrows H$ compose.

- Can the directed rule also be characterized as an adjunction? Yes!...

Directed equality as relative left adjoint

Symmetric equality is a left adjoint to identifying two variables together.

Directed equality as relative left adjoint

Symmetric equality is a left adjoint to identifying two variables together.

Idea: *Directed equality* is a *relative* left adjoint to identifying two *natural* variables together with a *dinatural* one.

Directed equality as relative left adjoint

Symmetric equality is a left adjoint to identifying two variables together.

Idea: *Directed equality* is a *relative* left adjoint to identifying two *natural* variables together with a *dinatural* one.

- The *relative* part imposes the syntactic restrictions of directed J : the relative functor is just reindexing along projections.

Directed equality as relative left adjoint

Symmetric equality is a left adjoint to identifying two variables together.

Idea: *Directed equality* is a *relative* left adjoint to identifying two *natural* variables together with a *dinatural* one.

- The *relative* part imposes the syntactic restrictions of directed J : the relative functor is just reindexing along projections.
- Unfortunately, for **Cat** we can only state this as a relative *para*-adjunction, because of non-compositionality of dinaturals.

Directed equality as relative left adjoint

Symmetric equality is a left adjoint to identifying two variables together.


Idea: *Directed equality* is a *relative* left adjoint to identifying two *natural* variables together with a *dinatural* one.

- The *relative* part imposes the syntactic restrictions of directed J : the relative functor is just reindexing along projections.
- Unfortunately, for **Cat** we can only state this as a relative *para*-adjunction, because of non-compositionality of dinaturals.
- *Para*- indicates that composition is partial: paracategories, parafunctors, para-adjunctions. [Hermida 2003]

Background: Relative adjunctions

Definition (Left relative adjunction [Arkor 2024])

Consider this situation of functors and categories:



We say that L is the J -relative left adjoint to R , written $L \dashv_J R$, if

$$\mathbb{D}(L(x), y) \cong \mathbb{X}(J(x), R(y))$$

is a bijection natural in both arguments $x : \mathbb{C}, y : \mathbb{D}$.

Directed equality as adjoint (1)

Theorem (Directed equality as relative left adjoint) ($\hookrightarrow \approx$)

► Let $[\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ the paracategory where morphisms are dinats natural in $\mathbb{A}^{\text{op}}, \mathbb{A}$ and dinatural in \mathbb{C} , and $[\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ as dinaturals.

Directed equality as adjoint (1)

Theorem (Directed equality as relative left adjoint) (\approx)

- Let $[\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ the paracategory where morphisms are dinats natural in $\mathbb{A}^{\text{op}}, \mathbb{A}$ and dinatural in \mathbb{C} , and $[\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ as dinaturals.
- Take the parafunctor $\pi_{\mathbb{A}}^* : [\mathbb{C}^{\diamond}, \mathbf{Set}] \rightarrow [\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ defined in the intuitive way by precomposing with projections.

Directed equality as adjoint (1)

Theorem (Directed equality as relative left adjoint) (\approx)

- Let $[\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ the paracategory where morphisms are dinats natural in $\mathbb{A}^{\text{op}}, \mathbb{A}$ and dinatural in \mathbb{C} , and $[\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ as dinaturals.
- Take the parafunctor $\pi_{\mathbb{A}}^* : [\mathbb{C}^{\diamond}, \mathbf{Set}] \rightarrow [\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ defined in the intuitive way by precomposing with projections.
- There is a dipresheaf $\text{hom}_{\mathbb{A}} \in [\mathbb{A}^{\text{op}} \times \mathbb{A}, \mathbf{Set}]$ such that the functor

$$\text{hom}_{\mathbb{A}} \times - : [\mathbb{C}^{\diamond}, \mathbf{Set}] \rightarrow [\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$$

$$\text{hom}_{\mathbb{A}} \times \Gamma := (\bar{a}, a, \bar{x}, x) \mapsto \text{hom}(\bar{a}, a) \times \Gamma(\bar{x}, x),$$

$$(\text{hom}_{\mathbb{A}} \times \alpha_x)_{abc} := \lambda(e \in \text{hom}(a, b), k \in \Gamma(c, c)).(e, \alpha_c(k))$$

Directed equality as adjoint (1)

Theorem (Directed equality as relative left adjoint) ($\hookrightarrow \approx$)

- Let $[\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ the paracategory where morphisms are dinats natural in $\mathbb{A}^{\text{op}}, \mathbb{A}$ and dinatural in \mathbb{C} , and $[\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ as dinaturals.
- Take the parafunctor $\pi_{\mathbb{A}}^* : [\mathbb{C}^{\diamond}, \mathbf{Set}] \rightarrow [\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$ defined in the intuitive way by precomposing with projections.
- There is a dipresheaf $\text{hom}_{\mathbb{A}} \in [\mathbb{A}^{\text{op}} \times \mathbb{A}, \mathbf{Set}]$ such that the functor

$$\text{hom}_{\mathbb{A}} \times - : [\mathbb{C}^{\diamond}, \mathbf{Set}] \rightarrow [\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$$

$$\text{hom}_{\mathbb{A}} \times \Gamma := (\bar{a}, a, \bar{x}, x) \mapsto \text{hom}(\bar{a}, a) \times \Gamma(\bar{x}, x),$$

$$(\text{hom}_{\mathbb{A}} \times \alpha_x)_{abc} := \lambda(e \in \text{hom}(a, b), k \in \Gamma(c, c)).(e, \alpha_c(k))$$

determines a $\pi_{\mathbb{A}}^*$ -relative left adjoint to the functor

$$\Delta_{\mathbb{A}} \times - : [\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}] \rightarrow [\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]$$

$$\Delta_{\mathbb{A}} \times P := P$$

$$(\Delta_{\mathbb{A}} \times \alpha_{abc})_{zx} := \alpha_{zzx}.$$

Directed equality as adjoint (2)

Theorem (Directed equality as relative left adjoint, cont.) ($\mathbb{U} \approx$)

► Thus the left relative adjointness situation

$$\text{hom}_{\mathbb{A}} \times - \dashv_{\pi_{\mathbb{A}}^*} \Delta_{\mathbb{A}} \times -$$

is as follows:

$$\begin{array}{ccc}
 & [\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}] & \\
 \text{hom}_{\mathbb{A}} \times - \nearrow & \dashv & \searrow \Delta_{\mathbb{A}} \times - \\
 [\mathbb{C}^{\diamond}, \mathbf{Set}] & \xrightarrow{\pi_{\mathbb{A}}^*} & [\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]
 \end{array}$$

Directed equality as adjoint (2)

Theorem (Directed equality as relative left adjoint, cont.) ($\mathbb{A} \approx$)

► Thus the left relative adjointness situation

$$\text{hom}_{\mathbb{A}} \times - \dashv_{\pi_{\mathbb{A}}^*} \Delta_{\mathbb{A}} \times -$$

is as follows:

$$\begin{array}{ccc}
 & [\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}] & \\
 \text{hom}_{\mathbb{A}} \times - \nearrow & \dashv & \searrow \Delta_{\mathbb{A}} \times - \\
 [\mathbb{C}^{\diamond}, \mathbf{Set}] & \xrightarrow{\pi_{\mathbb{A}}^*} & [\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]
 \end{array}$$

Proof. The required isomorphism is the following:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad \text{hom}_{\mathbb{A}}(a, b) \times \Gamma(\bar{x}, x) \vdash P(a, b, \bar{x}, x)} \quad (\text{hom-rel-adj})$$

which is an instance of directed J , where Γ is mute in $\bar{a} : \mathbb{A}, \bar{b} : \mathbb{A}^{\text{op}}$.

Directed equality as adjoint (2)

Theorem (Directed equality as relative left adjoint, cont.) ($\mathbb{A} \approx$)

► Thus the left relative adjointness situation

$$\text{hom}_{\mathbb{A}} \times - \dashv_{\pi_{\mathbb{A}}^*} \Delta_{\mathbb{A}} \times -$$

is as follows:

$$\begin{array}{ccc}
 & [\mathbb{A}^{\text{op}} \times \mathbb{A} \times \mathbb{C}^{\diamond}, \mathbf{Set}] & \\
 \text{hom}_{\mathbb{A}} \times - \nearrow & \dashv & \searrow \Delta_{\mathbb{A}} \times - \\
 [\mathbb{C}^{\diamond}, \mathbf{Set}] & \xrightarrow{\pi_{\mathbb{A}}^*} & [\mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond}, \mathbf{Set}]
 \end{array}$$

Proof. The required isomorphism is the following:

$$\frac{[z : \mathbb{A}, x : \mathbb{C}] \quad \Gamma(\bar{x}, x) \vdash P(\bar{z}, z, \bar{x}, x)}{[a : \mathbb{A}^{\text{op}}, b : \mathbb{A}, x : \mathbb{C}] \quad \text{hom}_{\mathbb{A}}(a, b) \times \Gamma(\bar{x}, x) \vdash P(a, b, \bar{x}, x)} \quad (\text{hom-rel-adj})$$

which is an instance of directed J , where Γ is mute in $\bar{a} : \mathbb{A}, \bar{b} : \mathbb{A}^{\text{op}}$.

$(\text{hom-rel-adj}) \Leftrightarrow (\text{hom})$: pick $P := \Gamma^{\text{op}}(b, a, x, \bar{x}) \Rightarrow P(a, b, \bar{x}, x)$, use (exp).

(Co)end calculus via dinaturality

- *What are the quantifiers of directed type theory?*
- Dinaturality allows us to view (co)ends as “adjoints” to weakening:

Theorem (Ends and coends as quantifiers)

Take $P : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, and the functor precomposing with projections

$$\pi_{\mathbb{A}}^*(P) : \mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond} \rightarrow \mathbf{Set}$$

$$\pi_{\mathbb{A}}^*(P) := (\bar{a}, a, \bar{x}, x) \mapsto P(\bar{x}, x),$$

(Co)end calculus via dinaturality

- What are the quantifiers of directed type theory?
- Dinaturality allows us to view (co)ends as “adjoints” to weakening:

Theorem (Ends and coends as quantifiers) (☞)

Take $P : \mathbb{C}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}$, and the functor precomposing with projections

$$\pi_{\mathbb{A}}^*(P) : \mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond} \rightarrow \mathbf{Set}$$

$$\pi_{\mathbb{A}}^*(P) := (\bar{a}, a, \bar{x}, x) \mapsto P(\bar{x}, x),$$

There are isos of sets of dinats, natural in $P : \mathbb{C}^{\diamond} \rightarrow \mathbf{Set}, Q : \mathbb{A}^{\diamond} \times \mathbb{C}^{\diamond} \rightarrow \mathbf{Set}$:

$$\frac{[a : \mathbb{A}, x : \mathbb{C}] \ P(\bar{x}, x) \vdash Q(\bar{a}, a, \bar{x}, x)}{[x : \mathbb{C}] \ P(\bar{x}, x) \vdash \int_{a:\mathbb{A}} Q(\bar{a}, a, \bar{x}, x)} \text{ (end)}$$
$$\frac{[x : \mathbb{C}] \ \int^{a:\mathbb{A}} Q(\bar{a}, a, \bar{x}, x) \vdash P(\bar{x}, x)}{[a : \mathbb{A}, x : \mathbb{C}] \ Q(\bar{a}, a, \bar{x}, x) \vdash P(\bar{x}, x)} \text{ (coend)}$$

(Co)ends as quantifiers

Theorem (Beck-Chevalley and Frobenius condition for (co)ends)

(Co)ends satisfy a Beck-Chevalley condition: for $F : \mathbb{C}^\diamond \rightarrow \mathbb{D}$ there is a strict isomorphism in the (large) functor category $[[\mathbb{A}^\diamond \times \mathbb{D}^\diamond, \mathbf{Set}], [\mathbb{D}^\diamond, \mathbf{Set}]]$

$$\int_{\mathbb{A}[\mathbb{D}]} ; F^* \cong (\mathrm{id}_{\mathbb{A}^\diamond} \times F)^* ; \int_{\mathbb{A}[\mathbb{C}]}$$

where $\int_{\mathbb{A}[\mathbb{C}]}, \int^{\mathbb{A}[\mathbb{C}]} : [\mathbb{A}^\diamond \times \mathbb{C}^\diamond, \mathbf{Set}] \rightarrow [\mathbb{C}^\diamond, \mathbf{Set}]$ are parametric (co)ends, and $F^ : [\mathbb{D}^\diamond, \mathbf{Set}] \rightarrow [\mathbb{C}^\diamond, \mathbf{Set}]$ is precomposition with F^\diamond .*

(Co)ends as quantifiers

Theorem (Beck-Chevalley and Frobenius condition for (co)ends)

(Co)ends satisfy a Beck-Chevalley condition: for $F : \mathbb{C}^\diamond \rightarrow \mathbb{D}$ there is a strict isomorphism in the (large) functor category $[[\mathbb{A}^\diamond \times \mathbb{D}^\diamond, \mathbf{Set}], [\mathbb{D}^\diamond, \mathbf{Set}]]$

$$\int_{\mathbb{A}[\mathbb{D}]} ; F^* \cong (\text{id}_{\mathbb{A}^\diamond} \times F)^* ; \int_{\mathbb{A}[\mathbb{C}]}$$

where $\int_{\mathbb{A}[\mathbb{C}]}, \int^{\mathbb{A}[\mathbb{C}]} : [\mathbb{A}^\diamond \times \mathbb{C}^\diamond, \mathbf{Set}] \rightarrow [\mathbb{C}^\diamond, \mathbf{Set}]$ are parametric (co)ends, and $F^ : [\mathbb{D}^\diamond, \mathbf{Set}] \rightarrow [\mathbb{C}^\diamond, \mathbf{Set}]$ is precomposition with F^\diamond .*

Moreover, a Frobenius condition for coends holds: there is an isomorphism natural in $\Gamma : \mathbb{A}^\diamond \times \mathbb{C}^\diamond \rightarrow \mathbf{Set}, P : \mathbb{C}^\diamond \rightarrow \mathbf{Set}$,

$$\int^{\mathbb{A}[\mathbb{C}]} (\pi_{\mathbb{A}[\mathbb{C}]}^*(P) \wedge \Gamma) \cong \pi_{\mathbb{A}[\mathbb{C}]}^*(P) \wedge \int^{\mathbb{A}[\mathbb{C}]} (\Gamma),$$

where $- \wedge - : [\mathbb{C}, \mathbf{Set}] \times [\mathbb{C}, \mathbf{Set}] \rightarrow [\mathbb{C}, \mathbf{Set}]$ is the pointwise product.

Proof. Beck-Chevalley is easy. For Frobenius, we can use logical rules to mirror the argument in [Jacobs 1999, 1.9.12(i)] with exponentials.

(Co)end calculus using our rules

- Using dinaturals as semantics, we can prove useful theorems “logically”.

Rules for (co)ends as quantifiers + directed equality for proofs of:

- (Co)Yoneda,
- Adjointness of Kan extensions via (co)ends,
- Presheaves are closed under exponentials,
- Associativity of composition of profunctors,
- Right lifts in profunctors,
- (Co)ends preserve limits,
- Adjointness of (co)ends in natural transformations,
- Characterization of dinaturals as certain ends,
- Frobenius property of (co)ends using exponentials.

(Co)end calculus with dinaturality (1)

Yoneda lemma: $(P, \Gamma : \mathbb{C} \rightarrow \mathbf{Set})$

$$\frac{\frac{[a : \mathbb{C}] \quad \Gamma(a) \vdash \int_{x:\mathbb{C}} \text{hom}_{\mathbb{C}}^{\text{op}}(a, \bar{x}) \Rightarrow P(x)}{\frac{[a : \mathbb{C}, x : \mathbb{C}] \quad \Gamma(a) \vdash \text{hom}_{\mathbb{C}}^{\text{op}}(a, \bar{x}) \Rightarrow P(x)}{\frac{[a : \mathbb{C}] \quad \text{hom}_{\mathbb{C}}(\bar{a}, x) \times \Gamma(a) \vdash P(x)}{[z : \mathbb{C}] \quad \Gamma(z) \vdash P(z)} \text{ (hom)}} \text{ (exp)} \quad \text{ (end)}$$

(Co)end calculus with dinaturality (1)

Yoneda lemma: $(P, \Gamma : \mathbb{C} \rightarrow \mathbf{Set})$

$$\begin{array}{c} [a : \mathbb{C}] \Gamma(a) \vdash \int_{x:\mathbb{C}} \text{hom}_{\mathbb{C}}^{\text{op}}(a, \bar{x}) \Rightarrow P(x) \\ \hline \hline [a : \mathbb{C}, x : \mathbb{C}] \Gamma(a) \vdash \text{hom}_{\mathbb{C}}^{\text{op}}(a, \bar{x}) \Rightarrow P(x) \quad (\text{end}) \\ \hline \hline [a : \mathbb{C}] \text{hom}_{\mathbb{C}}(\bar{a}, x) \times \Gamma(a) \vdash P(x) \quad (\text{exp}) \\ \hline \hline [z : \mathbb{C}] \Gamma(z) \vdash P(z) \quad (\text{hom}) \end{array}$$

CoYoneda lemma:

$$\begin{array}{c} [a : \mathbb{C}] \int_{x:\mathbb{C}} \text{hom}_{\mathbb{C}}(\bar{x}, a) \times P(x) \vdash \Gamma(a) \\ \hline \hline [a : \mathbb{C}, x : \mathbb{C}] \text{hom}_{\mathbb{C}}(\bar{a}, x) \times P(a) \vdash \Gamma(x) \quad (\text{coend}) \\ \hline \hline [z : \mathbb{C}] P(z) \vdash \Gamma(z) \quad (\text{hom}) \end{array}$$

(Co)end calculus with dinaturality (2)

Presheaves are cartesian closed: $(\Gamma, A, B : \mathbb{C} \rightarrow \mathbf{Set})$

$$\begin{array}{c}
 [x : \mathbb{C}] \Gamma(x) \vdash (A \Rightarrow B)(x) \\
 \quad := \text{Nat}(\text{hom}_{\mathbb{C}}(x, -) \times A, B) \\
 \quad \cong \int_{y:\mathbb{C}} \text{hom}_{\mathbb{C}}^{\text{op}}(x, \bar{y}) \times A^{\text{op}}(\bar{y}) \Rightarrow B(y) \\
 \hline \hline
 [x : \mathbb{C}, y : \mathbb{C}] \Gamma(x) \vdash \text{hom}_{\mathbb{C}}^{\text{op}}(x, \bar{y}) \times A^{\text{op}}(\bar{y}) \Rightarrow B(y) \quad (\text{end}) \\
 \hline \hline
 [x : \mathbb{C}, y : \mathbb{C}] A(y) \times \text{hom}_{\mathbb{C}}(\bar{x}, y) \times \Gamma(x) \vdash B(y) \quad (\text{exp}) \\
 \hline \hline
 [x : \mathbb{C}, y : \mathbb{C}] A(y) \times \text{hom}_{\mathbb{C}}(\bar{x}, y) \times \Gamma(x) \vdash B(y) \quad (\text{coend+frob.}) \\
 \hline \hline
 [y : \mathbb{C}] A(y) \times \left(\int^{x:\mathbb{C}} \text{hom}_{\mathbb{C}}(\bar{x}, y) \times \Gamma(x) \right) \vdash B(y) \\
 \hline \hline
 [y : \mathbb{C}] A(y) \times \Gamma(y) \vdash B(y) \quad (\text{coYoneda})
 \end{array}$$

(Co)end calculus with dinaturality (3)

Right Kan extensions via ends are right adjoints to precomposition with $F : \mathbb{C} \rightarrow \mathbb{D}$ ($P : \mathbb{C} \rightarrow \mathbf{Set}, \Gamma : \mathbb{D} \rightarrow \mathbf{Set}$):

$$\begin{array}{c}
 [y : \mathbb{D}] \Gamma(y) \vdash (\mathbf{Ran}_F P)(y) \\
 \quad := \int_{x:\mathbb{C}} \mathbf{hom}_{\mathbb{D}}^{\mathbf{op}}(y, F^{\mathbf{op}}(\bar{x})) \Rightarrow P(x) \\
 \hline \hline
 [x : \mathbb{C}, y : \mathbb{D}] \Gamma(y) \vdash \mathbf{hom}_{\mathbb{D}}^{\mathbf{op}}(y, F^{\mathbf{op}}(\bar{x})) \Rightarrow P(x) \quad (\text{end}) \\
 \hline \hline
 [x : \mathbb{C}, y : \mathbb{D}] \mathbf{hom}_{\mathbb{D}}(\bar{y}, F(x)) \times \Gamma(y) \vdash P(x) \quad (\text{exp}) \\
 \hline \hline
 [x : \mathbb{C}, y : \mathbb{D}] \mathbf{hom}_{\mathbb{D}}(\bar{y}, F(x)) \times \Gamma(y) \vdash P(x) \quad (\text{coend}) \\
 \hline \hline
 [x : \mathbb{C}] \int^{y:\mathbb{D}} \mathbf{hom}_{\mathbb{D}}(\bar{y}, F(x)) \times \Gamma(y) \vdash P(x) \quad (\text{coYoneda}) \\
 \hline \hline
 [x : \mathbb{C}] \Gamma(F(x)) \vdash P(x)
 \end{array}$$

(Co)end calculus with dinaturality (4)

Composition (on both sides) in **Prof** has a right adjoint (right lifts).

$(P : \mathbb{C}^{\text{op}} \times \mathbb{A} \rightarrow \mathbf{Set}, Q : \mathbb{A}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set}, \Gamma : \mathbb{C}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set})$:

$$\begin{array}{c}
 [x : \mathbb{C}^{\text{op}}, z : \mathbb{D}] \quad (P ; -)(Q)(x, z) := \\
 \int^{y:\mathbb{A}} P(x, y) \times Q(\bar{y}, z) \vdash \Gamma(x, z) \\
 \hline \hline
 [x : \mathbb{C}^{\text{op}}, y : \mathbb{A}, z : \mathbb{D}] P(x, y) \times Q(\bar{y}, z) \vdash \Gamma(x, z) \quad (\text{coend}) \\
 \hline \hline
 [x : \mathbb{C}^{\text{op}}, y : \mathbb{A}, z : \mathbb{D}] Q(\bar{y}, z) \vdash P^{\text{op}}(\bar{x}, \bar{y}) \Rightarrow \Gamma(x, z) \quad (\text{exp}) \\
 \hline \hline
 [x : \mathbb{C}^{\text{op}}, y : \mathbb{A}, z : \mathbb{D}] Q(\bar{y}, z) \vdash P^{\text{op}}(\bar{x}, \bar{y}) \Rightarrow \Gamma(x, z) \quad (\text{end}) \\
 \hline \hline
 [y : \mathbb{A}, z : \mathbb{D}] Q(\bar{y}, z) \vdash \int_{x:\mathbb{C}} P^{\text{op}}(\bar{x}, \bar{y}) \Rightarrow \Gamma(x, z) \\
 \hline \hline
 [y : \mathbb{A}^{\text{op}}, z : \mathbb{D}] Q(y, z) \vdash \int_{x:\mathbb{C}} P^{\text{op}}(\bar{x}, y) \Rightarrow \Gamma(x, z) \quad (\cong) \\
 \hline \hline
 [y : \mathbb{A}^{\text{op}}, z : \mathbb{D}] Q(y, z) \vdash \int_{x:\mathbb{C}} P^{\text{op}}(\bar{x}, y) \Rightarrow \Gamma(x, z) \\
 \hline \hline
 := \text{Rift}_P(\Gamma)(y, z)
 \end{array}$$

(Co)end calculus with dinaturality (5)

Fubini for ends ($\Gamma : \mathbf{Set}, P : (\mathbb{C}^{\text{op}} \times \mathbb{C}) \times (\mathbb{D}^{\text{op}} \times \mathbb{D}) \rightarrow \mathbf{Set}$)

$$\begin{array}{c}
 [] \Gamma \vdash \int_{x:\mathbb{C}} \int_{y:\mathbb{D}} P(\bar{x}, x, \bar{y}, y) \\
 \hline \hline
 \text{(end)} \\
 [x : \mathbb{C}] \Gamma \vdash \int_{y:\mathbb{D}} P(\bar{x}, x, \bar{y}, y) \\
 \hline \hline
 \text{(end)} \\
 [x : \mathbb{C}, y : \mathbb{D}] \Gamma \vdash P(\bar{x}, x, \bar{y}, y) \\
 \hline \hline
 \text{(structural property)} \\
 [y : \mathbb{D}, x : \mathbb{C}] \Gamma \vdash P(\bar{x}, x, \bar{y}, y) \\
 \hline \hline
 \text{(end)} \\
 [y : \mathbb{D}] \Gamma \vdash \int_{x:\mathbb{C}} P(\bar{x}, x, \bar{y}, y) \\
 \hline \hline
 \text{(end)} \\
 [] \Gamma \vdash \int_{y:\mathbb{D}} \int_{x:\mathbb{C}} P(\bar{x}, x, \bar{y}, y)
 \end{array}$$

(Co)end calculus with dinaturality (6)

Composition of profunctors is associative: $(\Gamma : \mathbb{A}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set},$
 $P : \mathbb{A}^{\text{op}} \times \mathbb{B} \rightarrow \mathbf{Set}, Q : \mathbb{B}^{\text{op}} \times \mathbb{C} \rightarrow \mathbf{Set}, R : \mathbb{C}^{\text{op}} \times \mathbb{D} \rightarrow \mathbf{Set})$

$$\frac{[a : \mathbb{A}, d : \mathbb{D}] \int^{b:\mathbb{B}} P(\bar{a}, b) \times \left(\int^{c:\mathbb{C}} Q(\bar{b}, c) \times R(\bar{c}, d) \right) \vdash \Gamma(\bar{a}, d)}{\text{(coend)}}$$

$$\frac{[a : \mathbb{A}, b : \mathbb{B}, d : \mathbb{D}] P(\bar{a}, b) \times \left(\int^{c:\mathbb{C}} Q(\bar{b}, c) \times R(\bar{c}, d) \right) \vdash \Gamma(\bar{a}, d)}{\text{(coend)}}$$

$$\frac{[a : \mathbb{A}, b : \mathbb{B}, c : \mathbb{C}, d : \mathbb{D}] P(\bar{a}, b) \times (Q(\bar{b}, c) \times R(\bar{c}, d)) \vdash \Gamma(\bar{a}, d)}{\text{(struct. prop.)}}$$

$$\frac{[a : \mathbb{A}, b : \mathbb{B}, c : \mathbb{C}, d : \mathbb{D}] (P(\bar{a}, b) \times Q(\bar{b}, c)) \times R(\bar{c}, d) \vdash \Gamma(\bar{a}, d)}{\text{(coend)}}$$

$$\frac{[a : \mathbb{A}, c : \mathbb{C}, d : \mathbb{D}] \left(\int^{b:\mathbb{B}} P(\bar{a}, b) \times Q(\bar{b}, c) \right) \times R(\bar{c}, d) \vdash \Gamma(\bar{a}, d)}{\text{(coend)}}$$

$$[a : \mathbb{A}, d : \mathbb{D}] \int^{c:\mathbb{C}} \left(\int^{b:\mathbb{B}} P(\bar{a}, b) \times Q(\bar{b}, c) \right) \times R(\bar{c}, d) \vdash \Gamma(\bar{a}, d)$$

Conclusion and future work

*We have seen how dinaturality allows us to give a semantic interpretation towards a directed type theory in **Cat** with quantifiers, where directed equality is given by hom-functors and quantifiers by (co)ends.*

Conclusion and future work

*We have seen how dinaturality allows us to give a semantic interpretation towards a directed type theory in **Cat** with quantifiers, where directed equality is given by hom-functors and quantifiers by (co)ends.*

Future work:

- 1 Big piece missing from the story: compositionality of dinaturals.

Conclusion and future work

*We have seen how dinaturality allows us to give a semantic interpretation towards a directed type theory in **Cat** with quantifiers, where directed equality is given by hom-functors and quantifiers by (co)ends.*

Future work:

- ① Big piece missing from the story: compositionality of dinaturals.
 - *Claim:* non-compositionality is intrinsic to **Cat**, like failure of UIP.

Conclusion and future work

*We have seen how dinaturality allows us to give a semantic interpretation towards a directed type theory in **Cat** with quantifiers, where directed equality is given by hom-functors and quantifiers by (co)ends.*

Future work:

- ① Big piece missing from the story: compositionality of dinaturals.
 - ▶ *Claim:* non-compositionality is intrinsic to **Cat**, like failure of UIP.
 - ▶ Find suitable structures axiomatizing composition of dinaturality (e.g., operads/multicategories but with explicit variances of variables.).

Conclusion and future work

*We have seen how dinaturality allows us to give a semantic interpretation towards a directed type theory in **Cat** with quantifiers, where directed equality is given by hom-functors and quantifiers by (co)ends.*

Future work:

- 1 Big piece missing from the story: compositionality of dinaturals.
 - ▶ *Claim:* non-compositionality is intrinsic to **Cat**, like failure of UIP.
 - ▶ Find suitable structures axiomatizing composition of dinaturality (e.g., operads/multicategories but with explicit variances of variables.).
- 2 Everything shown so far can be done in **Pos**, where all dinats compose.

Conclusion and future work

*We have seen how dinaturality allows us to give a semantic interpretation towards a directed type theory in **Cat** with quantifiers, where directed equality is given by hom-functors and quantifiers by (co)ends.*

Future work:

- ① Big piece missing from the story: compositionality of dinaturals.
 - ▶ *Claim:* non-compositionality is intrinsic to **Cat**, like failure of UIP.
 - ▶ Find suitable structures axiomatizing composition of dinaturality (e.g., operads/multicategories but with explicit variances of variables.).
- ② Everything shown so far can be done in **Pos**, where all dinats compose.
 - ▶ This can be used to characterize *the directed logic of posets*.

Conclusion and future work

*We have seen how dinaturality allows us to give a semantic interpretation towards a directed type theory in **Cat** with quantifiers, where directed equality is given by hom-functors and quantifiers by (co)ends.*

Future work:

- ① Big piece missing from the story: compositionality of dinaturals.
 - ▶ *Claim:* non-compositionality is intrinsic to **Cat**, like failure of UIP.
 - ▶ Find suitable structures axiomatizing composition of dinaturality (e.g., operads/multicategories but with explicit variances of variables.).
- ② Everything shown so far can be done in **Pos**, where all dinats compose.
 - ▶ This can be used to characterize *the directed logic of posets*.
 - ▶ The left relative adjunction characterizes directed equality in **Pos**.
 - ▶ Axiomatize this logic of directed equality using *directed doctrines*.

Conclusion and future work

*We have seen how dinaturality allows us to give a semantic interpretation towards a directed type theory in **Cat** with quantifiers, where directed equality is given by hom-functors and quantifiers by (co)ends.*

Future work:

- ① Big piece missing from the story: compositionality of dinaturals.
 - ▶ *Claim:* non-compositionality is intrinsic to **Cat**, like failure of UIP.
 - ▶ Find suitable structures axiomatizing composition of dinaturality (e.g., operads/multicategories but with explicit variances of variables.).
- ② Everything shown so far can be done in **Pos**, where all dinats compose.
 - ▶ This can be used to characterize *the directed logic of posets*.
 - ▶ The left relative adjunction characterizes directed equality in **Pos**.
 - ▶ Axiomatize this logic of directed equality using *directed doctrines*.
- ③ Long-term future: now that types are categories,
 - ▶ Internalize semantics of type theory inside type theory (e.g., dQIIT).
 - ▶ Naturality for free, since it can be proven internally.

The \int .

Paper: *“Directed equality with dinaturality”* (arXiv:2409.10237)

Agda code: github.com/iwilare/dinaturality

Thank you for the attention!

Directed doctrine (WIP)

- Take a category \mathbb{C} with binary products and terminal.
- The category $\mathbb{C}^{N\Delta P}$ is defined as follows:
 - Objects: $\mathbb{C}_0 \times \mathbb{C}_0 \times \mathbb{C}_0$
 - Morphisms $(N, \Delta, P) \rightarrow (N', \Delta', P')$: triples (n, p, d) where
 - $n : N \times \Delta \rightarrow N'$
 - $p : P \times \Delta \rightarrow P'$
 - $d : \Delta \rightarrow \Delta'$
 - Identities: $(\pi_1, \text{id}_\Delta, \pi_1)$
- **Definition:** a *directed doctrine* is a pseudofunctor $\mathcal{P} : \mathbb{C}^{N\Delta P} \rightarrow \mathbf{Pos}$.
- \mathcal{P} has **products** when each fiber has products.
- \mathcal{P} has **exponentials** when certain left/right-relative adjunctions hold.
- \mathcal{P} has **directed equality** when a certain left relative adjunction holds.
- \mathcal{P} has **involutions** when \mathbb{C} has a functor $-^{\text{op}} : \mathbb{C} \rightarrow \mathbb{C}$ s.t. $\text{op} ; \text{op} = \text{id}_{\mathbb{C}}$ and an iso $\mathcal{P}(P, \Delta, N) \cong \mathcal{P}(N^{\text{op}}, \Delta^{\text{op}}, N^{\text{op}})$ natural in P, Δ, N .

Directed doctrines - relative adj. for exponentials (WIP)

- Downgrade to dinaturality:

$$\frac{[a : ^+ \mathbb{C}, x : \Gamma] \quad A(a, x) \times \Gamma(a, x) \vdash B(a, x)}{[a : ^\Delta \mathbb{C}, x : \Gamma] \quad \uparrow_a^\Delta \Gamma(a, x) \vdash A^{\text{op}}(\bar{a}, x) \Rightarrow \uparrow_a^\Delta B(a, x)} \\ \frac{[a : ^- \mathbb{C}, x : \Gamma] \quad A(\bar{a}, x) \times \Gamma(\bar{a}, x) \vdash B(\bar{a}, x)}{[a : ^\Delta \mathbb{C}, x : \Gamma] \quad \uparrow_a^\Delta \Gamma(\bar{a}, x) \vdash A^{\text{op}}(a, x) \Rightarrow \uparrow_a^\Delta B(\bar{a}, x)}$$

- Dinatural to positive:

$$\frac{[a : ^\Delta \mathbb{C}, x : \Gamma] \quad A(\bar{a}, x) \times \uparrow_a^\Delta \Gamma(a, x) \vdash \uparrow_a^\Delta B(a, x)}{[a : ^+ \mathbb{C}, x : \Gamma] \quad \Gamma(a, x) \vdash A^{\text{op}}(a, x) \Rightarrow B(a, x)}$$

- Inversion:

$$\frac{[a : ^+ \mathbb{C}, x : \Gamma] \quad A(a, x) \times \Gamma(x) \vdash B(x)}{[a : ^- \mathbb{C}, x : \Gamma] \quad \pi_a^- \Gamma(x) \vdash A^{\text{op}}(\bar{a}, x) \Rightarrow \pi_a^- B(x)} \\ \frac{[a : ^- \mathbb{C}, x : \Gamma] \quad A(\bar{a}, x) \times \Gamma(x) \vdash B(x)}{[a : ^+ \mathbb{C}, x : \Gamma] \quad \pi_a^+ \Gamma(x) \vdash A^{\text{op}}(a, x) \Rightarrow \pi_a^+ B(x)}$$

- Dinatural to negative:

$$\frac{[a : ^\Delta \mathbb{C}, x : \Gamma] \quad A(a, x) \times \uparrow_a^\Delta \Gamma(\bar{a}, x) \vdash \uparrow_a^\Delta B(\bar{a}, x)}{[a : ^- \mathbb{C}, x : \Gamma] \quad \Gamma(\bar{a}, x) \vdash A^{\text{op}}(\bar{a}, x) \Rightarrow B(\bar{a}, x)}$$

Homotopical interpretation of dinaturality

We have maps both ways between these entailments:

$$\frac{[] \quad \top \vdash P}{[x : C] \ x = x \vdash P}$$

but in MLTT they are not isomorphic.

In the directed case, we do not even have both maps!

$$\frac{[] \quad \top \vdash P}{[x : \mathbb{C}] \ \text{hom}(\bar{x}, x) \vdash P}$$

We only have a map from top to bottom.

Related works

- **[Caccamo-Winskel 2001]**: axiomatic system to manipulate (co)ends; quantifier exchange is postulated, no type theory presented.
- **[Licata 2011]**: a model in **Cat** with directed equality defined judgementally and not propositionally.
- **[Nuyts 2015]**: a preliminary system of contexts variances is given, with no formal syntax nor models.
- **[North 2018]**: dependent type theory, but uses groupoid structure to type the refl rule. We use dinaturality precisely to avoid this problem.
- **[Riehl-Shulman 2017]**: a synthetic theory of $(\infty, 1)$ -categories with a directed interval type, no model in **Cat**.
- **[New-Licata 2022]**: a DTT with models in virtual equipments, with directed equality and quantifiers, but very different syntax w.r.t. MLTT
- **[Ahrens 2023]**: judgemental structure for directed equality reminiscent of a bicategorical model.
- **[Neumann 2024]**: groupoids are used in contexts rather than in the type of the conclusion.