

MA2001 LINEAR ALGEBRA

MATRICES

Goh Jun Le / Wang Fei

gohjunle@nus.edu.sg / matwf@nus.edu.sg

Department of Mathematics
Office: S17-06-25 / S17-06-16
Tel: 6601-1355 / 6516-2937

Introduction to Matrices	2
Definition of Matrix	3
Special Matrices	7
Matrix Operations	20
Identical Matrices	21
Addition, Subtraction & Scalar Multiplication	23
Properties	25
Matrix Multiplication	28
Properties	32
Powers of Square Matrices	35
Matrix Representation	37
Representation of Linear System	46
Transpose	52
Inverses of Square Matrices	55
Definition	56
Properties	61
Elementary Matrices	67
Elementary Operations	68
Elementary Matrices	71
Connection to Matrix Multiplication	72
Invertibility	85
Main Theorem for Invertible Matrices	90
Find Inverse	93
Column Operations	99
Determinant	107
Determinant of 2×2 Matrix	108
Determinant of 3×3 Matrix	111

Elementary Row Operation	120
Determinant	126
Properties	128
Cofactor Expansion	136
Finding Determinant	140
Adjoint Matrix	144
Cramer's Rule	148

Definition of Matrix

- **Definition.** A **matrix** (plural **matrices**) is a rectangular array of numbers.

$$\circ \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

- m is the number of **rows** in the matrix.
- n is the number of **columns** in the matrix.
- The **size** of the matrix is given by $m \times n$.
- The **(i, j) -entry** is the entry in i th row & j th column.
- In the given matrix, the (i, j) -entry is a_{ij} .
- **Remark.** Some books use $[\cdots]$ instead of (\cdots) .
- Notations $|\cdots|$ and $\|\cdots\|$ are reserved to use later.

3 / 150

Examples

1. $\begin{pmatrix} 1 & 2 \\ 3 & 4 \\ 0 & -1 \end{pmatrix}$ is a 3×2 matrix.
 - The $(1, 2)$ -entry is 2 and the $(3, 1)$ -entry is 0.
2. $\begin{pmatrix} \sqrt{2} & 3.1 & -2 \\ 3 & \frac{1}{2} & 0 \\ 0 & \pi & 0 \end{pmatrix}$ is a 3×3 matrix.
3. $(2 \ 1 \ 0)$ is a 1×3 matrix.
4. $\begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$ is a 3×1 matrix.
5. (4) is a 1×1 matrix.
 - A 1×1 matrix is usually treated as a real number in computation. For instance, $(4) = 4$.

4 / 150

Notation of Matrices

- A matrix is usually denoted by capital letters A, B, C, \dots .

- $m \times n$ matrix $A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}.$

- a_{ij} is the (i, j) -entry of A .
- It is denoted by $A = (a_{ij})_{m \times n}$

Sometimes, if the size of A is known (or not important)

- simple notation: $A = (a_{ij})$

5 / 150

Examples

- Write down the following matrices explicitly.

1. $A = (a_{ij})_{2 \times 3}$, where $a_{ij} = i + j$.

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} = \begin{pmatrix} 1+1 & 1+2 & 1+3 \\ 2+1 & 2+2 & 2+3 \end{pmatrix} \\ = \begin{pmatrix} 2 & 3 & 4 \\ 3 & 4 & 5 \end{pmatrix}$$

2. $B = (b_{ij})_{3 \times 2}$, where $b_{ij} = \begin{cases} 1 & \text{if } i + j \text{ is even} \\ -1 & \text{if } i + j \text{ is odd} \end{cases}$

$$B = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ -1 & 1 \\ 1 & -1 \end{pmatrix}$$

6 / 150

Special Matrices

- A **row matrix** (**row vector**) is a matrix with only one row.
 - $(2 \ 1 \ 0)$ is a row matrix (row vector).
- A **column matrix** (**column vector**) is a matrix with only one column.
 - $\begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$ is a column matrix (column vector).
- A **square matrix** is a matrix with the same number of rows and columns.
 - An $n \times n$ matrix is called a **square matrix** of **order** n .
 - $\begin{pmatrix} 8 & 1 & 6 \\ 3 & 5 & 7 \\ 4 & 9 & 2 \end{pmatrix}$ is a square matrix of order 3.

7 / 150

Special Matrices

- Let $A = (a_{ij})$ be a **square matrix** of order n .
 - The **diagonal** of A is the sequence of entries
 - $a_{11}, a_{22}, \dots, a_{nn}$
 - $A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}$.
 - $a_{ii}, i = 1, \dots, n$, are the **diagonal entries**.
 - $a_{ij}, i \neq j$, are called the **non-diagonal entries**.
 - a_{ij} is $\begin{cases} \text{a diagonal entry} & \text{if } i = j, \\ \text{a non-diagonal entry} & \text{if } i \neq j. \end{cases}$

8 / 150

Special Matrices

- Let $A = (a_{ij})$ be a **square matrix** of order n .
 - The **diagonal** of A is the sequence of entries
 - $a_{11}, a_{22}, \dots, a_{nn}$
 - $$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}.$$
- Remark.** In some textbook,
 - The **diagonal** is also called the **principle diagonal** or **major diagonal**.
 - The **anti-diagonal** or **minor diagonal** refers to the diagonal from the right top to the left bottom.
 - $a_{1n}, a_{2,n-1}, \dots, a_{n1}$.

9 / 150

Special Matrices

- A **square matrix** is called a **diagonal matrix** if all its non-diagonal entries are zero.

$$A = \begin{pmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{pmatrix}.$$

$A = (a_{ij})_{n \times n}$ is diagonal $\Leftrightarrow a_{ij} = 0$ for all $i \neq j$.

Example.

- $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ is a diagonal matrix.
- $\begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix}$, a, b, c are (possibly zero) numbers.

10 / 150

Special Matrices

- A **diagonal matrix** is called a **scalar matrix** if all its diagonal entries are the same.

- $A = \begin{pmatrix} c & 0 & \cdots & 0 \\ 0 & c & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & c \end{pmatrix}$, where c is a constant.

- $A = (a_{ij})_{n \times n}$ is scalar $\Leftrightarrow a_{ij} = \begin{cases} c & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$

Example.

- $\begin{pmatrix} 2001 & 0 \\ 0 & 2001 \end{pmatrix}$ is a scalar matrix.

- $\begin{pmatrix} c & 0 & 0 \\ 0 & c & 0 \\ 0 & 0 & c \end{pmatrix}$, where c is a (possibly zero) number.

11 / 150

Special Matrices

- A **scalar matrix** is called an **identity matrix** if all its diagonal entries are 1.

- Denote the identity matrix of order n (size $n \times n$) by I_n .

- If no confusion in order, write I instead of I_n .

- $A = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}$.

- $A = (a_{ij})_{n \times n}$ is identity $\Leftrightarrow a_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$

- **Note:** There is exactly one identity matrix in order n .

- $I_1 = (1) = 1$; $I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

12 / 150

Special Matrices

- A matrix with all entries equal to zero is a **zero matrix**.

- Denote the zero matrix of size $m \times n$ by $\mathbf{0}_{m \times n}$.
 - If no confusion in size, write $\mathbf{0}$ instead of $\mathbf{0}_{m \times n}$.

- $\mathbf{A} = \begin{pmatrix} 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 0 \end{pmatrix}.$

- $\mathbf{A} = (a_{ij})_{m \times n}$ is zero $\Leftrightarrow a_{ij} = 0$ for all i, j .

Note: There is exactly one zero matrix in size $m \times n$.

- $\mathbf{0}_{1 \times 1} = (0) = 0$; $\mathbf{0}_{1 \times 3} = \begin{pmatrix} 0 & 0 & 0 \end{pmatrix}.$

- $\mathbf{0}_{3 \times 2} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}; \quad \mathbf{0}_{3 \times 4} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$

13 / 150

Special Matrices

- A **square matrix** is called **symmetric** if it is symmetric with respect to the diagonal.

- $\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & a_{24} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & a_{34} & \cdots & a_{3n} \\ a_{41} & a_{42} & a_{43} & a_{44} & \cdots & a_{4n} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & a_{n4} & \cdots & a_{nn} \end{pmatrix}.$

- $\mathbf{A} = (a_{ij})_{n \times n}$ is symmetric $\Leftrightarrow a_{ij} = a_{ji}$ for all i, j .

- (There is no restriction to the diagonal entries.)

- **Examples.**

- (2); $\begin{pmatrix} a & b & c \\ b & d & e \\ c & e & f \end{pmatrix}; \quad \begin{pmatrix} a & b & c & d \\ b & e & f & g \\ c & f & h & i \\ d & g & i & j \end{pmatrix}$

14 / 150

Special Matrices

- A **square matrix** is called **upper triangular** if all the entries **below** the diagonal are zero.

$$\circ \quad A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} & \cdots & a_{1n} \\ 0 & a_{22} & a_{23} & a_{24} & \cdots & a_{2n} \\ 0 & 0 & a_{33} & a_{34} & \cdots & a_{3n} \\ 0 & 0 & 0 & a_{44} & \cdots & a_{4n} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & a_{nn} \end{pmatrix}.$$

$$\circ \quad A = (a_{ij})_{n \times n} \text{ is upper triangular} \Leftrightarrow a_{ij} = 0 \text{ if } i > j.$$

- (There is no restriction to the diagonal entries.)

- Examples.**

$$\circ \quad (2); \quad \begin{pmatrix} a & b & c \\ 0 & d & e \\ 0 & 0 & f \end{pmatrix}; \quad \begin{pmatrix} a & b & c & d \\ 0 & e & f & g \\ 0 & 0 & h & i \\ 0 & 0 & 0 & j \end{pmatrix}$$

15 / 150

Special Matrices

- A **square matrix** is called **lower triangular** if all the entries **above** the diagonal are zero.

$$\circ \quad A = \begin{pmatrix} a_{11} & 0 & 0 & 0 & \cdots & 0 \\ a_{21} & a_{22} & 0 & 0 & \cdots & 0 \\ a_{31} & a_{32} & a_{33} & 0 & \cdots & 0 \\ a_{41} & a_{42} & a_{43} & a_{44} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & a_{n4} & \cdots & a_{nn} \end{pmatrix}.$$

$$\circ \quad A = (a_{ij})_{n \times n} \text{ is lower triangular} \Leftrightarrow a_{ij} = 0 \text{ if } i < j.$$

- (There is no restriction to the diagonal entries.)

- Examples.**

$$\circ \quad (2); \quad \begin{pmatrix} a & 0 & 0 \\ b & d & 0 \\ c & e & f \end{pmatrix}; \quad \begin{pmatrix} a & 0 & 0 & 0 \\ b & e & 0 & 0 \\ c & f & h & 0 \\ d & g & i & j \end{pmatrix}$$

16 / 150

Special Matrices

- Both **upper triangular matrices** and **lower triangular matrices** are called **triangular matrices**.

- (A matrix is both upper and lower triangular

\Leftrightarrow it is diagonal.)

- More Examples.**

- Square matrices:

$$\bullet \begin{pmatrix} 4 \end{pmatrix} \begin{pmatrix} 0 & 4 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ -1 & 3 & 2 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & 1 & 6 & 2 \\ 0 & 3 & 9 & -1 \\ 1 & 0 & 0 & 0 \\ -2 & 3 & 0 & 1 \end{pmatrix}$$

- Diagonal matrices:

$$\bullet \begin{pmatrix} 4 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

17 / 150

Special Matrices

- More Examples.**

- Scalar matrices:

$$\bullet \begin{pmatrix} 4 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix}$$

- Identity matrices:

$$\bullet \begin{pmatrix} 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

- Zero matrices:

$$\bullet \begin{pmatrix} 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

18 / 150

Special Matrices

- **More Examples.**

- Symmetric matrices:

- $\begin{pmatrix} 4 \end{pmatrix} \begin{pmatrix} 0 & 4 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} 1 & -1 & 0 \\ -1 & 3 & 2 \\ 0 & 2 & 2 \end{pmatrix} \begin{pmatrix} 2 & 1 & 6 & -2 \\ 1 & 3 & 0 & -1 \\ 6 & 0 & 0 & 0 \\ -2 & -1 & 0 & 1 \end{pmatrix}$

- Upper triangular matrices:

- $\begin{pmatrix} 4 \end{pmatrix} \begin{pmatrix} 0 & 4 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & -1 & 0 \\ 0 & 3 & 2 \\ 0 & 0 & 2 \end{pmatrix}$

- Lower triangular matrix:

- $\begin{pmatrix} 2 & 0 & 0 & 0 \\ 1 & 3 & 0 & 0 \\ 6 & 0 & 0 & 0 \\ -2 & -1 & 0 & 1 \end{pmatrix}$

19 / 150

Matrix Operations

20 / 150

Identical Matrices

- A matrix is completely determined by its size and entries.
- **Definition.** Two matrices are **equal** if
 - they have the same size (same number of rows, same number of columns), and
 - all the corresponding entries are the same.

Let $A = (a_{ij})_{m \times n}$ and $B = (b_{ij})_{p \times q}$. Then

- $A = B \Leftrightarrow m = p \text{ \& } n = q \text{ \& } a_{ij} = b_{ij} \text{ for all } i, j$

- **Examples.**

- $\mathbf{0}_{2 \times 2} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}; \quad \mathbf{0}_{2 \times 3} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$

- $\mathbf{0}_{2 \times 2}$ and $\mathbf{0}_{2 \times 3}$ have different size $\Rightarrow \mathbf{0}_{2 \times 2} \neq \mathbf{0}_{2 \times 3}.$

21 / 150

Identical Matrices

- A matrix is completely determined by its size and entries.
- **Definition.** Two matrices are **equal** if
 - they have the same size (same number of rows, same number of columns), and
 - all the corresponding entries are the same.

Let $A = (a_{ij})_{m \times n}$ and $B = (b_{ij})_{p \times q}$. Then

$$\circ A = B \Leftrightarrow m = p \ \& \ n = q \ \& \ a_{ij} = b_{ij} \text{ for all } i, j$$

- **Examples.**

$$\circ A = \begin{pmatrix} x & y \\ z & w \end{pmatrix}; \quad B = \begin{pmatrix} 1 & -1 \\ 2 & 4 \end{pmatrix}.$$

$$\bullet A = B \Leftrightarrow \begin{cases} x = 1 \\ y = -1 \\ z = 2 \\ w = 4 \end{cases}$$

22 / 150

Addition, Subtraction & Scalar Multiplication

- Let $A = (a_{ij})_{m \times n}$ and $B = (b_{ij})_{m \times n}$ be matrices.
 - **Addition:** $A + B = (a_{ij} + b_{ij})_{m \times n}$.
 - **Subtraction:** $A - B = (a_{ij} - b_{ij})_{m \times n}$.

- **Example.**

$$\circ \text{ Let } A = \begin{pmatrix} 2 & 3 & 4 \\ 4 & 5 & 6 \end{pmatrix}, B = \begin{pmatrix} 1 & 2 & 3 \\ -1 & -1 & -1 \end{pmatrix}.$$

$$\begin{aligned} A + B &= \begin{pmatrix} 2+1 & 3+2 & 4+3 \\ 4+(-1) & 5+(-1) & 6+(-1) \end{pmatrix} \\ &= \begin{pmatrix} 3 & 5 & 7 \\ 3 & 4 & 5 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} A - B &= \begin{pmatrix} 2-1 & 3-2 & 4-3 \\ 4-(-1) & 5-(-1) & 6-(-1) \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 & 1 \\ 5 & 6 & 7 \end{pmatrix} \end{aligned}$$

23 / 150

Addition, Subtraction & Scalar Multiplication

- Let $A = (a_{ij})_{m \times n}$ be a matrix, and c a constant.

- **Scalar multiplication:** $cA = (ca_{ij})_{m \times n}$.

- **Example.**

- Let $A = \begin{pmatrix} 2 & 3 & 4 \\ 4 & 5 & 6 \end{pmatrix}$. Then

$$4A = \begin{pmatrix} 4 \cdot 2 & 4 \cdot 3 & 4 \cdot 4 \\ 4 \cdot 4 & 4 \cdot 5 & 4 \cdot 6 \end{pmatrix} = \begin{pmatrix} 8 & 12 & 16 \\ 16 & 20 & 24 \end{pmatrix}$$

- **Remarks.**

- $(-1)A$ is usually denoted by $-A$.
- It can be proved that $A - B = A + (-B)$.
 - In the discussion we usually only consider addition and scalar multiplication.

24 / 150

Properties

- **Theorem.** Let A, B, C be matrices of the same size.

- $A - B = A + (-B)$.
- Commutative Law for Matrix Addition:
 - $A + B = B + A$.
- Associative Law for Matrix Addition:
 - $(A + B) + C = A + (B + C)$.
- Let 0 be the zero matrix of the same size as A .
 - $0 + A = A$; $A - A = 0$; $0A = 0$; $c0 = 0$.
- Distributive Law for Scalar Multiplication over Addition:
 - $c(A + B) = cA + cB$.
 - $(c + d)A = cA + dA$.
- $c(dA) = (cd)A$, $1A = A$.

25 / 150

Properties

- Let A and B be matrices of the same size.

- Prove that $A - B = A + (-B)$.

Proof. Suppose that $A = (a_{ij})_{m \times n}$ and $B = (b_{ij})_{m \times n}$.

1. Verify that LHS and RHS have the same size.

- A is $m \times n$ & B is $m \times n \Rightarrow A - B$ is $m \times n$.
- B is $m \times n \Rightarrow -B$ is $m \times n$.
- $A, -B$ are $m \times n \Rightarrow A + (-B)$ is $m \times n$.

2. Verify: “ (i, j) -entry of LHS” = “ (i, j) -entry of RHS”.

$$\begin{aligned}(i, j)\text{-entry of } (A - B) &= a_{ij} - b_{ij} \\ &= a_{ij} + (-b_{ij}) \\ &= a_{ij} + (i, j)\text{-entry of } (-B) \\ &= (i, j)\text{-entry of } [A + (-B)]\end{aligned}$$

26 / 150

Properties

- Let A, B, C be matrices of the same size.

- Prove that $A + (B + C) = (A + B) + C$.

Proof. Let $A = (a_{ij})_{m \times n}$, $B = (b_{ij})_{m \times n}$, $C = (c_{ij})_{m \times n}$.

1. It is clear that LHS & RHS are both $m \times n$. (Why?)
2. Verify: “ (i, j) -entry of LHS” = “ (i, j) -entry of RHS”.

$$\begin{aligned}(i, j)\text{-entry of } A + (B + C) &= a_{ij} + [(i, j)\text{-entry of } B + C] \\ &= a_{ij} + (b_{ij} + c_{ij}) \\ &= (a_{ij} + b_{ij}) + c_{ij} \\ &= [(i, j)\text{-entry of } A + B] + c_{ij} \\ &= (i, j)\text{-entry of } (A + B) + C.\end{aligned}$$

$$\therefore A + (B + C) = (A + B) + C.$$

- **Exercise:** Prove the remaining properties. (Question 2.18)

27 / 150

Matrix Multiplication

- Consider the following linear systems:

$$\circ \begin{cases} a_{11}y_1 + a_{12}y_2 = z_1 \\ a_{21}y_1 + a_{22}y_2 = z_2 \\ a_{31}y_1 + a_{32}y_2 = z_3 \end{cases} \quad \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}$$

$$\circ \begin{cases} b_{11}x_1 + b_{12}x_2 + b_{13}x_3 = y_1 \\ b_{21}x_1 + b_{22}x_2 + b_{23}x_3 = y_2 \end{cases} \quad \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix}$$

$$\begin{cases} (a_{11}b_{11} + a_{12}b_{21})x_1 + (a_{11}b_{12} + a_{12}b_{22})x_2 + (a_{11}b_{13} + a_{12}b_{23})x_3 = z_1 \\ (a_{21}b_{11} + a_{22}b_{21})x_1 + (a_{21}b_{12} + a_{22}b_{22})x_2 + (a_{21}b_{13} + a_{22}b_{23})x_3 = z_2 \\ (a_{31}b_{11} + a_{32}b_{21})x_1 + (a_{31}b_{12} + a_{32}b_{22})x_2 + (a_{31}b_{13} + a_{32}b_{23})x_3 = z_3 \end{cases}$$

Can we use the coefficient matrices of the first two linear systems to obtain the coefficient matrix of their composite?

$$\circ \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} & a_{11}b_{13} + a_{12}b_{23} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} & a_{21}b_{13} + a_{22}b_{23} \\ a_{31}b_{11} + a_{32}b_{21} & a_{31}b_{12} + a_{32}b_{22} & a_{31}b_{13} + a_{32}b_{23} \end{pmatrix}$$

28 / 150

Matrix Multiplication

- Define matrix multiplication so that the **product** of

$$\circ \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix} \text{ and } \begin{pmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{pmatrix} \text{ is given by}$$

$$\circ \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} & a_{11}b_{13} + a_{12}b_{23} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} & a_{21}b_{13} + a_{22}b_{23} \\ a_{31}b_{11} + a_{32}b_{21} & a_{31}b_{12} + a_{32}b_{22} & a_{31}b_{13} + a_{32}b_{23} \end{pmatrix}$$

The **(3, 2)**-entry of the product is $a_{31}b_{12} + a_{32}b_{22}$.

- a_{31} and a_{32} are from the **3rd row** of the first matrix.
- b_{12} and b_{22} are from the **2nd column** of the second.

In order to get the **(i, j)**-entry of the **product** matrix:

- Find the **i**th **row** of the first matrix;
- Find the **j**th **column** of the second matrix;
- Multiply the corresponding entries.
- Add the products together.

29 / 150

Matrix Multiplication

- **Definition.** Let $A = (a_{ij})_{m \times p}$ and $B = (b_{ij})_{p \times n}$.
 - AB is the $m \times n$ matrix such that its (i, j) -entry is

$$a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ip}b_{pj} = \sum_{k=1}^p a_{ik}b_{kj}$$

Note: No. of columns of A = the no. of rows of B .

1. i th row of A : $\begin{pmatrix} \vdots \\ a_{i1} & a_{i2} & \cdots & a_{ip} \\ \vdots \end{pmatrix}$

2. j th column of B : $\begin{pmatrix} b_{1j} \\ b_{2j} \\ \vdots \\ b_{pj} \end{pmatrix}$

3. Multiply componentwise and add the products.

30 / 150

Examples

- $\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 2 & 3 \\ -1 & -2 \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 8 & 7 \end{pmatrix}$
 - $\begin{pmatrix} 1 \cdot 1 + 2 \cdot 2 + 3 \cdot (-1) & 1 \cdot 1 + 2 \cdot 3 + 3 \cdot (-2) \\ 4 \cdot 1 + 5 \cdot 2 + 6 \cdot (-1) & 4 \cdot 1 + 5 \cdot 3 + 6 \cdot (-2) \end{pmatrix}$
- $\begin{pmatrix} 1 & 1 \\ 2 & 3 \\ -1 & -2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} = \begin{pmatrix} 5 & 7 & 9 \\ 14 & 19 & 24 \\ -9 & -12 & -15 \end{pmatrix}$
 - $\begin{pmatrix} 1 \cdot 1 + 1 \cdot 4 & 1 \cdot 2 + 1 \cdot 5 & 1 \cdot 3 + 1 \cdot 6 \\ 2 \cdot 1 + 3 \cdot 4 & 2 \cdot 2 + 3 \cdot 5 & 2 \cdot 3 + 3 \cdot 6 \\ (-1) \cdot 1 + (-2) \cdot 4 & (-1) \cdot 2 + (-2) \cdot 5 & (-1) \cdot 3 + (-2) \cdot 6 \end{pmatrix}$
- **Remark.** Matrix multiplication is **NOT commutative**.
 - AB is the **pre-multiplication** of A to B (to B by A).
 - BA is the **post-multiplication** of A to B (to B by A).

31 / 150

Properties

- **Theorem.**

- Let A, B, C be $m \times p, p \times q, q \times n$ matrices, respectively.
 - Associative Law: $A(BC) = (AB)C$.
- Let A be $m \times p$ matrix, B_1, B_2 be $p \times n$ matrices.
 - Distributive Law: $A(B_1 + B_2) = AB_1 + AB_2$.
- Let A_1, A_2 be $m \times p$ matrices, B be $p \times n$ matrix.
 - Distributive Law: $(A_1 + A_2)B = A_1B + A_2B$.
- Let A be $m \times p$ and B be $p \times n$. For constant c ,
 - $c(AB) = (cA)B = A(cB)$.
- Let A be an $m \times n$ matrix.
 - $A\mathbf{0}_{n \times p} = \mathbf{0}_{m \times p}$; $\mathbf{0}_{p \times m}A = \mathbf{0}_{p \times n}$.
 - $AI_n = A$; $I_mA = A$.

32 / 150

Properties

- Let A be an $m \times p$ matrix and B_1, B_2 be $p \times n$ matrices.
 - Prove that $A(B_1 + B_2) = AB_1 + AB_2$.

Proof. Verify that LHS and RHS have the same size.

- B_1 and B_2 are $p \times n \Rightarrow B_1 + B_2$ is $p \times n$.
 $\Rightarrow A(B_1 + B_2)$ is $m \times n$.
- AB_1 is $m \times n$, and AB_2 is $m \times n$
 $\Rightarrow AB_1 + AB_2$ is $m \times n$.

Let $A = (a_{ij})_{m \times p}$, $B_1 = (b_{ij})_{p \times n}$ and $B_2 = (b'_{ij})_{p \times n}$.

- We shall verify that the following are equal:
 - “the (i, j) -entry of $A(B_1 + B_2)$ ” and
 - “the (i, j) -entry of $AB_1 + AB_2$ ”,for all $i = 1, \dots, m, j = 1, \dots, n$.

33 / 150

Properties

- Let A be an $m \times p$ matrix and B_1, B_2 be $p \times n$ matrices.

- Prove that $A(B_1 + B_2) = AB_1 + AB_2$.

Proof. Let $A = (a_{ij})$, $B_1 = (b_{ij})$, $B_2 = (b'_{ij})$.

- $B_1 + B_2 = (b_{ij} + b'_{ij})$. Let $b''_{ij} = b_{ij} + b'_{ij}$.

$$\begin{aligned}
 & (i, j)\text{-entry of } A(B_1 + B_2) \\
 &= a_{i1}b''_{1j} + a_{i2}b''_{2j} + \cdots + a_{ip}b''_{pj} \\
 &= a_{i1}(b_{1j} + b'_{1j}) + a_{i2}(b_{2j} + b'_{2j}) + \cdots + a_{ip}(b_{pj} + b'_{pj}) \\
 &= (a_{i1}b_{1j} + a_{i1}b'_{1j}) + (a_{i2}b_{2j} + a_{i2}b'_{2j}) + \cdots + (a_{ip}b_{pj} + a_{ip}b'_{pj}) \\
 &= (a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ip}b_{pj}) + (a_{i1}b'_{1j} + a_{i2}b'_{2j} + \cdots + a_{ip}b'_{pj}) \\
 &= (i, j)\text{-entry of } AB_1 + (i, j)\text{-entry of } AB_2 \\
 &= (i, j)\text{-entry of } (AB_1 + AB_2).
 \end{aligned}$$

$$\therefore A(B_1 + B_2) = AB_1 + AB_2.$$

34 / 150

Powers of Square Matrices

- Let A be an $m \times n$ matrix.
 - AA is well-defined $\Leftrightarrow m = n \Leftrightarrow A$ is square.

Definition. Let A be a **square** matrix of order n . For nonnegative integers k , the **powers** of A are defined as

$$A^k = \begin{cases} I_n & \text{if } k = 0, \\ \underbrace{AA \cdots A}_{k \text{ times}} & \text{if } k \geq 1. \end{cases}$$

- Example.** Let $A = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix}$. Then
 - $A^2 = AA = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} = \begin{pmatrix} 3 & 8 \\ 4 & 11 \end{pmatrix}$.
 - $A^3 = AAA = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} = \begin{pmatrix} 11 & 30 \\ 15 & 41 \end{pmatrix}$.

35 / 150

Properties

- Let A be a square matrix, and m, n nonnegative integers.
 - $A^m A^n = A^{m+n}$, $(A^m)^n = A^{mn}$.
- Recall that matrix multiplication is NOT commutative.
 - In general, $(AB)^n \neq A^n B^n$ for $n = 2, 3, \dots$.
 - For example, let $A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.
 - $(AB)^2 = (AB)(AB) = ABAB = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$.
 - $A^2 B^2 = (AA)(BB) = AAB B = \begin{pmatrix} 1 & 0 \\ 0 & 4 \end{pmatrix}$.
- **Exercise.** Let A, B be square matrices of the same size.
 - Suppose that $AB = BA$. Prove that
 - $(AB)^n = A^n B^n$ for all nonnegative integers n .

36 / 150

Matrix Representation

- Let $A = (a_{ij})_{m \times n} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$.
 - Let \mathbf{a}_i denote the i th row of A , $i = 1, \dots, m$.
 - $\mathbf{a}_1 = (a_{11} \ a_{12} \ \cdots \ a_{1n})$
 - $\mathbf{a}_2 = (a_{21} \ a_{22} \ \cdots \ a_{2n})$
 - $\dots\dots\dots$
 - $\mathbf{a}_m = (a_{m1} \ a_{m2} \ \cdots \ a_{mn})$

Then each \mathbf{a}_i is a $1 \times n$ matrix (row vector).

- $A = \begin{pmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \vdots \\ \mathbf{a}_m \end{pmatrix}$.

37 / 150

Matrix Representation

- Let $A = (a_{ij})_{m \times n} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$.
 - Let b_j denote the j th column of A , $j = 1, \dots, n$.
 - $b_1 = \begin{pmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{pmatrix}, b_2 = \begin{pmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{pmatrix}, \dots, b_n = \begin{pmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{pmatrix}$.

Then each b_j is an $m \times 1$ matrix (column vector).

- $A = (b_1 \ b_2 \ \cdots \ b_n)$.

38 / 150

Matrix Representation

- Let $a = (a_1 \ a_2 \ \cdots \ a_n)$ and $b = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$.
 - Then ab is a 1×1 matrix, i.e., a real number.
 - $ab = a_1b_1 + a_2b_2 + \cdots + a_nb_n$.
 - Note:** ba is an $n \times n$ matrix.
 - (i, j) -entry is the i th entry of b times the j th entry of a .
 - $ba = \begin{pmatrix} b_1a_1 & b_1a_2 & \cdots & b_1a_n \\ b_2a_1 & b_2a_2 & \cdots & b_2a_n \\ \vdots & \vdots & \ddots & \vdots \\ b_na_1 & b_na_2 & \cdots & b_na_n \end{pmatrix}$.

39 / 150

Matrix Representation

- Suppose $A = (a_{ij})_{m \times p}$.
 - Let $a_i = (a_{i1} \ a_{i2} \ \cdots \ a_{ip})$ be the i th row of A .

Suppose $B = (b_{ij})_{p \times n}$.

- Let $b_j = \begin{pmatrix} b_{1j} \\ b_{2j} \\ \vdots \\ b_{pj} \end{pmatrix}$ be the j th column of B .

$$\begin{aligned} a_i b_j &= a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ip}b_{pj} \\ &= (i, j)\text{-entry of } AB. \end{aligned}$$

- $AB = \begin{pmatrix} a_1 b_1 & a_1 b_2 & \cdots & a_1 b_n \\ a_2 b_1 & a_2 b_2 & \cdots & a_2 b_n \\ \vdots & \vdots & & \vdots \\ a_m b_1 & a_m b_2 & \cdots & a_m b_n \end{pmatrix}.$

40 / 150

Matrix Representation

- The (i, j) -entry of AB is
 - The i th row of A times the j th column of B .

$$\begin{aligned} a_i B &= a_i (b_1 \ b_2 \ \cdots \ b_n) \\ &= (a_i b_1 \ a_i b_2 \ \cdots \ a_i b_n) \\ &= i\text{th row of } AB \end{aligned}$$

$$AB = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{pmatrix} B = \begin{pmatrix} a_1 B \\ a_2 B \\ \vdots \\ a_m B \end{pmatrix}.$$

41 / 150

Matrix Representation

- The (i, j) -entry of AB is
 - The i th row of A times the j th column of B .

$$Ab_j = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{pmatrix} b_j = \begin{pmatrix} a_1 b_j \\ a_2 b_j \\ \vdots \\ a_m b_j \end{pmatrix} = j\text{th column of } AB.$$

$$\begin{aligned} AB &= A (b_1 \quad b_2 \quad \cdots \quad b_n) \\ &= (Ab_1 \quad Ab_2 \quad \cdots \quad Ab_n) \end{aligned}$$

- **Remark.** Matrices can be multiplied in blocks (provided that the sizes are matched).
 - Reference: Question 2.23.

42 / 150

Example

- Let $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 1 \\ 2 & 3 \\ -1 & -2 \end{pmatrix}$.
 - Let $a_1 = (1 \quad 2 \quad 3)$, $a_2 = (4 \quad 5 \quad 6)$. Then

$$\begin{aligned} AB &= \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} B = \begin{pmatrix} a_1 B \\ a_2 B \end{pmatrix} \\ &= \begin{pmatrix} (1 \quad 2 \quad 3) \begin{pmatrix} 1 & 1 \\ 2 & 3 \\ -1 & -2 \end{pmatrix} \\ (4 \quad 5 \quad 6) \begin{pmatrix} 1 & 1 \\ 2 & 3 \\ -1 & -2 \end{pmatrix} \end{pmatrix} \\ &= \begin{pmatrix} (2 \quad 1) \\ (8 \quad 7) \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 8 & 7 \end{pmatrix} \end{aligned}$$

43 / 150

Example

- Let $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 1 \\ 2 & 3 \\ -1 & -2 \end{pmatrix}$.

- Let $b_1 = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$ and $b_2 = \begin{pmatrix} 1 \\ 3 \\ -2 \end{pmatrix}$. Then

$$\begin{aligned}
 AB &= A(b_1 \ b_2) = (Ab_1 \ Ab_2) \\
 &= \left(\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \begin{pmatrix} 1 \\ 3 \\ -2 \end{pmatrix} \right) \\
 &= \left(\begin{pmatrix} 2 \\ 8 \end{pmatrix} \quad \begin{pmatrix} 1 \\ 7 \end{pmatrix} \right) = \begin{pmatrix} 2 & 1 \\ 8 & 7 \end{pmatrix}
 \end{aligned}$$

44 / 150

Example

- Let $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 1 \\ 2 & 3 \\ -1 & -2 \end{pmatrix}$.

$$\begin{aligned}
 AB &= \begin{pmatrix} a_1b_1 & a_1b_2 \\ a_2b_1 & a_2b_2 \end{pmatrix} \\
 &= \left(\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \begin{pmatrix} 1 \\ 3 \\ -2 \end{pmatrix} \right) \\
 &= \begin{pmatrix} 2 & 1 \\ 8 & 7 \end{pmatrix}
 \end{aligned}$$

45 / 150

Representation of Linear System

- Linear System of m equations in n variables x_1, \dots, x_n :

$$\circ \begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

$$b_i = a_{i1}x_1 + a_{i2}x_2 + \cdots + a_{in}x_n$$

$$= (a_{i1} \quad a_{i2} \quad \cdots \quad a_{in}) \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

46 / 150

Representation of Linear System

- Linear System of m equations in n variables x_1, \dots, x_n .

$$\circ \begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

$$\circ \mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}, \text{ coefficient matrix.}$$

$$\bullet \mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \text{ variable matrix.}$$

$$\bullet \mathbf{b} = \begin{pmatrix} b_1 \\ \vdots \\ b_m \end{pmatrix}, \text{ constant matrix. Then } \boxed{\mathbf{Ax} = \mathbf{b}}$$

47 / 150

Representation of Linear System

- Let $A = (a_{ij})_{m \times n}$, $\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} b_1 \\ \vdots \\ b_m \end{pmatrix}$.
 - Then $A\mathbf{x} = \mathbf{b}$ is the linear system of
 - m linear equations in n variables x_1, \dots, x_n ,
 - a_{ij} are the coefficients, and b_i are the constants.
 - Let $\mathbf{u} = \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix}$.
 - $x_1 = u_1, \dots, x_n = u_n$ is a solution to the system
 - $\Leftrightarrow A\mathbf{u} = \mathbf{b}$
 - $\Leftrightarrow \mathbf{u}$ is a solution to $A\mathbf{x} = \mathbf{b}$.

48 / 150

Representation of Linear System

- **Problem.** Suppose that a linear system has more than one solutions. Then it has infinitely many solutions.
Proof. Let the system be represented by $A\mathbf{x} = \mathbf{b}$.
 - Suppose that it has two solutions $\mathbf{u}_1 \neq \mathbf{u}_2$.
 - $A\mathbf{u}_1 = \mathbf{b}$ and $A\mathbf{u}_2 = \mathbf{b}$.Consider $c_1\mathbf{u}_1 + c_2\mathbf{u}_2$, where c_1, c_2 are constants.
 - $A(c_1\mathbf{u}_1 + c_2\mathbf{u}_2) = c_1A\mathbf{u}_1 + c_2A\mathbf{u}_2 = c_1\mathbf{b} + c_2\mathbf{b} = (c_1 + c_2)\mathbf{b}$.In particular,
 - If $c_1 + c_2 = 1$, then $A(c_1\mathbf{u}_1 + c_2\mathbf{u}_2) = 1\mathbf{b} = \mathbf{b}$.Therefore, $A\mathbf{x} = \mathbf{b}$ has infinitely many solutions:
 - $\mathbf{x} = c_1\mathbf{u}_1 + c_2\mathbf{u}_2$, where $c_1 + c_2 = 1$.

49 / 150

Representation of Linear System

- Consider $\begin{cases} 4x + 5y + 6z = 5 \\ x - y = 2 \\ y - z = 3. \end{cases}$

$$\begin{aligned} \begin{pmatrix} 5 \\ 2 \\ 3 \end{pmatrix} &= \begin{pmatrix} 4 & 5 & 6 \\ 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} \\ &= \begin{pmatrix} 4x + 5y + 6z \\ x - y \\ y - z \end{pmatrix} \\ &= \begin{pmatrix} 4x \\ x \\ 0 \end{pmatrix} + \begin{pmatrix} 5y \\ -y \\ y \end{pmatrix} + \begin{pmatrix} 6z \\ 0 \\ -z \end{pmatrix} \\ &= x \begin{pmatrix} 4 \\ 1 \\ 0 \end{pmatrix} + y \begin{pmatrix} 5 \\ -1 \\ 1 \end{pmatrix} + z \begin{pmatrix} 6 \\ 0 \\ -1 \end{pmatrix} \end{aligned}$$

50 / 150

Representation of Linear System

- $\mathbf{A} = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix}, \mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \mathbf{b} = \begin{pmatrix} b_1 \\ \vdots \\ b_m \end{pmatrix}.$

$$\begin{aligned} \begin{pmatrix} b_1 \\ \vdots \\ b_m \end{pmatrix} &= \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} a_{11}x_1 + \cdots + a_{1n}x_n \\ \vdots \\ a_{m1}x_1 + \cdots + a_{mn}x_n \end{pmatrix} \\ &= x_1 \begin{pmatrix} a_{11} \\ \vdots \\ a_{m1} \end{pmatrix} + \cdots + x_n \begin{pmatrix} a_{1n} \\ \vdots \\ a_{mn} \end{pmatrix} \end{aligned}$$

- Let \mathbf{a}_j denote the j th column of \mathbf{A} . Then

- $\mathbf{b} = \mathbf{Ax} = x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n = \sum_{j=1}^n x_j\mathbf{a}_j.$

51 / 150

Transpose

- Let $A = (a_{ij})_{m \times n}$ be a matrix.
 - The **transpose** of A is the $n \times m$ matrix A^T (or A^t)
 - whose (i, j) -entry is a_{ji} .
- **Example.** Let $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{pmatrix}$.
 - $A^T = \begin{pmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{pmatrix}$ and $(A^T)^T = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{pmatrix}$.
- **Remarks.**
 - The i th row of A^T is the i th column of A .
 - The j th column of A^T is the j th row of A .

52 / 150

Properties

- **Theorem.** Let A be an $m \times n$ matrix.
 - $(A^T)^T = A$.
 - A is symmetric $\Leftrightarrow A = A^T$.
 - Let c be a scalar. Then $(cA)^T = cA^T$.
 - Let B be $m \times n$. Then $(A + B)^T = A^T + B^T$.
 - Let B be $n \times p$. Then $(AB)^T = B^T A^T$.
 - **Proof.** We only prove the last statement.
 - Left-hand side:
 - AB is $m \times p \Rightarrow (AB)^T$ is $p \times m$.
 - Right-hand side:
 - B^T is $p \times n$ & A^T is $n \times m \Rightarrow B^T A^T$ is $p \times m$.
- So $(AB)^T$ and $B^T A^T$ have the same size.

53 / 150

Properties

- **Proof.** (Continued) Let $A = (a_{ij})_{m \times n}$ and $B = (b_{ij})_{n \times p}$.

$$\begin{aligned}(i, j)\text{-entry of } (AB)^T &= (j, i)\text{-entry of } AB \\ &= a_{j1}b_{1i} + a_{j2}b_{2i} + \cdots + a_{jn}b_{ni}.\end{aligned}$$

- Let $A^T = (a'_{ij})_{n \times m}$ and $B^T = (b'_{ij})_{p \times n}$.
 - $a'_{ij} = a_{ji}$ and $b'_{ij} = b_{ji}$.

$$\begin{aligned}(i, j)\text{-entry of } B^T A^T &= b'_{i1}a'_{1j} + b'_{i2}a'_{2j} + \cdots + b'_{in}a'_{nj} \\ &= b_{1i}a_{j1} + b_{2i}a_{j2} + \cdots + b_{ni}a_{jn} \\ &= a_{j1}b_{1i} + a_{j2}b_{2i} + \cdots + a_{jn}b_{ni}.\end{aligned}$$

$$\therefore (AB)^T = B^T A^T.$$

Note: In general, $(AB)^T \neq A^T B^T$.

54 / 150

Inverses of Square Matrices

55 / 150

Inverses of Numbers

- Let a and b be real numbers.
 - $a + x = b \Rightarrow x = b - a = b + (-a)$.
The number $-a$ is the **additive inverse** of a .
 - $ax = b \Rightarrow x = b/a = a^{-1}b$, provided that $a \neq 0$.
The number a^{-1} is the **multiplicative inverse** of a ($\neq 0$).

- Let A and B be matrices of the same size.
 - $A + X = B \Rightarrow X = B - A = B + (-A)$.
So $-A$ is the **additive inverse** of A .

- Let A be an $m \times n$ matrix and B be an $m \times p$ matrix.
 - $AX = B \Rightarrow X = \dots$
It is expected to have a matrix A^{-1} so that $X = A^{-1}B$.

56 / 150

Inverses of Square Matrices

- **Definition.** Let A be a **square matrix** of order n .
 - If there exists a square matrix B of order n so that
 - $AB = I_n$ and $BA = I_n$
 then A is called **invertible**, and B is an **inverse** of A .
 - If A is not **invertible**, A is called **singular**.

Note: Non-square matrix is neither invertible nor singular.

- **Example.** Suppose that A is invertible with inverse B .

$$\begin{aligned}
 AX &= C \Rightarrow B(AX) = BC \\
 &\Rightarrow (BA)X = BC \\
 &\Rightarrow IX = BC \\
 &\Rightarrow X = BC.
 \end{aligned}$$

(A and C must have the same number of rows.)

57 / 150

Examples

- Let $A = \begin{pmatrix} 2 & -5 \\ -1 & 3 \end{pmatrix}$ and $B = \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix}$.
 - $AB = \begin{pmatrix} 2 & -5 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I.$
 - $BA = \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 2 & -5 \\ -1 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I.$

Therefore, A is invertible and B is an inverse of A .

- Solve the equation $\begin{pmatrix} 2 & -5 \\ -1 & 3 \end{pmatrix} X = \begin{pmatrix} 4 \\ 0 \end{pmatrix}$.
 - Pre-multiply the equation by $\begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix}$.

$$\begin{pmatrix} 12 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 4 \\ 0 \end{pmatrix} = \begin{pmatrix} 3 & 5 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 2 & -5 \\ -1 & 3 \end{pmatrix} X = IX = X.$$

58 / 150

Examples

- Prove that $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ is invertible, and find its inverse.

- Let $B = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ such that $AB = BA = I$.

- $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = AB = \begin{pmatrix} a+2c & b+2d \\ 3a+4c & 3b+4d \end{pmatrix}.$

Solve a linear system in a, b, c, d :

- $\begin{cases} 1 = a + 2c \\ 0 = b + 2d \\ 0 = 3a + 4c \\ 1 = 3b + 4d \end{cases} \dots \Rightarrow \dots \begin{cases} a = -2 \\ b = 1 \\ c = 3/2 \\ d = -1/2. \end{cases}$

Moreover, one must verify that $BA = \dots = I$.

- A is invertible with an inverse $\begin{pmatrix} -2 & 1 \\ 3/2 & -1/2 \end{pmatrix}.$

59 / 150

Examples

- Prove that $A = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}$ is a singular matrix.

- Assume that A is invertible (prove by contradiction).

- Then A has an inverse B : $AB = BA = I$.

Let $B = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Then

- $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & b \\ a & b \end{pmatrix}.$

- $1 = a, 0 = b, 0 = a, 1 = b$, a contradiction!

So such B does not exist.

- Therefore, A is not invertible, i.e., it is singular.

Remark: One also gets a contradiction by checking $BA = I$.

60 / 150

Properties

- **Theorem.** Let A be a square matrix.
 - If A is invertible, then its inverse is unique.
- **Proof.** Suppose that B_1 and B_2 are both inverses of A .
 - $AB_1 = B_1A = I$ and $AB_2 = B_2A = I$.We need to verify that $B_1 = B_2$:
 - $B_1 = B_1I = B_1(AB_2) = (B_1A)B_2 = IB_2 = B_2$.
- **Notation.** The unique inverse of A , if exists, is denoted by A^{-1} .
 - $AA^{-1} = A^{-1}A = I$.
- Suppose that A is an invertible matrix. Then
 - If $AX = B$ (A and B have the same number of rows),
 - then $A^{-1}B = A^{-1}(AX) = (A^{-1}A)X = IX = X$.

61 / 150

Properties

- **Cancellation Law.** Let A be an invertible matrix.
 - $AB_1 = AB_2 \Rightarrow B_1 = B_2$.
 - $C_1A = C_2A \Rightarrow C_1 = C_2$.
- **Proof.** Suppose that $AB_1 = AB_2$. Then
 - B_1 is the solution to $AX = AB_2$.
 - $B_1 = A^{-1}(AB_2) = (A^{-1}A)B_2 = IB_2 = B_2$.The other statement is left as an exercise.
- **Remark.** The cancellation law fails if A is singular.
 - Recall that $A = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}$ is singular.
 - $A \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 2 & 0 \end{pmatrix}$ & $A \begin{pmatrix} 2 & 0 \\ 1 & 3 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 2 & 0 \end{pmatrix}$

62 / 150

Example

- Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Find the condition when A is invertible.

Let $B = \begin{pmatrix} x & y \\ z & w \end{pmatrix}$. Suppose that $AB = BA = I$.

$$\circ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = AB = \begin{pmatrix} ax + bz & ay + bw \\ cx + dz & cy + dw \end{pmatrix}.$$

Solve a linear system in x, y, z, w :

$$\circ \begin{cases} ax + bz = 1 \\ ay + bw = 0 \\ cx + dz = 0 \\ cy + dw = 1 \end{cases} \Rightarrow \begin{cases} ax + bz = 1 \\ cx + dz = 0 \\ ay + bw = 0 \\ cy + dw = 1 \end{cases}$$

- They are inconsistent $\Leftrightarrow a : c = b : d \Leftrightarrow ad = bc$.
- They are consistent $\Leftrightarrow ad \neq bc$.

63 / 150

Example

- Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Find the condition when A is invertible.

- If $ad = bc$, then A is singular. Suppose that $ad \neq bc$.

$$\bullet \begin{cases} ax + bz = 1 \\ cx + dz = 0 \end{cases} \Rightarrow x = \frac{d}{ad - bc}, z = \frac{-c}{ad - bc}.$$

$$\bullet \begin{cases} ay + bw = 0 \\ cy + dw = 1 \end{cases} \Rightarrow y = \frac{-b}{ad - bc}, w = \frac{a}{ad - bc}.$$

$$\text{Let } B = \begin{pmatrix} x & y \\ z & w \end{pmatrix} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

- One verifies that $AB = I$ and $BA = I$.

Conclusion: A is invertible $\Leftrightarrow ad - bc \neq 0$.

- If A is invertible, then $A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$.

64 / 150

Properties

- **Theorem.** Let A, B be invertible matrices of same size.

- Let $c \neq 0$. cA is invertible, and $(cA)^{-1} = \frac{1}{c}A^{-1}$.
- A^T is invertible, and $(A^T)^{-1} = (A^{-1})^T$.
- A^{-1} is invertible, and $(A^{-1})^{-1} = A$.
- AB is invertible, and $(AB)^{-1} = B^{-1}A^{-1}$.

- **Proof.** To prove that A is invertible with $A^{-1} = M$,

- Verify that $AM = MA = I$.

To prove that A^T is invertible with inverse $(A^{-1})^T$.

- We shall verify that $A^T(A^{-1})^T = (A^{-1})^T A^T = I$.

- $A^T(A^{-1})^T = (A^{-1}A)^T = I^T = I$.
- $(A^{-1})^T A^T = (AA^{-1})^T = I^T = I$.

Other properties are left as exercises.

65 / 150

Properties

- Let A_1, A_2, \dots, A_k be invertible matrices of same size.

- $(A_1 A_2 \cdots A_k)^{-1} = A_k^{-1} \cdots A_2^{-1} A_1^{-1}$.

In particular, $(\underbrace{AA \cdots A}_{k \text{ times}})^{-1} = \underbrace{A^{-1} \cdots A^{-1}}_{k \text{ times}} A^{-1}$.

- $(A^k)^{-1} = (A^{-1})^k$.

- **Definition.** Let A be an invertible matrix.

- For any positive integer k , $A^{-k} = (A^{-1})^k$

- **Exercise.** Let A be an invertible matrix.

- For any integers m and n ,
 - $A^{m+n} = A^m A^n$ and $(A^m)^n = A^{mn}$.

- **Note.** If A is singular, then A^{-1} is undefined.

66 / 150

Elementary Operations

- Recall the **elementary row operations** of matrices.
 - Multiply a row by a nonzero constant.
 - Interchange two rows.
 - Add a constant multiple of a row to another row.
- What is the resulting matrix by applying an elementary row operation to the identity matrix I ?

$$\circ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{cR_3} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & c & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

- cR_i , where $c \neq 0$:
 - Replace the i th diagonal entry by c :
 - $a_{ii} = 1 \mapsto c$.

68 / 150

Elementary Operations

- Recall the **elementary row operations** of matrices.
 - Multiply a row by a nonzero constant.
 - Interchange two rows.
 - Add a constant multiple of a row to another row.
- What is the resulting matrix by applying an elementary row operation to the identity matrix I ?

$$\circ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_2 \leftrightarrow R_4} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

- $R_i \leftrightarrow R_j$, where $i \neq j$:
 - The i th and j th diagonal entries become 0.
 - The (i, j) and (j, i) -entries become 1.
 - $a_{ii} = a_{jj} = 1 \mapsto 0$, $a_{ij} = a_{ji} = 0 \mapsto 1$.

69 / 150

Elementary Operations

- Recall the **elementary row operations** of matrices.
 - Multiply a row by a nonzero constant.
 - Interchange two rows.
 - Add a constant multiple of a row to another row.
- What is the resulting matrix by applying an elementary row operation to the identity matrix I ?

$$\circ \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_2 + cR_4} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & c \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

- $R_i + cR_j$, where $i \neq j$:
 - The (i, j) -entry becomes c .
 - $a_{ij} = 0 \mapsto c$.

70 / 150

Elementary Matrices

- Definition.** A square matrix is called an **elementary matrix** if it can be obtained from the identity matrix by performing a single elementary row operation.

$$\circ cR_i, \text{ where } c \neq 0: \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & c & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\circ R_i \leftrightarrow R_j, \text{ where } i \neq j, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

$$\circ R_i + cR_j, \text{ where } i \neq j, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & c \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

71 / 150

Connection to Matrix Multiplication

- Let E be an elementary matrix.
 - Suppose that it is obtained from I by
 - Multiplying the i th row by a nonzero number c .

$$\text{Let } E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & c & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{pmatrix}.$$

$$\circ EA = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ ca_{31} & ca_{32} & ca_{33} \\ a_{41} & a_{42} & a_{43} \end{pmatrix}.$$

Pre-multiplication by E to A

\Leftrightarrow Multiplying the i th row of A by number c .

72 / 150

Connection to Matrix Multiplication

- Let $E = (e_{ij})_{m \times m}$ be obtained from I_m by multiplying k th row by c .

$$\circ e_{ij} = \begin{cases} 0 & \text{if } i \neq j, \\ 1 & \text{if } i = j \neq k, \\ c & \text{if } i = j = k. \end{cases}$$

Let $A = (a_{ij})_{m \times n}$. Then EA is well-defined.

- Let $i \neq k$. Then (i, j) -entry of EA is

$$\begin{aligned} & e_{i1}a_{1j} + \cdots + e_{ii}a_{ij} + \cdots + e_{im}a_{mj} \\ &= 0 + \cdots + 1 \cdot a_{ij} + \cdots + 0 = a_{ij}. \end{aligned}$$

- Let $i = k$. Then (i, j) -entry of EA is

$$\begin{aligned} & e_{i1}a_{1j} + \cdots + e_{ii}a_{ij} + \cdots + e_{im}a_{mj} \\ &= 0 + \cdots + c \cdot a_{ij} + \cdots + 0 = ca_{ij}. \end{aligned}$$

Therefore, EA is the matrix obtained from A by multiplying the k th row by c .

73 / 150

Connection to Matrix Multiplication

- Let E be an elementary matrix.
 - Suppose that it is obtained from I by
 - Interchanging the i th row and the j th row.

$$\text{Let } E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{pmatrix}$$

$$\circ EA = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{41} & a_{42} & a_{43} \\ a_{31} & a_{32} & a_{33} \\ a_{21} & a_{22} & a_{23} \end{pmatrix}$$

Pre-multiplication by E to A

\Leftrightarrow Interchanging the i th row and j th row of A .

74 / 150

Connection to Matrix Multiplication

- Let $E = (e_{ij})$ be obtained from I_m by interchanging ℓ th and k th rows ($\ell < k$).
 - $e_{ij} = \begin{cases} 1 & \text{if } i = j \neq \ell, k \text{ or } (i, j) = (\ell, k), (k, \ell), \\ 0 & \text{otherwise.} \end{cases}$

Let $A = (a_{ij})_{m \times n}$. Then EA is well-defined.

- Let $i \neq \ell, k$. Then (i, j) -entry of EA is

$$\begin{aligned} & e_{i1}a_{1j} + \cdots + e_{ii}a_{ij} + \cdots + e_{im}a_{mj} \\ &= 0 + \cdots + 1 \cdot a_{ij} + \cdots + 0 = a_{ij}. \end{aligned}$$

- Let $i = \ell$ (the case $i = k$ is similar). Then (i, j) -entry of EA

$$\begin{aligned} & e_{i1}a_{1j} + \cdots + e_{ii}a_{ij} + \cdots + e_{ik}a_{kj} + \cdots + e_{im}a_{mj} \\ &= 0 + \cdots + 0 \cdot a_{ij} + \cdots + 1 \cdot a_{kj} + \cdots + 0 = a_{kj}. \end{aligned}$$

Therefore, EA is the matrix obtained from A by interchanging ℓ th and k th rows.

75 / 150

Connection to Matrix Multiplication

- Let E be an elementary matrix.
 - Suppose that it is obtained from I by
 - Adding c times of the j th row to the i th row.

$$\text{Let } E = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & c \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{pmatrix}$$

$$\circ EA = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} + ca_{41} & a_{22} + ca_{42} & a_{23} + ca_{43} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{pmatrix}$$

Pre-multiplication by E to A

\Leftrightarrow Adding c times the j th row to the i th row of A .

76 / 150

Connection to Matrix Multiplication

- Let $E = (e_{ij})$ be obtained from I_m by adding c times of ℓ th row to k th row.

$$\circ e_{ij} = \begin{cases} 1 & \text{if } i = j, \\ c & \text{if } (i, j) = (k, \ell), \\ 0 & \text{otherwise.} \end{cases}$$

Let $A = (a_{ij})_{m \times n}$. Then EA is well-defined.

- Let $i \neq k$. Then (i, j) -entry of EA is

$$\begin{aligned} & e_{i1}a_{1j} + \cdots + e_{ii}a_{ij} + \cdots + e_{im}a_{mj} \\ &= 0 + \cdots + 1 \cdot a_{ij} + \cdots + 0 = a_{ij}. \end{aligned}$$

- Let $i = k$. Then (i, j) -entry of EA is (w.l.o.g., assume $k < \ell$)

$$\begin{aligned} & e_{i1}a_{1j} + \cdots + e_{ii}a_{ij} + \cdots + e_{i\ell}a_{\ell j} + \cdots + e_{im}a_{mj} \\ &= 0 + \cdots + 1 \cdot a_{ij} + \cdots + c \cdot a_{\ell j} + \cdots + 0 = a_{ij} + ca_{\ell j}. \end{aligned}$$

Therefore, EA is obtained from A by adding c times of ℓ th row to the k th row.

77 / 150

Connection to Multiplication

- **Theorem.**

- Let E be the elementary matrix obtained
 - by performing an elementary row operation to I_m .

Then for any $m \times n$ matrix A , EA can be obtained

- by performing same elementary row operation to A .
- Let A be an $m \times n$ matrix.
 - $I_m \xrightarrow{cR_i} E \Rightarrow A \xrightarrow{cR_i} EA.$
 - $I_m \xrightarrow{R_i \leftrightarrow R_j} E \Rightarrow A \xrightarrow{R_i \leftrightarrow R_j} EA.$
 - $I_m \xrightarrow{R_i + cR_j} E \Rightarrow A \xrightarrow{R_i + cR_j} EA.$

78 / 150

Example

- Let $A = \begin{pmatrix} 0 & 4 & 2 \\ -2 & 1 & -3 \\ 1 & 0 & 2 \end{pmatrix}.$
 - $A = \begin{pmatrix} 0 & 4 & 2 \\ -2 & 1 & -3 \\ 1 & 0 & 2 \end{pmatrix} \xrightarrow{R_1 \leftrightarrow R_3} \begin{pmatrix} 1 & 0 & 2 \\ -2 & 1 & -3 \\ 0 & 4 & 2 \end{pmatrix} = A_1.$
 - $I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_1 \leftrightarrow R_3} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} = E_1.$

$$\begin{aligned} E_1 A &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 4 & 2 \\ -2 & 1 & -3 \\ 1 & 0 & 2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 2 \\ -2 & 1 & -3 \\ 0 & 4 & 2 \end{pmatrix} = A_1. \end{aligned}$$

79 / 150

Example

- Let $A = \begin{pmatrix} 0 & 4 & 2 \\ -2 & 1 & -3 \\ 1 & 0 & 2 \end{pmatrix}$.

- $A_1 = \begin{pmatrix} 1 & 0 & 2 \\ -2 & 1 & -3 \\ 0 & 4 & 2 \end{pmatrix} \xrightarrow{R_2+2R_1} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 4 & 2 \end{pmatrix} = A_2.$

- $I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_2+2R_1} \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = E_2.$

$$\begin{aligned} E_2 A_1 &= \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 2 \\ -2 & 1 & -3 \\ 0 & 4 & 2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 4 & 2 \end{pmatrix} = A_2. \end{aligned}$$

80 / 150

Example

- Let $A = \begin{pmatrix} 0 & 4 & 2 \\ -2 & 1 & -3 \\ 1 & 0 & 2 \end{pmatrix}$.

- $A_2 = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 4 & 2 \end{pmatrix} \xrightarrow{R_3+(-4)R_2} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & -2 \end{pmatrix} = A_3.$

- $I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_3+(-4)R_2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -4 & 1 \end{pmatrix} = E_3.$

$$\begin{aligned} E_3 A_2 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -4 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 4 & 2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & -2 \end{pmatrix} = A_3. \end{aligned}$$

81 / 150

Example

- Let $A = \begin{pmatrix} 0 & 4 & 2 \\ -2 & 1 & -3 \\ 1 & 0 & 2 \end{pmatrix}$.

- $A_3 = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & -2 \end{pmatrix} \xrightarrow{-\frac{1}{2}R_3} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} = A_4.$

- $I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{-\frac{1}{2}R_3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{2} \end{pmatrix} = E_4.$

$$\begin{aligned} E_4 A_3 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & -2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} = A_4. \end{aligned}$$

82 / 150

Example

- Let $A = \begin{pmatrix} 0 & 4 & 2 \\ -2 & 1 & -3 \\ 1 & 0 & 2 \end{pmatrix}$.

- $A_4 = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_1 + (-2)R_3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} = A_5.$

- $I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_1 + (-2)R_3} \begin{pmatrix} 1 & 0 & -2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = E_5.$

$$\begin{aligned} E_5 A_4 &= \begin{pmatrix} 1 & 0 & -2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} = A_5. \end{aligned}$$

83 / 150

Example

- Let $A = \begin{pmatrix} 0 & 4 & 2 \\ -2 & 1 & -3 \\ 1 & 0 & 2 \end{pmatrix}$.

- $A_5 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_2 + (-1)R_3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = A_6 = I.$

- $I = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_2 + (-1)R_3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} = E_6.$

$$\begin{aligned} E_6 A_5 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = A_6 = I. \end{aligned}$$

84 / 150

Invertibility

- Theorem.** Every elementary matrix is invertible.

- The inverse of an elementary matrix is elementary.

Proof. There are three types of elementary matrices.

- Suppose that $I \xrightarrow{cR_i} E$, where $c \neq 0$.

- Then $E \xrightarrow{\frac{1}{c}R_i} I$.

Let D denote the elementary matrix $I \xrightarrow{\frac{1}{c}R_i} D$.

- $I \xrightarrow[cR_i]{E} E \xrightarrow[\frac{1}{c}R_i]{D} I$. So $DE = I$.

- $I \xrightarrow[\frac{1}{c}R_i]{D} D \xrightarrow[cR_i]{E} I$. So $ED = I$.

It follows that E is invertible and $E^{-1} = D$.

The other two cases are similar and left as exercises.

85 / 150

Invertibility

- Let E be an elementary matrix.
 - $I \xrightarrow{cR_i} E \Rightarrow I \xrightarrow{\frac{1}{c}R_i} E^{-1}$.
 - $I \xrightarrow{R_i \leftrightarrow R_j} E \Rightarrow I \xrightarrow{R_i \leftrightarrow R_j} E^{-1}$. (So $E = E^{-1}$)
 - $I \xrightarrow{R_i + cR_j} E \Rightarrow I \xrightarrow{R_i + (-c)R_j} E^{-1}$.
- Suppose that matrices A and B are row equivalent.
 - $A = A_0 \xrightarrow{\text{ero1}} A_1 \xrightarrow{\text{ero2}} A_2 \cdots \rightarrow A_{k-1} \xrightarrow{\text{ero}k} A_k = B$.

Let E_i be the elementary matrix corresponding to the i th elementary row operation: $I \xrightarrow{\text{ero}i} E_i$.

$$A = A_0 \xrightarrow[E_1]{\text{ero1}} A_1 \xrightarrow[E_2]{\text{ero2}} A_2 \cdots \rightarrow A_{k-1} \xrightarrow[E_k]{\text{ero}k} A_k = B.$$

Then $B = E_k E_{k-1} \cdots E_2 E_1 A$.

86 / 150

Invertibility

- Theorem.** Two matrices A and B are row equivalent
 - \Leftrightarrow there exist elementary matrices E_1, E_2, \dots, E_k such that $B = E_k E_{k-1} \cdots E_2 E_1 A$.

- Remarks.** Suppose for elementary matrices E_i ,

$$B = E_k E_{k-1} \cdots E_2 E_1 A.$$

$$\begin{aligned} & \bullet A \xrightarrow{E_1} \bullet \xrightarrow{E_2} \bullet \rightarrow \cdots \rightarrow \bullet \xrightarrow{E_{k-1}} \bullet \xrightarrow{E_k} B. \\ & \bullet A \xleftarrow{E_1^{-1}} \bullet \xleftarrow{E_2^{-1}} \bullet \leftarrow \cdots \leftarrow \bullet \xleftarrow{E_{k-1}^{-1}} \bullet \xleftarrow{E_k^{-1}} B. \end{aligned}$$

$$\therefore A = E_1^{-1} E_2^{-1} \cdots E_{k-1}^{-1} E_k^{-1} B.$$

We can now prove the theorem stated in Chapter 1:

- Theorem.** Suppose that the augmented matrices of two linear systems are row equivalent.
 - Then the two systems have the same solution set.

87 / 150

Invertibility

- **Proof.** Let the two linear systems be

- $Ax = b$ and $Cx = d$.

Then the associated augmented matrices are

- $(A \mid b)$ and $(C \mid d)$.

There exist elementary matrices E_1, E_2, \dots, E_k so that

- $E_k \cdots E_1 (A \mid b) = (C \mid d)$.

- $E_k \cdots E_1 A = C$ and $E_k \cdots E_1 b = d$.

Let u be a solution to $Ax = b$, i.e., $Au = b$.

- $E_k \cdots E_1 Au = E_k \cdots E_1 b \Rightarrow Cu = d$.

So u is also a solution to $Cx = d$.

88 / 150

Invertibility

- **Proof.** Let the two linear systems be

- $Ax = b$ and $Cx = d$.

Then the associated augmented matrices are

- $(A \mid b)$ and $(C \mid d)$.

There exist elementary matrices E_1, E_2, \dots, E_k so that

- $E_k \cdots E_1 (A \mid b) = (C \mid d)$.

- $E_k \cdots E_1 A = C$ and $E_k \cdots E_1 b = d$.

- $A = E_1^{-1} \cdots E_k^{-1} C$ and $b = E_1^{-1} \cdots E_k^{-1} d$.

Let v be a solution to $Cx = d$, i.e., $Cv = d$.

- $E_1^{-1} \cdots E_k^{-1} Cv = E_1^{-1} \cdots E_k^{-1} d \Rightarrow Av = b$.

So v is also a solution to $Ax = b$.

89 / 150

Main Theorem for Invertible Matrices

• **Theorem.** Let A be a square matrix. Then the followings are equivalent:

1. A is an invertible matrix.
2. Linear system $Ax = b$ has a unique solution.
3. Linear system $Ax = 0$ has only the trivial solution.
4. The reduced row-echelon form of A is I .
5. A is the product of elementary matrices.

90 / 150

Main Theorem for Invertible Matrices

• **Proof.** We prove $1 \Rightarrow 2 \Rightarrow 3 \Rightarrow 4 \Rightarrow 5 \Rightarrow 1$:

$1 \Rightarrow 2$: Suppose A is invertible.

- $Ax = b \Rightarrow x = Ix = A^{-1}Ax = A^{-1}b$.

$2 \Rightarrow 3$: Suppose $Ax = b$ has a unique solution u .

- If $Ax = 0$ has a solution v , then $Av = 0$.
 - $A(u - v) = Au - Av = b - 0 = b$.
 - $u - v$ is also a solution to $Ax = b$.
- By uniqueness, $u = u - v$; so $v = 0$.

$3 \Rightarrow 4$: Suppose $Ax = 0$ has only the trivial solution.

- Let R be the reduced row-echelon form of A .
 - Except the last column, all other columns of $(R \mid 0)$ are pivot columns.
- Note that R is a square matrix. So $R = I$.

91 / 150

Main Theorem for Invertible Matrices

- **Proof.** We prove $1 \Rightarrow 2 \Rightarrow 3 \Rightarrow 4 \Rightarrow 5 \Rightarrow 1$:

4 \Rightarrow 5: Suppose that the reduced row-echelon form of A is I .

- I can be obtained from A by elementary row operations.
 - $A \xrightarrow{\text{ero1}} \bullet \dots \rightarrow \dots \bullet \xrightarrow{\text{ero}k} I$.
- Let E_i be the elementary matrix corresponding to the i th elementary row operation.
 - $I = E_k E_{k-1} \dots E_2 E_1 A$.
- Then $A = E_1^{-1} E_2^{-1} \dots E_{k-1}^{-1} E_k^{-1}$.
 - (Recall: each E_i^{-1} is also an elementary matrix.)

5 \Rightarrow 1: Suppose that $A = E_1 E_2 \dots E_k$,

- where E_1, \dots, E_k are elementary matrices.

Each E_i is invertible $\Rightarrow A$ is invertible.

92 / 150

Find Inverse

- Let A be an invertible matrix of order n . Its reduced row-echelon form is I_n .
 - There exist elementary matrices E_i such that

$$\bullet \quad E_k \dots E_2 E_1 A = I_n.$$

$$\text{Then } E_k \dots E_2 E_1 = A^{-1}.$$

Consider the $n \times 2n$ matrix $(A \mid I)$.

- Apply the ele. row oper. corresponding to E_1, \dots, E_k :

$$\begin{aligned} (A \mid I_n) &\xrightarrow{E_1} (E_1 A \mid E_1) \\ &\xrightarrow{E_2} (E_2 E_1 A \mid E_2 E_1) \\ &\rightarrow \dots \rightarrow \dots \\ &\xrightarrow{E_k} (E_k \dots E_2 E_1 A \mid E_k \dots E_2 E_1) \\ &= (I_n \mid A^{-1}). \end{aligned}$$

93 / 150

Find Inverse

- **Theorem.** Let A be an invertible matrix.

◦ The reduced row-echelon form of $(A \mid I)$ is $(I \mid A^{-1})$

- **Example.** Let $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 5 & 3 \\ 1 & 0 & 8 \end{pmatrix}$. Find A^{-1} .

$$\begin{aligned} \left(\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 5 & 3 & 0 & 1 & 0 \\ 1 & 0 & 8 & 0 & 0 & 1 \end{array} \right) &\xrightarrow[R_3+(-1)R_1]{R_2+(-2)R_1} \left(\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & -2 & 5 & -1 & 0 & 1 \end{array} \right) \\ &\xrightarrow{R_3+2R_2} \left(\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & 0 & -1 & -5 & 2 & 1 \end{array} \right) \\ &\xrightarrow{(-1)R_2} \left(\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & 0 & 1 & 5 & -2 & -1 \end{array} \right) \end{aligned}$$

94 / 150

Find Inverse

- **Theorem.** Let A be an invertible matrix.

◦ The reduced row-echelon form of $(A \mid I)$ is $(I \mid A^{-1})$

- **Example.** Let $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 5 & 3 \\ 1 & 0 & 8 \end{pmatrix}$. Find A^{-1} .

$$\begin{aligned} \left(\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & -3 & -2 & 1 & 0 \\ 0 & 0 & 1 & 5 & -2 & -1 \end{array} \right) &\xrightarrow[R_2+3R_3]{R_1+(-3)R_3} \left(\begin{array}{ccc|ccc} 1 & 2 & 0 & -14 & 6 & 3 \\ 0 & 1 & 0 & 13 & -5 & -3 \\ 0 & 0 & 1 & 5 & -2 & -1 \end{array} \right) \\ &\xrightarrow{R_1+(-2)R_2} \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & -40 & 16 & 9 \\ 0 & 1 & 0 & 13 & -5 & -3 \\ 0 & 0 & 1 & 5 & -2 & -1 \end{array} \right) \end{aligned}$$

Therefore, $A^{-1} = \begin{pmatrix} -40 & 16 & 9 \\ 13 & -5 & -3 \\ 5 & -2 & -1 \end{pmatrix}$.

95 / 150

Find Inverse

- A square matrix is **invertible**

- ⇔ Its reduced row-echelon form is I
- ⇔ All the columns in its row-echelon form are pivot.
- ⇔ All the rows in its row-echelon form are nonzero.

A square matrix is **singular**

- ⇔ Its reduced row-echelon form is not I
- ⇔ Some columns in its row-echelon form are non-pivot.
- ⇔ Some rows in its row-echelon form are zero.

Example. Let $A = \begin{pmatrix} 1 & 2 & 0 \\ 2 & 4 & 4 \\ 3 & 6 & 3 \end{pmatrix}$. Then A is singular.

$$\begin{pmatrix} 1 & 2 & 0 \\ 2 & 4 & 4 \\ 3 & 6 & 3 \end{pmatrix} \xrightarrow[R_3+(-3)R_1]{R_2+(-2)R_1} \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 4 \\ 0 & 0 & 3 \end{pmatrix} \xrightarrow{R_3+(-\frac{3}{4})R_2} \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 4 \\ 0 & 0 & 0 \end{pmatrix}$$

96 / 150

Find Inverse

- **Theorem.** Let A and B be square matrices of the same size. If $AB = I$, then
 - A and B are invertible, and $A^{-1} = B$, $B^{-1} = A$.

Proof. Consider the linear system $Bx = 0$.

- $Bx = 0 \Rightarrow ABx = A0 \Rightarrow x = 0$.
- $Bx = 0$ has only the trivial solution $\Rightarrow B$ is invertible.

B^{-1} exists such that $BB^{-1} = B^{-1}B = I$.

- $AB = I \Rightarrow ABB^{-1} = IB^{-1} \Rightarrow A = B^{-1}$.

Therefore, A is invertible, and $A^{-1} = (B^{-1})^{-1} = B$.

- **Corollary and Exercise.** Let A_1, A_2, \dots, A_k be square matrices of the same size.
 - $A_1 A_2 \cdots A_k$ is invertible \Leftrightarrow all A_i are invertible.
 - $A_1 A_2 \cdots A_k$ is singular \Leftrightarrow some A_i are singular.

97 / 150

Find Inverse

- Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $B = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$, $ad \neq bc$.
 - One verifies that $AB = I$.
 - Using the theorem, $A^{-1} = B$.
 - The verification that $BA = I$ is not necessary.
- Let A be a square matrix such that $A^2 - 3A - 4I = 0$.
 - Prove that A is invertible, and find A^{-1} .

Wrong Proof. $0 = (A - 4I)(A + I)$.

- $A - 4I = 0$ or $A + I = 0 \Rightarrow A = 4I$ or $A = -I$.

Proof. $4I = A^2 - 3A = A(A - 3I)$.

- $I = \frac{1}{4}A(A - 3I) = A \left[\frac{1}{4}(A - 3I) \right]$.
- So A is invertible with $A^{-1} = \frac{1}{4}(A - 3I)$.

98 / 150

Column Operations

- Recall that the pre-multiplication of an elementary matrix \Leftrightarrow corresponding elementary row operation.

Question. What is the effect of **post-multiplication** of an elementary matrix?

- Answer: Corresponding elementary column operation.
- **Elementary column operations:**
 - kC_i : multiply i th column by a nonzero constant k .
 - $C_i \leftrightarrow C_j$: interchange i th and j th columns.
 - $C_i + kC_j$: add k times j th column to i th column.

Let E be the matrix obtained from I by a single elementary column operation.

- Then E is an **elementary matrix**.
 - (i.e., E can be obtained from I by a single elementary row operation.)

99 / 150

Column Operations

- If E is obtained from I_n by a single elementary column operation, then E is an elementary matrix.

- $I \xrightarrow{kC_i} E \Leftrightarrow I \xrightarrow{kR_i} E.$
- $I \xrightarrow{C_i \leftrightarrow C_j} E \Leftrightarrow I \xrightarrow{R_i \leftrightarrow R_j} E.$
- $I \xrightarrow{C_i + kC_j} E \Leftrightarrow I \xrightarrow{R_j + kR_i} E.$

- Let A be an $m \times n$ matrix. Then A^T is $n \times m$.

- Suppose that $I_n \xrightarrow{kC_i} E$. Note that $E = E^T$.
 - $I_n \xrightarrow{kR_i} E^T \Rightarrow A^T \xrightarrow{kR_i} E^T A^T = (AE)^T.$

Then $A \xrightarrow{kC_i} AE$.

100 / 150

Column Operations

- If E is obtained from I_n by a single elementary column operation, then E is an elementary matrix.

- $I \xrightarrow{kC_i} E \Leftrightarrow I \xrightarrow{kR_i} E.$
- $I \xrightarrow{C_i \leftrightarrow C_j} E \Leftrightarrow I \xrightarrow{R_i \leftrightarrow R_j} E.$
- $I \xrightarrow{C_i + kC_j} E \Leftrightarrow I \xrightarrow{R_j + kR_i} E.$

- Let A be an $m \times n$ matrix. Then A^T is $n \times m$.

- Suppose that $I_n \xrightarrow{C_i \leftrightarrow C_j} E$. Note that $E = E^T$.
 - $I \xrightarrow{R_i \leftrightarrow R_j} E^T \Rightarrow A^T \xrightarrow{R_i \leftrightarrow R_j} E^T A^T = (AE)^T.$

Then $A \xrightarrow{C_i \leftrightarrow C_j} AE$.

101 / 150

Column Operations

- If E is obtained from I_n by a single elementary column operation, then E is an elementary matrix.
 - $I \xrightarrow{kC_i} E \Leftrightarrow I \xrightarrow{kR_i} E$.
 - $I \xrightarrow{C_i \leftrightarrow C_j} E \Leftrightarrow I \xrightarrow{R_i \leftrightarrow R_j} E$.
 - $I \xrightarrow{C_i + kC_j} E \Leftrightarrow I \xrightarrow{R_j + kR_i} E$.
 - Let A be an $m \times n$ matrix. Then A^T is $n \times m$.
 - Suppose that $I \xrightarrow{C_i + kC_j} E$.
 - Then $I \xrightarrow{R_i + kR_j} E^T$.
 - $A^T \xrightarrow{R_i + kR_j} E^T A^T = (AE)^T$.
- Then $A \xrightarrow{C_i + kC_j} AE$.

102 / 150

Column Operations

- If E is obtained from I_n by a single elementary column operation, then E is an elementary matrix.
 - $I \xrightarrow{kC_i} E \Leftrightarrow I \xrightarrow{kR_i} E$.
 - $I \xrightarrow{C_i \leftrightarrow C_j} E \Leftrightarrow I \xrightarrow{R_i \leftrightarrow R_j} E$.
 - $I \xrightarrow{C_i + kC_j} E \Leftrightarrow I \xrightarrow{R_j + kR_i} E$.
- Let A be an $m \times n$ matrix, and E an $n \times n$ elementary matrix. Then
 - The post-multiplication of E to A
 - \Leftrightarrow Corresponding elementary column operation to A .
 - $I \xrightarrow{kC_i} E \Rightarrow A \xrightarrow{kC_i} AE$.
 - $I \xrightarrow{C_i \leftrightarrow C_j} E \Rightarrow A \xrightarrow{C_i \leftrightarrow C_j} AE$.
 - $I \xrightarrow{C_i + kC_j} E \Rightarrow A \xrightarrow{C_i + kC_j} AE$.

103 / 150

Examples

- Let $A = \begin{pmatrix} 1 & 0 & 2 & 3 \\ 2 & -1 & 3 & 6 \\ 1 & 4 & 4 & 0 \end{pmatrix}$.
- $I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{2C_3} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = E_1.$
- $AE_1 = \dots = \begin{pmatrix} 1 & 0 & 4 & 3 \\ 2 & -1 & 6 & 6 \\ 1 & 4 & 8 & 0 \end{pmatrix}$
- $A \xrightarrow{2C_3} AE_1.$

104 / 150

Examples

- Let $A = \begin{pmatrix} 1 & 0 & 2 & 3 \\ 2 & -1 & 3 & 6 \\ 1 & 4 & 4 & 0 \end{pmatrix}$.
- $I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{C_2 \leftrightarrow C_4} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} = E_2.$
- $AE_2 = \dots = \begin{pmatrix} 1 & 3 & 2 & 0 \\ 2 & 6 & 3 & -1 \\ 1 & 0 & 4 & 4 \end{pmatrix}$
- $A \xrightarrow{C_2 \leftrightarrow C_4} AE_2.$

105 / 150

Examples

- Let $A = \begin{pmatrix} 1 & 0 & 2 & 3 \\ 2 & -1 & 3 & 6 \\ 1 & 4 & 4 & 0 \end{pmatrix}$.
 - $I = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \xrightarrow{C_1+2C_3} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 2 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = E_3.$
 - $AE_3 = \dots = \begin{pmatrix} 5 & 0 & 2 & 3 \\ 8 & -1 & 3 & 6 \\ 9 & 4 & 4 & 0 \end{pmatrix}.$
- $A \xrightarrow{C_1+2C_3} AE_3.$

106 / 150

Determinant

107 / 150

Determinant of 2×2 Matrix

- Recall that $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is invertible $\Leftrightarrow ad - bc \neq 0$.
- **Definition.** Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$.
 - The **determinant** of A is $\det(A) = |A| = ad - bc$

Therefore, A is invertible $\Leftrightarrow \det(A) \neq 0$.
- **Definition.** If $A = (a)$, it is natural to set $\det(A) = a$.
- **Properties & Exercises.** Let A, B be 2×2 matrices.
 - $\det(I_2) = 1$.
 - $A \xrightarrow{cR_i} B \Rightarrow \det(B) = c \det(A)$.
 - $A \xrightarrow{R_i \leftrightarrow R_j} B \Rightarrow \det(B) = -\det(A)$.
 - $A \xrightarrow{R_i + cR_j} B \Rightarrow \det(B) = \det(A)$.

108 / 150

Determinant of 2×2 Matrix

- Consider linear system $\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}$.
 - Suppose $\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ is invertible.
 - $\mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} \begin{pmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{pmatrix}$.

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \mathbf{A}^{-1} \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = \dots = \frac{1}{\det(\mathbf{A})} \begin{pmatrix} a_{22}b_1 - a_{12}b_2 \\ -a_{21}b_1 + a_{11}b_2 \end{pmatrix}$$
- $x_1 = \frac{\begin{vmatrix} b_1 & a_{12} \\ b_2 & a_{22} \end{vmatrix}}{\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}}$ and $x_2 = \frac{\begin{vmatrix} a_{11} & b_1 \\ a_{21} & b_2 \end{vmatrix}}{\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}}$.

109 / 150

Determinant of 2×2 Matrix

- One can verify that

$$\begin{vmatrix} a_{11} + a'_{11} & a_{12} + a'_{12} \\ a_{21} & a_{22} \end{vmatrix} = (a_{11} + a'_{11})a_{22} - (a_{12} + a'_{12})a_{21}$$

$$= (a_{11}a_{22} - a_{12}a_{21}) + (a'_{11}a_{22} - a'_{12}a_{21})$$

$$= \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} + \begin{vmatrix} a'_{11} & a'_{12} \\ a_{21} & a_{22} \end{vmatrix}$$

In particular,

$$\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = \begin{vmatrix} a_{11} & 0 \\ a_{21} & a_{22} \end{vmatrix} + \begin{vmatrix} 0 & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$$

$$= a_{11} \begin{vmatrix} 1 & 0 \\ a_{21} & a_{22} \end{vmatrix} + a_{12} \begin{vmatrix} 0 & 1 \\ a_{21} & a_{22} \end{vmatrix}$$

$$= a_{11} \cdot a_{22} - a_{12} \cdot a_{21}$$

$$= a_{11} \cdot \det(a_{22}) - a_{12} \cdot \det(a_{21}).$$

110 / 150

Determinant of 3×3 Matrix

- Let A be a square matrix. It is expected that
 - $\det(I) = 1$.
 - A is invertible $\Leftrightarrow \det(A) \neq 0$.
 - Equivalently, A is singular $\Leftrightarrow \det(A) = 0$.
 - $A \xrightarrow{cR_i} B \Rightarrow \det(B) = c \det(A)$.
 - $A \xrightarrow{R_i \leftrightarrow R_j} B \Rightarrow \det(B) = -\det(A)$.
 - $A \xrightarrow{R_i + cR_j} B \Rightarrow \det(B) = \det(A)$.
- Suppose A is invertible. Then there exist elementary row operations
 - $A \xrightarrow{\text{ero1}} A_1 \xrightarrow{\text{ero2}} A_2 \rightarrow \dots \rightarrow A_{k-1} \xrightarrow{\text{erok}} A_k = I$.
 Then $\det(A)$ can be evaluated backwards.
- Example.** $A \xrightarrow{R_1 \leftrightarrow R_3} \bullet \xrightarrow{3R_2} \bullet \xrightarrow{R_2 + 2R_4} I \Rightarrow \det(A) = -\frac{1}{3}$.

111 / 150

Determinant of 3×3 Matrix

- It is also expected that
 - $\det \begin{pmatrix} R_1 \\ R_2 \\ \vdots \\ R_n \end{pmatrix} + \det \begin{pmatrix} R'_1 \\ R_2 \\ \vdots \\ R_n \end{pmatrix} = \det \begin{pmatrix} R_1 + R'_1 \\ R_2 \\ \vdots \\ R_n \end{pmatrix}$
- Consider 3×3 matrix $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$
 - It is expected to have $\det(A)$:
 - $\begin{vmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} + \begin{vmatrix} 0 & a_{12} & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} + \begin{vmatrix} 0 & 0 & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}.$

112 / 150

Determinant of 3×3 Matrix

- If $a_{11} = 0$, the determinant is supposed to be 0. Suppose $a_{11} \neq 0$.

$$\begin{vmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11} \begin{vmatrix} 1 & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & a_{32} & a_{33} \end{vmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}$$

$$\begin{pmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \xrightarrow{\frac{1}{a_{11}}R_1} \begin{pmatrix} 1 & 0 & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \xrightarrow{\begin{matrix} R_2 + (-a_{21})R_1 \\ R_3 + (-a_{31})R_1 \end{matrix}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & a_{32} & a_{33} \end{pmatrix}.$$

We use the same elementary row operations:

$$\circ \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & a_{32} & a_{33} \end{pmatrix} \rightarrow \text{RREF} \text{ and } \begin{pmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{pmatrix} \rightarrow \text{RREF}.$$

113 / 150

Determinant of 3×3 Matrix

- If $a_{12} = 0$, the determinant is supposed to be 0. Suppose $a_{12} \neq 0$.

$$\begin{vmatrix} 0 & a_{12} & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = -a_{12} \begin{vmatrix} 1 & 0 & 0 \\ 0 & a_{21} & a_{23} \\ 0 & a_{31} & a_{33} \end{vmatrix} = -a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix}$$

$$\begin{pmatrix} 0 & a_{12} & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \xrightarrow{\frac{1}{a_{12}}R_1} \begin{pmatrix} 0 & 1 & 0 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \xrightarrow{\begin{matrix} R_2 + (-a_{21})R_1 \\ R_3 + (-a_{31})R_1 \end{matrix}} \begin{pmatrix} 0 & 1 & 0 \\ a_{21} & 0 & a_{23} \\ a_{31} & 0 & a_{33} \end{pmatrix} \xrightarrow{C_1 \leftrightarrow C_2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_{21} & a_{23} \\ 0 & a_{31} & a_{33} \end{pmatrix}$$

We use the same elementary row operations:

$$\circ \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_{21} & a_{23} \\ 0 & a_{31} & a_{33} \end{pmatrix} \rightarrow \text{RREF} \text{ and } \begin{pmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{pmatrix} \rightarrow \text{RREF}.$$

114 / 150

Determinant of 3×3 Matrix

- If $a_{13} = 0$, the determinant is supposed to be 0. Suppose $a_{13} \neq 0$.

$$\begin{vmatrix} 0 & 0 & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{13} \begin{vmatrix} 1 & 0 & 0 \\ 0 & a_{21} & a_{22} \\ 0 & a_{31} & a_{32} \end{vmatrix} = a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

$$\begin{aligned} & \begin{pmatrix} 0 & 0 & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \xrightarrow{\frac{1}{a_{13}}R_1} \begin{pmatrix} 0 & 0 & 1 \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \\ & \xrightarrow[\begin{matrix} R_2 + (-a_{23})R_1 \\ R_3 + (-a_{33})R_1 \end{matrix}]{\begin{matrix} R_2 + (-a_{23})R_1 \\ R_3 + (-a_{33})R_1 \end{matrix}} \begin{pmatrix} 0 & 0 & 1 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & 0 \end{pmatrix} \xrightarrow{C_1 \leftrightarrow C_3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_{21} & a_{22} \\ 0 & a_{31} & a_{32} \end{pmatrix} \end{aligned}$$

We use the same elementary row operations:

$$\circ \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_{21} & a_{22} \\ 0 & a_{31} & a_{32} \end{pmatrix} \rightarrow \text{RREF} \text{ and } \begin{pmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix} \rightarrow \text{RREF}.$$

115 / 150

Determinant of 3×3 Matrix

- Definition.** Let $\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$.

$$\circ \det(\mathbf{A}) = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

- Notation.** Let $\mathbf{A} = (a_{ij})_{n \times n}$.
 - Let \mathbf{M}_{ij} be the **submatrix** obtained from \mathbf{A} by deleting the i th row and j th column.
 - If $\mathbf{A} = (a_{ij})_{3 \times 3}$, then $\det(\mathbf{A})$ is given by
 - $a_{11} \det(\mathbf{M}_{11}) - a_{12} \det(\mathbf{M}_{12}) + a_{13} \det(\mathbf{M}_{13})$
 - Let $A_{ij} = (-1)^{i+j} \det(\mathbf{M}_{ij})$, the **(i, j) -cofactor** of \mathbf{A} .
 - If $\mathbf{A} = (a_{ij})_{3 \times 3}$, then
 - $\det(\mathbf{A}) = a_{11}A_{11} + a_{12}A_{12} + a_{13}A_{13}$

116 / 150

Example

- Let $B = (b_{ij})_{3 \times 3} = \begin{pmatrix} -3 & -2 & 4 \\ 4 & 3 & 1 \\ 0 & 2 & 4 \end{pmatrix}$.
 - (1, 1)-cofactor: $(-1)^{1+1} \begin{vmatrix} 3 & 1 \\ 2 & 4 \end{vmatrix} = 3 \cdot 4 - 1 \cdot 2 = 10$.
 - (1, 2)-cofactor:
 - $(-1)^{1+2} \begin{vmatrix} 4 & 1 \\ 0 & 4 \end{vmatrix} = -(4 \cdot 4 - 1 \cdot 0) = -16$.
 - (1, 3)-cofactor: $(-1)^{1+3} \begin{vmatrix} 4 & 3 \\ 0 & 2 \end{vmatrix} = 4 \cdot 2 - 3 \cdot 0 = 8$.

$$\begin{aligned} \det(B) &= (-3) \cdot 10 + (-2) \cdot (-16) + 4 \cdot 8 \\ &= 34. \end{aligned}$$

117 / 150

An Alternative Formula

- Let $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$.

$$\begin{aligned} \det(A) &= a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \\ &= a_{11}(a_{22}a_{33} - a_{23}a_{32}) - a_{12}(a_{21}a_{33} - a_{23}a_{31}) \\ &\quad + a_{13}(a_{21}a_{32} - a_{22}a_{31}) \\ &= (a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32}) \\ &\quad - (a_{11}a_{23}a_{32} + a_{12}a_{21}a_{33} + a_{13}a_{22}a_{31}). \end{aligned}$$

- The positive terms come from the
 - 3 (broken) diagonals from the top left to bottom right.The negative terms come from the
 - 3 (broken) diagonals from the top right to bottom left.

118 / 150

Example

- Let $B = (b_{ij})_{3 \times 3} = \begin{pmatrix} -3 & -2 & 4 \\ 4 & 3 & 1 \\ 0 & 2 & 4 \end{pmatrix}$.

$$\begin{aligned}\det(B) &= [(-3) \cdot 3 \cdot 4 + (-2) \cdot 1 \cdot 0 + 4 \cdot 4 \cdot 2] \\ &\quad - [(-3) \cdot 1 \cdot 2 + (-2) \cdot 4 \cdot 4 + 4 \cdot 3 \cdot 0] \\ &= (-36 + 0 + 32) - (-6 - 32 + 0) = 34.\end{aligned}$$

- Find the determinant of $I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$.

$$\begin{aligned}\det(I_3) &= (1 \cdot 1 \cdot 1 + 0 \cdot 0 \cdot 0 + 0 \cdot 0 \cdot 0) \\ &\quad - (1 \cdot 0 \cdot 0 + 0 \cdot 0 \cdot 1 + 0 \cdot 1 \cdot 0) \\ &= 1 - 0 = 1.\end{aligned}$$

119 / 150

Elementary Row Operation

- Let $A = (a_{ij})_{3 \times 3} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$.

- $\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \xrightarrow{cR_2} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ ca_{21} & ca_{22} & ca_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = B.$

$$\begin{aligned}\det(B) &= (a_{11}ca_{22}a_{33} + a_{12}ca_{23}a_{31} + a_{13}ca_{21}a_{31}) \\ &\quad - (a_{11}ca_{23}a_{32} + a_{12}ca_{21}a_{33} + a_{13}ca_{22}a_{31}) \\ &= c \cdot [(a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{31}) \\ &\quad - (a_{11}a_{23}a_{32} + a_{12}a_{21}a_{33} + a_{13}a_{22}a_{31})] \\ &= c \det(A).\end{aligned}$$

In particular, $I \xrightarrow{cR_i} E \Rightarrow \det(E) = c \cdot \det(I) = c$.

- $A \xrightarrow{cR_i} EA \Rightarrow \det(EA) = c \det(A) = \det(E) \det(A).$

120 / 150

Elementary Row Operation

- Let $\mathbf{A} = (a_{ij})_{3 \times 3} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$.

- $\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \xrightarrow{R_1 \leftrightarrow R_3} \begin{pmatrix} a_{31} & a_{32} & a_{33} \\ a_{21} & a_{22} & a_{23} \\ a_{11} & a_{12} & a_{13} \end{pmatrix} = \mathbf{B}.$

$$\begin{aligned} \det(\mathbf{B}) &= (a_{31}a_{22}a_{13} + a_{32}a_{23}a_{11} + a_{33}a_{21}a_{11}) \\ &\quad - (a_{31}a_{23}a_{12} + a_{32}a_{21}a_{13} + a_{33}a_{22}a_{11}) \\ &= (-1) \cdot [(a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32}) \\ &\quad - (a_{11}a_{23}a_{32} + a_{12}a_{21}a_{33} + a_{13}a_{22}a_{31})] \\ &= -\det(\mathbf{A}). \end{aligned}$$

In particular, $\mathbf{I} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{E} \Rightarrow \det(\mathbf{E}) = -\det(\mathbf{I}) = -1.$

- $\mathbf{A} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{EA} \Rightarrow \det(\mathbf{EA}) = \det(\mathbf{E}) \det(\mathbf{A}).$

121 / 150

Elementary Row Operation

- Let $\mathbf{A} = (a_{ij})_{3 \times 3} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$.

- $\mathbf{A} \xrightarrow{R_2 + cR_1} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} + ca_{11} & a_{22} + ca_{12} & a_{23} + ca_{13} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \mathbf{B}.$

$$\begin{aligned} \det(\mathbf{B}) &= [a_{11}(a_{22} + ca_{12})a_{33} + a_{12}(a_{23} + ca_{13})a_{31} \\ &\quad + a_{13}(a_{21} + ca_{11})a_{32}] \\ &\quad - [a_{11}(a_{23} + ca_{13})a_{32} + a_{12}(a_{21} + ca_{11})a_{33} \\ &\quad + a_{13}(a_{22} + ca_{12})a_{31}] \\ &= \overset{\text{Exercise}}{\dots\dots\dots} = \det(\mathbf{A}). \end{aligned}$$

In particular, $\mathbf{I} \xrightarrow{R_i + cR_j} \mathbf{E} \Rightarrow \det(\mathbf{E}) = \det(\mathbf{I}) = 1.$

- $\mathbf{A} \xrightarrow{R_i + cR_j} \mathbf{EA} \Rightarrow \det(\mathbf{EA}) = \det(\mathbf{E}) \det(\mathbf{A}).$

122 / 150

Elementary Row Operation

- Let A be a square matrix of order 3.
 - For any elementary matrix E of order 3,
 - $\det(EA) = \det(E) \det(A)$.

This property can be used to find $\det(A)$.

- Let R be the reduced row-echelon form of A .
 - Then $R = E_k E_{k-1} \cdots E_2 E_1 A$, E_i elementary.

$$\begin{aligned} \det(R) &= \det(E_k) \det(E_{k-1} \cdots E_2 E_1 A) \\ &= \cdots \cdots \cdots \\ &= \det(E_k) \det(E_{k-1}) \cdots \det(E_2) \det(E_1 A) \\ &= \det(E_k) \det(E_{k-1}) \cdots \det(E_2) \det(E_1) \det(A). \end{aligned}$$

If A is **invertible**, $R = I$, and $\det(R) = 1$.

- $\det(A) = [\det(E_k) \det(E_{k-1}) \cdots \det(E_2) \det(E_1)]^{-1}$

123 / 150

Elementary Row Operation

- Let A be a square matrix of order 3.
 - For any elementary matrix E of order 3,
 - $\det(EA) = \det(E) \det(A)$.
- Let R be the reduced row-echelon form of A .
 - Then $R = E_k E_{k-1} \cdots E_2 E_1 A$, E_i elementary.

$$\det(R) = \det(E_k) \det(E_{k-1}) \cdots \det(E_2) \det(E_1) \det(A).$$

If A is **singular**, then the last row of R is zero.

- $R \xrightarrow{2R_3} R \Rightarrow 2 \det(R) = \det(R) \Rightarrow \det(R) = 0$.

Note that $\det(E) \neq 0$ for any elementary matrix E .

- We must have $\det(A) = 0$.

124 / 150

Example

- Let $A = \begin{pmatrix} -3 & -2 & 4 \\ 4 & 3 & 1 \\ 0 & 2 & 4 \end{pmatrix}$.

$$A \xrightarrow[E_1]{R_2 + \frac{4}{3}R_1} \bullet \xrightarrow[E_2]{R_3 + (-6)R_2} \bullet \xrightarrow[E_3]{-\frac{1}{3}R_1} \bullet \xrightarrow[E_4]{3R_2} \bullet \\ \xrightarrow[E_5]{-\frac{1}{34}R_3} \bullet \xrightarrow[E_6]{R_1 + \frac{4}{3}R_3} \bullet \xrightarrow[E_7]{R_2 + (-19)R_3} \bullet \xrightarrow[E_8]{R_1 + (-\frac{2}{3}R_2)} I.$$

- $\det(E_i) = 1$ if $i = 1, 2, 6, 7, 8$.
 - $\det(E_3) = -\frac{1}{3}$, $\det(E_4) = 3$, $\det(E_5) = -\frac{1}{34}$.
- $\det(A) = [\det(E_1) \cdots \det(E_8)]^{-1} = (\frac{1}{34})^{-1} = 34$.
- We will show that in order to find $\det(A)$ it suffices to use the Gaussian elimination to get a row-echelon form of A .

125 / 150

Determinant

- Definition.** Let $A = (a_{ij})_{n \times n}$. Its **determinant** is:

- If $n = 1$, define $\det(A) = a_{11}$.
 - If $n > 1$, let A_{ij} be its (i, j) -cofactor, define
 - $\det(A) = a_{11}A_{11} + a_{12}A_{12} + \cdots + a_{1n}A_{1n}$.

- Example.** Let $A = (a_{ij})_{4 \times 4} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}$.

$$\det(A) = a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix} \\ + a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} - a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}.$$

126 / 150

Determinant

- **Example.** Find $\det(\mathbf{C})$ if $\mathbf{C} = \begin{pmatrix} 0 & -1 & 2 & 0 \\ 2 & -3 & 3 & -2 \\ 0 & 2 & 4 & 0 \\ 0 & 0 & 2 & -1 \end{pmatrix}$.

$$\begin{aligned} \det(\mathbf{C}) &= 0 \cdot \begin{vmatrix} -3 & 3 & -2 \\ 2 & 4 & 0 \\ 0 & 2 & -1 \end{vmatrix} - (-1) \cdot \begin{vmatrix} 2 & 3 & -2 \\ 0 & 4 & 0 \\ 0 & 2 & -1 \end{vmatrix} \\ &\quad + 2 \cdot \begin{vmatrix} 2 & -3 & -2 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{vmatrix} - 0 \cdot \begin{vmatrix} 2 & -3 & 3 \\ 0 & 2 & 4 \\ 0 & 0 & 2 \end{vmatrix} \\ &= 0 \cdot 10 - (-1) \cdot (-8) + 2 \cdot (-4) - 0 \cdot 8 = -16. \end{aligned}$$

- **Warning:** The “diagonal expansion” of $\det(\mathbf{A})$ for 2×2 or 3×3 matrices is no longer valid if the order ≥ 4 .

127 / 150

Properties

- **Theorem.** $\det(\mathbf{I}) = 1$. For any square matrices,
 - If $\mathbf{A} \xrightarrow{cR_i} \mathbf{B}$, then $\det(\mathbf{B}) = c \det(\mathbf{A})$.
 - In particular, if $\mathbf{I} \xrightarrow{cR_i} \mathbf{E}$, then $\det(\mathbf{E}) = c$.
 - If $\mathbf{A} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{B}$, then $\det(\mathbf{B}) = -\det(\mathbf{A})$.
 - In particular, if $\mathbf{I} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{E}$, then $\det(\mathbf{E}) = -1$.
 - If $\mathbf{A} \xrightarrow{R_i + cR_j} \mathbf{B}$, then $\det(\mathbf{B}) = \det(\mathbf{A})$.
 - In particular, if $\mathbf{I} \xrightarrow{R_i + cR_j} \mathbf{E}$, then $\det(\mathbf{E}) = 1$.
- **Theorem.** Let \mathbf{A} be a square matrix.
 - For any elementary matrix \mathbf{E} of the same order,
 - $\det(\mathbf{EA}) = \det(\mathbf{E}) \det(\mathbf{A})$.

128 / 150

Properties

- **Theorem.** Suppose a square matrix A has a **zero row**.

- Then $\det(A) = 0$.

Proof. Suppose the i th row of square matrix A is 0 .

- $A \xrightarrow{2R_i} A \Rightarrow \det(A) = 2 \det(A) \Rightarrow \det(A) = 0$.

- Suppose square matrices A and B are **row equivalent**.

- There exist elementary matrices E_1, E_2, \dots, E_k s.t.

- $B = E_k \cdots E_2 E_1 A$.

$$\det(B) = \det(E_k) \cdots \det(E_2) \det(E_1) \det(A).$$

Note that $\det(E) \neq 0$ for every elementary matrix E .

- $\det(A) = 0 \Leftrightarrow \det(B) = 0$.
- Equivalently, $\det(A) \neq 0 \Leftrightarrow \det(B) \neq 0$.

129 / 150

Properties

- **Theorem.** Suppose a square matrix A has a **zero row**.

- Then $\det(A) = 0$.

Proof. Suppose the i th row of square matrix A is 0 .

- $A \xrightarrow{2R_i} A \Rightarrow \det(A) = 2 \det(A) \Rightarrow \det(A) = 0$.

- **Theorem.** $\det(A) = 0 \Leftrightarrow A$ is **singular**.

- Equivalently, $\det(A) \neq 0 \Leftrightarrow A$ is **invertible**.

Proof. Suppose A is invertible. Then

- A is row equivalent to I .

- $\det(I) = 1 \neq 0 \Rightarrow \det(A) \neq 0$.

Suppose A is singular. Then the RREF of A is not I .

- The RREF of A has a zero row $\Rightarrow \det(A) = 0$.

130 / 150

Properties

- **Theorem.** Let A, B be square matrices of the same size.

- Then $\det(AB) = \det(A) \det(B)$.

Proof. Suppose that A is invertible. Then

- $A = E_1 E_2 \cdots E_k$ for elementary matrices E_i .

$$\begin{aligned}\det(AB) &= \det(E_1 E_2 \cdots E_k B) \\ &= \det(E_1) \det(E_2) \cdots \det(E_k) \det(B) \\ &= \det(E_1 E_2 \cdots E_k) \det(B) \\ &= \det(A) \det(B).\end{aligned}$$

Suppose that A is singular. Then AB is singular.

- $\det(A) = 0$ and $\det(AB) = 0$.

Then $\det(AB) = 0 = \det(A) \det(B)$.

131 / 150

Properties

- **Theorem.** For any square A , $\det(A) = \det(A^T)$.

Proof. Suppose A is singular.

- Then A^T is also singular, because
 - A^T is invertible $\Rightarrow A = (A^T)^T$ is invertible.

For this case, $\det(A) = 0 = \det(A^T)$.

Suppose A is invertible. Then

- $A = E_1 E_2 \cdots E_k$ for elementary matrices E_i .

Note that $\det(E) = \det(E^T)$ for elementary matrix E .

$$\begin{aligned}\det(A^T) &= \det(E_k^T \cdots E_2^T E_1^T) \\ &= \det(E_k^T) \cdots \det(E_2^T) \det(E_1^T) \\ &= \det(E_k) \cdots \det(E_2) \det(E_1) \\ &= \det(E_1) \det(E_2) \cdots \det(E_k) \\ &= \det(E_1 E_2 \cdots E_k) = \det(A).\end{aligned}$$

132 / 150

Properties

- **Theorem.** Suppose $A = (a_{ij})_{n \times n}$ is **upper triangular**.

- Then $\det(A) = a_{11}a_{22} \cdots a_{nn}$.

- **Example.**

- $$\begin{vmatrix} 1 & 2 & 3 & 4 & 5 \\ 0 & 6 & 7 & 8 & 9 \\ 0 & 0 & 10 & 11 & 12 \\ 0 & 0 & 0 & 13 & 14 \\ 0 & 0 & 0 & 0 & 15 \end{vmatrix} = 1 \cdot 6 \cdot 10 \cdot 13 \cdot 15 = 11700.$$

- **Remarks.**

- The result is also true for **lower triangular matrices**.
- Note that a row-echelon form of a square matrix is always upper triangular.
 - To find the determinant using elementary row operation, it suffices to use Gaussian elimination to get a row-echelon form.

133 / 150

Properties

- **Theorem.** Suppose $A = (a_{ij})_{n \times n}$ is **upper triangular**.

- Then $\det(A) = a_{11}a_{22} \cdots a_{nn}$.

Idea of Proof. Suppose that $a_{ii} \neq 0$ for all $i = 1, \dots, n$.

- Illustration:

$$A = \begin{pmatrix} a_{11} & * & * & * \\ 0 & a_{22} & * & * \\ 0 & 0 & a_{33} & * \\ 0 & 0 & 0 & a_{44} \end{pmatrix} \xrightarrow{\text{multiply } a_{ii}^{-1} \text{ to } R_i} \begin{pmatrix} 1 & * & * & * \\ 0 & 1 & * & * \\ 0 & 0 & 1 & * \\ 0 & 0 & 0 & 1 \end{pmatrix} = B$$

$$\xrightarrow{\text{convert to RREF}} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = I.$$

Therefore, $\det(A) = a_{11} \cdots a_{nn} \det(I) = a_{11} \cdots a_{nn}$.

134 / 150

Properties

- **Theorem.** Suppose $A = (a_{ij})_{n \times n}$ is **upper triangular**.

- Then $\det(A) = a_{11}a_{22} \cdots a_{nn}$.

Idea of Proof. Suppose that $a_{ii} = 0$ for some i .

- Illustration: Assume that $a_{22} = 0$ but $a_{33} \neq 0, a_{44} \neq 0$.

$$\begin{aligned}
 \begin{pmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & \otimes & * \\ 0 & 0 & 0 & \otimes \end{pmatrix} &\xrightarrow[\text{to the 3rd and 4th rows}]{\text{row operations}} \begin{pmatrix} * & * & * & * \\ 0 & 0 & * & * \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\
 &\xrightarrow[\text{to the 2nd row}]{\text{add multiples of 3rd and 4th rows}} \begin{pmatrix} * & * & * & * \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.
 \end{aligned}$$

A is row equivalent to a singular matrix; $\det(A) = 0 = a_{11} \cdots a_{nn}$.

135 / 150

Cofactor Expansion

- **Theorem.** Let A be a square matrix of order n .

- Let A_{ij} denote the (i, j) -cofactor of A .

Then for any i and j ,

- $\det(A) = a_{i1}A_{i1} + a_{i2}A_{i2} + \cdots + a_{in}A_{in}$.
- $\det(A) = a_{1j}A_{1j} + a_{2j}A_{2j} + \cdots + a_{nj}A_{nj}$.

Idea of Proof. Suppose A is 4×4 . Let $B = (b_{ij})$ be obtained from A by

- Moving the 4th row to the top. Then $b_{1j} = a_{4j}$.

$$\bullet \quad A = \begin{pmatrix} R_1 \\ R_2 \\ R_3 \\ R_4 \end{pmatrix} \xrightarrow{R_3 \leftrightarrow R_4, R_2 \leftrightarrow R_3, R_1 \leftrightarrow R_2} \begin{pmatrix} R_4 \\ R_1 \\ R_2 \\ R_3 \end{pmatrix} = B.$$

So $\det(B) = (-1)^{4-1} \det(A)$.

136 / 150

Cofactor Expansion

- **Idea of Proof.** Suppose A is 4×4 . Let $B = (b_{ij})$ be obtained from A by

- Moving the 4th row to the top. Then $b_{1j} = a_{4j}$.

$$\bullet \quad A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix} \dashrightarrow \begin{pmatrix} a_{41} & a_{42} & a_{43} & a_{44} \\ a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{pmatrix} = B.$$

So $\det(B) = (-1)^{4-1} \det(A)$.

The submatrix obtained from B by deleting its 1st row and j th column is the same as that obtained from A by deleting its 4th row and j th column, say M_{ij} .

- Let B_{ij} be the (i, j) -cofactor of B .
 - $B_{1j} = (-1)^{1+j} \det(M_{ij})$, $A_{4j} = (-1)^{4+j} \det(M_{ij})$.

So $B_{1j} = (-1)^{4-1} A_{4j}$, $j = 1, \dots, n$.

137 / 150

Cofactor Expansion

- **Proof.** Fix i . Let $B = (b_{ij})$ be obtained from A by

- Moving the i th row to the top.

Then $b_{1j} = a_{ij}$, $B_{1j} = (-1)^{i-1} A_{ij}$ for any j .

$$\begin{aligned} \det(A) &= (-1)^{i-1} \det(B) \\ &= (-1)^{i-1} \cdot (b_{11}B_{11} + b_{12}B_{12} + \dots + b_{1n}B_{1n}) \\ &= (-1)^{i-1} \cdot [a_{i1}(-1)^{i-1}A_{i1} + a_{i2}(-1)^{i-1}A_{i2} \\ &\quad + \dots + a_{in}(-1)^{i-1}A_{in}] \\ &= (-1)^{2i-2}(a_{i1}A_{i1} + a_{i2}A_{i2} + \dots + a_{in}A_{in}) \\ &= a_{i1}A_{i1} + a_{i2}A_{i2} + \dots + a_{in}A_{in}. \end{aligned}$$

This is called the **cofactor expansion along the i th row**.

- The proof for the **cofactor expansion along the j th column** is left as exercises. (*Hint: Consider A^T .*)

138 / 150

Cofactor Expansion

- In evaluating the determinant using cofactor expansion,
 - expand along the row or column with the most zeros.

• **Example.** $A = \begin{pmatrix} 0 & -1 & 2 & 0 \\ 2 & -3 & 3 & -2 \\ 0 & 2 & 4 & 0 \\ 0 & 0 & 2 & -1 \end{pmatrix}.$

$$\det(\mathbf{A}) = a_{11}A_{11} + a_{21}A_{21} + a_{31}A_{31} + a_{41}A_{41}$$

$$= 2 \cdot (-1)^{2+1} \begin{vmatrix} -1 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 2 & -1 \end{vmatrix}$$

$$= -2 \cdot (-1) \cdot (-1)^{3+3} \begin{vmatrix} -1 & 2 \\ 2 & 4 \end{vmatrix}$$

$$= -16.$$

139 / 150

Finding Determinant

- Find $\det(\mathbf{A})$ if \mathbf{A} is a square matrix of order n .
 - If \mathbf{A} has a zero row/column, then $\det(\mathbf{A}) = 0$.
 - If \mathbf{A} is triangular, $\det(\mathbf{A}) = a_{11} \cdots a_{nn}$.
 - Suppose that \mathbf{A} is not triangular.
 - If $n = 2$, use formula $\det(\mathbf{A}) = a_{11}a_{22} - a_{12}a_{21}$.
 - If a row/coln has many 0, use cofactor expansion.
 - Otherwise, use ele. row operations to get REF:
 - $\det(\mathbf{EA}) = \det(\mathbf{E}) \det(\mathbf{A})$.
- Note the following formulas (exercises for the last two):
 - $\det(\mathbf{A}) = \det(\mathbf{A}^T)$.
 - $\det(\mathbf{AB}) = \det(\mathbf{A}) \det(\mathbf{B})$.
 - $\det(c\mathbf{A}) = c^n \det(\mathbf{A})$, where \mathbf{A} is $n \times n$.
 - $\det(\mathbf{A}^{-1}) = \det(\mathbf{A})^{-1}$ if \mathbf{A} is invertible.

140 / 150

Examples

- Find $\det(\mathbf{A})$, where $\mathbf{A} = \begin{pmatrix} 3 & -1 & 1 & 1 \\ 3 & -1 & 2 & 1 \\ 0 & 2 & 4 & 0 \\ 0 & 0 & 2 & -1 \end{pmatrix}$.

$$\begin{aligned} & \begin{pmatrix} 3 & -1 & 1 & 1 \\ 3 & -1 & 2 & 1 \\ 0 & 2 & 4 & 0 \\ 0 & 0 & 2 & -1 \end{pmatrix} \xrightarrow[\mathbf{E}_1]{R_2 + (-1)R_1} \begin{pmatrix} 3 & -1 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 2 & 4 & 0 \\ 0 & 0 & 2 & -1 \end{pmatrix} \\ & \xrightarrow[\mathbf{E}_2]{R_2 \leftrightarrow R_3} \begin{pmatrix} 3 & -1 & 1 & 1 \\ 0 & 2 & 4 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 2 & -1 \end{pmatrix} \xrightarrow[\mathbf{E}_3]{R_4 + (-2)R_3} \begin{pmatrix} 3 & -1 & 1 & 1 \\ 0 & 2 & 4 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} = \mathbf{B}. \end{aligned}$$

- $\det(\mathbf{B}) = 3 \cdot 2 \cdot 1 \cdot (-1) = -6$.
- $\det(\mathbf{A}) = [\det(\mathbf{E}_1) \det(\mathbf{E}_2) \det(\mathbf{E}_3)]^{-1} \det(\mathbf{B}) = 6$.

141 / 150

Examples

- Find $\det(\mathbf{A})$, where $\mathbf{A} = \begin{pmatrix} 3 & -1 & 1 & 1 \\ 3 & -1 & 2 & 1 \\ 0 & 2 & 4 & 0 \\ 0 & 0 & 2 & -1 \end{pmatrix}$.

- Expand along the 4th row: $\det(\mathbf{A}) = (-1)(-6) = 6$.

$$\bullet \quad 2 \cdot (-1)^{4+3} \begin{vmatrix} 3 & -1 & 1 \\ 3 & -1 & 1 \\ 0 & 2 & 0 \end{vmatrix} + (-1) \cdot (-1)^{4+4} \begin{vmatrix} 3 & -1 & 1 \\ 3 & -1 & 2 \\ 0 & 2 & 4 \end{vmatrix}$$

$$\bullet \quad \begin{vmatrix} 3 & -1 & 1 \\ 3 & -1 & 1 \\ 0 & 2 & 0 \end{vmatrix} \xrightarrow{R_2 + (-1)R_1} \begin{vmatrix} 3 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 2 & 0 \end{vmatrix} = 0$$

$$\bullet \quad \begin{vmatrix} 3 & -1 & 1 \\ 3 & -1 & 2 \\ 0 & 2 & 4 \end{vmatrix} = 3 \cdot \begin{vmatrix} -1 & 2 \\ 2 & 4 \end{vmatrix} - 3 \cdot \begin{vmatrix} -1 & 1 \\ 2 & 4 \end{vmatrix} = -6.$$

142 / 150

Examples

- Let $\mathbf{A} = \begin{pmatrix} -3 & -2 & 4 \\ 4 & 3 & 1 \\ 0 & 2 & 4 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 2 \end{pmatrix}$.
 - It is given that $\det(\mathbf{A}) = 34$ and $\det(\mathbf{B}) = -1$.
 - $\det(\mathbf{A}^T) = \det(\mathbf{A}) = 34$.
 - $\det(2\mathbf{A}) = 2^3 \det(\mathbf{A}) = 8 \cdot 34 = 272$.
 - $\det(\mathbf{A}^{-1}) = [\det(\mathbf{A})]^{-1} = \frac{1}{34}$.
 - $\det(\mathbf{AB}) = \det(\mathbf{A}) \det(\mathbf{B}) = 34 \cdot (-1) = -34$.
 - $\det(\mathbf{BA}) = \det(\mathbf{B}) \det(\mathbf{A}) = (-1) \cdot 34 = -34$.

143 / 150

Adjoint Matrix

- Definition.** Let \mathbf{A} be a square matrix of order n . The **(classical) adjoint** (or **adjugate**, or **adjunct**) of \mathbf{A} is

- $\text{adj}(\mathbf{A}) = (A_{ji})_{n \times n}$

where A_{ij} is the (i, j) -cofactor of \mathbf{A} .

- Example.** Let $\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$. Then
 - $A_{11} = a_{22}, A_{12} = -a_{21}, A_{21} = -a_{12}, A_{22} = a_{11}$.
 - $\text{adj}(\mathbf{A}) = \begin{pmatrix} A_{11} & A_{21} \\ A_{12} & A_{22} \end{pmatrix} = \begin{pmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{pmatrix}$.

Recall that if \mathbf{A} is invertible, then

- $\mathbf{A}^{-1} = \frac{1}{a_{11}a_{22} - a_{12}a_{21}} \begin{pmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{pmatrix}$.

It is conjectured that $\mathbf{A}^{-1} = [\det(\mathbf{A})]^{-1} \text{adj}(\mathbf{A})$.

144 / 150

Adjoint Matrix

- **Theorem.** Let A be a square matrix. Then

- $A[\text{adj}(A)] = \det(A)I$.

Proof. Let $A[\text{adj}(A)] = (c_{ij})$. Then

- $c_{ij} = a_{i1}A_{j1} + a_{i2}A_{j2} + \cdots + a_{in}A_{jn}$.

Let $i = j$. Then $c_{ii} = a_{i1}A_{i1} + \cdots + a_{in}A_{in} = \det(A)$.

Suppose that $i \neq j$. Let B be the matrix obtained from A by replacing j th row by the i th row.

- $B \xrightarrow{R_i \leftrightarrow R_j} B \Rightarrow \det(B) = -\det(B) \Rightarrow \det(B) = 0$.

$$\begin{aligned} c_{ij} &= a_{i1}A_{j1} + a_{i2}A_{j2} + \cdots + a_{in}A_{jn} \\ &= b_{j1}B_{j1} + b_{j2}B_{j2} + \cdots + b_{jn}B_{jn} \\ &= \det(B) = 0. \end{aligned}$$

Therefore, $A[\text{adj}(A)] = \det(A)I$.

145 / 150

Adjoint Matrix

- **Theorem.** Let A be a square matrix. Then

- $A[\text{adj}(A)] = \det(A)I$.

- $[\text{adj}(A)]A = \det(A)I$. (Exercise!)

If A is invertible, then $A^{-1} = \frac{1}{\det(A)} \text{adj}(A)$.

- **Exercises.** Let A, B be invertible matrices of order n .

- Find $[\text{adj}(A)]^{-1}$ and $\text{adj}(A^{-1})$.
- Find $\det(\text{adj}(A))$ and $\text{adj}(\text{adj}(A))$.
- Prove that $\text{adj}(AB) = \text{adj}(B) \text{adj}(A)$.

- **Challenging Problems.** Suppose A and B are not necessarily invertible.

- Find $\det(\text{adj}(A))$ and $\text{adj}(\text{adj}(A))$.
- Is it true that $\text{adj}(AB) = \text{adj}(B) \text{adj}(A)$?

146 / 150

Example

- Let $\mathbf{A} = \begin{pmatrix} 1 & -1 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 3 \end{pmatrix}$. $\det(\mathbf{A}) = (-1) \begin{vmatrix} 1 & 1 \\ 1 & 3 \end{vmatrix} = -2$.

$$\begin{aligned} \text{adj}(\mathbf{A}) &= \begin{pmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{pmatrix} \\ &= \begin{pmatrix} \begin{vmatrix} -1 & 0 \\ 0 & 3 \end{vmatrix} & -\begin{vmatrix} -1 & 1 \\ 0 & 3 \end{vmatrix} & \begin{vmatrix} -1 & 1 \\ -1 & 0 \end{vmatrix} \\ -\begin{vmatrix} 0 & 0 \\ 1 & 3 \end{vmatrix} & \begin{vmatrix} 1 & 1 \\ 1 & 3 \end{vmatrix} & -\begin{vmatrix} 1 & 1 \\ 0 & 0 \end{vmatrix} \\ \begin{vmatrix} 0 & -1 \\ 1 & 0 \end{vmatrix} & -\begin{vmatrix} 1 & -1 \\ 1 & 0 \end{vmatrix} & \begin{vmatrix} 1 & -1 \\ 0 & -1 \end{vmatrix} \end{pmatrix} = \begin{pmatrix} -3 & 3 & 1 \\ 0 & 2 & 0 \\ 1 & -1 & -1 \end{pmatrix} \\ \mathbf{A}^{-1} &= \frac{1}{\det(\mathbf{A})} \text{adj}(\mathbf{A}) = \begin{pmatrix} \frac{3}{2} & -\frac{3}{2} & -\frac{1}{2} \\ 0 & -1 & 0 \\ -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} \end{pmatrix} \end{aligned}$$

147 / 150

Cramer's Rule

- Let $\mathbf{A} = (a_{ij})_{n \times n}$ be an invertible matrix.
 - The linear system $\mathbf{A}\mathbf{x} = \mathbf{b}$ has a unique solution
 - $\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$.

Recall that $\mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} [\text{adj}(\mathbf{A})]$. Let $\mathbf{b} = (b_i)_{n \times 1}$.

$$x_j = \frac{1}{\det(\mathbf{A})} (A_{1j}b_1 + A_{2j}b_2 + \cdots + A_{nj}b_n).$$

Fix j , and let \mathbf{A}_j be the matrix obtained by replacing the j th column of \mathbf{A} by \mathbf{b} . Then

- b_i is the (i, j) -entry of \mathbf{A}_j .
- A_{ij} is the (i, j) -cofactor of \mathbf{A}_j .

Therefore, $x_j = \frac{\det(\mathbf{A}_j)}{\det(\mathbf{A})}$, $j = 1, \dots, n$.

148 / 150

Cramer's Rule

- **Cramer's Rule.** Let A be an invertible matrix of order n .
 - For every column matrix b of size $n \times 1$, the linear system $Ax = b$ has a unique solution

- $$x = \frac{1}{\det(A)} \begin{pmatrix} \det(A_1) \\ \vdots \\ \det(A_n) \end{pmatrix},$$

A_j is obtained from A by replacing its j th coln by b .

- **Example.** Let $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ and $b = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}$.

- Suppose that A is invertible. $Ax = b$ implies

- $$x = \frac{1}{\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}} \begin{pmatrix} \begin{vmatrix} b_1 & a_{12} \\ b_2 & a_{22} \end{vmatrix} \\ \begin{vmatrix} a_{11} & b_1 \\ a_{21} & b_2 \end{vmatrix} \end{pmatrix}$$

149 / 150

Example

- $$\begin{pmatrix} 1 & 1 & 3 \\ 2 & -2 & 2 \\ 3 & 9 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 4 \\ 3 \end{pmatrix}. \quad \begin{vmatrix} 1 & 1 & 3 \\ 2 & -2 & 2 \\ 3 & 9 & 0 \end{vmatrix} = 60.$$

- $$x = \frac{\begin{vmatrix} 0 & 1 & 3 \\ 4 & -2 & 2 \\ 3 & 9 & 0 \end{vmatrix}}{60} = \frac{132}{60} = 2.2$$

- $$y = \frac{\begin{vmatrix} 1 & 0 & 3 \\ 2 & 4 & 2 \\ 3 & 3 & 0 \end{vmatrix}}{60} = \frac{-24}{60} = -0.4$$

- $$z = \frac{\begin{vmatrix} 1 & 1 & 0 \\ 2 & -2 & 4 \\ 3 & 9 & 3 \end{vmatrix}}{60} = \frac{-36}{60} = -0.6$$

150 / 150