

MA2001 LINEAR ALGEBRA

VECTOR SPACES

Goh Jun Le / Wang Fei

gohjunle@nus.edu.sg / matwf@nus.edu.sg


Department of Mathematics
Office: S17-06-25 / S17-06-16
Tel: 6601-1355 / 6516-2937

Euclidean n-Spaces	2
Vectors in xy -Plane	3
Operations on Vectors	5
Vectors in xyz -Space	11
Euclidean Spaces	12
Implicit and Explicit Forms	15
Linear Combinations and Linear Spans	24
Linear Combination	25
Linear Span	28
Criterion for $\text{span}(S) = \mathbb{R}^n$	32
Properties of Linear Spans	37
Subspaces	45
Subspaces	46
Subspaces of $\mathbb{R}^1, \mathbb{R}^2, \mathbb{R}^3$	49
Solution Space	52
Linear Independence	56
Linear Independence	57
Properties	61
Bases	67
Motivation	68
Properties	74
Coordinate Vector	76
Properties	80
Dimensions	85
Criterion for Bases	86
Dimension	88
Properties	91

Transition Matrices	98
Coordinate Vector	99
Transition Matrix.....	101
Properties	104

Vectors in xy -Plane

- Recall the xy -plane:




- Every point P on the plane is represented by (a, b) .
 - a is the x -coordinate and b is the y -coordinate.
- The arrow from the origin O to the point P is called a **vector**, denoted by $\overrightarrow{OP} = \mathbf{v} = (a, b)$.

3 / 106

Vectors in xy -Plane

- A **vector** represents the change from the **initial point** to the **end point**.




If \overrightarrow{PQ} is **parallel** shifted to $\overrightarrow{P'Q'}$, then

- $\overrightarrow{PQ} = \overrightarrow{P'Q'}$,
 - that is, $(a_2 - a_1, b_2 - b_1) = (a'_2 - a'_1, b'_2 - b'_1)$,
 - that is, $a_2 - a_1 = a'_2 - a'_1$ & $b_2 - b_1 = b'_2 - b'_1$.

4 / 106

Length

- Let $\mathbf{v} = (v_1, v_2)$ be a vector in xy -plane.



- Its **length** is $\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2}$
- If \mathbf{v} is the vector from $P(a_1, b_1)$ to $Q(a_2, b_2)$.
 - $\mathbf{v} = (a_2 - a_1, b_2 - b_1)$.
 - $\|\mathbf{v}\| = \sqrt{(a_2 - a_1)^2 + (b_2 - b_1)^2}$.

5 / 106

Scalar Multiplication

- Scalar Multiplication.** Let $\mathbf{v} = (v_1, v_2)$ and $c \in \mathbb{R}$.

- Then $c\mathbf{v} = (cv_1, cv_2)$

Geometric interpretation:

- $c\mathbf{v}$ is a vector parallel to \mathbf{v} such that
 - its length is $|c|$ times the length of \mathbf{v} .
 - 1. If $c = 0$, then $c\mathbf{v} = 0\mathbf{v} = \mathbf{0}$ is the **zero vector**.
 - 2. If $c > 0$, then $c\mathbf{v}$ has the same direction as \mathbf{v} .
 - 3. If $c < 0$, then $c\mathbf{v}$ has the opposite direction of \mathbf{v} .

In particular, $(-1)\mathbf{v}$ is the **negative** of \mathbf{v} , denoted by $-\mathbf{v}$.


- Example.** Let $\mathbf{v} = (2, 1)$. Then
 - $0\mathbf{v} = (0, 0)$, $-\mathbf{v} = (-2, -1)$, $2\mathbf{v} = (4, 2)$.
 - $(-2)\mathbf{v} = (-4, -2) = -(2\mathbf{v}) = 2(-\mathbf{v})$.

6 / 106

Scalar Multiplication

- **Scalar Multiplication.** Let $\mathbf{v} = (v_1, v_2)$ and $c \in \mathbb{R}$.

- Then $c\mathbf{v} = (cv_1, cv_2)$



- **Properties & Exercises.**

- $c(d\mathbf{v}) = (cd)\mathbf{v} = d(c\mathbf{v})$.
- In particular, $-c\mathbf{v} = (-c)\mathbf{v} = c(-\mathbf{v})$, $-(-\mathbf{v}) = \mathbf{v}$.
- $\|c\mathbf{v}\| = |c| \|\mathbf{v}\|$, $\mathbf{v} = \mathbf{0} \Leftrightarrow \|\mathbf{v}\| = 0$.

7 / 106


Addition and Subtraction

- **Addition.** Let $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$.

- Then $\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2)$

Geometric interpretation:

- Parallel shift \mathbf{v} so that its initial point is the same as the end of \mathbf{u} . Then $\mathbf{u} + \mathbf{v}$ is the vector from the initial point of \mathbf{u} to the end point of \mathbf{v} .



8 / 106

Addition and Subtraction


- **Subtraction.** Let $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$.

- Then $\mathbf{u} - \mathbf{v} = (u_1 - v_1, u_2 - v_2)$

Note that $\mathbf{u} - \mathbf{v} = \mathbf{u} + (-\mathbf{v})$.

Geometric interpretation.

- Parallel shift \mathbf{v} so that \mathbf{u} and \mathbf{v} have the same initial point. Then $\mathbf{u} - \mathbf{v}$ is the vector from the end point of \mathbf{v} to the end point of \mathbf{u} .



9 / 106

Addition and Subtraction

- **Example.** Let $\mathbf{u} = (2, 3)$ and $\mathbf{v} = (4, -5)$.

- $\mathbf{u} + \mathbf{v} = (2 + 4, 3 + (-5)) = (6, -2)$.
- $\mathbf{u} - \mathbf{v} = (2 - 4, 3 - (-5)) = (-2, 8)$.


- **Properties & Exercises.** Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be vectors in the xy -plane and $c, d \in \mathbb{R}$.

- $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$.
- $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$.
- $\mathbf{0} + \mathbf{v} = \mathbf{v}$, where $\mathbf{0}$ is the zero vector.
- $\mathbf{v} + (-\mathbf{v}) = \mathbf{0}$.
- $c(d\mathbf{v}) = (cd)\mathbf{v} = d(c\mathbf{v})$.
- $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$.
- $(c + d)\mathbf{v} = c\mathbf{v} + d\mathbf{v}$.
- $1\mathbf{v} = \mathbf{v}$.

10 / 106

Vectors in xyz -Space

- Consider the xyz -space:



The **vector** $\mathbf{v} = \overrightarrow{OP}$ is the arrow from the origin O to P , denoted by $\mathbf{v} = (a, b, c)$.

11 / 106

Euclidean Spaces

- Definition.** An **n -vector** or **ordered n -tuple** of real numbers is $\mathbf{v} = (v_1, v_2, \dots, v_i, \dots, v_n)$.

- $v_i \in \mathbb{R}$ is the **i th component** or **i th coordinate** of \mathbf{v} .

Let $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$.

- \mathbf{u} and \mathbf{v} are **equal** if $u_i = v_i$ for all $i = 1, \dots, n$.
- The n -vector $\mathbf{0} = (0, 0, \dots, 0)$ is the **zero vector**.
- Let $c \in \mathbb{R}$. The **scalar multiple** $c\mathbf{v}$ is
 - $c\mathbf{v} = (cv_1, cv_2, \dots, cv_n)$.
- The **negative** of \mathbf{v} is $(-1)\mathbf{v}$, denoted by $-\mathbf{v}$.
- The **addition** $\mathbf{u} + \mathbf{v}$ is
 - $\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2, \dots, u_n + v_n)$.
- The **subtraction** $\mathbf{u} - \mathbf{v}$ is
 - $\mathbf{u} - \mathbf{v} = (u_1 - v_1, u_2 - v_2, \dots, u_n - v_n)$.

12 / 106

Euclidean Spaces

- **Notation.** An n -vector (v_1, v_2, \dots, v_n) can be viewed as

- a **row matrix** (**row vector**) $(v_1 \ v_2 \ \cdots \ v_n)$,
- a **column matrix** (**column vector**) $\begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix}$.

- **Properties.** Let u, v, w be n -vectors and $c, d \in \mathbb{R}$.

- $u + v = v + u$.
- $(u + v) + w = u + (v + w)$.
- $v + \mathbf{0} = v$ and $v + (-v) = \mathbf{0}$.
- $c(u + v) = cu + cv$.
- $(c + d)v = cv + dv$.
- $c(dv) = (cd)v$.
- $1v = v$. (Verification is left as exercise.)

13 / 106

Euclidean Spaces

- The **Euclidean n -space** (or simply **n -space**) is the set of all n -vectors of real numbers.

- $\mathbb{R}^n = \{(v_1, v_2, \dots, v_n) \mid v_1, v_2, \dots, v_n \in \mathbb{R}\}$.

$v \in \mathbb{R}^n$ if and only if v is of the form

- $v = (v_1, v_2, \dots, v_n)$ for real numbers v_1, v_2, \dots, v_n .

- In particular,

- If $n = 1$, then $\mathbb{R} = \mathbb{R}^1$ is the real line.
- If $n = 2$, then \mathbb{R}^2 is the xy -plane.
- If $n = 3$, then \mathbb{R}^3 is the xyz -space.

- Linear system $Ax = b$ in m equations and n variables.

- x can be viewed as an n -vector, i.e., $x \in \mathbb{R}^n$.

Then the solution set of $Ax = b$ is a subset of \mathbb{R}^n .

14 / 106

Implicit and Explicit Forms

- A linear system is given in the **implicit form**:

$$\circ \begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

- **Example.** $\begin{cases} x + y + z = 0, \\ x - y + 2z = 1. \end{cases}$

- An implicit form of the solution set is

- $\{(x, y, z) \mid x + y + z = 0 \text{ and } x - y + 2z = 1\}.$

Geometrically, the solution set is the intersection of two non-parallel planes, which is a straight line in \mathbb{R}^3 .

15 / 106

Implicit and Explicit Forms

- A linear system is given in the **implicit form**:

$$\circ \begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

Its general solution is in the **explicit form**.

- **Example.** $\begin{cases} x + y + z = 0, \\ x - y + 2z = 1. \end{cases}$

$$\circ \left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 1 & -1 & 2 & 1 \end{array} \right) \xrightarrow{R_2 + (-1)R_1} \left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 0 & -2 & 1 & 1 \end{array} \right).$$

- $x = \frac{1}{2} - \frac{3}{2}t, y = -\frac{1}{2} + \frac{1}{2}t, z = t, \text{ where } t \in \mathbb{R}.$

- An explicit form of the solution set is

- $\{(\frac{1}{2} - \frac{3}{2}t, -\frac{1}{2} + \frac{1}{2}t, t) \mid t \in \mathbb{R}\}.$

16 / 106

Lines in \mathbb{R}^2

- A **straight line** in \mathbb{R}^2 is of the form
 - $ax + by = c$, where a and b are not both zero.

Implicit form: $\{(x, y) \mid ax + by = c\}$.

Explicit form:

- If $a \neq 0$, then $y = t$ and $x = \frac{c - bt}{a}$.
 - $\left\{ \left(\frac{c - bt}{a}, t \right) \mid t \in \mathbb{R} \right\}$.
- If $b \neq 0$, then $x = s$ and $y = \frac{c - as}{b}$.
 - $\left\{ \left(s, \frac{c - as}{b} \right) \mid s \in \mathbb{R} \right\}$.

17 / 106

Lines in \mathbb{R}^2

- A **straight line** in \mathbb{R}^2 is determined by a point (x_0, y_0) on the line, and its direction vector $(a, b) \neq \mathbf{0}$.
 - A point on the line is of the form $(x_0, y_0) + t(a, b)$.

Explicit form of the line:

- $\{(x_0 + ta, y_0 + tb) \mid t \in \mathbb{R}\}$.
- **Example.** Suppose a line has an explicit form:
 - $\{(2 + 5t, 3 - 2t) \mid t \in \mathbb{R}\}$.

Find an implicit form of the line.

Solution. $x = 2 + 5t$ and $y = 3 - 2t$.

- $t = \frac{x - 2}{5}$ and $t = \frac{3 - y}{2}$.
- $\frac{x - 2}{5} = \frac{3 - y}{2} \Rightarrow \{(x, y) \mid 2x + 5y = 19\}$.

18 / 106

Planes in \mathbb{R}^3

- A **plane** in \mathbb{R}^3 is of the form
 - $ax + by + cz = d$, where a, b, c are not all zero.

Implicit form: $\{(x, y, z) \mid ax + by + cz = d\}$.


Explicit form:

- If $a \neq 0$, $\left\{ \left(\frac{d - bs - ct}{a}, s, t \right) \mid s, t \in \mathbb{R} \right\}$.
- If $b \neq 0$, $\left\{ \left(s, \frac{d - as - ct}{b}, t \right) \mid s, t \in \mathbb{R} \right\}$.
- If $c \neq 0$, $\left\{ \left(s, t, \frac{d - as - bt}{c} \right) \mid s, t \in \mathbb{R} \right\}$.

19 / 106

Planes in \mathbb{R}^3

- Three **non-collinear** points A, B, C determines a plane.



$$\begin{aligned} \mathbf{r} - \mathbf{a} &= \overrightarrow{AR} = \overrightarrow{AB'} + \overrightarrow{AC'} \\ &= s\overrightarrow{AB} + t\overrightarrow{AC} \\ &= s\mathbf{u} + t\mathbf{v}. \end{aligned}$$

- $\boxed{\mathbf{r} = \mathbf{a} + s\mathbf{u} + t\mathbf{v}, \quad s, t \in \mathbb{R}}$

20 / 106

Planes in \mathbb{R}^3

- A **plane** in \mathbb{R}^3 can be explicitly represented as
 - $\{(x_0, y_0, z_0) + s(a_1, b_1, c_1) + t(a_2, b_2, c_2) \mid s, t \in \mathbb{R}\}$,
 (x_0, y_0, z_0) is a point on the plane, and (a_1, b_1, c_1) & (a_2, b_2, c_2) are non-parallel vectors parallel to the plane.
- **Example.** A plane is given by
 - $\{(1 + s - t, 2 + s - 2t, 4 - s - 3t) \mid s, t \in \mathbb{R}\}$.Let $x = 1 + s - t$, $y = 2 + s - 2t$, $z = 4 - s - 3t$.
 - $\left(\begin{array}{cc|c} 1 & -1 & x-1 \\ 1 & -2 & y-2 \\ -1 & -3 & z-4 \end{array} \right) \xrightarrow[\text{Elimination}]{\text{Gaussian}} \left(\begin{array}{cc|c} 1 & -1 & x-1 \\ 0 & -1 & -x+y-1 \\ 0 & 0 & 5x-4y+z-1 \end{array} \right)$.The system is consistent. So $5x - 4y + z - 1 = 0$.
 - Implicit form:
 - $\{(x, y, z) \mid 5x - 4y + z = 1\}$.

21 / 106

Lines in \mathbb{R}^3

- A **straight line** in \mathbb{R}^3 is the intersection of two non-parallel planes. An implicit form is
 - $\{(x, y, z) \mid a_1x + b_1y + c_1z = d_1 \text{ \& } a_2x + b_2y + c_2z = d_2\}$,
 a_i, b_i, c_i not all zero, and the planes are not parallel.
- **Example.** Suppose a line is the intersection of
 - $x + 2y + 3z = 4$ and $2x + 3y + 4z = 5$.Solve the system to have
 - $x = t - 2$, $y = -2t + 3$ and $z = t$.An explicit form of the line:
 - $\{(t - 2, -2t + 3, t) \mid t \in \mathbb{R}\}$.Note that $(t - 2, -2t + 3, t) = (-2, 3, 0) + t(1, -2, 1)$.

22 / 106

Lines in \mathbb{R}^3

- A **straight line** in \mathbb{R}^3 is determined by a point (x_0, y_0, z_0) on the line, and its direction vector $(a, b, c) \neq \mathbf{0}$.
 - A point on the line: $(x_0, y_0, z_0) + t(a, b, c)$.

Explicit form: $\{(x_0 + ta, y_0 + tb, z_0 + tc) \mid t \in \mathbb{R}\}$.

In order to have an implicit form, we need to find two non-parallel planes $ax + by + cz = d$ containing the line.

- **Example.** $\{(t - 2, -2t + 3, t + 1) \mid t \in \mathbb{R}\}$.
 - $x = t - 2$ and $y = -2t + 3$
 - $y = -2(2 + x) + 3 \Rightarrow 2x + y = -1$.
 - $x = t - 2$ and $z = t + 1$.
 - $x - z = -3$.

An implicit form of the line is

- $\{(x, y, z) \mid 2x + y = -1 \ \& \ x - z = -3\}$.

23 / 106

Linear Combinations and Linear Spans

24 / 106

Linear Combination

- Recall the operations on vectors in \mathbb{R}^n .
 - If $\mathbf{v} = (v_1, \dots, v_n) \in \mathbb{R}^n$ and $c \in \mathbb{R}$,
 - $c\mathbf{v} = (cv_1, \dots, cv_n)$.
 - If $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{v} = (v_1, \dots, v_n) \in \mathbb{R}^n$,
 - $\mathbf{u} + \mathbf{v} = (u_1 + v_1, \dots, u_n + v_n)$.
- **Definition.** Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ be vectors in \mathbb{R}^n .
 - A **linear combination** of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ has the form
 - $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k$,where $c_1, c_2, \dots, c_k \in \mathbb{R}$.
 - In particular, $\mathbf{0}$ is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$:
 - $\mathbf{0} = 0\mathbf{v}_1 + 0\mathbf{v}_2 + \dots + 0\mathbf{v}_k$.

25 / 106

Examples

- Let $v_1 = (2, 1, 3)$, $v_2 = (1, -1, 2)$ and $v_3 = (3, 0, 5)$.
 - Is $v = (3, 3, 4)$ a linear combination of v_1, v_2, v_3 ?

Suppose that $v = av_1 + bv_2 + cv_3$, i.e.,

$$\begin{aligned}(3, 3, 4) &= a(2, 1, 3) + b(1, -1, 2) + c(3, 0, 5) \\ &= (2a + b + 3c, a - b, 3a + 2b + 5c).\end{aligned}$$

Solve the linear system
$$\begin{cases} 2a + b + 3c = 3 \\ a - b = 3 \\ 3a + 2b + 5c = 4. \end{cases}$$

$$\circ \left(\begin{array}{ccc|c} 2 & 1 & 3 & 3 \\ 1 & -1 & 0 & 3 \\ 3 & 2 & 5 & 4 \end{array} \right) \xrightarrow[\text{elimination}]{\text{Gaussian}} \left(\begin{array}{ccc|c} 2 & 1 & 3 & 3 \\ 0 & -\frac{3}{2} & -\frac{3}{2} & \frac{3}{2} \\ 0 & 0 & 0 & 0 \end{array} \right)$$

The system is consistent.

- Therefore, v is a linear combination of v_1, v_2, v_3 .

26 / 106

Examples

- Let $v_1 = (2, 1, 3)$, $v_2 = (1, -1, 2)$ and $v_3 = (3, 0, 5)$.
 - Is $v = (1, 2, 4)$ a linear combination of v_1, v_2, v_3 ?

Suppose that $v = av_1 + bv_2 + cv_3$, i.e.,

$$\begin{aligned}(1, 2, 4) &= a(2, 1, 3) + b(1, -1, 2) + c(3, 0, 5) \\ &= (2a + b + 3c, a - b, 3a + 2b + 5c).\end{aligned}$$

Solve the linear system
$$\begin{cases} 2a + b + 3c = 1 \\ a - b = 2 \\ 3a + 2b + 5c = 4. \end{cases}$$

$$\circ \left(\begin{array}{ccc|c} 2 & 1 & 3 & 1 \\ 1 & -1 & 0 & 2 \\ 3 & 2 & 5 & 4 \end{array} \right) \xrightarrow[\text{elimination}]{\text{Gaussian}} \left(\begin{array}{ccc|c} 2 & 1 & 3 & 1 \\ 0 & -\frac{3}{2} & -\frac{3}{2} & \frac{3}{2} \\ 0 & 0 & 0 & 3 \end{array} \right)$$

The system is inconsistent.

- Therefore, v is not a linear combination of v_1, v_2, v_3 .

27 / 106

Linear Span

- **Definition.** Let $S = \{v_1, v_2, \dots, v_k\}$ be a subset of \mathbb{R}^n .

- The **set of all linear combinations** of v_1, v_2, \dots, v_k
 - $\{c_1 v_1 + c_2 v_2 + \dots + c_k v_k \mid c_1, c_2, \dots, c_k \in \mathbb{R}\}$

is called the **linear span** of S (or v_1, v_2, \dots, v_k).

- It is denoted by $\text{span}(S)$ or $\text{span}\{v_1, v_2, \dots, v_k\}$.

v is a linear combination of v_1, v_2, \dots, v_k

$$\Leftrightarrow v \in \text{span}\{v_1, v_2, \dots, v_k\}.$$

- **Example.** Let $S = \{(2, 1, 3), (1, -1, 2), (3, 0, 5)\}$.

- $(3, 3, 4) \in \text{span}(S)$ but $(1, 2, 4) \notin \text{span}(S)$.

Example. Let $S = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$.

- $(x, y, z) = x(1, 0, 0) + y(0, 1, 0) + z(0, 0, 1)$.

Therefore, $\text{span}(S) = \mathbb{R}^3$.

28 / 106

Examples

- Let $S = \{(1, 0, 0, -1), (0, 1, 1, 0)\}$ be a subset of \mathbb{R}^4 .

- Every vector in $\text{span}(S)$ is of the form
 - $a(1, 0, 0, -1) + b(0, 1, 1, 0) = (a, b, b, -a)$,
where $a, b \in \mathbb{R}$.
- $\text{span}(S) = \{(a, b, b, -a) \mid a, b \in \mathbb{R}\}$.

- Let $V = \{(2a + b, a, 3b - a) \mid a, b \in \mathbb{R}\} \subseteq \mathbb{R}^3$.

- Every vector in V is of the form
 - $(2a + b, a, 3b - a) = a(2, 1, -1) + b(1, 0, 3)$,
where $a, b \in \mathbb{R}$.
- $V = \text{span}\{(2, 1, -1), (1, 0, 3)\}$.

29 / 106

Examples

- Prove that $\text{span}\{(1, 0, 1), (1, 1, 0), (0, 1, 1)\} = \mathbb{R}^3$.
 - It is clear: $\text{span}\{(1, 0, 1), (1, 1, 0), (0, 1, 1)\} \subseteq \mathbb{R}^3$.
Is $\mathbb{R}^3 \subseteq \text{span}\{(1, 0, 1), (1, 1, 0), (0, 1, 1)\}$?
- Let $(x, y, z) \in \mathbb{R}^3$. We shall find $a, b, c \in \mathbb{R}$ such that
 - $(x, y, z) = a(1, 0, 1) + b(1, 1, 0) + c(0, 1, 1)$.
 - Equivalently,
$$\begin{cases} a + b = x \\ b + c = y \\ a + c = z \end{cases}$$
 - $$\left(\begin{array}{ccc|c} 1 & 1 & 0 & x \\ 0 & 1 & 1 & y \\ 1 & 0 & 1 & z \end{array} \right) \xrightarrow[\text{Elimination}]{\text{Gaussian}} \left(\begin{array}{ccc|c} 1 & 1 & 0 & x \\ 0 & 1 & 1 & y \\ 0 & 0 & 2 & z - x + y \end{array} \right)$$

The system is always consistent for any $(x, y, z) \in \mathbb{R}^3$.
Therefore, $\text{span}\{(1, 0, 1), (1, 1, 0), (0, 1, 1)\} = \mathbb{R}^3$.

30 / 106

Examples

- Prove that $\text{span}\{(1, 1, 1), (1, 2, 0), (2, 1, 3), (2, 3, 1)\} \neq \mathbb{R}^3$.
 - Clear: $\text{span}\{(1, 1, 1), (1, 2, 0), (2, 1, 3), (2, 3, 1)\} \subseteq \mathbb{R}^3$.
Is $\mathbb{R}^3 \not\subseteq \text{span}\{(1, 1, 1), (1, 2, 0), (2, 1, 3), (2, 3, 1)\}$?
- Let $(x, y, z) \in \mathbb{R}^3$. Can we find $a, b, c, d \in \mathbb{R}$ such that
 - $(x, y, z) = a(1, 1, 1) + b(1, 2, 0) + c(2, 1, 3) + d(2, 3, 1)$?
 - Equivalently,
$$\begin{cases} a + b + 2c + 2d = x \\ a + 2b + c + 3d = y \\ a + 3c + d = z \end{cases}$$
 - $$\left(\begin{array}{cccc|c} 1 & 1 & 2 & 2 & x \\ 1 & 2 & 1 & 3 & y \\ 1 & 0 & 3 & 1 & z \end{array} \right) \xrightarrow{\text{G.E.}} \left(\begin{array}{cccc|c} 1 & 1 & 2 & 2 & x \\ 0 & 1 & -1 & 1 & y - x \\ 0 & 0 & 0 & 0 & y + z - 2x \end{array} \right)$$

The system is consistent $\Leftrightarrow y + z - 2x = 0$.
Therefore, $\text{span}\{(1, 1, 1), (1, 2, 0), (2, 1, 3), (2, 3, 1)\} \neq \mathbb{R}^3$.
◦ $\text{span}\{(1, 1, 1), (1, 2, 0), (2, 1, 3), (2, 3, 1)\} = \{(x, y, z) \mid y + z - 2x = 0\}$.

31 / 106

Criterion for $\text{span}(S) = \mathbb{R}^n$

- Let $S = \{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n$. Is $\text{span}(S) = \mathbb{R}^n$?
 - For arbitrary $v \in \mathbb{R}^n$, we shall check the consistency of
 - $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = v$.

View each v_j as a column vector $v_j = \begin{pmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{nj} \end{pmatrix}$.

$$\begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} = v = c_1 \begin{pmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{pmatrix} + c_2 \begin{pmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{n2} \end{pmatrix} + \dots + c_k \begin{pmatrix} a_{1k} \\ a_{2k} \\ \vdots \\ a_{nk} \end{pmatrix}$$

$$= \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1k} \\ a_{21} & a_{22} & \dots & a_{2k} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nk} \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_k \end{pmatrix}.$$

32 / 106

Criterion for $\text{span}(S) = \mathbb{R}^n$

- Let $S = \{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n$. Is $\text{span}(S) = \mathbb{R}^n$?
 - For arbitrary $v \in \mathbb{R}^n$, we shall check the consistency of
 - $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = v$.

View v_j as column vectors. Let $A = (v_1 \ v_2 \ \dots \ v_k)$.

- The system can be written as $Ax = v$.

Let R be a row-echelon form of A .

- $(A \mid v) \xrightarrow[\text{Elimination}]{\text{Gaussian}} (R \mid v')$.

- Since $v \in \mathbb{R}^n$ is arbitrary, $v' \in \mathbb{R}^n$ is also arbitrary.

$$\begin{aligned} \text{span}(S) = \mathbb{R}^n &\Leftrightarrow Ax = v \text{ consistent for every } v \in \mathbb{R}^n \\ &\Leftrightarrow \text{last column of } (R \mid v') \text{ non-pivot for any } v' \in \mathbb{R}^n \\ &\Leftrightarrow R \text{ has no zero row.} \end{aligned}$$

33 / 106

Criterion for $\text{span}(S) = \mathbb{R}^n$

- Let $S = \{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n$.
 1. View each v_j as a column vector.
 2. Let $A = (v_1 \ v_2 \ \cdots \ v_k)$.
 3. Find a row-echelon form R of A .
 - If R has a zero row, then $\text{span}(S) \neq \mathbb{R}^n$.
 - If R has no zero row, then $\text{span}(S) = \mathbb{R}^n$.

- **Example.**

- $\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix} \xrightarrow[\text{Elimination}]{\text{Gaussian}} \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \end{pmatrix}.$

$\therefore \text{span}\{(1, 0, 1), (1, 1, 0), (0, 1, 1)\} = \mathbb{R}^3.$

34 / 106

Criterion for $\text{span}(S) = \mathbb{R}^n$

- Let $S = \{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n$.
 1. View each v_j as a column vector.
 2. Let $A = (v_1 \ v_2 \ \cdots \ v_k)$.
 3. Find a row-echelon form R of A .
 - If R has a zero row, then $\text{span}(S) \neq \mathbb{R}^n$.
 - If R has no zero row, then $\text{span}(S) = \mathbb{R}^n$.

- **Example.**

- $\begin{pmatrix} 1 & 1 & 2 & 2 \\ 1 & 2 & 1 & 3 \\ 1 & 0 & 3 & 1 \end{pmatrix} \xrightarrow[\text{Elimination}]{\text{Gaussian}} \begin{pmatrix} 1 & 1 & 2 & 2 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$

$\therefore \text{span}\{(1, 1, 1), (1, 2, 0), (2, 1, 3), (2, 3, 1)\} \neq \mathbb{R}^3.$

35 / 106

Criterion for $\text{span}(S) = \mathbb{R}^n$

- **Theorem.** Let $S = \{v_1, v_2, \dots, v_k\}$ be a subset of \mathbb{R}^n .

- If $k < n$, then $\text{span}(S) \neq \mathbb{R}^n$.

Proof. View each v_j as a column vector. Then

- $A = (v_1 \ v_2 \ \cdots \ v_k)$ is an $n \times k$ matrix.

Let R be a row-echelon form of A . Then R is $n \times k$.

1. R has at most k pivot columns.
2. R has at most k nonzero rows.
3. R has at least $n - k > 0$ zero rows.

Therefore, $\text{span}(S) \neq \mathbb{R}^n$.

- **Examples.**

- One vector cannot span \mathbb{R}^2 .
- Two vectors cannot span \mathbb{R}^3 .

36 / 106

Properties of Linear Spans

- Let $S = \{u_1, u_2, \dots, u_k\} \subseteq \mathbb{R}^n$.

- Then $0 = 0u_1 + 0u_2 + \cdots + 0u_k \in \text{span}(S)$.

Suppose $v_1, v_2, \dots, v_r \in \text{span}(S)$.

- Each v_i is a linear combination of u_1, u_2, \dots, u_k .

- $v_1 = a_{11}u_1 + a_{12}u_2 + \cdots + a_{1k}u_k$.
- $v_2 = a_{21}u_1 + a_{22}u_2 + \cdots + a_{2k}u_k$.
- \vdots
- $v_r = a_{r1}u_1 + a_{r2}u_2 + \cdots + a_{rk}u_k$.

Then for any $c_1, c_2, \dots, c_r \in \mathbb{R}$,

$$\begin{aligned} c_1v_1 + \cdots + c_rv_r &= c_1(a_{11}u_1 + \cdots + a_{1k}u_k) \\ &\quad + \cdots + c_r(a_{r1}u_1 + \cdots + a_{rk}u_k) \\ &= (c_1a_{11} + \cdots + c_ra_{r1})u_1 \\ &\quad + \cdots + (c_1a_{1k} + \cdots + c_ra_{rk})u_k. \end{aligned}$$

37 / 106

Properties of Linear Spans

- **Theorem.** Let $S = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$ be a subset of \mathbb{R}^n .
 - $\mathbf{0} \in \text{span}(S)$, where $\mathbf{0}$ is the zero vector in \mathbb{R}^n .
 - Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r \in \text{span}(S)$, $c_1, c_2, \dots, c_r \in \mathbb{R}$.
 - $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_r\mathbf{v}_r \in \text{span}(S)$.
- **Remarks.** In particular,
 - Since $\mathbf{0} \in \text{span}(S)$, $\text{span}(S) \neq \emptyset$.
 - $\mathbf{v} \in \text{span}(S)$ and $c \in \mathbb{R} \Rightarrow c\mathbf{v} \in \text{span}(S)$.
 - $\text{span}(S)$ is **closed** under scalar multiplication.
 - $\mathbf{u} \in \text{span}(S)$ and $\mathbf{v} \in \text{span}(S) \Rightarrow \mathbf{u} + \mathbf{v} \in \text{span}(S)$.
 - $\text{span}(S)$ is **closed** under addition.

38 / 106

Properties of Linear Spans

- **Theorem.** Given two subsets of \mathbb{R}^n :
 - $S_1 = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$, $S_2 = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$.Then $\text{span}(S_1) \subseteq \text{span}(S_2)$
 - \Leftrightarrow Every \mathbf{u}_i is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$.**Proof.** \Rightarrow : Suppose that $\text{span}(S_1) \subseteq \text{span}(S_2)$.
 - $\mathbf{u}_i = 0\mathbf{u}_1 + \dots + 1\mathbf{u}_i + \dots + 0\mathbf{u}_k \in \text{span}(S_1)$.
 - Then $\mathbf{u}_i \in \text{span}(S_2)$ by assumption.That is, \mathbf{u}_i is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$.
 - \Leftarrow : Suppose each \mathbf{u}_i is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$.
 - Let $\mathbf{w} \in \text{span}(S_1)$. There exist $c_1, c_2, \dots, c_k \in \mathbb{R}$ s.t.
 - $\mathbf{w} = c_1\mathbf{u}_1 + c_2\mathbf{u}_2 + \dots + c_k\mathbf{u}_k \in \text{span}(S_2)$.Therefore, $\text{span}(S_1) \subseteq \text{span}(S_2)$.

39 / 106

Properties of Linear Spans

- **Theorem.** Let $v_1, v_2, \dots, v_{k-1}, v_k \in \mathbb{R}^n$.
 - If v_k is a linear combination of v_1, v_2, \dots, v_{k-1} , then
 - $\text{span}\{v_1, \dots, v_{k-1}\} = \text{span}\{v_1, \dots, v_{k-1}, v_k\}$.

Proof. It follows from the definition of linear span that

$$\text{span}\{v_1, \dots, v_{k-1}\} \subseteq \text{span}\{v_1, \dots, v_{k-1}, v_k\}.$$

Since v_k is a linear combination of v_1, \dots, v_{k-1} ,

$$\text{span}\{v_1, \dots, v_{k-1}, v_k\} \subseteq \text{span}\{v_1, \dots, v_{k-1}\}.$$

$$\therefore \text{span}\{v_1, \dots, v_{k-1}\} = \text{span}\{v_1, \dots, v_{k-1}, v_k\}.$$

- **Example.** Let $v_1 = (1, 1, 0, 2)$, $v_2 = (1, 0, 0, 1)$.
 - Let $v_3 = v_1 - v_2 = (0, 1, 0, 1)$.
 Then $\text{span}\{v_1, v_2\} = \text{span}\{v_1, v_2, v_3\}$.

40 / 106

Properties of Linear Spans

- Let $S = \{v_1, v_2, \dots, v_k\}$ be a subset of \mathbb{R}^n .

$$v \in \text{span}(S) \Leftrightarrow v = c_1 v_1 + \dots + c_k v_k \text{ for some } c_i \in \mathbb{R}$$

$$\Leftrightarrow (v_1 \quad \dots \quad v_k) \begin{pmatrix} c_1 \\ \vdots \\ c_k \end{pmatrix} = v.$$

1. View each v_j as a column vector.
2. Let $A = (v_1 \quad \dots \quad v_k)$.
3. Check if the linear system $Ax = v$ is consistent.
 - If $Ax = v$ is consistent, then $v \in \text{span}(S)$.
 - If $Ax = v$ is inconsistent, then $v \notin \text{span}(S)$.

41 / 106

Examples

- Let $\mathbf{u}_1 = (1, 0, 1)$, $\mathbf{u}_2 = (1, 1, 2)$, $\mathbf{u}_3 = (-1, 2, 1)$, and
 - $\mathbf{v}_1 = (1, 2, 3)$, $\mathbf{v}_2 = (2, -1, 1)$.

Prove that $\text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\} = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$.

Solution. Step 1: $\text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\} \subseteq \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$.

- Show that $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3 \in \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$.

$$\left(\begin{array}{cc|c|c|c} 1 & 2 & 1 & 1 & -1 \\ 2 & -1 & 0 & 1 & 2 \\ 3 & 1 & 1 & 2 & 1 \end{array} \right) \xrightarrow{\text{G.E.}} \left(\begin{array}{cc|c|c|c} 1 & 2 & 1 & 1 & -1 \\ 0 & -5 & -2 & -1 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

- The systems $(\mathbf{v}_1 \quad \mathbf{v}_2) \mathbf{x} = \mathbf{u}_j$ are all consistent.

- Then $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3 \in \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$.

Therefore, $\text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\} \subseteq \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$.

42 / 106

Examples

- Let $\mathbf{u}_1 = (1, 0, 1)$, $\mathbf{u}_2 = (1, 1, 2)$, $\mathbf{u}_3 = (-1, 2, 1)$, and
 - $\mathbf{v}_1 = (1, 2, 3)$, $\mathbf{v}_2 = (2, -1, 1)$.

Prove that $\text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\} = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$.

Solution. Step 2: $\text{span}\{\mathbf{v}_1, \mathbf{v}_2\} \subseteq \text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$.

- Show that $\mathbf{v}_1, \mathbf{v}_2 \in \text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$.

$$\left(\begin{array}{ccc|c|c} 1 & 1 & -1 & 1 & 2 \\ 0 & 1 & 2 & 2 & -1 \\ 1 & 2 & 1 & 3 & 1 \end{array} \right) \xrightarrow{\text{G.E.}} \left(\begin{array}{ccc|c|c} 1 & 1 & -1 & 1 & 2 \\ 0 & 1 & 2 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

- The systems $(\mathbf{u}_1 \quad \mathbf{u}_2 \quad \mathbf{u}_3) \mathbf{x} = \mathbf{v}_j$ are all consistent.

- Then $\mathbf{v}_1, \mathbf{v}_2 \in \text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$.

Therefore, $\text{span}\{\mathbf{v}_1, \mathbf{v}_2\} \subseteq \text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$.

We can conclude that $\text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\} = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$.

43 / 106

Examples

- Let $\mathbf{u}_1 = (1, 0, 0, 1)$, $\mathbf{u}_2 = (0, 1, -1, 2)$, $\mathbf{u}_3 = (2, 1, -1, 4)$.
 $\mathbf{v}_1 = (1, 1, 1, 1)$, $\mathbf{v}_2 = (-1, 1, -1, 1)$, $\mathbf{v}_3 = (-1, 1, 1, -1)$.

- $(\mathbf{v}_1 \quad \mathbf{v}_2 \quad \mathbf{v}_3 \mid \mathbf{u}_1 \mid \mathbf{u}_2 \mid \mathbf{u}_3)$

$$\xrightarrow{\text{G.E.}} \left(\begin{array}{ccc|c|c|c} 1 & -1 & -1 & 1 & 0 & 2 \\ 0 & 2 & 2 & -1 & 1 & -1 \\ 0 & 0 & 2 & -1 & -1 & -3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

- The systems $(\mathbf{v}_1 \quad \mathbf{v}_2 \quad \mathbf{v}_3) \mathbf{x} = \mathbf{u}_j$ are consistent.

- Then $\text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\} \subseteq \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

- $(\mathbf{u}_1 \quad \mathbf{u}_2 \quad \mathbf{u}_3 \mid \mathbf{v}_1 \mid \mathbf{v}_2 \mid \mathbf{v}_3)$

$$\xrightarrow{\text{G.E.}} \left(\begin{array}{ccc|c|c|c} 1 & 0 & 2 & 1 & -1 & -1 \\ 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 2 & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

- $(\mathbf{u}_1 \quad \mathbf{u}_2 \quad \mathbf{u}_3) \mathbf{x} = \mathbf{v}_j$ are inconsistent for $j = 1, 3$.

- Then $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} \not\subseteq \text{span}\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$.

44 / 106

Subspaces

45 / 106

Subspaces

- Definition.** Let V be a subset of \mathbb{R}^n . Then V is called a **subspace** of \mathbb{R}^n if there exist $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$ s.t.

- $V = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$.

More precisely,

- V is the **subspace spanned** by $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$;
- $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ **spans** the subspace V .

- Remark.**

- Let $\mathbf{0} \in \mathbb{R}^n$ be the zero vector. Then

- $\{\mathbf{0}\} = \text{span}\{\mathbf{0}\}$ is the **zero space**.

- Let \mathbf{e}_i denote the n -vector whose i th coordinate is 1 and elsewhere 0, e.g., $\mathbf{e}_2 = (0, 1, 0, \dots, 0)$.

- Then for every $\mathbf{v} = (v_1, v_2, \dots, v_n) \in \mathbb{R}^n$

- $\mathbf{v} = v_1 \mathbf{e}_1 + v_2 \mathbf{e}_2 + \dots + v_n \mathbf{e}_n$.

$\mathbb{R}^n = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is a subspace of \mathbb{R}^n .

46 / 106

Examples

- In order to show that a subset V of \mathbb{R}^n is a subspace:
 - Find $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$.
 - Show that every $\mathbf{v} \in V$ is of the form
 - $\mathbf{v} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k, c_1, c_2, \dots, c_k \in \mathbb{R}$.
- Let $V_1 = \{(a + 4b, a) \mid a, b \in \mathbb{R}\}$.
 - $(a + 4b, a) = a(1, 1) + b(4, 0)$ for all $a, b \in \mathbb{R}$.Then $V_1 = \text{span}\{(1, 1), (4, 0)\}$ is a subspace of \mathbb{R}^2 .
- Let $V_2 = \{(x, y, z) \mid x + y - z = 0\}$.
 - $x + y - z = 0$ can be explicitly solved:
 - $(x, y, z) = (-s + t, s, t)$, where $s, t \in \mathbb{R}$.
 - $(-s + t, s, t) = s(-1, 1, 0) + t(1, 0, 1)$. $V_2 = \text{span}\{(-1, 1, 0), (1, 0, 1)\}$ is a subspace of \mathbb{R}^3 .

47 / 106

Examples

- Recall that a subspace V is of the form $\text{span}(S)$. Then
 - $\mathbf{0} \in V$,
 - $c \in \mathbb{R} \ \& \ \mathbf{v} \in V \Rightarrow c\mathbf{v} \in V$,
 - $\mathbf{u} \in V \ \& \ \mathbf{v} \in V \Rightarrow \mathbf{u} + \mathbf{v} \in V$.If any of the above fails, then V is not a subspace (of \mathbb{R}^n).
- Let $V_3 = \{(1, a) \mid a \in \mathbb{R}\}$.
 - $\mathbf{0} = (0, 0) \notin V_3$. So V is not a subspace of \mathbb{R}^2 .
- Let $V_4 = \{(x, y, z) \mid x^2 \leq y^2 \leq z^2\}$.
 - $(1, 2, 3) \in V_4$ because $1^2 \leq 2^2 \leq 3^2$.
 - $(1, 2, -3) \in V_4$ because $1^2 \leq 2^2 \leq (-3)^2$.
 - $(1, 2, 3) + (1, 2, -3) = (2, 4, 0) \notin V_4$.Therefore, V_4 is not a subspace of \mathbb{R}^3 .

48 / 106

Subspaces of $\mathbb{R}^1, \mathbb{R}^2, \mathbb{R}^3$

- Let \mathbf{v} be a nonzero vector in \mathbb{R}^n , $n = 1, 2, 3$.

- $V = \text{span}\{\mathbf{v}\} = \{c\mathbf{v} \mid c \in \mathbb{R}\}$.

This is a line through the origin.

1. $n = 1$: $\mathbf{v} = v \in \mathbb{R}$, and V is the whole $\mathbb{R}^1 = \mathbb{R}$.

2. $n = 2$: $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$.

- $V = \{(x, y) \mid v_2x - v_1y = 0\}$.

3. $n = 3$: $\mathbf{v} = (v_1, v_2, v_3) \in \mathbb{R}^3$.

- $V = \{(cv_1, cv_2, cv_3) \mid c \in \mathbb{R}\}$.

- If $v_1 \neq 0$, then V is the intersection of planes

- $v_2x - v_1y = 0$ and $v_3x - v_1z = 0$.

- $V = \{(x, y, z) \mid v_2x - v_1y = 0 \text{ \& } v_3x - v_1z = 0\}$

49 / 106

Subspaces of $\mathbb{R}^1, \mathbb{R}^2, \mathbb{R}^3$

- Let \mathbf{u}, \mathbf{v} be nonzero vectors in \mathbb{R}^n , $n = 2, 3$.

- $V = \text{span}\{\mathbf{u}, \mathbf{v}\} = \{s\mathbf{u} + t\mathbf{v} \mid s, t \in \mathbb{R}\}$.

If \mathbf{u} and \mathbf{v} are parallel, then $V = \text{span}\{\mathbf{u}\} = \text{span}\{\mathbf{v}\}$.

Suppose \mathbf{u} and \mathbf{v} are not parallel. Then

- $V = \text{span}\{\mathbf{u}, \mathbf{v}\}$ is a plane containing the origin.

1. $n = 2$: Then V is the whole \mathbb{R}^2 .

2. $n = 3$: Let $\mathbf{u} = (u_1, u_2, u_3)$ and $\mathbf{v} = (v_1, v_2, v_3)$.

- $V = \{s(u_1, u_2, u_3) + t(v_1, v_2, v_3) \mid s, t \in \mathbb{R}\}$.

We can find an implicit form of V :

- $V = \{(x, y, z) \mid ax + by + cz = 0\}$,

where a, b, c are not all zero.

50 / 106

Subspaces of $\mathbb{R}^1, \mathbb{R}^2, \mathbb{R}^3$

- Subspaces of \mathbb{R}^1 :
 - $\{0\}$,
 - \mathbb{R} .
- Subspaces of \mathbb{R}^2 :
 - $\{0\} = \{(0, 0)\}$,
 - A straight line passing through the origin $(0, 0)$,
 - \mathbb{R}^2 .
- Subspaces of \mathbb{R}^3 :
 - $\{0\} = \{(0, 0, 0)\}$,
 - A straight line passing through the origin $(0, 0, 0)$,
 - A plane containing the origin $(0, 0, 0)$,
 - \mathbb{R}^3 .

A subspace of $\mathbb{R}^i, i = 1, 2, 3$, is always the solution set of a homogeneous linear system.

51 / 106

Solution Space

- **Theorem.** The **solution set** of a **homogeneous** linear system of n variables is a **subspace** of \mathbb{R}^n .

Proof. Recall that a homogeneous system is consistent.

- If the system has only the trivial solution,
 - then the solution set $\{0\}$ is a subspace of \mathbb{R}^n .
- Suppose that the system has infinitely many solutions.
 - Use Gauss-Jordan elimination to find RREF. By setting the variables corresponding to non-pivot columns as arbitrary parameters t_1, \dots, t_k , solve the variables corresponding to pivot columns.
 - $x_1 = r_{11}t_1 + r_{12}t_2 + \dots + r_{1k}t_k$.
 - $x_2 = r_{21}t_1 + r_{22}t_2 + \dots + r_{2k}t_k$.
 -
 - $x_n = r_{n1}t_1 + r_{n2}t_2 + \dots + r_{nk}t_k$.

52 / 106

Solution Space

- **Theorem.** The **solution set** of a **homogeneous** linear system of n variables is a **subspace** of \mathbb{R}^n .

Proof. Recall that a homogeneous system is consistent.

- If the system has only the trivial solution,
 - then the solution set $\{\mathbf{0}\}$ is a subspace of \mathbb{R}^n .
- Suppose that the system has infinitely many solutions.

$$\bullet \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = t_1 \begin{pmatrix} r_{11} \\ r_{21} \\ \vdots \\ r_{n1} \end{pmatrix} + t_2 \begin{pmatrix} r_{12} \\ r_{22} \\ \vdots \\ r_{n2} \end{pmatrix} + \cdots + t_k \begin{pmatrix} r_{1k} \\ r_{2k} \\ \vdots \\ r_{nk} \end{pmatrix}.$$

The solution set is spanned by

$$\bullet (r_{11}, r_{21}, \dots, r_{n1}), \dots, (r_{1k}, r_{2k}, \dots, r_{nk}).$$

So the solution set is a subspace of \mathbb{R}^n .

53 / 106

Examples

- The **solution set** of a **homogeneous** linear system is called the **solution space** of the system.

We will see later that a subspace of \mathbb{R}^n is always the solution space of a homogeneous linear system.

- $$\begin{cases} x - 2y + 3z = 0 \\ 2x - 4y + 6z = 0 \\ 3x - 6y + 9z = 0 \end{cases}$$
 - $$\left(\begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ 2 & -4 & 6 & 0 \\ 3 & -6 & 9 & 0 \end{array} \right) \xrightarrow[R_3+(-3)R_1]{R_2+(-2)R_1} \left(\begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$
 - $(x, y, z) = (2s - 3t, s, t) = s(2, 1, 0) + t(-3, 0, 1).$

The solution space is $\text{span}\{(2, 1, 0), (-3, 0, 1)\}$.

54 / 106

Examples

- $$\begin{cases} x - 2y + 3z = 0 \\ -3x + 7y - 8z = 0 \\ -2x + 4y - 6z = 0 \end{cases}$$
 - $\left(\begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ -3 & 7 & -8 & 0 \\ -2 & 4 & -6 & 0 \end{array} \right) \xrightarrow{\text{G-J.E.}} \left(\begin{array}{ccc|c} 1 & 0 & 5 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$
 - $(x, y, z) = (-5t, -t, t) = t(-5, -1, 1)$.

The solution space is $\text{span}\{(-5, -1, 1)\}$.

- $$\begin{cases} x - 2y + 3z = 0 \\ -3x + 7y - 8z = 0 \\ 4x + y + 2z = 0 \end{cases}$$
 - $\left(\begin{array}{ccc|c} 1 & -2 & 3 & 0 \\ -3 & 7 & -8 & 0 \\ 4 & 1 & 2 & 0 \end{array} \right) \xrightarrow{\text{G-J.E.}} \left(\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right)$
 - $(x, y, z) = (0, 0, 0)$. The solution space is $\{(0, 0, 0)\}$.

55 / 106

Linear Independence

56 / 106

Linear Independence

- In \mathbb{R}^3 , a plane containing the origin can be spanned by two non-parallel vectors: $V = \text{span}\{\mathbf{u}, \mathbf{v}\}$.
 - If a plane is spanned by more than two vectors, then
 - some vectors in the spanning set is redundant.
- Suppose that $V = \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$.
 - Recall that if \mathbf{v}_k is a linear combination of $\mathbf{v}_1, \dots, \mathbf{v}_{k-1}$,
 - $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_{k-1}, \mathbf{v}_k\} = \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_{k-1}\}$.

Continuing this procedure, we can remove the redundant vectors in the spanning set to obtain

- $V = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$,

so that any \mathbf{v}_i is NOT a linear combination of the other vectors.

57 / 106

Linear Independence

• **Definition.** Let $S = \{v_1, \dots, v_k\}$ be a subset of \mathbb{R}^n .

◦ The equation $c_1v_1 + c_2v_2 + \dots + c_kv_k = \mathbf{0}$

has a **trivial solution** $c_1 = c_2 = \dots = c_k = 0$.

1. If the equation has a **non-trivial solution**, then

- S is a **linearly dependent set**,
- v_1, v_2, \dots, v_k are **linearly dependent**.

There exist $c_1, c_2, \dots, c_k \in \mathbb{R}$ not all zero such that

- $c_1v_1 + c_2v_2 + \dots + c_kv_k = \mathbf{0}$.

2. If the equation has **only the trivial solution**, then

- S is a **linearly independent set**,
- v_1, v_2, \dots, v_k are **linearly independent**.

$$c_1v_1 + c_2v_2 + \dots + c_kv_k = \mathbf{0} \Rightarrow c_1 = \dots = c_k = 0.$$

58 / 106

Examples

• Let $S = \{(1, -2, 3), (5, 6, -1), (3, 2, 1)\}$.

◦ $c_1(1, -2, 3) + c_2(5, 6, -1) + c_3(3, 2, 1) = (0, 0, 0)$.

- $c_1 \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} + c_2 \begin{pmatrix} 5 \\ 6 \\ -1 \end{pmatrix} + c_3 \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$

- $\begin{pmatrix} 1 & 5 & 3 \\ -2 & 6 & 2 \\ 3 & -1 & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$

- $\begin{pmatrix} 1 & 5 & 3 \\ -2 & 6 & 2 \\ 3 & -1 & 1 \end{pmatrix} \xrightarrow{\text{G.E.}} \begin{pmatrix} 1 & 5 & 3 \\ 0 & 16 & 8 \\ 0 & 0 & 0 \end{pmatrix}.$

The 3rd column is non-pivot.

- The system has infinitely many solutions.

Therefore, S is a linearly dependent set.

59 / 106

Examples

- Let $S = \{(1, 0, 0, 1), (0, 2, 1, 0), (1, -1, 1, 1)\}$.
 - $c_1(1, 0, 0, 1) + c_2(0, 2, 1, 0) + c_3(1, -1, 1, 1) = \mathbf{0}$.

$$\bullet \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & -1 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

$$\bullet \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & -1 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{pmatrix} \xrightarrow[\text{elimination}]{\text{Gaussian}} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & -1 \\ 0 & 0 & \frac{3}{2} \\ 0 & 0 & 0 \end{pmatrix}.$$

All the columns are pivot.

- The system has only the trivial solution.

Therefore, S is a linearly independent set.

60 / 106

Properties

- Let S_1, S_2 be finite subsets of \mathbb{R}^n such that $S_1 \subseteq S_2$.
 - S_1 linearly dependent $\Rightarrow S_2$ linearly dependent.
 - S_2 linearly independent $\Rightarrow S_1$ linearly independent.
- $c\mathbf{0} = \mathbf{0}$ has infinitely many solutions $c \in \mathbb{R}$.
 - $\{\mathbf{0}\}$ is linearly dependent.
 - If $\mathbf{0} \in S (\subseteq \mathbb{R}^n)$ then S is linearly dependent.
- Let $\mathbf{v} \in \mathbb{R}^n$. Then $c\mathbf{v} = \mathbf{0} \Leftrightarrow c = 0$ or $\mathbf{v} = \mathbf{0}$.
 - $\{\mathbf{v}\}$ is linearly independent $\Leftrightarrow \mathbf{v} \neq \mathbf{0}$.
- Let $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$. Then

$$\{\mathbf{u}, \mathbf{v}\} \text{ is linearly dependent } \Leftrightarrow \mathbf{u} = a\mathbf{v} \text{ for some } a \in \mathbb{R} \\ \text{or } \mathbf{v} = a\mathbf{u} \text{ for some } a \in \mathbb{R}$$

61 / 106

Properties

- **Theorem.** Let $S = \{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n, k \geq 2$.
 - S is linearly dependent
 - \Leftrightarrow there exists v_i such that it is a linear combination of other vectors $v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k$.
- **Proof.** \Rightarrow : Suppose S is linearly dependent.
 - There exist $c_1, c_2, \dots, c_k \in \mathbb{R}$ not all zero such that
 - $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = \mathbf{0}$.
 - Suppose that $c_i \neq 0$. Then
 - $v_i = -\frac{c_1}{c_i} v_1 - \dots - \frac{c_{i-1}}{c_i} v_{i-1} - \frac{c_{i+1}}{c_i} v_{i+1} - \dots - \frac{c_k}{c_i} v_k$.
 - \Leftarrow : Suppose v_i is a linear combination of other vectors. Then there exist $c_1, \dots, c_{i-1}, c_{i+1}, \dots, c_k \in \mathbb{R}$ such that
 - $v_i = c_1 v_1 + \dots + c_{i-1} v_{i-1} + c_{i+1} v_{i+1} + \dots + c_k v_k$.
$$c_1 v_1 + \dots + c_{i-1} v_{i-1} + (-1)v_i + c_{i+1} v_{i+1} + \dots + c_k v_k = \mathbf{0}$$

$\therefore S = \{v_1, \dots, v_{i-1}, v_i, v_{i+1}, \dots, v_k\}$ is linearly dept.

62 / 106

Properties

- **Theorem.** Let $S = \{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n, k \geq 2$.
 - S is linearly dependent
 - \Leftrightarrow there exists v_i such that it is a linear combination of other vectors $v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k$.
 - S is linearly independent
 - \Leftrightarrow no vector in S can be written as a linear combination of other vectors.
- **Remarks.** Suppose $S = \{v_1, v_2, \dots, v_k\}$ is linearly dependent. Let $V = \text{span}(S)$.
 - Some $v_i \in S$ is a linear combination of other vectors.
 - Remove v_i from S and repeat the procedure until we obtain a linearly independent set S' .
 - Then $\text{span}(S') = V$ and S' has no "redundant vector" to span V .

63 / 106

Examples

- Let $S_1 = \{(1, 0), (0, 4), (2, 4)\}$.
 - Note that $(2, 4) = 2(1, 0) + 1(0, 4)$.
Then S_1 is linearly dependent. Moreover,
 - $\text{span}(S_1) = \text{span}\{(1, 0), (0, 4)\}$.
- Let $S_2 = \{(-1, 0, 0), (0, 3, 0), (0, 0, 7)\}$.
 - $(-1, 0, 0)$ is the only vector whose 1st component $\neq 0$
 - $(0, 3, 0)$ is the only vector whose 2nd component $\neq 0$.
 - $(0, 0, 7)$ is the only vector whose 3rd component $\neq 0$.Any vector is NOT a linear combination of the other two vectors.
 $\therefore S_2$ is linearly independent.

64 / 106

Properties

- **Theorem.** Let $S = \{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n$.
 - If $k > n$, then S is linearly dependent.
- **Proof.** Consider $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = \mathbf{0}$.
 - View each v_j as a column vector.
 - Let $A = (v_1 \ v_2 \ \dots \ v_k)$.
 - Determine if $Ax = \mathbf{0}$ has only the trivial solution.
 - A is of size $n \times k$; so is the RREF R .
 - R has at most n nonzero rows.
 - R has at most n pivot columns.
 - R has at least $k - n > 0$ non-pivot columns.
 - Then $Ax = \mathbf{0}$ has non-trivial solutions.
- $\therefore S$ is linearly dependent.

65 / 106

Properties

- **Theorem.** Suppose $\{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n$ is linearly independent.

- If $v_{k+1} \in \mathbb{R}^n$ is not in $\text{span}\{v_1, v_2, \dots, v_k\}$.
 - then $\{v_1, v_2, \dots, v_k, v_{k+1}\}$ is linearly independent.

Proof. Suppose $c_1 v_1 + \dots + c_k v_k + c_{k+1} v_{k+1} = \mathbf{0}$.

- If $c_{k+1} \neq 0$, then
 - $v_{k+1} = -\frac{c_1}{c_{k+1}} v_1 - \dots - \frac{c_k}{c_{k+1}} v_k$,
 - $v_{k+1} \in \text{span}\{v_1, \dots, v_k\}$, contradiction!
- So $c_{k+1} = 0$. This implies
 - $c_1 v_1 + \dots + c_k v_k = \mathbf{0}$.
 - $\{v_1, \dots, v_k\}$ is linearly independent.
 $\Rightarrow c_1 = c_2 = \dots = c_k = 0$.

Therefore, $\{v_1, \dots, v_k, v_{k+1}\}$ is linearly independent.

66 / 106

Bases

67 / 106

Motivation

- Let $\{v_1, \dots, v_k\} \subseteq \mathbb{R}^n$ be linearly independent.
 1. Suppose $\text{span}\{v_1, \dots, v_k\} \neq \mathbb{R}^n$.
 2. Pick $v_{k+1} \in \mathbb{R}^n$ but $v_{k+1} \notin \text{span}\{v_1, \dots, v_k\}$.
 3. Then $\{v_1, \dots, v_k, v_{k+1}\}$ is linearly independent.
 4. Repeat this procedure until
 - $\{v_1, \dots, v_k, v_{k+1}, \dots, v_m\}$ is linearly independent &
 - $\text{span}\{v_1, \dots, v_k, v_{k+1}, \dots, v_m\} = \mathbb{R}^n$.
- If $m > n$, then $\{v_1, \dots, v_m\}$ is linearly dependent.
If $m < n$, then $\{v_1, \dots, v_m\}$ cannot span \mathbb{R}^n .
 - We must have $n = m$. $\{v_1, \dots, v_n\}$ is linearly independent, and spans \mathbb{R}^n .

68 / 106

Vector Spaces

- **Definition.** A set V is called a **vector space** if
 - V is a subspace of \mathbb{R}^n for some positive integer n .If W and V are vector spaces such that $W \subseteq V$,
 - then W is a **subspace** of V .
- **Examples.**
 - Let $U = \text{span}\{(1, 1, 1)\}$, $V = \text{span}\{(1, 1, -1)\}$ and $W = \text{span}\{(1, 0, 0), (0, 1, 1)\}$.
Then U, V, W are vector spaces (subspace of \mathbb{R}^3).
 - $(1, 1, 1) = (1, 0, 0) + (0, 1, 1) \in W$.
 - Then $U \subseteq W$; so U is a subspace of W .
 - $(1, 1, -1) \notin \text{span}\{(1, 0, 0), (0, 1, 1)\}$.
 - Then $V \not\subseteq W$; so V is NOT a subspace of W .

69 / 106

Bases

- **Definition.** Let $S = \{v_1, \dots, v_k\}$ be a subset of a vector space V . Then S is called a **basis** (plural **bases**) for V if
 - S is **linearly independent**, and $\text{span}(S) = V$.
- **Example.** Show that $S = \{(1, 2, 1), (2, 9, 0), (3, 3, 4)\}$ is a basis for \mathbb{R}^3 .
 1. Prove that S is linearly independent.
 - Let $c_1(1, 2, 1) + c_2(2, 9, 0) + c_3(3, 3, 4) = \mathbf{0}$.
 - $$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 9 & 3 \\ 1 & 0 & 4 \end{pmatrix} \xrightarrow[\text{Elimination}]{\text{Gaussian}} \begin{pmatrix} 1 & 2 & 3 \\ 0 & 5 & -3 \\ 0 & 0 & -\frac{1}{5} \end{pmatrix}.$$
 - All the three columns are pivot.
 - The system has only the trivial solution.Therefore, S is linearly independent.

70 / 106

Bases

- **Definition.** Let $S = \{v_1, \dots, v_k\}$ be a subset of a vector space V . Then S is called a **basis** (plural **bases**) for V if

- S is **linearly independent**, and $\text{span}(S) = V$.

- **Example.** Show that $S = \{(1, 2, 1), (2, 9, 0), (3, 3, 4)\}$ is a basis for \mathbb{R}^3 .

2. Prove that $\text{span}(S) = \mathbb{R}^3$.

- $$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 9 & 3 \\ 1 & 0 & 4 \end{pmatrix} \xrightarrow[\text{Elimination}]{\text{Gaussian}} \begin{pmatrix} 1 & 2 & 3 \\ 0 & 5 & -3 \\ 0 & 0 & -\frac{1}{5} \end{pmatrix}.$$

- A row-echelon form has no zero row.

Therefore, $\text{span}(S) = \mathbb{R}^3$.

We can conclude that S is a basis for \mathbb{R}^3 .

71 / 106

Examples

- Let $V = \text{span}\{(1, 1, 1, 1), (1, -1, -1, 1), (1, 0, 0, 1)\}$.
- $S = \{(1, 1, 1, 1), (1, -1, -1, 1)\}$. Is S a basis for V ?

1.
$$\begin{pmatrix} 1 & 1 \\ 1 & -1 \\ 1 & -1 \\ 1 & 1 \end{pmatrix} \xrightarrow{\text{Gaussian elimination}} \begin{pmatrix} 1 & 1 \\ 0 & -2 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

- All the two columns are pivot.
- $c_1(1, 1, 1, 1) + c_2(1, -1, -1, 1) = \mathbf{0}$

has only the trivial solution.

So S is linearly independent.

2. $(1, 0, 0, 1) = \frac{1}{2}(1, 1, 1, 1) + \frac{1}{2}(1, -1, -1, 1).$

- So $(1, 0, 0, 1) \in \text{span}(S)$.

So $\text{span}(S) = V$.

Therefore, S is a basis for V .

72 / 106

Examples

- Let $S = \{(1, 1, 1, 1), (0, 0, 1, 2), (-1, 0, 0, 1)\}$.
 - Let $|S|$ be the number of vectors in S . Then $|S| = 3$.
 - So $\text{span}(S) \neq \mathbb{R}^4$; thus S is NOT a basis for \mathbb{R}^4 .
- Let $V = \text{span}(S)$, $S = \{(1, 1, 1), (0, 0, 1), (1, 1, 0)\}$.
 - $(1, 1, 1) = (0, 0, 1) + (1, 1, 0)$.
 - So S is linearly dependent; thus S is not a basis for V .
- **Remarks.**
 - A basis for a vector space V contains
 - smallest possible number of vectors that spans V ,
 - largest possible number of vectors that is linearly independent.
 - For convenience, \emptyset is said to be the **basis** for $\{0\}$.
 - Other than $\{0\}$, any vector space has infinitely many different bases.

73 / 106

Properties

- **Theorem.** Let $S = \{v_1, v_2, \dots, v_k\}$ be a subset of a vector space V . Then the following are equivalent:
 - S is a basis for V .
 - Every vector $v \in V$ can be **uniquely** expressed as
 - $v = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$, $c_i \in \mathbb{R}$.
 - **Proof.** \Rightarrow : Suppose S is a basis. Then $\text{span}(S) = V$.
 - For every $v \in V$, there exist $c_1, c_2, \dots, c_k \in \mathbb{R}$ s.t.
 - $v = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$.
 - Suppose $v = d_1 v_1 + d_2 v_2 + \dots + d_k v_k$, $d_i \in \mathbb{R}$.
 - $0 = (c_1 - d_1)v_1 + \dots + (c_k - d_k)v_k$.
- Since S is linearly independent,
- $c_1 - d_1 = c_2 - d_2 = \dots = c_k - d_k = 0$;
 - that is, $c_1 = d_1, c_2 = d_2, \dots, c_k = d_k$.

74 / 106

Properties

- **Theorem.** Let $S = \{v_1, v_2, \dots, v_k\}$ be a subset of a vector space V . Then the following are equivalent:
 - S is a basis for V .
 - Every vector $v \in V$ can be **uniquely** expressed as
 - $v = c_1 v_1 + c_2 v_2 + \dots + c_k v_k, c_i \in \mathbb{R}$.
- **Proof.** \Leftarrow : Suppose that every vector $v \in V$ can be uniquely expressed as
 - $v = c_1 v_1 + c_2 v_2 + \dots + c_k v_k, c_i \in \mathbb{R}$.Then by definition $\text{span}(S) = V$.
Let $0 \in V$. Suppose $0 = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$.
 - Note that $0 = 0v_1 + 0v_2 + \dots + 0v_k$.By the uniqueness, $c_1 = 0, c_2 = 0, \dots, c_k = 0$.
 - So S is linearly independent.

75 / 106

Coordinate Vector

- **Definition.** Let $S = \{v_1, v_2, \dots, v_k\}$ be a basis for a vector space V .
 - For every $v \in V$, there exist unique $c_1, \dots, c_k \in \mathbb{R}$ s.t.
 - $v = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$.
 - c_1, c_2, \dots, c_k are the **coordinates** of v **relative** to S .
 - (c_1, c_2, \dots, c_k) is the **coordinate vector** of v **relative** to the basis S , denoted by $(v)_S$.
- **Remark.** The order of v_1, v_2, \dots, v_k is fixed.
 - Let $S_1 = \{(1, 1), (-1, 1)\}$ be a basis for \mathbb{R}^2 . (Check!)
 - Let $v = 2(1, 1) + 3(-1, 1) = (-1, 5)$.
 - Then $(v)_{S_1} = (2, 3)$.Let $S_2 = \{(-1, 1), (1, 1)\}$. Then $(v)_{S_2} = (3, 2)$.

76 / 106

Examples

- Let $S = \{(1, 2, 1), (2, 9, 0), (3, 3, 4)\}$.
 - One can check that S is a basis for \mathbb{R}^3 . (Exercise!)

Let $\mathbf{v} = (5, -1, 9)$. Solve

- $\mathbf{v} = a(1, 2, 1) + b(2, 9, 0) + c(3, 3, 4)$.

$$\bullet \left(\begin{array}{ccc|c} 1 & 2 & 3 & 5 \\ 2 & 9 & 3 & -1 \\ 1 & 0 & 4 & 9 \end{array} \right) \xrightarrow{\text{G.-J.E.}} \left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 2 \end{array} \right).$$

$$(\mathbf{v})_S = (a, b, c) = (1, -1, 2).$$

Suppose that $(\mathbf{w})_S = (-1, 3, 2)$.

$$\begin{aligned} \mathbf{w} &= (-1)(1, 2, 1) + 3(2, 9, 0) + 2(3, 3, 4) \\ &= (11, 31, 7). \end{aligned}$$

77 / 106

Examples

- Let $\mathbf{v} = (2, 3) \in \mathbb{R}^2$.
 - Let $S_1 = \{(1, 0), (0, 1)\}$ be a basis for \mathbb{R}^2 .
 - $\mathbf{v} = 2(1, 0) + 3(0, 1)$; so $(\mathbf{v})_{S_1} = (2, 3)$.

- Let $S_2 = \{(1, -1), (1, 1)\}$ be a basis for \mathbb{R}^2 .

$$\bullet \left(\begin{array}{cc|c} 1 & 1 & 2 \\ -1 & 1 & 3 \end{array} \right) \xrightarrow{\text{G.-J.E.}} \left(\begin{array}{cc|c} 1 & 0 & -\frac{1}{2} \\ 0 & 1 & \frac{5}{2} \end{array} \right).$$

$$\bullet (\mathbf{v})_{S_2} = \left(-\frac{1}{2}, \frac{5}{2}\right).$$

- Let $S_3 = \{(1, 0), (1, 1)\}$ be a basis for \mathbb{R}^2 .

$$\bullet \left(\begin{array}{cc|c} 1 & 1 & 2 \\ 0 & 1 & 3 \end{array} \right) \xrightarrow{R_1 + (-1)R_2} \left(\begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & 3 \end{array} \right).$$

$$\bullet (\mathbf{v})_{S_3} = (-1, 3).$$

78 / 106

Standard Basis

• **Definition.** Let $E = \{e_1, e_2, \dots, e_n\}$ be a subset of \mathbb{R}^n ,

◦ $e_1 = (1, 0, \dots, 0), e_2 = (0, 1, \dots, 0), \dots, e_n = (0, 0, \dots, 1).$

1. Let $v = (v_1, v_2, \dots, v_n) \in \mathbb{R}^n$. Then

◦ $v = v_1 e_1 + v_2 e_2 + \dots + v_n e_n.$

So $\text{span}(E) = \mathbb{R}^n$.

2. Suppose that $c_1 e_1 + c_2 e_2 + \dots + c_n e_n = \mathbf{0}$. Then

◦ $(c_1, c_2, \dots, c_n) = (0, 0, \dots, 0).$

So E is linearly independent.

E is called the **standard basis** for \mathbb{R}^n .

◦ For any $v = (v_1, v_2, \dots, v_n) \in \mathbb{R}^n$,

• $(v)_E = (v_1, v_2, \dots, v_n) = v.$

79 / 106

Properties

• **Theorem.** Let S be a basis for a vector space V .

◦ $(v)_S = \mathbf{0} \Leftrightarrow v = \mathbf{0}.$

◦ For any $c \in \mathbb{R}$ and $v \in \mathbb{R}$, $(cv)_S = c(v)_S.$

◦ For any $u, v \in V$, $(u + v)_S = (u)_S + (v)_S.$

Proof. Let $S = \{v_1, \dots, v_k\}.$

◦ \Rightarrow : Suppose $(v)_S = (\overbrace{0, 0, \dots, 0}^k) = \mathbf{0} \in \mathbb{R}^k.$

• $v = 0v_1 + \dots + 0v_k = \mathbf{0} \in V.$

\Leftarrow : Let $v = \mathbf{0} \in V$. Then $v = 0v_1 + \dots + 0v_k.$

• $(v)_S = (\overbrace{0, \dots, 0}^k) = \mathbf{0} \in \mathbb{R}^k.$

80 / 106

Properties

- **Theorem.** Let S be a basis for a vector space V

- $(\mathbf{v})_S = \mathbf{0} \Leftrightarrow \mathbf{v} = \mathbf{0}$.
- For any $c \in \mathbb{R}$ and $\mathbf{v} \in \mathbb{R}$, $(c\mathbf{v})_S = c(\mathbf{v})_S$.
- For any $\mathbf{u}, \mathbf{v} \in V$, $(\mathbf{u} + \mathbf{v})_S = (\mathbf{u})_S + (\mathbf{v})_S$.

Proof. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$.

- Let $(\mathbf{v})_S = (c_1, \dots, c_k)$. Then
 - $\mathbf{v} = c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k$.
 - $c\mathbf{v} = cc_1\mathbf{v}_1 + \dots + cc_k\mathbf{v}_k$.
$$(c\mathbf{v})_S = (cc_1, \dots, cc_k) = c(c_1, \dots, c_k) = c(\mathbf{v})_S.$$
- Let $(\mathbf{u})_S = (c_1, \dots, c_k)$, $(\mathbf{v})_S = (d_1, \dots, d_k)$.
 - $\mathbf{u} = c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k$, $\mathbf{v} = d_1\mathbf{v}_1 + \dots + d_k\mathbf{v}_k$.
 - $\mathbf{u} + \mathbf{v} = (c_1 + d_1)\mathbf{v}_1 + \dots + (c_k + d_k)\mathbf{v}_k$.
$$(\mathbf{u} + \mathbf{v})_S = (c_1 + d_1, \dots, c_k + d_k) = (\mathbf{u})_S + (\mathbf{v})_S.$$

81 / 106

Properties

- **Theorem.** Let S be a basis for a vector space V .

- For any $\mathbf{u}, \mathbf{v} \in V$, $\mathbf{u} = \mathbf{v} \Leftrightarrow (\mathbf{u})_S = (\mathbf{v})_S$.
- For any $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r \in V$ and $c_1, c_2, \dots, c_r \in \mathbb{R}$,
 - $(c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r)_S = c_1(\mathbf{v}_1)_S + \dots + c_r(\mathbf{v}_r)_S$.

Proof. Left as exercises.

- **Theorem.** Let S be a basis for a vector space V .

- Suppose $|S| = k$. Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r \in V$.
 1. $\mathbf{v}_1, \dots, \mathbf{v}_r$ are linearly independent

$$\Leftrightarrow (\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S \text{ are linearly independent.}$$
 2. $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\} = V$

$$\Leftrightarrow \text{span}\{(\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S\} = \mathbb{R}^k.$$

82 / 106

Properties

- **Proof.** 1. \Rightarrow : Suppose $\mathbf{v}_1, \dots, \mathbf{v}_r$ are linearly independent.
 - Consider equation $c_1(\mathbf{v}_1)_S + \dots + c_r(\mathbf{v}_r)_S = \mathbf{0} \in \mathbb{R}^k$.
 - $(c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r)_S = (\mathbf{0})_S$, where $\mathbf{0} \in V$.
 Then $c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r = \mathbf{0}$.
 - $\mathbf{v}_1, \dots, \mathbf{v}_r$ linearly independent $\Rightarrow c_1 = \dots = c_r = 0$.
 Therefore, $(\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S$ are linearly independent.
 - \Leftarrow : Suppose $(\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S$ are linearly independent.
 - Consider equation $c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r = \mathbf{0} \in V$.
 - $(c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r)_S = (\mathbf{0})_S$.
 Then $c_1(\mathbf{v}_1)_S + \dots + c_r(\mathbf{v}_r)_S = \mathbf{0} \in \mathbb{R}^k$.
 - $(\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S$ are linearly independent
 - $\Rightarrow c_1 = \dots = c_r = 0$.
 Therefore, $\mathbf{v}_1, \dots, \mathbf{v}_r$ are linearly independent.

83 / 106

Properties

- **Proof.** 2. \Rightarrow : Suppose $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\} = V$.
 - Let $\mathbf{w} = (c_1, \dots, c_k) \in \mathbb{R}^k$. If $S = \{\mathbf{u}_1, \dots, \mathbf{u}_k\}$,
 - then $\mathbf{v} = c_1\mathbf{u}_1 + \dots + c_k\mathbf{u}_k \in V$, $(\mathbf{v})_S = \mathbf{w}$.
 Since $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\} = V$, there exist $d_i \in \mathbb{R}$ s.t.
 - $\mathbf{v} = d_1\mathbf{v}_1 + \dots + d_r\mathbf{v}_r$.
 Then $\mathbf{w} = (\mathbf{v})_S = d_1(\mathbf{v}_1)_S + \dots + d_r(\mathbf{v}_r)_S$.
 Therefore, $\text{span}\{(\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S\} = \mathbb{R}^k$.
 - 2. \Leftarrow : Suppose $\text{span}\{(\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S\} = \mathbb{R}^k$.
 - Let $\mathbf{v} \in V$. Then $(\mathbf{v})_S \in \mathbb{R}^k$. There exist $c_i \in \mathbb{R}^k$ s.t.
 - $(\mathbf{v})_S = c_1(\mathbf{v}_1)_S + \dots + c_r(\mathbf{v}_r)_S$.
 - $(\mathbf{v})_S = (c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r)_S$.
 Then $\mathbf{v} = c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r$.
 Therefore, $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\} = V$.

84 / 106

Criterion for Bases

- Let $S = \{v_1, \dots, v_k\}$ be a subset of \mathbb{R}^n .
 - If $k > n$, then S is linearly dependent.
 - If $k < n$, then $\text{span}(S) \neq \mathbb{R}^n$.

If S is a basis, then $k = n$.
- **Theorem.** Let V be a vector space having a basis with k vectors.
 - Any subset of V of $> k$ vectors is linearly dependent.
 - Any subset of V of $< k$ vectors cannot span V .
- **Corollary.** All bases of a vector space have same size.
 - To be more precise, if S_1 and S_2 are two bases of a vector space V ,
 - then $|S_1| = |S_2|$.

86 / 106

Criterion for Bases

- **Proof.** Let S be a basis of V with $|S| = k$.
 - Let $T = \{v_1, \dots, v_r\}$ be a subset of V .
 - Then $\{(v_1)_S, \dots, (v_r)_S\}$ is a subset of \mathbb{R}^k .
 1. Suppose $r > k$.
 - $\{(v_1)_S, \dots, (v_r)_S\}$ is linearly dependent in \mathbb{R}^k .

Then $\{v_1, \dots, v_r\}$ is linearly dependent in V .
 2. Suppose $r < k$.
 - $\text{span}\{(v_1)_S, \dots, (v_r)_S\} \neq \mathbb{R}^k$.

Then $\text{span}\{v_1, \dots, v_r\} \neq V$.

87 / 106

Dimension

- **Definition.** Let V be a vector space and S a basis for V .
 - The **dimension** of V is $\dim(V) = |S|$.
- **Examples.**
 - \emptyset is a (the) basis for $\{\mathbf{0}\}$.
 - Then $\dim(\{\mathbf{0}\}) = |\emptyset| = 0$.
 - \mathbb{R}^n has the standard basis $E = \{e_1, e_2, \dots, e_n\}$.
 - Then $\dim(\mathbb{R}^n) = n$.
 - In \mathbb{R}^2 and \mathbb{R}^3 , every straight line through the origin is of the form $\text{span}\{v\}$ with $v \neq \mathbf{0}$.
 - The dimension of such a straight line is 1.
 - In \mathbb{R}^3 , every plane containing the origin is of the form $\text{span}\{u, v\}$, where u, v are linearly independent.
 - The dimension of such a plane is 2.

88 / 106

Dimension of Solution Space

- Let $Ax = \mathbf{0}$ be a homogeneous linear system.
 - Recall that the solution set is a vector space V .
- Let R be a row-echelon form of A .
- $$\begin{aligned} & \text{no. of non-pivot cols of } R \\ &= \text{no. of arbitrary parameters in soln} \\ &= \text{the dimension of } V. \end{aligned}$$
- **Example.** $x + y + z = 0$.
 - $(1 \ 1 \ 1) \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 0$.
 - The 2nd and 3rd columns are non-pivot.
 - The dimension of the solution space is 2.

89 / 106

Example

$$\bullet \begin{cases} 2v + 2w - x + z = 0 \\ -v - w + 2x - 3y + z = 0 \\ x + y + z = 0 \\ v + w - 2x - z = 0 \end{cases}$$

$$\circ \begin{pmatrix} 2 & 2 & -1 & 0 & 1 \\ -1 & -1 & 2 & -3 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & -2 & 0 & -1 \end{pmatrix}$$

$$\xrightarrow[\text{Elimination}]{\text{Gaussian}} \begin{pmatrix} 2 & 2 & -1 & 0 & 1 \\ 0 & 0 & \frac{3}{2} & -3 & \frac{3}{2} \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

- The 2nd and 5th columns are non-pivot.
- The solution space has dimension 2.

90 / 106

Properties

- **Theorem.** Let S be a subset of a vector space V . The following are equivalent:
 1. S is a basis for V .
 2. S is linearly independent, and $|S| = \dim(V)$.
 3. S spans V , and $|S| = \dim(V)$.
- To check whether a subset S is a basis for a vector space V , simply check any **two** of the following three conditions:
 - S is linearly independent,
 - S spans V ,
 - $|S| = \dim(V)$.
- **Example.** Let $S = \{(2, 0, -1), (4, 0, 7), (-1, 1, 4)\}$.
 - One can check that S is linearly independent.

Since $|S| = 3$, S is a basis for \mathbb{R}^3 .

91 / 106

Properties

- **Proof.** “ $1 \Rightarrow 2$ ” and “ $1 \Rightarrow 3$ ” are clear.

“ $2 \Rightarrow 1$ ”: Suppose S is linearly indept. & $|S| = \dim(V)$.

- It suffices to show that $\text{span}(S) = V$.

Assume $\text{span}(S) \neq V$. Pick $v \in V$ but $v \notin \text{span}(S)$.

- Then $S \cup \{v\}$ is linearly independent.

But $|S \cup \{v\}| = \dim(V) + 1 > \dim(V)$.

- Then $S \cup \{v\}$ is linearly dependent.

Therefore, we must have $\text{span}(S) = V$.

- Since S is linearly independent, S is a basis for V .

92 / 106

Properties

- **Proof.** “ $1 \Rightarrow 2$ ” and “ $1 \Rightarrow 3$ ” are clear.

“ $3 \Rightarrow 1$ ”: Suppose S spans V & $|S| = \dim(V)$.

- It suffices to show that S is linearly independent.

Assume that S is linearly dependent.

- Then there exists $v \in S$ such that v is a linear combination of other vectors in S .

- Hence, $\text{span}(S - \{v\}) = \text{span}(S) = V$.

On the other hand, $|S - \{v\}| = \dim(V) - 1 < \dim(V)$.

- Then $\text{span}(S - \{v\}) \neq V$.

Therefore, S must be linearly independent.

- Since S spans V , S is a basis for V .

93 / 106

Properties

- **Theorem.** Let U be a subspace of a vector space V .

- $U = V \Leftrightarrow \dim(U) = \dim(V)$.

Proof. \Rightarrow : Clear!

\Leftarrow : Suppose $\dim(U) = \dim(V)$.

- Let S be a basis for U . Then
 - S is linearly independent (in U , and thus) in V .
 - $|S| = \dim(U) = \dim(V)$.

Then S is also a basis for V .

- Therefore, $V = \text{span}(S) = U$.

- **Corollary.** Let U be a subspace of a vector space V .

- $U \neq V \Leftrightarrow \dim(U) < \dim(V)$.

94 / 106

Properties

- **Theorem.** Let A be a square matrix of order n . Then the following are equivalent:

1. A is invertible.
2. $Ax = b$ has a unique solution.
3. $Ax = 0$ has only the trivial solution.
4. The reduced row-echelon form of A is I_n .
5. A is a product of elementary matrices.
6. $\det(A) \neq 0$.
7. The rows of A form a basis for \mathbb{R}^n .
8. The columns of A form a basis for \mathbb{R}^n .

- We have proved the equivalence of 1 to 6.

- It remains to show that " $1 \Leftrightarrow 7$ " & " $1 \Leftrightarrow 8$ ".

95 / 106

Properties

- **Proof.** “1 \Leftrightarrow 8”: Let \mathbf{a}_j be the j th column of \mathbf{A} .

- $\mathbf{A} = (\mathbf{a}_1 \quad \mathbf{a}_2 \quad \cdots \quad \mathbf{a}_n)$.

$\{\mathbf{a}_1, \dots, \mathbf{a}_n\}$ is a basis for \mathbb{R}^n

$\Leftrightarrow \text{span}\{\mathbf{a}_1, \dots, \mathbf{a}_n\} = \mathbb{R}^n$

\Leftrightarrow a row-echelon form of \mathbf{A} has no zero row

$\Leftrightarrow \mathbf{A}$ is invertible.

“1 \Leftrightarrow 7”:

rows of \mathbf{A} form a basis for \mathbb{R}^n

\Leftrightarrow columns of \mathbf{A}^T form a basis for \mathbb{R}^n

$\Leftrightarrow \mathbf{A}^T$ is invertible

$\Leftrightarrow \mathbf{A}$ is invertible.

96 / 106

Examples

- Let $\mathbf{v}_1 = (1, 1, 1)$, $\mathbf{v}_2 = (-1, 1, 2)$ and $\mathbf{v}_3 = (1, 0, 1)$.

- Let $\mathbf{A} = \begin{pmatrix} \mathbf{v}_1 \\ \mathbf{v}_2 \\ \mathbf{v}_3 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ -1 & 1 & 2 \\ 1 & 0 & 1 \end{pmatrix}$. Then $\det(\mathbf{A}) = 3$.

- \mathbf{A} is invertible.

- So $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is a basis for \mathbb{R}^3 .

- $\mathbf{v}_1 = (1, 1, 1, 1)$, $\mathbf{v}_2 = (1, -1, 1, -1)$, $\mathbf{v}_3 = (0, 1, -1, 0)$,

$\mathbf{v}_4 = (2, 1, 1, 0)$.

- $\begin{pmatrix} 1 & 1 & 0 & 2 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 0 & 0 \end{pmatrix} \xrightarrow{\text{Gaussian elimination}} \begin{pmatrix} 1 & 1 & 0 & 2 \\ 0 & -2 & 1 & -1 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$.

- So $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ is NOT a basis for \mathbb{R}^4 .

97 / 106

Coordinate Vector

- Let $S = \{v_1, \dots, v_k\}$ be a basis for a vector space V .
 - Then every vector $v \in V$ can be uniquely expressed as a linear combination of v_1, \dots, v_k :
 - $v = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$, where $c_i \in \mathbb{R}$.

Then $(c_1, c_2, \dots, c_k) = (v)_S$ is the **coordinate vector** of v relative to the basis S .

View each v_i as a column vector. Then

$$\circ \begin{pmatrix} v_1 & v_2 & \cdots & v_k \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_k \end{pmatrix} = v.$$

99 / 106

Coordinate Vector

- Let $S = \{v_1, \dots, v_k\}$ be a basis for a vector space V .
 - Then every vector $v \in V$ can be uniquely expressed as a linear combination of v_1, \dots, v_k :
 - $v = c_1 v_1 + c_2 v_2 + \dots + c_k v_k$, where $c_i \in \mathbb{R}$.

The **column vector** $[v]_S = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_k \end{pmatrix}$ is also called

- the **coordinate vector** of v relative to S .
- View v_1, \dots, v_k as column vectors.
 - Let $A = \begin{pmatrix} v_1 & v_2 & \cdots & v_k \end{pmatrix}$. Then
 - $A[v]_S = v$ for every $v \in V$.

100 / 106

Transition Matrix

- Let S and T be bases for a vector space V .
 - $S = \{u_1, u_2, \dots, u_k\}$ and $T = \{v_1, v_2, \dots, v_k\}$.
- Let $w \in V$. What is the relation between $[w]_S$ and $[w]_T$?

- Suppose all u_j, v_j and w are viewed as column vectors.
 - Let $A = (u_1 \ \dots \ u_k)$ and $B = (v_1 \ \dots \ v_k)$.

Let $w \in V$. Then

$$\begin{aligned} w &= A[w]_S = (u_1 \ \dots \ u_k) [w]_S \\ &= (B[u_1]_T \ \dots \ B[u_k]_T) [w]_S \\ &= B([u_1]_T \ \dots \ [u_k]_T) [w]_S \end{aligned}$$

So $([u_1]_T \ \dots \ [u_k]_T) [w]_S$ is the coordinate vector of w relative to the basis T ; that is,

- $([u_1]_T \ \dots \ [u_k]_T) [w]_S = [w]_T$.

101 / 106

Transition Matrix

- Definition.** Let V be a vector space, and
 - $S = \{u_1, \dots, u_k\}$ and T be bases for V .

$([u_1]_T \ \dots \ [u_k]_T)$ is the **transition matrix** from S to T .

 - Denote it by P . Then $P[w]_S = [w]_T$ for all $w \in V$.
- Example.** Let $S = \{u_1, u_2, u_3\}$, $T = \{v_1, v_2, v_3\}$.
 - $u_1 = (1, 0, -1)$, $u_2 = (0, -1, 0)$, $u_3 = (1, 0, 2)$.
 - $v_1 = (1, 1, 1)$, $v_2 = (1, 1, 0)$, $v_3 = (-1, 0, 0)$.

View all vectors as column vectors.

- $(v_1 \ v_2 \ v_3 \mid u_1 \mid u_2 \mid u_3)$

$$\xrightarrow{\text{G.-J.E.}} \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & 0 & 2 \\ 0 & 1 & 0 & 1 & -1 & -2 \\ 0 & 0 & 1 & -1 & -1 & -1 \end{array} \right) = (I \mid P).$$

102 / 106

Transition Matrix

- **Definition.** Let V be a vector space, and
 - $S = \{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ and T be bases for V .
 - $([\mathbf{u}_1]_T \ \cdots \ [\mathbf{u}_k]_T)$ is the **transition matrix** from S to T .
 - Denote it by P . Then $P[\mathbf{w}]_S = [\mathbf{w}]_T$ for all $\mathbf{w} \in V$.
- **Example.** Let $S = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$, $T = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.
 - $\mathbf{u}_1 = (1, 0, -1)$, $\mathbf{u}_2 = (0, -1, 0)$, $\mathbf{u}_3 = (1, 0, 2)$.
 - $\mathbf{v}_1 = (1, 1, 1)$, $\mathbf{v}_2 = (1, 1, 0)$, $\mathbf{v}_3 = (-1, 0, 0)$.

Transition matrix from S to T is $P = \begin{pmatrix} -1 & 0 & 2 \\ 1 & -1 & -2 \\ -1 & -1 & -1 \end{pmatrix}$.

 - Suppose $(\mathbf{w})_S = (2, -1, 2)$.
 - $[\mathbf{w}]_T = P[\mathbf{w}]_S = \begin{pmatrix} 2 \\ -1 \\ -3 \end{pmatrix}$. $(\mathbf{w})_T = (2, -1, -3)$.

103 / 106

Properties

- **Theorem.** Let S and T be bases for a vector space V .
 - Let P be the transition matrix from S to T . Then
 - P is an invertible matrix.
 - P^{-1} is the transition matrix from T to S .
- **Proof.** Let Q be the transition matrix from T to S .
 - It suffices to show that $QP = I$.

Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$. Then

 - $[\mathbf{v}_1]_S = \mathbf{e}_1, [\mathbf{v}_2]_S = \mathbf{e}_2, \dots, [\mathbf{v}_k]_S = \mathbf{e}_k$.
$$\begin{aligned}
 QP &= QPI = QP(\mathbf{e}_1 \ \cdots \ \mathbf{e}_k) = (QP\mathbf{e}_1 \ \cdots QP\mathbf{e}_k) \\
 &= (QP[\mathbf{v}_1]_S \ \cdots QP[\mathbf{v}_k]_S) \\
 &= (Q[\mathbf{v}_1]_T \ \cdots Q[\mathbf{v}_k]_T) \\
 &= ([\mathbf{v}_1]_S \ \cdots [\mathbf{v}_k]_S) \\
 &= (\mathbf{e}_1 \ \cdots \ \mathbf{e}_k) = I.
 \end{aligned}$$

104 / 106

Examples

- Let $S = \{\mathbf{u}_1, \mathbf{u}_2\}$, $\mathbf{u}_1 = (1, 1)$, $\mathbf{u}_2 = (1, -1)$.

$$T = \{\mathbf{v}_1, \mathbf{v}_2\}, \mathbf{v}_1 = (1, 0), \mathbf{v}_2 = (1, 1).$$

Note that both S and T are bases for \mathbb{R}^2 .

$$\circ \left(\begin{array}{cc|cc} \mathbf{v}_1 & \mathbf{v}_2 & \mathbf{u}_1 & \mathbf{u}_2 \end{array} \right) \xrightarrow{R_1 + (-1)R_2} \left(\begin{array}{cc|cc} 1 & 0 & 0 & 2 \\ 0 & 1 & 1 & -1 \end{array} \right)$$

- Transition matrix from S to T : $\mathbf{P} = \begin{pmatrix} 0 & 2 \\ 1 & -1 \end{pmatrix}$.

$$\circ \left(\begin{array}{cc|cc} \mathbf{u}_1 & \mathbf{u}_2 & \mathbf{v}_1 & \mathbf{v}_2 \end{array} \right) \xrightarrow{\text{G.J.E.}} \left(\begin{array}{cc|cc} 1 & 0 & \frac{1}{2} & 1 \\ 0 & 1 & \frac{1}{2} & 0 \end{array} \right)$$

- Transition matrix from T to S : $\mathbf{Q} = \begin{pmatrix} \frac{1}{2} & 1 \\ \frac{1}{2} & 0 \end{pmatrix}$.

- One checks easily that $\mathbf{PQ} = \mathbf{QP} = \mathbf{I}$.

105 / 106

Examples

- Let $S = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ and $T = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

$$\circ S = \{(1, 0, -1), (0, -1, 0), (1, 0, 2)\};$$

$$\circ T = \{(1, 1, 1), (1, 1, 0), (-1, 0, 0)\}.$$

We have computed the transition matrix from S to T :

$$\circ \mathbf{P} = \begin{pmatrix} -1 & 0 & 2 \\ 1 & -1 & -2 \\ -1 & -1 & -1 \end{pmatrix}.$$

Then the transition matrix from T to S is

$$\circ \mathbf{P}^{-1} = \dots = \begin{pmatrix} \frac{1}{3} & \frac{2}{3} & -\frac{2}{3} \\ -1 & -1 & 0 \\ \frac{2}{3} & \frac{1}{3} & -\frac{1}{3} \end{pmatrix}.$$

For any $\mathbf{w} \in \mathbb{R}^3 = \text{span}(S) = \text{span}(T)$,

$$\circ \mathbf{P}[\mathbf{w}]_S = [\mathbf{w}]_T \text{ and } \mathbf{P}^{-1}[\mathbf{w}]_T = [\mathbf{w}]_S.$$

106 / 106