

MA2001 LINEAR ALGEBRA

DIAGONALIZATION

Goh Jun Le / Wang Fei

gohjunle@nus.edu.sg / matwf@nus.edu.sg

Department of Mathematics
Office: S17-06-25 / S17-06-16
Tel: 6601-1355 / 6516-2937

Eigenvalues and Eigenvectors	2
Motivations	3
Definitions	9
Characteristic Equation	11
Eigenspace	19
Diagonalization	26
Diagonalizable Matrices	27
Criterion of Diagonalizability	30
Diagonalization	33
Examples	38
Application	47
Orthogonal Diagonalization	53
Introduction	54
Definition	56
Algorithm	58
Examples	61
Quadratic Forms and Conic Sections	70
Quadratic Form	71
Simplification	75
Quadratic Equation	78
Classification	89

Motivations

- Let A be a square matrix. Then

- $$A^m = \underbrace{AA \cdots AA}_{m \text{ times}}$$

In general, the matrix multiplication is complicated.

- Is there a shortcut?
- Suppose A and B are **diagonal matrices** of order n .

- $$A = \begin{pmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{pmatrix}, B = \begin{pmatrix} b_{11} & 0 & \cdots & 0 \\ 0 & b_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & b_{nn} \end{pmatrix}$$
- $$AB = \begin{pmatrix} a_{11}b_{11} & 0 & \cdots & 0 \\ 0 & a_{22}b_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn}b_{nn} \end{pmatrix}.$$

3 / 96

Motivations

- Let A be a square matrix. Then

- $$A^m = \underbrace{AA \cdots AA}_{m \text{ times}}$$

In general, the matrix multiplication is complicated.

- Is there a shortcut?
- Suppose A is a **diagonal matrix** of order n .

- $$A = \begin{pmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{pmatrix}.$$
- $$A^m = \begin{pmatrix} a_{11}^m & 0 & \cdots & 0 \\ 0 & a_{22}^m & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn}^m \end{pmatrix}.$$

4 / 96

Motivations

- Let A be a square matrix.
 - Suppose there exists an invertible matrix P such that
 - $P^{-1}AP = D$ is a diagonal matrix.

Then $A = PDP^{-1}$.

$$\begin{aligned}
 A^m &= (PDP^{-1})^m \\
 &= \underbrace{(PDP^{-1})(PDP^{-1}) \cdots (PDP^{-1})}_{m \text{ times}} \\
 &= PD(P^{-1}P)D(P^{-1}P) \cdots (P^{-1}P)DP^{-1} \\
 &= P \underbrace{DD \cdots DD}_{m \text{ times}} P^{-1} \\
 &= PD^m P^{-1}.
 \end{aligned}$$

5 / 96

Motivations

- **Example.** Suppose that each year
 - 4% of the rural population moves to the urban district.
 - 1% of the urban populations moves to the rural district.

After n years,

- Let a_n be the rural population;
- Let b_n be the urban population.

$$a_n = 0.96a_{n-1} + 0.01b_{n-1}, \quad b_n = 0.04a_{n-1} + 0.99b_{n-1}.$$

$$\begin{pmatrix} a_n \\ b_n \end{pmatrix} = \begin{pmatrix} 0.96 & 0.01 \\ 0.04 & 0.99 \end{pmatrix} \begin{pmatrix} a_{n-1} \\ b_{n-1} \end{pmatrix}.$$

$$\text{Let } \mathbf{x}_n = \begin{pmatrix} a_n \\ b_n \end{pmatrix} \text{ and } \mathbf{A} = \begin{pmatrix} 0.96 & 0.01 \\ 0.04 & 0.99 \end{pmatrix}.$$

$$\mathbf{x}_n = \mathbf{A}\mathbf{x}_{n-1} = \mathbf{A}^2\mathbf{x}_{n-2} = \cdots = \mathbf{A}^n\mathbf{x}_0.$$

6 / 96

Motivations

- Let $\mathbf{x}_n = \begin{pmatrix} a_n \\ b_n \end{pmatrix}$ and $\mathbf{A} = \begin{pmatrix} 0.96 & 0.01 \\ 0.04 & 0.99 \end{pmatrix}$.

- $\mathbf{x}_n = \mathbf{A}\mathbf{x}_{n-1} = \mathbf{A}^2\mathbf{x}_{n-2} = \dots = \mathbf{A}^n\mathbf{x}_0$.

Let $\mathbf{P} = \begin{pmatrix} 1 & 1 \\ 4 & -1 \end{pmatrix}$. Then $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D} = \begin{pmatrix} 1 & 0 \\ 0 & 0.95 \end{pmatrix}$.

- $\mathbf{A}^n = \mathbf{P}\mathbf{D}^n\mathbf{P}^{-1}$

- $\mathbf{A}^n = \begin{pmatrix} 0.2 + 0.8 \cdot 0.95^n & 0.2 - 0.2 \cdot 0.95^n \\ 0.8 - 0.8 \cdot 0.95^n & 0.8 + 0.2 \cdot 0.95^n \end{pmatrix}$

- $\begin{pmatrix} a_n \\ b_n \end{pmatrix} = \mathbf{x}_n = \mathbf{A}^n\mathbf{x}_0 = \mathbf{A}^n \begin{pmatrix} a_0 \\ b_0 \end{pmatrix}$.

- $\begin{pmatrix} a_n \\ b_n \end{pmatrix} = \begin{pmatrix} 0.2a_0 + 0.2b_0 + (0.8a_0 - 0.2b_0) \cdot 0.95^n \\ 0.8a_0 + 0.8b_0 - (0.8a_0 - 0.2b_0) \cdot 0.95^n \end{pmatrix}$.

In particular, $\begin{pmatrix} a_n \\ b_n \end{pmatrix} \xrightarrow{n \rightarrow \infty} \begin{pmatrix} 0.2(a_0 + b_0) \\ 0.8(a_0 + b_0) \end{pmatrix}$.

7 / 96

Motivations

- Let \mathbf{A} be a square matrix of order 3.
- Suppose $\mathbf{P} = (\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3)$ is invertible such that

- $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D} = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$.

Then $\mathbf{A}\mathbf{P} = \mathbf{P}\mathbf{D}$.

- $\mathbf{A}(\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3) = (\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3) \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$

- $(\mathbf{A}\mathbf{v}_1 \ \mathbf{A}\mathbf{v}_2 \ \mathbf{A}\mathbf{v}_3) = (\lambda_1\mathbf{v}_1 \ \lambda_2\mathbf{v}_2 \ \lambda_3\mathbf{v}_3)$.

- Hence, $\mathbf{A}\mathbf{v}_1 = \lambda_1\mathbf{v}_1$, $\mathbf{A}\mathbf{v}_2 = \lambda_2\mathbf{v}_2$, $\mathbf{A}\mathbf{v}_3 = \lambda_3\mathbf{v}_3$.

8 / 96

Definitions

- **Definition.** Let A be a square matrix of order n .
 - Suppose that for some $\lambda \in \mathbb{R}$ and **nonzero** $v \in \mathbb{R}^n$
 - $Av = \lambda v$
 - λ is called an **eigenvalue** of A .
 - v is called an **eigenvector** of A associated with λ .
- **Example.** Let $A = \begin{pmatrix} 0.96 & 0.01 \\ 0.04 & 0.99 \end{pmatrix}$.
 - Let $u = \begin{pmatrix} 1 \\ 4 \end{pmatrix}$. Then $Au = \begin{pmatrix} 1 \\ 4 \end{pmatrix} = 1u$.
 - u is an eigenvector of A associated to eigenvalue 1.
 - Let $v = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$. $Av = \begin{pmatrix} 0.95 \\ -0.95 \end{pmatrix} = 0.95v$.
 - v is an eigenvector associated to eigenvalue 0.95.

9 / 96

Example

- Let $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$.
 - Let $u = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$. Then $Bu = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} = 3u$.
 - u is an eigenvector of B associated to eigenvalue 3.
 - Let $v = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$. Then $Bv = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} = 0v$.
 - v is an eigenvector of B associated to eigenvalue 0.
 - Let $w = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$. Then $Bw = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} = 0w$.
 - w is an eigenvector of B associated to eigenvalue 0.

10 / 96

Characteristic Equation

- Let A be a square matrix. How to find its eigenvalues?

- $\lambda \in \mathbb{R}$ is an eigenvalue of A
 - $\Leftrightarrow Av = \lambda v$ for some nonzero column vector v
 - $\Leftrightarrow \lambda v - Av = 0$ for some nonzero column vector v
 - $\Leftrightarrow (\lambda I - A)v = 0$ for some nonzero column vector v
 - $\Leftrightarrow (\lambda I - A)x = 0$ has non-trivial solution
 - $\Leftrightarrow \lambda I - A$ a singular matrix
 - $\Leftrightarrow \det(\lambda I - A) = 0$.

If A is of order n , then $\det(\lambda I - A)$ is a monic polynomial in λ of degree n :

$$\lambda^n + c_{n-1}\lambda^{n-1} + \cdots + c_1\lambda + c_0.$$

- Definition.** Let A be a square matrix.
 - $\det(\lambda I - A)$ is the **characteristic polynomial** of A .
 - $\det(\lambda I - A) = 0$ is the **characteristic equation** of A .

11 / 96

Characteristic Equation

- Theorem.** Let A be a square matrix.

- Then the eigenvalues of A are precisely all the roots to the characteristic equation $\det(\lambda I - A) = 0$.

- Examples.**

- Let $A = \begin{pmatrix} 0.96 & 0.01 \\ 0.04 & 0.99 \end{pmatrix}$. Characteristic polynomial is

$$\begin{aligned} \det(\lambda I - A) &= \det \left(\lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 0.96 & 0.01 \\ 0.04 & 0.99 \end{pmatrix} \right) \\ &= \det \begin{pmatrix} \lambda - 0.96 & -0.01 \\ -0.04 & \lambda - 0.99 \end{pmatrix} \\ &= (\lambda - 0.96)(\lambda - 0.99) - (-0.01)(-0.04) \\ &= \lambda^2 - 1.95\lambda + 0.95 \\ &= (\lambda - 0.95)(\lambda - 1). \end{aligned}$$

Hence, A has two eigenvalues 0.95 and 1.

12 / 96

Characteristic Equation

- **Theorem.** Let A be a square matrix.
 - Then the eigenvalues of A are precisely all the roots to the characteristic equation $\det(\lambda I - A) = 0$.

- **Examples.**

- Let $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$. Characteristic polynomial:

$$\begin{aligned}\det(\lambda I - B) &= \begin{vmatrix} \lambda - 1 & -1 & -1 \\ -1 & \lambda - 1 & -1 \\ -1 & -1 & \lambda - 1 \end{vmatrix} \\ &= \lambda^3 - 3\lambda^2 \\ &= \lambda^2(\lambda - 3).\end{aligned}$$

Hence, B has two eigenvalues 0 and 3.

13 / 96

Characteristic Equation

- **Theorem.** Let A be a square matrix.
 - Then the eigenvalues of A are precisely all the roots to the characteristic equation $\det(\lambda I - A) = 0$.

- **Examples.**

- Let $C = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 2 \\ 1 & 1 & 1 \end{pmatrix}$. Characteristic polynomial:

$$\begin{aligned}\det(\lambda I - C) &= \begin{vmatrix} \lambda & 1 & 0 \\ 0 & \lambda & -2 \\ -1 & -1 & \lambda - 1 \end{vmatrix} \\ &= \lambda^3 - \lambda^2 - 2\lambda + 2 \\ &= (\lambda - 1)(\lambda - \sqrt{2})(\lambda + \sqrt{2}).\end{aligned}$$

Hence, C has three eigenvalues 1, $\sqrt{2}$ and $-\sqrt{2}$.

14 / 96

Main Theorem for Invertible Matrices

- **Theorem.** Let A be a square matrix of order n . Then the following are equivalent:

1. A is invertible.
2. The reduced row-echelon form of A is I_n .
3. The homogeneous linear system $Ax = 0$ has only the trivial solution.
4. The linear system $Ax = b$ has exactly one solution.
5. A is the product of elementary matrices.
6. $\det(A) \neq 0$.
7. The rows of A form a basis for \mathbb{R}^n .
8. The columns of A form a basis for \mathbb{R}^n .
9. $\text{rank}(A) = n$.
10. 0 is not an eigenvalue of A .

15 / 96

Main Theorem for Invertible Matrices

- **Proof.** It remains to show that “10” is equivalent to “6”:

- 0 is not an eigenvalue of A
 - $\Leftrightarrow 0$ is not a root to $\det(\lambda I - A) = 0$
 - $\Leftrightarrow \det(0I - A) \neq 0$
 - $\Leftrightarrow \det(-A) \neq 0$
 - $\Leftrightarrow (-1)^n \det(A) \neq 0$
 - $\Leftrightarrow \det(A) \neq 0$.

16 / 96

Upper Triangular Matrices

- Let A be an **upper triangular** matrix of order n :

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a_{22} & a_{23} & \cdots & a_{2n} \\ 0 & 0 & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a_{nn} \end{pmatrix}.$$

Its characteristic polynomial is $\det(\lambda I - A)$:

$$\begin{aligned} \det(\lambda I - A) &= \begin{vmatrix} \lambda - a_{11} & -a_{12} & -a_{13} & \cdots & -a_{1n} \\ 0 & \lambda - a_{22} & -a_{23} & \cdots & -a_{2n} \\ 0 & 0 & \lambda - a_{33} & \cdots & -a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda - a_{nn} \end{vmatrix} \\ &= (\lambda - a_{11})(\lambda - a_{22})(\lambda - a_{33}) \cdots (\lambda - a_{nn}). \end{aligned}$$

17 / 96

Upper Triangular Matrices

- Theorem.** Let A be an upper (or lower) triangular matrix. Then its eigenvalues are all the diagonal entries of A .
 - More precisely, if $A = (a_{ij})_{n \times n}$ is upper triangular ($a_{ij} = 0$ if $i > j$) or lower triangular ($a_{ij} = 0$ if $i < j$),
 - then the eigenvalues of A are $a_{11}, a_{22}, \dots, a_{nn}$.

- Examples.**

$$\circ \begin{pmatrix} -1 & 3.5 & 14 \\ 0 & 5 & -26 \\ 0 & 0 & 2 \end{pmatrix}. \text{ Eigenvalues: } -1, 5 \text{ and } 2.$$

$$\circ \begin{pmatrix} -2 & 0 & 0 \\ 99 & 0 & 0 \\ 10 & -4.5 & 10 \end{pmatrix}. \text{ Eigenvalues: } -2, 0 \text{ and } 10.$$

18 / 96

Eigenspace

- Let A be a square matrix of order n .
 - Let λ be an eigenvalue of A .

Let $\mathbf{0} \neq \mathbf{v} \in \mathbb{R}^n$. Then

- \mathbf{v} is an eigenvector of A associated to λ
 - $\Leftrightarrow A\mathbf{v} = \lambda\mathbf{v}$
 - $\Leftrightarrow (\lambda I - A)\mathbf{v} = \mathbf{0}$
 - $\Leftrightarrow \mathbf{v}$ is a nonzero vector in the nullspace of $\lambda I - A$.

- Definition.** Let A be a square matrix and λ an eigenvalue of A . (Then $\lambda I - A$ is singular.)
 - The **eigenspace** of A associated to λ is the nullspace of $\lambda I - A$, denoted by E_λ (or $E_{A,\lambda}$).
 - E_λ consists of all the eigenvectors of A associated to λ , and the zero vector $\mathbf{0}$. Note that $\dim E_\lambda \geq 1$.

19 / 96

Examples

- $A = \begin{pmatrix} 0.96 & 0.01 \\ 0.04 & 0.99 \end{pmatrix}$ has eigenvalues 1 and 0.95.
 - The eigenspace E_1 is the nullspace of $1I - A$.
 - $1I - A = \begin{pmatrix} 0.04 & -0.01 \\ -0.04 & 0.01 \end{pmatrix}$.
 - $(1I - A)\mathbf{x} = \mathbf{0} \Leftrightarrow \mathbf{x} = t \begin{pmatrix} 0.25 \\ 1 \end{pmatrix}, t \in \mathbb{R}$.
- Then $E_1 = \text{span} \left\{ \begin{pmatrix} 0.25 \\ 1 \end{pmatrix} \right\}$, and $\dim(E_1) = 1$.
- The eigenspace $E_{0.95}$ is the nullspace of $0.95I - A$.
 - $0.95I - A = \begin{pmatrix} -0.01 & -0.01 \\ -0.04 & -0.04 \end{pmatrix}$.
 - $(0.95I - A)\mathbf{x} = \mathbf{0} \Leftrightarrow \mathbf{x} = t \begin{pmatrix} -1 \\ 1 \end{pmatrix}, t \in \mathbb{R}$.
- Then $E_{0.95} = \text{span} \left\{ \begin{pmatrix} -1 \\ 1 \end{pmatrix} \right\}$, and $\dim(E_{0.95}) = 1$.

20 / 96

Examples

- $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ has eigenvalues 3 and 0.

- The eigenspace E_3 is the nullspace of $3I - B$.

- $3I - B = \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}.$

- $(3I - B)x = 0 \Leftrightarrow x = t \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, t \in \mathbb{R}.$

Then $E_3 = \text{span} \left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$, and $\dim(E_3) = 1$.

21 / 96

Examples

- $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ has eigenvalues 3 and 0.

- The eigenspace E_0 is the nullspace of $0I - B$.

- $0I - B = -B = \begin{pmatrix} -1 & -1 & -1 \\ -1 & -1 & -1 \\ -1 & -1 & -1 \end{pmatrix}.$

- $(0I - B)x = 0 \Leftrightarrow x = s \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, s, t \in \mathbb{R}.$

$E_0 = \text{span} \left\{ \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right\}$, and $\dim(E_0) = 2$.

- **Note:** If A is singular, then 0 is an eigenvalue of A .

- The eigenspace E_0 is the nullspace of A .

22 / 96

Examples

- $C = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 2 \\ 1 & 1 & 1 \end{pmatrix}$ has eigenvalues 1 , $\sqrt{2}$ and $-\sqrt{2}$.

◦ The eigenspace E_1 is the nullspace of $1I - C$.

- $1I - C = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & -2 \\ -1 & -1 & 0 \end{pmatrix}.$

- $(1I - C)x = 0 \Leftrightarrow x = t \begin{pmatrix} -2 \\ 2 \\ 1 \end{pmatrix}, t \in \mathbb{R}.$

$$E_1 = \text{span} \left\{ \begin{pmatrix} -2 \\ 2 \\ 1 \end{pmatrix} \right\}, \text{ and } \dim(E_1) = 1.$$

23 / 96

Examples

- $C = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 2 \\ 1 & 1 & 1 \end{pmatrix}$ has eigenvalues 1 , $\sqrt{2}$ and $-\sqrt{2}$.

◦ The eigenspace $E_{\sqrt{2}}$ is the nullspace of $\sqrt{2}I - C$.

- $\sqrt{2}I - C = \begin{pmatrix} \sqrt{2} & 1 & 0 \\ 0 & \sqrt{2} & -2 \\ -1 & -1 & \sqrt{2} - 1 \end{pmatrix}.$

- $(\sqrt{2}I - C)x = 0 \Leftrightarrow x = t \begin{pmatrix} -1 \\ \sqrt{2} \\ 1 \end{pmatrix}, t \in \mathbb{R}.$

$$E_{\sqrt{2}} = \text{span} \left\{ \begin{pmatrix} -1 \\ \sqrt{2} \\ 1 \end{pmatrix} \right\}, \text{ and } \dim(E_{\sqrt{2}}) = 1.$$

24 / 96

Examples

- $C = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 2 \\ 1 & 1 & 1 \end{pmatrix}$ has eigenvalues 1 , $\sqrt{2}$ and $-\sqrt{2}$.
 - The eigenspace $E_{-\sqrt{2}}$ is the nullspace of $-\sqrt{2}I - C$.
 - $-\sqrt{2}I - C = \begin{pmatrix} -\sqrt{2} & 1 & 0 \\ 0 & -\sqrt{2} & -2 \\ -1 & -1 & -\sqrt{2}-1 \end{pmatrix}$.
 - $(-\sqrt{2}I - C)x = 0 \Leftrightarrow x = t \begin{pmatrix} -1 \\ -\sqrt{2} \\ 1 \end{pmatrix}, t \in \mathbb{R}$.
- $$E_{-\sqrt{2}} = \text{span} \left\{ \begin{pmatrix} -1 \\ -\sqrt{2} \\ 1 \end{pmatrix} \right\}, \text{ and } \dim(E_{-\sqrt{2}}) = 1.$$

25 / 96

Diagonalization

26 / 96

Diagonalizable Matrices

- **Definition.** Let A be a square matrix.
 - A is called **diagonalizable** if there exists an **invertible** matrix P such that $P^{-1}AP$ is a **diagonal** matrix.
 - **Examples.**
 - $A = \begin{pmatrix} 0.96 & 0.01 \\ 0.04 & 0.99 \end{pmatrix}$ and $P = \begin{pmatrix} 0.25 & -1 \\ 1 & 1 \end{pmatrix}$
 - Then $P^{-1}AP = D = \begin{pmatrix} 1 & 0 \\ 0 & 0.95 \end{pmatrix}$.
- Then A is diagonalizable.
- Note that the diagonal entries of D are the eigenvalues of A .
 - The columns of P are eigenvectors of A associated to these eigenvalues.

27 / 96

Diagonalizable Matrices

- **Definition.** Let A be a square matrix.
 - A is called **diagonalizable** if there exists an **invertible** matrix P such that $P^{-1}AP$ is a **diagonal** matrix.
- **Examples.**
 - $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ and $P = \begin{pmatrix} 1 & -1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$.
 - $P^{-1}BP = D = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$. So B is diagonalizable.
 - Note that the diagonal entries of D are the eigenvalues of B .
 - The columns of P are eigenvectors of B associated to these eigenvalues.

28 / 96

Examples

- Prove that $M = \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix}$ is not diagonalizable.
 - Suppose there exists invertible $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ such that
 - $\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix}$.
 i.e., $\begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix}$.
 - $\begin{pmatrix} 2a & 2b \\ a + 2c & b + 2d \end{pmatrix} = \begin{pmatrix} \lambda a & \mu b \\ \lambda c & \mu d \end{pmatrix}$.
 If $a \neq 0$, then $\lambda = 2$, and $a + 2c = 2c \Rightarrow a = 0$; so $a = 0$.
 If $b \neq 0$, then $\mu = 2$, and $b + 2d = 2d \Rightarrow b = 0$; so $b = 0$.
 - Then $\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ c & d \end{pmatrix}$ is singular.
 - Therefore, M is not diagonalizable.

29 / 96

Criterion of Diagonalizability

- Let A be a square matrix of order n .
 - Suppose that A is diagonalizable.
 - There exist invertible matrices P such that $P^{-1}AP$ is a diagonal matrix D , i.e., $AP = PD$.

Let $P = (v_1 \ \cdots \ v_n)$ and $D = \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix}$.

- $A(v_1 \ \cdots \ v_n) = (v_1 \ \cdots \ v_n) \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix}$.

- $(Av_1 \ \cdots \ Av_n) = (\lambda_1 v_1 \ \cdots \ \lambda_n v_n)$.

- Then $Av_i = \lambda_i v_i$, $i = 1, \dots, n$.

λ_i is an eigenvalue of A , v_i is an eigenvector associated to λ_i .

- P is invertible $\Rightarrow v_1, \dots, v_n$ are linearly independent.

30 / 96

Criterion of Diagonalizability

- Let A be a square matrix of order n .
 - Suppose A has n linearly independent eigenvectors.
 - $Av_1 = \lambda_1 v_1, Av_2 = \lambda_2 v_2, \dots, Av_n = \lambda_n v_n$,
 - where v_1, \dots, v_n are linearly independent.

Let $P = (v_1 \ \cdots \ v_n)$. Then P is invertible.

- Let $D = \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix}$.

$$\begin{aligned} AP &= A(v_1 \ \cdots \ v_n) = (Av_1 \ \cdots \ Av_n) \\ &= (\lambda_1 v_1 \ \cdots \ \lambda_n v_n) \\ &= (v_1 \ \cdots \ v_n) \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix} = PD. \end{aligned}$$

- $P^{-1}AP = D$; so A is diagonalizable.

31 / 96

Criterion of Diagonalizability

- **Theorem.** Let A be a square matrix of order n .
 - A is diagonalizable
 - $\Leftrightarrow A$ has n linearly independent eigenvectors.
- **Remark.** Suppose that $P^{-1}AP = D$ is diagonal.
 - The diagonal entries of D are eigenvalues of A :
 - $\lambda_1, \dots, \lambda_n$, which may be repeated.

D is not unique unless A has only one eigenvalue.
 - The columns of P are eigenvectors of A :
 - v_1, \dots, v_n , which are linearly independent.
 - v_i is an eigenvector of A associated to λ_i .

P is not unique. For instance,

 - v_i can be replaced by a nonzero multiple of v_i .

32 / 96

Diagonalization

- **Algorithm of Diagonalization**
 - Let A be a square matrix of order n .
 1. Solve $\det(\lambda I - A) = 0$ to find eigenvalues of A .
 2. For each eigenvalue λ_i of A ,
 - find a basis S_i for the eigenspace E_{λ_i} .

A is diagonalizable $\Leftrightarrow |S_1| + \dots + |S_k| = n$,
 A is not diagonalizable $\Leftrightarrow |S_1| + \dots + |S_k| < n$.

Suppose A is diagonalizable. Then

 - $S_1 \cup \dots \cup S_k = \{v_1, \dots, v_n\}$ is a basis for \mathbb{R}^n .
 - A is diagonalized by $P = (v_1 \ \dots \ v_n)$.

33 / 96

Remarks

- $\det(\lambda \mathbf{I} - \mathbf{A})$ is a polynomial of λ in degree n .
 - It can be completely factorized as
 - $(\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n)$, $\lambda_i \in \mathbb{C}$.
 But $\lambda_1, \dots, \lambda_n$ are not necessarily real numbers.
 - If some λ_i is not real,
 - then \mathbf{A} is not diagonalizable (over \mathbb{R}).
- **Example.** Let $\mathbf{A} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.
 - $\det(\lambda \mathbf{I} - \mathbf{A}) = \lambda^2 + 1 = (\lambda - i)(\lambda + i)$.
 - \mathbf{A} is not diagonalizable over \mathbb{R} .
 - $\begin{pmatrix} -i & i \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} -i & i \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$.

34 / 96

Remarks

- Suppose that $\det(\lambda \mathbf{I} - \mathbf{A})$ can be completely factorized:
 - $(\lambda - \lambda_1)^{r_1} (\lambda - \lambda_2)^{r_2} \cdots (\lambda - \lambda_k)^{r_k}$,
 - where $\lambda_1, \lambda_2, \dots, \lambda_k$ are all distinct.
 Then r_i is the **algebraic multiplicity** $a(\lambda_i)$ of λ_i .
 - Let E_i be the eigenspace of \mathbf{A} associated to λ_i .
 - $\dim E_i$ is the **geometric multiplicity** $g(\lambda_i)$ of λ_i .
 - One can prove (MA2101) that $g(\lambda_i) \leq a(\lambda_i)$.
 Note that $a(\lambda_1) + a(\lambda_2) + \cdots + a(\lambda_k) = n$.
 - If $\dim E_i < a(\lambda_i)$ for some i ,
 - then $\dim E_1 + \dim E_2 + \cdots + \dim E_k < n$;
 consequently, \mathbf{A} is not diagonalizable.

35 / 96

Remarks

- Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be distinct eigenvalues of A .
 - and v_i be an eigenvector of A associated to λ_i .

Then v_1, v_2, \dots, v_k are linearly independent.

- **Proof.** Let $k = 2$. Suppose $c_1 v_1 + c_2 v_2 = 0$.

$$\begin{aligned} 0 &= A0 = A(c_1 v_1 + c_2 v_2) \\ &= c_1 (A v_1) + c_2 (A v_2) \\ &= (c_1 \lambda_1) v_1 + (c_2 \lambda_2) v_2, \\ 0 &= \lambda_1 0 = \lambda_1 (c_1 v_1 + c_2 v_2) \\ &= (c_1 \lambda_1) v_1 + (c_2 \lambda_1) v_2. \end{aligned}$$

- Then $c_2 \lambda_2 v_2 = c_2 \lambda_1 v_2$, i.e., $c_2 (\lambda_2 - \lambda_1) v_2 = 0$.

- $v_2 \neq 0, \lambda_1 \neq \lambda_2$; so $c_2 = 0$ & $c_1 = 0$.

The general case can be proved by mathematical induction. (Exercise.)

36 / 96

Diagonalization

- **Algorithm of Diagonalization**

- Let A be a square matrix of order n .

Case 1. If $\det(\lambda I - A)$ cannot be completely factorized,

- then A is not diagonalizable.

Case 2. If $\det(\lambda I - A)$ can be completely factorized,

- for each λ_i , find a basis S_i for its eigenspace.

2a. If $|S_i| < a(\lambda_i)$ for some i ,

- then A is not diagonalizable.

2b. If $|S_i| = a(\lambda_i)$ for all i ,

- then A is diagonalizable.
- $S_1 \cup \dots \cup S_k = \{v_1, \dots, v_n\}$ is a basis for \mathbb{R}^n .
- $P = (v_1 \ \dots \ v_n)$ diagonalizes A .

37 / 96

Examples

- Let $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$.

Step 1. $\det(\lambda I - B) = (\lambda - 3)\lambda^2$.

- B has eigenvalues $\lambda = 3$ and $\lambda = 0$.

Step 2. Find bases for eigenspaces:

- E_3 : $\{(1, 1, 1)^T\}$.
- E_0 : $\{(-1, 1, 0)^T, (-1, 0, 1)^T\}$.

Step 3. $P = \begin{pmatrix} 1 & -1 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$. Then $P^{-1}BP = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$.

38 / 96

Examples

- Let $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$.

Step 1. $\det(\lambda I - B) = (\lambda - 3)\lambda^2$.

- B has eigenvalues $\lambda = 3$ and $\lambda = 0$.

Step 2. Find bases for eigenspaces:

- E_3 : $\{(1, 1, 1)^T\}$.
- E_0 : $\{(-1, 1, 0)^T, (-1, 0, 1)^T\}$.

Step 3. $P = \begin{pmatrix} -1 & 1 & -1 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}$. Then $P^{-1}BP = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 0 \end{pmatrix}$.

- The i th column of P is an eigenvector of B associated to the i th diagonal entry (eigenvalue) of $P^{-1}BP$.

39 / 96

Examples

- Let $C = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 2 \\ 1 & 1 & 1 \end{pmatrix}$.

Step 1. $\det(\lambda I - C) = (\lambda - 1)(\lambda - \sqrt{2})(\lambda + \sqrt{2})$.

- C has eigenvalues $\lambda = 1, \sqrt{2}$ and $-\sqrt{2}$.

Step 2. Find bases for eigenspaces:

- E_1 : $\{(-2, 2, 1)^T\}$.
- $E_{\sqrt{2}}$: $\{(-1, \sqrt{2}, 1)^T\}$.
- $E_{-\sqrt{2}}$: $\{(-1, -\sqrt{2}, 1)^T\}$.

Step 3. $P = \begin{pmatrix} -2 & -1 & -1 \\ 2 & \sqrt{2} & -\sqrt{2} \\ 1 & 1 & 1 \end{pmatrix}, P^{-1}CP = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sqrt{2} & 0 \\ 0 & 0 & -\sqrt{2} \end{pmatrix}$.

40 / 96

Examples

- Let $A = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 2 & 0 \\ -3 & 5 & 2 \end{pmatrix}$.

Step 1. $\det(\lambda I - A) = (\lambda - 1)(\lambda - 2)^2$.

- A has eigenvalues $\lambda = 1$ and 2 .

Step 2. Find bases for eigenspaces:

- $1I - A = \begin{pmatrix} 0 & 0 & 0 \\ -1 & -1 & 0 \\ 3 & -5 & -1 \end{pmatrix}$.
- $(1I - A)x = 0 \Leftrightarrow x = t \begin{pmatrix} 1 \\ -1 \\ 8 \end{pmatrix}, t \in \mathbb{R}$.
- $E_1 = \text{span} \left\{ \begin{pmatrix} 1 \\ -1 \\ 8 \end{pmatrix} \right\}$.

41 / 96

Examples

- Let $\mathbf{A} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 2 & 0 \\ -3 & 5 & 2 \end{pmatrix}$.

Step 1. $\det(\lambda \mathbf{I} - \mathbf{A}) = (\lambda - 1)(\lambda - 2)^2$.

- \mathbf{A} has eigenvalues $\lambda = 1$ and 2 .

Step 2. Find bases for eigenspaces:

- $2\mathbf{I} - \mathbf{A} = \begin{pmatrix} 1 & 0 & 0 \\ -1 & 0 & 0 \\ 3 & -5 & 0 \end{pmatrix}$.

- $(2\mathbf{I} - \mathbf{A})\mathbf{x} = \mathbf{0} \Leftrightarrow \mathbf{x} = t \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, t \in \mathbb{R}$.

- $E_2 = \text{span} \left\{ \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$.

42 / 96

Examples

- Let $\mathbf{A} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 2 & 0 \\ -3 & 5 & 2 \end{pmatrix}$.

Step 1. $\det(\lambda \mathbf{I} - \mathbf{A}) = (\lambda - 1)(\lambda - 2)^2$.

- \mathbf{A} has eigenvalues $\lambda = 1$ and 2 .

Step 2. Find bases for eigenspaces:

- $E_1 = \text{span} \left\{ \begin{pmatrix} 1 \\ -1 \\ 8 \end{pmatrix} \right\}$,

- $E_2 = \text{span} \left\{ \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$.

Step 3. Since there are only two linearly independent eigenvectors, \mathbf{A} is not diagonalizable.

43 / 96

Examples

- Let $A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$.

Step 1. $\det(\lambda I - A) = \lambda^2 - \lambda - 1$.

- A has eigenvalues $\frac{1+\sqrt{5}}{2}$ and $\frac{1-\sqrt{5}}{2}$.

Step 2. Find eigenspaces:

- $\left(\frac{1+\sqrt{5}}{2}I - A\right)x = 0 \Leftrightarrow x = t \begin{pmatrix} 1 \\ \frac{1+\sqrt{5}}{2} \end{pmatrix}, t \in \mathbb{R}$.

- $E_{\frac{1+\sqrt{5}}{2}} = \text{span} \left\{ \begin{pmatrix} 1 \\ \frac{1+\sqrt{5}}{2} \end{pmatrix} \right\}$.

- $\left(\frac{1-\sqrt{5}}{2}I - A\right)x = 0 \Leftrightarrow x = t \begin{pmatrix} 1 \\ \frac{1-\sqrt{5}}{2} \end{pmatrix}, t \in \mathbb{R}$.

- $E_{\frac{1-\sqrt{5}}{2}} = \text{span} \left\{ \begin{pmatrix} 1 \\ \frac{1-\sqrt{5}}{2} \end{pmatrix} \right\}$.

44 / 96

Examples

- Let $A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$.

Step 1. $\det(\lambda I - A) = \lambda^2 - \lambda - 1$.

- A has eigenvalues $\frac{1+\sqrt{5}}{2}$ and $\frac{1-\sqrt{5}}{2}$.

Step 2. Find eigenspaces:

- $E_{\frac{1+\sqrt{5}}{2}} = \text{span} \left\{ \begin{pmatrix} 1 \\ \frac{1+\sqrt{5}}{2} \end{pmatrix} \right\}$.

- $E_{\frac{1-\sqrt{5}}{2}} = \text{span} \left\{ \begin{pmatrix} 1 \\ \frac{1-\sqrt{5}}{2} \end{pmatrix} \right\}$.

Step 3. $P = \begin{pmatrix} 1 & 1 \\ \frac{1+\sqrt{5}}{2} & \frac{1-\sqrt{5}}{2} \end{pmatrix}$. $P^{-1}AP = \begin{pmatrix} \frac{1+\sqrt{5}}{2} & 0 \\ 0 & \frac{1-\sqrt{5}}{2} \end{pmatrix}$.

45 / 96

Examples

- **Theorem.** Let A be a square matrix of order n .

- If A has n distinct eigenvalues,
 - then A is diagonalizable.

Proof. Suppose A has distinct eigenvalues $\lambda_1, \dots, \lambda_n$.

- Let v_i be an eigenvector of A associated to λ_i .
- It is known that v_1, \dots, v_n are linearly independent.

Therefore, A is diagonalizable.

- **Example.** Let $A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 2 & 2 & 2 \\ 0 & 0 & 3 & 3 \\ 0 & 0 & 0 & 4 \end{pmatrix}$.

- A has eigenvalues 1, 2, 3, 4; so A is diagonalizable.
- Can you diagonalize it? (Exercise.)

46 / 96

Application

- Suppose that A is diagonalizable.
 - There exists an invertible matrix P such that

- $P^{-1}AP = \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix}$ is diagonal.

- $A = P \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix} P^{-1}$.

- Let m be a nonnegative integer. Then

- $A^m = P \begin{pmatrix} \lambda_1^m & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n^m \end{pmatrix} P^{-1}$.

47 / 96

Application

- Suppose that A is diagonalizable.
 - There exists an invertible matrix P such that
 - $P^{-1}AP = \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix}$ is diagonal.
 - $A = P \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix} P^{-1}$.
 - Suppose that A is invertible. Then for any integer m ,
 - $A^m = P \begin{pmatrix} \lambda_1^m & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n^m \end{pmatrix} P^{-1}$.

48 / 96

Examples

- Let $A = \begin{pmatrix} -4 & 0 & -6 \\ 2 & 1 & 2 \\ 3 & 0 & 5 \end{pmatrix}$.
 - $\det(\lambda I - A) = (\lambda + 1)(\lambda - 1)(\lambda - 2)$.
 - $(-1I - A)x = 0 \Leftrightarrow x = t \begin{pmatrix} -2 \\ 1 \\ 1 \end{pmatrix}, t \in \mathbb{R}$.
 - $(1I - A)x = 0 \Leftrightarrow x = t \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, t \in \mathbb{R}$.
 - $(2I - A)x = 0 \Leftrightarrow x = t \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, t \in \mathbb{R}$.
 - $P = \begin{pmatrix} -2 & 0 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$. $P^{-1}AP = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$.

49 / 96

Examples

- Let $A = \begin{pmatrix} -4 & 0 & -6 \\ 2 & 1 & 2 \\ 3 & 0 & 5 \end{pmatrix}$.

- $P = \begin{pmatrix} -2 & 0 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$. $P^{-1}AP = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$.

- $A^m = P \begin{pmatrix} (-1)^m & 0 & 0 \\ 0 & 1^m & 0 \\ 0 & 0 & 2^m \end{pmatrix} P^{-1}$.

$$A^{10} = P \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1024 \end{pmatrix} P^{-1}$$

$$= \dots = \begin{pmatrix} -1022 & 0 & -2046 \\ 0 & 1 & 0 \\ 1023 & 0 & 2047 \end{pmatrix}.$$

50 / 96

Examples

- The **Fibonacci numbers** a_n are defined by
 - $a_0 = 0, a_1 = 1$ and $a_n = a_{n-1} + a_{n-2}$ for $n \geq 2$.

Note that $a_{n+1} = a_{n-1} + a_n$ for $n \geq 1$.

- $\begin{pmatrix} a_n \\ a_{n+1} \end{pmatrix} = \begin{pmatrix} a_n \\ a_{n-1} + a_n \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} a_{n-1} \\ a_n \end{pmatrix}$.

Let $x_n = \begin{pmatrix} a_n \\ a_{n+1} \end{pmatrix}$ and $A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$.

- $x_n = Ax_{n-1} = A^2x_{n-2} = \dots = A^n x_0, x_0 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

We have diagonalized $A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$.

- $P = \begin{pmatrix} 1 & 1 \\ \frac{1+\sqrt{5}}{2} & \frac{1-\sqrt{5}}{2} \end{pmatrix}$. $P^{-1}AP = \begin{pmatrix} \frac{1+\sqrt{5}}{2} & 0 \\ 0 & \frac{1-\sqrt{5}}{2} \end{pmatrix}$.

51 / 96

Examples

- The **Fibonacci numbers** F_n are defined by
 - $a_0 = 0, a_1 = 1$ and $a_n = a_{n-1} + a_{n-2}$ for $n \geq 2$.

Let $x_n = \begin{pmatrix} a_n \\ a_{n+1} \end{pmatrix}$ and $A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$.

$$\begin{aligned} x_n &= A^n x_0 = P \begin{pmatrix} \frac{1+\sqrt{5}}{2} & 0 \\ 0 & \frac{1-\sqrt{5}}{2} \end{pmatrix}^n P^{-1} x_0 \\ &= P \begin{pmatrix} \left(\frac{1+\sqrt{5}}{2}\right)^n & 0 \\ 0 & \left(\frac{1-\sqrt{5}}{2}\right)^n \end{pmatrix} P^{-1} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\ &= \begin{pmatrix} \frac{1}{\sqrt{5}} \left(\frac{1+\sqrt{5}}{2}\right)^n - \frac{1}{\sqrt{5}} \left(\frac{1-\sqrt{5}}{2}\right)^n \\ \frac{1}{\sqrt{5}} \left(\frac{1+\sqrt{5}}{2}\right)^{n+1} - \frac{1}{\sqrt{5}} \left(\frac{1-\sqrt{5}}{2}\right)^{n+1} \end{pmatrix} \end{aligned}$$

Therefore, $a_n = \frac{1}{\sqrt{5}} \left(\frac{1+\sqrt{5}}{2}\right)^n - \frac{1}{\sqrt{5}} \left(\frac{1-\sqrt{5}}{2}\right)^n$.

52 / 96

Orthogonal Diagonalization

53 / 96

Introduction

- Recall that an $n \times n$ matrix A is **diagonalizable**
 - $\Leftrightarrow A$ has n **linearly independent eigenvectors**

v_1, \dots, v_n (associated to $\lambda_1, \dots, \lambda_n$).

Then $P^{-1}AP = D$, where

- $P = (v_1 \ v_2 \ \dots \ v_n), D = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix}$

- In order to find P^{-1} , we may need:
 - Gauss-Jordan elimination: $(P \mid I) \dashrightarrow (I \mid P^{-1})$.
 - Adjoint matrix: $P^{-1} = \frac{1}{\det(P)} \text{adj}(P)$.
- Note:** If P is **orthogonal**, then $P^{-1} = P^T$.

54 / 96

Introduction

- Let $B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$. Then it can be diagonalized by
 - $P = \begin{pmatrix} 1 & -1 & -1 \\ 1 & 1 & -1 \\ 1 & 0 & 2 \end{pmatrix}$.

We can verify that the columns of P , which are eigenvectors of B , form an **orthogonal** basis for \mathbb{R}^3 .

- Normalizing:

- $R = \begin{pmatrix} \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & 0 & \frac{2}{\sqrt{6}} \end{pmatrix}$.

- R is an orthogonal matrix, which also diagonalizes B .

55 / 96

Definition

- **Definition.** A square matrix A is called **orthogonally diagonalizable** if it can be diagonalized by an **orthogonal** matrix. That is,
 - there exists an **orthogonal** matrix P such that
 - $P^T A P (= P^{-1} A P)$ is a **diagonal** matrix.

P is said to **orthogonally diagonalize** A .

- **Remarks.** For any eigenvalue λ of A , we can always choose an orthonormal basis for the associated eigenspace E_λ .

Suppose further that A is **orthogonally diagonalizable**.

- Then A is diagonalizable, and A has n linearly independent eigenvectors.
- For distinct eigenvalues $\lambda \neq \mu$,
 - Every eigenvector of λ is orthogonal to that of μ .

56 / 96

Classification

- **Theorem.** A square matrix is **orthogonally diagonalizable**

\Leftrightarrow it is a **symmetric** matrix.

- **Proof.** (\Rightarrow) Suppose A is orthogonally diagonalizable.

- There is an orthogonal matrix P & a diagonal matrix D

- such that $D = P^T A P$.

Since D is diagonal, it is also symmetric.

- $D = D^T = (P^T A P)^T = P^T A^T P$.

Therefore, $P^T A P = P^T A^T P$.

- Note that both P and P^T are invertible.

- By Cancellation Law: $A = A^T$.

That is, A is symmetric.

(\Leftarrow) is left in MA2101 Linear Algebra II.

57 / 96

Algorithm

- **Algorithm.** (Orthogonally diagonalize symmetric matrix).

Let A be a **symmetric** matrix of order n .

1. Find all distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_k$.
2. For each eigenvalue λ_i , find an **orthonormal** basis for the eigenspace E_{λ_i} .
 - (i) Find a basis S_{λ_i} for E_{λ_i} .
 - (ii) Use Gram-Schmidt process to transfer S_{λ_i} to an orthonormal basis T_{λ_i} for E_{λ_i} .
3. Let $T = T_{\lambda_1} \cup T_{\lambda_2} \cup \dots \cup T_{\lambda_k}$,
 - $T = \{v_1, \dots, v_n\}$ is an orthonormal basis for \mathbb{R}^n . $P = (v_1 \ \dots \ v_n)$ orthogonally diagonalizes A .

58 / 96

Algorithm

- Compare with the **algorithm for diagonalization**:

Let A be a square matrix of order n .

1. Find all distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_k$.
2. For each eigenvalue λ_i , find a basis for the eigenspace E_{λ_i} .
3. Let $S = S_{\lambda_1} \cup S_{\lambda_2} \cup \dots \cup S_{\lambda_k}$.
 - (i) If $|S| < n$, then A is not diagonalizable.
 - (ii) If $|S| = n$, say $S = \{v_1, v_2, \dots, v_n\}$,
 - $P = (v_1 \ v_2 \ \dots \ v_n)$ diagonalizes A .

59 / 96

Algorithm

- **Remarks.** Let A be a **symmetric** matrix of order n .

1. Every eigenvalue of A is a real number.
2. Write the characteristic polynomial
 - $\det(\lambda I - A) = (\lambda - \lambda_1)^{r_1} \dots (\lambda - \lambda_k)^{r_k}$,
 - where $\lambda_1, \dots, \lambda_k$ are distinct eigenvalues.

Then $\dim E_{\lambda_1} = r_1, \dots, \dim E_{\lambda_k} = r_k$.

$\therefore \dim E_{\lambda_1} + \dots + \dim E_{\lambda_k} = r_1 + \dots + r_k = n$.

3. If each basis S_{λ_i} for E_{λ_i} is orthonormal, then
 - $T = S_{\lambda_1} \cup \dots \cup S_{\lambda_k} = \{v_1, \dots, v_n\}$ is an orthonormal set. (Exercise.)
 - $P = (v_1 \ \dots \ v_n)$ is an orthogonal matrix.

60 / 96

Examples

- Let $\mathbf{A} = \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix}$.
 1. Find eigenvalues: For 2×2 matrix,
 - $\det(\lambda \mathbf{I} - \mathbf{A}) = \lambda^2 - \text{tr}(\mathbf{A})\lambda + \det(\mathbf{A})$.
 - $\lambda^2 - 2\lambda + \frac{3}{4} = (\lambda - \frac{1}{2})(\lambda - \frac{3}{2})$.
 - $\therefore \lambda = \frac{1}{2}$ and $\lambda = \frac{3}{2}$.
 2. Find eigenvectors. For $\lambda = \frac{1}{2}$,
 - Solve $(\lambda \mathbf{I} - \mathbf{A})\mathbf{x} = \mathbf{0}$:
 - $\begin{pmatrix} -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$.
 - $\begin{pmatrix} x \\ y \end{pmatrix} = t \begin{pmatrix} 1 \\ 1 \end{pmatrix} \xrightarrow{\text{normalizing}} \mathbf{v}_1 = \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix}$.

61 / 96

Examples

- Let $\mathbf{A} = \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix}$.
 1. Find eigenvalues: For 2×2 matrix,
 - $\det(\lambda \mathbf{I} - \mathbf{A}) = \lambda^2 - \text{tr}(\mathbf{A})\lambda + \det(\mathbf{A})$.
 - $\lambda^2 - 2\lambda + \frac{3}{4} = (\lambda - \frac{1}{2})(\lambda - \frac{3}{2})$.
 - $\therefore \lambda = \frac{1}{2}$ and $\lambda = \frac{3}{2}$.
 2. Find eigenvectors. For $\lambda = \frac{3}{2}$,
 - Solve $(\lambda \mathbf{I} - \mathbf{A})\mathbf{x} = \mathbf{0}$:
 - $\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$.
 - $\begin{pmatrix} x \\ y \end{pmatrix} = t \begin{pmatrix} -1 \\ 1 \end{pmatrix} \xrightarrow{\text{normalizing}} \mathbf{v}_2 = \begin{pmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix}$.

62 / 96

Examples

- Let $\mathbf{A} = \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix}$.
 1. Find eigenvalues: For 2×2 matrix,
 - $\det(\lambda \mathbf{I} - \mathbf{A}) = \lambda^2 - \text{tr}(\mathbf{A})\lambda + \det(\mathbf{A})$.
 - $\lambda^2 - 2\lambda + \frac{3}{4} = (\lambda - \frac{1}{2})(\lambda - \frac{3}{2})$.
 - $\therefore \lambda = \frac{1}{2}$ and $\lambda = \frac{3}{2}$.
 3. Let $\mathbf{P} = (\mathbf{v}_1 \ \mathbf{v}_2) = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$. Then
 - $\mathbf{P}^T \mathbf{A} \mathbf{P} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{3}{2} \end{pmatrix}$.

63 / 96

Examples

- Let $\mathbf{B} = \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & 3 & 1 \\ -1 & 1 & 1 & 3 \end{pmatrix}$.
 1. Find the eigenvalues. The characteristic polynomial

$$\begin{aligned} & \det(\lambda \mathbf{I} - \mathbf{B}) \\ &= \det \begin{pmatrix} \lambda - 1 & 1 & -1 & 1 \\ 1 & \lambda - 1 & 1 & -1 \\ -1 & 1 & \lambda - 3 & -1 \\ 1 & -1 & -1 & \lambda - 3 \end{pmatrix} \\ &= \dots\dots\dots \\ &= \lambda^4 - 8\lambda^3 + 16\lambda^2 = \lambda^2(\lambda - 4)^2. \end{aligned}$$
 - The eigenvalues are $\lambda = 0$ and $\lambda = 4$.

64 / 96

Examples

- Let $B = \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & 3 & 1 \\ -1 & 1 & 1 & 3 \end{pmatrix}$.

2. Find the eigenvectors. Let $\lambda = 0$. Solve

- $(\lambda I - B)x = 0$.

$$\begin{pmatrix} -1 & 1 & -1 & 1 \\ 1 & -1 & 1 & -1 \\ -1 & 1 & -3 & -1 \\ 1 & -1 & -1 & -3 \end{pmatrix} \begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

- $\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = s \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + t \begin{pmatrix} 2 \\ 0 \\ -1 \\ 1 \end{pmatrix}, s, t \in \mathbb{R}.$

65 / 96

Examples

- Let $B = \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & 3 & 1 \\ -1 & 1 & 1 & 3 \end{pmatrix}$.

2. Find the eigenvectors. Let $\lambda = 0$. Set

- $u_1 = (1, 1, 0, 0)$ and $u_2 = (2, 0, -1, 1)$.

$$v_1 = u_1 = (1, 1, 0, 0)$$

$$v_2 = u_2 - \frac{u_2 \cdot v_1}{v_1 \cdot v_2} v_1 = (1, -1, -1, 1).$$

- Normalizing:

$$w_1 = \frac{v_1}{\|v_1\|} = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}, 0, 0\right)$$

$$w_2 = \frac{v_2}{\|v_2\|} = \left(\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, \frac{1}{2}\right).$$

66 / 96

Examples

- Let $B = \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & 3 & 1 \\ -1 & 1 & 1 & 3 \end{pmatrix}$.

2. Find the eigenvectors. Let $\lambda = 4$. Solve

- $(\lambda I - B)x = 0$.

$$\begin{pmatrix} 3 & 1 & -1 & 1 \\ 1 & 3 & 1 & -1 \\ -1 & 1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

- $\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = s \begin{pmatrix} \frac{1}{2} \\ -\frac{1}{2} \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ 0 \\ 1 \end{pmatrix}, s, t \in \mathbb{R}.$

67 / 96

Examples

- Let $B = \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & 3 & 1 \\ -1 & 1 & 1 & 3 \end{pmatrix}$.

2. Find the eigenvectors. Let $\lambda = 4$. Set

- $\mathbf{u}_3 = (\frac{1}{2}, -\frac{1}{2}, 1, 0)$ and $\mathbf{u}_4 = (-\frac{1}{2}, \frac{1}{2}, 0, 1)$.

$$\mathbf{v}_3 = \mathbf{u}_3 = (\frac{1}{2}, -\frac{1}{2}, 1, 0)$$

$$\mathbf{v}_4 = \mathbf{u}_4 - \frac{\mathbf{u}_4 \cdot \mathbf{v}_3}{\mathbf{v}_3 \cdot \mathbf{v}_3} \mathbf{v}_3 = (-\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 1).$$

- Normalizing:

$$\mathbf{w}_3 = \frac{\mathbf{v}_3}{\|\mathbf{v}_3\|} = (\frac{1}{\sqrt{6}}, -\frac{1}{\sqrt{6}}, \frac{2}{\sqrt{6}}, 0)$$

$$\mathbf{w}_4 = \frac{\mathbf{v}_4}{\|\mathbf{v}_4\|} = (-\frac{1}{\sqrt{12}}, \frac{1}{\sqrt{12}}, \frac{1}{\sqrt{12}}, \frac{3}{\sqrt{12}}).$$

68 / 96

Examples

- Let $B = \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & 3 & 1 \\ -1 & 1 & 1 & 3 \end{pmatrix}$.

3. Let $P = (w_1 \ w_2 \ w_3 \ w_4)$.

- $P = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{2} & \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{12}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{2} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{12}} \\ 0 & -\frac{1}{2} & \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{12}} \\ 0 & \frac{1}{2} & 0 & \frac{3}{\sqrt{12}} \end{pmatrix}$.

Then $P^T A P = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 4 \end{pmatrix}$ is diagonal.

69 / 96

Quadratic Forms and Conic Sections

70 / 96

Quadratic Form

- A **homogeneous** polynomial in degree 2 in variables x, y :
 - $f(x, y) = ax^2 + bxy + cy^2$, a, b, c are real constants.

It is known as a **quadratic form** in variables x, y .

- Definition.** A **quadratic form** in n variables x_1, \dots, x_n is

- $Q(x_1, \dots, x_n) = \sum_{i=1}^n q_{ii}x_i^2 + \sum_{i < j} q_{ij}x_i x_j$.

- Examples.**

- $Q(x, y) = x^2 + y^2 - xy$.
- $Q(x, y, z) = x^2 + 2y^2 + 3z^2 + 4xy + 5xz + 6yz$.
- $Q(x_1, \dots, x_n) = x_1^2 + x_2^2 + \dots + x_n^2$.

71 / 96

Quadratic Form

- Let $\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n$, $\mathbf{A} = (a_{ij})_{n \times n}$ a symmetric matrix.
 - $\mathbf{Ax} = \begin{pmatrix} a_{11}x_1 + \cdots + a_{1n}x_n \\ \vdots \\ a_{n1}x_1 + \cdots + a_{nn}x_n \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^n a_{1j}x_j \\ \vdots \\ \sum_{j=1}^n a_{nj}x_j \end{pmatrix}.$

$$\begin{aligned} \mathbf{x}^T \mathbf{Ax} &= (x_1, \dots, x_n) \begin{pmatrix} \sum_{j=1}^n a_{1j}x_j \\ \vdots \\ \sum_{j=1}^n a_{nj}x_j \end{pmatrix} \\ &= \sum_{i=1}^n x_i \left(\sum_{j=1}^n a_{ij}x_j \right) \\ &= \sum_{i=1}^n \sum_{j=1}^n a_{ij}x_i x_j \\ &= \sum_{i=1}^n a_{ii}x_i^2 + \sum_{i < j} 2a_{ij}x_i x_j. \end{aligned}$$

72 / 96

Quadratic Form

- $Q(x_1, \dots, x_n) = \sum_{i=1}^n q_{ii}x_i^2 + \sum_{i < j} q_{ij}x_i x_j$ is a quadratic form.
 - Let $\mathbf{x} = (x_1, \dots, x_n)^T$ and $\mathbf{A} = (a_{ij})_{n \times n}$ be defined by
 - $a_{ii} = q_{ii}$ and $a_{ij} = a_{ji} = \frac{1}{2}q_{ij}$ for $i < j$.
 Then $Q(\mathbf{x}) = \mathbf{x}^T \mathbf{Ax}$, $\mathbf{x} \in \mathbb{R}^n$.
- Examples.**
 - $Q(x, y) = 2x^2 + 3y^2$ is a quadratic form in x and y .
 - Let $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$ and $\mathbf{A} = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$.
 - Then $Q(x, y) = \mathbf{x}^T \mathbf{Ax}$.
 - $Q(x, y) = x^2 + y^2 - xy$ is a quadratic form in x and y .
 - Let $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$ and $\mathbf{A} = \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix}$.
 - Then $Q(x, y) = \mathbf{x}^T \mathbf{Ax}$.

73 / 96

Quadratic Form

- $Q(x_1, \dots, x_n) = \sum_{i=1}^n q_{ii}x_i^2 + \sum_{i < j} q_{ij}x_i x_j$ is a quadratic form.
 - Let $\mathbf{x} = (x_1, \dots, x_n)^T$ and $\mathbf{A} = (a_{ij})_{n \times n}$ be defined by
 - $a_{ii} = q_{ii}$ and $a_{ij} = a_{ji} = \frac{1}{2}q_{ij}$ for $i < j$.
 Then $Q(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$, $\mathbf{x} \in \mathbb{R}^n$.
- **Examples.**
 - $Q(x, y, z) = x^2 + 2y^2 + 3z^2 + 4xy + 5xz + 6yz$.
 - It is a quadratic form in variables x, y, z .
 - Let $\mathbf{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ and $\mathbf{A} = \begin{pmatrix} 1 & 2 & \frac{5}{2} \\ 2 & 2 & 3 \\ \frac{5}{2} & 3 & 3 \end{pmatrix}$.
 - Then $Q(x, y, z) = \mathbf{x}^T \mathbf{A} \mathbf{x}$.

74 / 96

Simplification

- Suppose the quadratic form is presented as
 - $Q(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$, $\mathbf{x} = (x_1, \dots, x_n)^T \in \mathbb{R}^n$,
where \mathbf{A} is a **symmetric** matrix of order n .
- Recall that \mathbf{A} is **orthogonally diagonalizable**.
 - There exists an orthogonal matrix \mathbf{P} such that

$$\bullet \quad \mathbf{P}^T \mathbf{A} \mathbf{P} = \mathbf{D} = \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix}.$$

Let $\mathbf{y} = \mathbf{P}^T \mathbf{x} = (y_1, \dots, y_n)^T \in \mathbb{R}^n$. Then $\mathbf{x} = \mathbf{P} \mathbf{y}$.

$$\begin{aligned} Q(\mathbf{x}) &= (\mathbf{P} \mathbf{y})^T \mathbf{A} (\mathbf{P} \mathbf{y}) = \mathbf{y}^T (\mathbf{P}^T \mathbf{A} \mathbf{P}) \mathbf{y} \\ &= (y_1 \quad \cdots \quad y_n) \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix} \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} \\ &= \lambda_1 y_1^2 + \cdots + \lambda_n y_n^2. \end{aligned}$$

75 / 96

Examples

- Let $Q(x, y) = x^2 - xy + y^2$.
 - $Q(x, y) = \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$.
 - Orthogonally diagonalize $\begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix}$.
 - $\begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}^T \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{3}{2} \end{pmatrix}$.
 - Let $\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}^T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}}(x + y) \\ \frac{1}{\sqrt{2}}(-x + y) \end{pmatrix}$.

$$\begin{aligned} Q(x, y) &= \frac{1}{2}(x')^2 + \frac{3}{2}(y')^2 \\ &= \frac{1}{4}(x + y)^2 + \frac{3}{4}(-x + y)^2. \end{aligned}$$

76 / 96

Examples

- Let $Q(x, y, z) = x^2 + 2y^2 + z^2 + 2xz$.
 - $Q(x, y, z) = \begin{pmatrix} x & y & z \end{pmatrix} \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$.
 - Orthogonally diagonalize $\mathbf{A} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 1 \end{pmatrix}$.
 - $\mathbf{P}^T \mathbf{A} \mathbf{P} = \mathbf{D} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \mathbf{P} = \begin{pmatrix} \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{pmatrix}$.
 - Let $\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \mathbf{P}^T \mathbf{x} = \begin{pmatrix} \frac{1}{\sqrt{2}}(x + z) \\ y \\ \frac{1}{\sqrt{2}}(-x + z) \end{pmatrix}$.
 - $Q(x, y, z) = 2(x')^2 + 2(y')^2 + 0(z')^2 = (x + z)^2 + 2y^2$.

77 / 96

Quadratic Equation

- A **quadratic equation** in variable x is of the form
 - $ax^2 + bx = c$.
- **Definition.** A **quadratic equation** in variables x and y is
 - $ax^2 + bxy + cy^2 + dx + ey = f$.

The graph of a quadratic equation is a **conic section**.

- **Note.** Let $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$, $\mathbf{A} = \begin{pmatrix} a & \frac{1}{2}b \\ \frac{1}{2}b & c \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} d \\ e \end{pmatrix}$.
 - $\mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} = f$.
- **Definition.** $ax^2 + bxy + cy^2 = \mathbf{x}^T \mathbf{A} \mathbf{x}$ is the quadratic form **associated** with the quadratic equation.
 - $ax^2 + bxy + cy^2 + dx + ey = f$.

78 / 96

Classification of Conics

- Classification of **conic sections**.
 - **Degenerated** conic sections.
 - The whole plane \mathbb{R}^2 : $0 = 0$.
 - Empty set: $x^2 + y^2 = -1$.
 - A point: $x^2 + y^2 = 0$.
 - A line: $x = 0$ or $x^2 = 0$.
 - A pair of distinct lines: $x^2 - y^2 = 0$.
 - **Non-degenerated** conic sections.
 - Circle: $x^2 + y^2 = 1$.
 - Ellipse: $x^2 + 2y^2 = 1$.
 - Hyperbola: $x^2 - y^2 = 1$.
 - Parabola: $x^2 - y = 0$.

79 / 96


Standard Forms

- Standard form of **circle** or **ellipse**:

- $\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1, \quad \alpha > 0, \beta > 0.$

- $(x \ y) \begin{pmatrix} \frac{1}{\alpha^2} & 0 \\ 0 & \frac{1}{\beta^2} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 1.$

If $\alpha = \beta$, it is a circle of radius $r = \alpha = \beta$.



80 / 96


Standard Forms

- Standard form of **circle** or **ellipse**:

- $\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1, \quad \alpha > 0, \beta > 0.$

- $(x \ y) \begin{pmatrix} \frac{1}{\alpha^2} & 0 \\ 0 & \frac{1}{\beta^2} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 1.$

If $\alpha > \beta$, ellipse of major radius α , minor radius β :



81 / 96


Standard Forms

- Standard form of **circle** or **ellipse**:

- $\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1, \quad \alpha > 0, \beta > 0.$

- $(x \ y) \begin{pmatrix} \frac{1}{\alpha^2} & 0 \\ 0 & \frac{1}{\beta^2} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 1.$

If $\alpha < \beta$, ellipse of major radius β , minor radius α :



82 / 96


Standard Forms

- Standard form of **hyperbola**:

- Case 1: $\frac{x^2}{\alpha^2} - \frac{y^2}{\beta^2} = 1, \quad \alpha > 0, \beta > 0.$

- $(x \ y) \begin{pmatrix} \frac{1}{\alpha^2} & 0 \\ 0 & -\frac{1}{\beta^2} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 1.$

Semi-major axis α and semi-minor axis β .



83 / 96


Standard Forms

- Standard form of **hyperbola**:

- Case 2: $-\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1, \alpha > 0, \beta > 0.$

- $(x \ y) \begin{pmatrix} -\frac{1}{\alpha^2} & 0 \\ 0 & \frac{1}{\beta^2} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 1.$

Semi-major axis β and semi-minor axis α .



84 / 96


Standard Forms

- Standard form of **parabola**:

- Case 1: $x^2 = \alpha y, |\alpha|/4 \neq 0$ is the focal length.

- $(x \ y) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + (0 \ -\alpha) \begin{pmatrix} x \\ y \end{pmatrix} = 0.$

Suppose that $\alpha > 0$.



85 / 96


Standard Forms

- Standard form of **parabola**:

- Case 1: $x^2 = \alpha y$, $|\alpha|/4 \neq 0$ is the focal length.

- $$\begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 & -\alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 0.$$

Suppose that $\alpha < 0$.



86 / 96


Standard Forms

- Standard form of **parabola**:

- Case 2: $y^2 = \alpha x$, $|\alpha|/4 \neq 0$ is the focal length.

- $$\begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} -\alpha & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 0.$$

Suppose that $\alpha > 0$.




87 / 96

Standard Forms

- Standard form of **parabola**:
 - Case 2: $y^2 = \alpha x$, $|\alpha|/4 \neq 0$ is the focal length.
 - $(x \ y) \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + (-\alpha \ 0) \begin{pmatrix} x \\ y \end{pmatrix} = 0.$

Suppose that $\alpha < 0$.



88 / 96

Classification

- Classify $\mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} = f$, $\mathbf{x} \in \mathbb{R}^2$.
 1. Orthogonally diagonalize \mathbf{A} .
 - $\mathbf{P}^T \mathbf{A} \mathbf{P} = \begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix}$, \mathbf{P} an orthogonal matrix.
 2. Let $\mathbf{y} = \mathbf{P}^T \mathbf{x}$. Then
 - $\mathbf{y}^T \begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix} \mathbf{y} + \mathbf{b}^T \mathbf{P} \mathbf{y} = f.$
 3. Complete the squares.
- **Remark.** λ and μ are eigenvalues of \mathbf{A} ; $\lambda\mu = \det(\mathbf{A})$.
 - Suppose the conic section is **non-degenerate**.
 - $\det(\mathbf{A}) > 0 \Leftrightarrow$ ellipse (or circle).
 - $\det(\mathbf{A}) < 0 \Leftrightarrow$ hyperbola.
 - $\det(\mathbf{A}) = 0 \Leftrightarrow$ parabola.

89 / 96

Examples

- $x^2 - xy + y^2 - x - y = 1.$

Let $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$, $\mathbf{A} = \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} -1 \\ -1 \end{pmatrix}.$

- $\mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} = 1.$

1. Orthogonally diagonalize $\mathbf{A}.$

- $\mathbf{P}^T \mathbf{A} \mathbf{P} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{3}{2} \end{pmatrix},$ where $\mathbf{P} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}.$

2. Let $\begin{pmatrix} x' \\ y' \end{pmatrix} = \mathbf{y} = \mathbf{P}^T \mathbf{x} = \begin{pmatrix} \frac{1}{\sqrt{2}}(x + y) \\ \frac{1}{\sqrt{2}}(-x + y) \end{pmatrix}.$

- $\mathbf{y}^T \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{3}{2} \end{pmatrix} \mathbf{y} + \mathbf{b}^T \mathbf{P} \mathbf{y} = 1.$

90 / 96

Examples

- $x^2 - xy + y^2 - x - y = 1.$

Let $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$, $\mathbf{A} = \begin{pmatrix} 1 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} -1 \\ -1 \end{pmatrix}.$

- $\mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} = 1.$

3. $\frac{1}{2}(x')^2 + \frac{3}{2}(y')^2 - \sqrt{2}(x') = 1.$

- $\frac{1}{2}(x' - \sqrt{2})^2 + \frac{3}{2}(y')^2 = 1 + \frac{1}{2}(\sqrt{2})^2 = 2.$

- $\frac{(x' - \sqrt{2})^2}{2^2} + \frac{(y')^2}{(2/\sqrt{3})^2} = 1.$

Note that $\mathbf{P} = \begin{pmatrix} \cos \frac{\pi}{4} & -\sin \frac{\pi}{4} \\ \sin \frac{\pi}{4} & \cos \frac{\pi}{4} \end{pmatrix}, \begin{pmatrix} x' \\ y' \end{pmatrix} = \mathbf{P}^T \begin{pmatrix} x \\ y \end{pmatrix}.$

- The x' - and y' -axis is obtained by rotating the x - and y -axis about the origin O anticlockwise by $\pi/4.$


91 / 96

Examples

- $x^2 - xy + y^2 - x - y = 1.$

- $\frac{(x' - \sqrt{2})^2}{2^2} + \frac{(y')^2}{(2/\sqrt{3})^2} = 1.$

The x' - and y' -axis is obtained by rotating the x - and y -axis about the origin O anticlockwise by $\pi/4$.



92 / 96

Examples

- $2x^2 + 24xy + 9y^2 + 20x - 6y = 5.$

Let $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$, $\mathbf{A} = \begin{pmatrix} 2 & 12 \\ 12 & 9 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 20 \\ -6 \end{pmatrix}$.

- $\mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} = 5.$

1. Orthogonally diagonalize \mathbf{A} (Exercise).

- $\mathbf{P}^T \mathbf{A} \mathbf{P} = \begin{pmatrix} 18 & 0 \\ 0 & -7 \end{pmatrix}$, where $\mathbf{P} = \begin{pmatrix} \frac{3}{5} & -\frac{4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{pmatrix}.$

2. Let $\begin{pmatrix} x' \\ y' \end{pmatrix} = \mathbf{y} = \mathbf{P}^T \mathbf{x} = \begin{pmatrix} \frac{3}{5}x + \frac{4}{5}y \\ -\frac{4}{5}x + \frac{3}{5}y \end{pmatrix}.$

- $\mathbf{y}^T \begin{pmatrix} 18 & 0 \\ 0 & -7 \end{pmatrix} \mathbf{y} + \mathbf{b}^T \mathbf{P} \mathbf{y} = 5.$

93 / 96

Examples

- $2x^2 + 24xy + 9y^2 + 20x - 6y = 5.$

Let $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$, $\mathbf{A} = \begin{pmatrix} 2 & 12 \\ 12 & 9 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 20 \\ -6 \end{pmatrix}.$

- $\mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} = 5.$

3. $18(x')^2 - 7(y')^2 + \frac{36}{5}x' - \frac{98}{5}y' = 5.$

- $18(x' + \frac{1}{5})^2 - 7(y' + \frac{7}{5})^2 = -8.$

- $-\frac{(x' + \frac{1}{5})^2}{(2/3)^2} + \frac{(y' + \frac{7}{5})^2}{(\sqrt{8/7})^2} = 1.$

Note that $\mathbf{P} = \begin{pmatrix} \frac{3}{5} & -\frac{4}{5} \\ \frac{4}{5} & \frac{3}{5} \end{pmatrix}$ and $\mathbf{y} = \mathbf{P}^T \mathbf{x}.$

- The x' - and y' -axis is obtained by rotating the x - and y -axis about the origin O anticlockwise by $\cos^{-1}(\frac{3}{5}).$


94 / 96

Examples

- $2x^2 + 24xy + 9y^2 + 20x - 6y = 5.$

- $-\frac{(x' + \frac{1}{5})^2}{(2/3)^2} + \frac{(y' + \frac{7}{5})^2}{(\sqrt{8/7})^2} = 1.$

The x' - and y' -axis is obtained by rotating the x - and y -axis about the origin O anticlockwise by $\cos^{-1}(\frac{3}{5}).$



95 / 96

Remark

- Let \mathbf{P} be orthogonal of order 2. Then $\det(\mathbf{P}) = \pm 1$.
 - $\det(\mathbf{P}) = 1 \Rightarrow \mathbf{P} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$.
 - Let $\mathbf{y} = \mathbf{P}^T \mathbf{x}$. Then the new axes are obtained by rotating the original axes about O anticlockwise by θ .
 - $\det(\mathbf{P}) = -1 \Rightarrow \mathbf{P} = \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}$.
 - $\mathbf{P} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.
 - Let $\mathbf{y} = \mathbf{P}^T \mathbf{x}$. Then the new axes are obtained by first rotating the original axes about O anticlockwise by θ , then reflecting w.r.t. the x' -axis.

By multiplying the 2nd column of \mathbf{P} by -1 if necessary, we can always diagonalize a symmetric \mathbf{A} by an orthogonal matrix with determinant 1.

96 / 96