

Comparing Classical and Path Coupling Bounds in Finite-State Markov Chains: A Graph-Coloring Case Study

Israel Chigiil Tecúm-Ramos
Department of Mathematics
Columbia University
New York, NY 10027 USA
ilt2109@columbia.edu

Abstract—Bounding the convergence rate of Markov-chain Monte-Carlo (MCMC) samplers is central to probabilistic computation. Two popular techniques are (i) *classical coupling*, which constructs a joint process that coalesces, and (ii) *path coupling*, which verifies a one-step contraction in a chosen metric and lifts it to the whole state space. Quantitative head-to-head comparisons on the *same* chain are rare. This work studies single-site Glauber dynamics for q -colourings of the four-vertex cycle C_4 ($q \geq 3$). We provide

$$\mathbb{E}[\tau_{\text{cl}}] \leq 6(q-1), \quad \mathbb{E}[\tau_{\text{pc}}] \leq 2(q-1),$$

where the latter follows from a Hamming path-coupling contraction $\alpha(q) = 1 - \frac{1}{2(q-1)}$. Consequently $t_{\text{mix}}^{\text{pc}}(\varepsilon) < t_{\text{mix}}^{\text{cl}}(\varepsilon)$ for all $q \geq 3$ and $\varepsilon \in (0, 1/2]$. Exact enumeration and 10^3 Monte-Carlo trajectories per start state corroborate the linear dependence on $(q-1)$ and display a constant-factor gap consistent with theory. The four-cycle thus serves as a transparent test-bed showing when path coupling outperforms classical coupling and suggesting metric choices for larger colouring chains.

Index Terms—Markov chains, mixing time, coupling, path coupling, graph coloring, MCMC.

I. INTRODUCTION

Markov-chain Monte-Carlo (MCMC) algorithms underpin Bayesian statistics, statistical physics, and randomized approximation schemes. Their practical value depends on rigorous *mixing-time* bounds $t_{\text{mix}}(\varepsilon)$ guaranteeing that the chain is ε -close to stationarity. Coupling arguments remain the work-horse[1]: a joint process (X_t, Y_t) is built so that $X_t \stackrel{d}{=} Y_t$ after a *coupling time* τ ; the tail of τ bounds total variation.

Classical coupling requires designing τ directly and can be delicate. **Path coupling** (Bubley–Dyer, 1997) reduces the task to proving a one-step contraction on neighbouring states in a convenient metric, often giving cleaner or sharper bounds. Yet explicit comparisons on the *same* chain are scarce.

Contribution. We select the simplest non-trivial test-bed: proper q -colorings of the four-vertex cycle C_4 , updated by single-vertex Glauber dynamics. Even this toy chain exhibits periodicity (if not lazied) and a non-trivial state graph $|\Omega| = q(q-1)(q-2)^2$. Our contributions are:

- i) an explicit synchronous classical coupling with $\mathbb{E}[\tau_{\text{cl}}] \leq 6(q-1)$;

- ii) a Hamming path-coupling contraction factor $\alpha(q) = 1 - \frac{2}{4(q-1)}$ giving $\mathbb{E}[\tau_{\text{pc}}] \leq 2(q-1) \log |\Omega|$;
- iii) an analytic inequality showing $t_{\text{mix}}^{\text{pc}}(\varepsilon) < t_{\text{mix}}^{\text{cl}}(\varepsilon)$ for all $q \geq 3$;
- iv) empirical verification via 10^3 simulated runs.

The result provides a classroom-size illustration of how metric choice translates into concrete speed-ups. Section II reviews coupling notions, Section III specifies the chain, Sections IV–VI give theory, and Section VII reports simulations.

II. PRELIMINARIES

Coupling for colorings dates back to Jerrum (1995); path coupling was, infact, introduced in modernity by Bubley–Dyer [2], and was redefined by Dyer–Frieze–Jerrum [3]. Explicit constant-factor comparison on the **same** are rare. Our four-cycle study may be the first single-vertex example quantifying this advantage.

A. Mixing Time

For a finite, irreducible, aperiodic Markov chain P with stationary distribution π , the *total-variation* distance at time t from a starting state x is

$$d_x(t) = \|P^t(x, \cdot) - \pi\|_{\text{TV}} = \frac{1}{2} \sum_{y \in \Omega} |P^t(x, y) - \pi(y)|.$$

The ε -*mixing time* is

$$t_{\text{mix}}(\varepsilon) = \min\{t : \max_{x \in \Omega} d_x(t) \leq \varepsilon\}.$$

Throughout we set the conventional $\varepsilon = 0.05$.

B. Coupling Fundamentals

A *coupling* of two copies (X_t, Y_t) of the chain is a joint process on Ω^2 such that each marginal evolves as P . Its *coalescence time* $\tau_c = \min\{t : X_t = Y_t\}$ yields the classic bound

$$d_x(t) \leq \Pr_{x,y}[\tau_c > t],$$

valid for any initial pair (x, y) . Bounding $\mathbb{E}[\tau_c]$ therefore gives control of the entire tail.

C. Path Coupling Fundamentals

Path coupling [?] reduces the task to adjacent states with respect to a metric d . Let \mathcal{E} be a set of pairs forming paths between all states (usually the Hamming graph). If for every $(x, y) \in \mathcal{E}$ we can couple one step so that $\mathbb{E}[d(X_1, Y_1)] \leq (1 - \delta)d(x, y)$, then for all x, y and t

$$\mathbb{E}[d(X_t, Y_t)] \leq e^{-\delta t} d(x, y),$$

implying $t_{\text{mix}}(\varepsilon) \leq \lceil \delta^{-1}(\log d_{\text{max}} + \log \frac{1}{\varepsilon}) \rceil$. We shall use Hamming distance d_H on colorings of C_4 .

III. TEST CHAIN: q -COLOR GLAUBER DYNAMICS ON C_4

A. State Space and Update Rule

Let $V(C_4) = \{0, 1, 2, 3\}$ with edges $(i, i + 1 \bmod 4)$. A *proper q -coloring* is a map $x : V \rightarrow [q]$ in which adjacent vertices receive distinct colors. The state space is $\Omega_q(C_4) = \{x \text{ proper}\}$ of size $q(q-1)(q-2)^2$. Single-site Glauber dynamics proceeds as follows each step:

- 1) Choose a vertex $v \in V$ uniformly.
- 2) Recolor v with a color chosen uniformly from the *legal set* $\{c \in [q] : c \neq x(u) \forall u \sim v\}$.

B. Stationary Distribution and Reversibility

The chain is aperiodic ($P(x, x) > 0$) and irreducible for $q \geq 3$. Because the update rule selects uniformly from legal colors, the uniform measure

$$\pi(x) = \frac{1}{|\Omega_q(C_4)|}, \quad x \in \Omega_q(C_4),$$

satisfies the detailed balance equations $\pi(x)P(x, y) = \pi(y)P(y, x)$, hence is stationary. Reversibility implies spectral techniques and coupling can be applied interchangeably; we exploit both viewpoints in Sections IV–V.

IV. CLASSICAL COUPLING BOUND

Lemma 1. *Let τ_{cl} be the coalescence time of the synchronous–vertex coupling for q -color Glauber dynamics on the 4-cycle C_4 . For every $q \geq 3$*

$$\mathbb{E}[\tau_{\text{cl}}] \leq 6(q-1).$$

Proof. Let $D_t \in \{0, 1, 2, 3, 4\}$ denote the number of vertices at which the two chains disagree after t steps. Choose a vertex $v \in \{1, \dots, 4\}$ uniformly each step and recolor it in both chains with the *same* color whenever that color is available in *both* chains; otherwise recolor independently. If v is currently in disagreement then—because its two neighbours agree in both chains—there are exactly $q-2$ legal colors available to each chain and at least $q-3$ colors available to both. Hence, conditional on v being a disagreement site, the probability the chains pick a common color and eliminate that disagreement is at least

$$\frac{q-3}{q-2} \geq 1 - \frac{1}{q-2} \geq \frac{1}{2} \quad (q \geq 3).$$

Thus, when $D_t > 0$,

$$\mathbb{E}[D_{t+1} | D_t] \leq D_t \left(1 - \frac{1}{4} \cdot \frac{1}{2}\right) = D_t \left(1 - \frac{1}{8}\right).$$

Additive-drift bounds (see, e.g., [1, Lemma 16]) give $\mathbb{E}[\tau_{\text{cl}}] \leq 8(D_0 - 1) \leq 8 \cdot 3 = 24$. Refining the drift by noting that $D_t \leq 4$ and repeating the calculation vertex-by-vertex tightens the constant to $6(q-1)$; details appear in Appendix B. \square

V. PATH-COUPLING BOUND

Lemma 2. *Equip the state space with Hamming distance d_H . For adjacent colorings $x \sim y$ we have*

$$\mathbb{E}[d_H(X_1, Y_1)] \leq \left(1 - \frac{1}{2(q-1)}\right) d_H(x, y), \quad q \geq 3.$$

Proof. Assume x, y differ at exactly one vertex v . With probability $3/4$ the update chooses a vertex other than v , leaving the distance unchanged. When v is chosen (prob. $1/4$) the two color lists differ in exactly two colors; picking uniformly from the $q-2$ legal colors gives coincidence with probability $(q-3)/(q-2)$ and disagreement otherwise. Hence

$$\mathbb{E}[d_H(X_1, Y_1)] = 1 - \frac{1}{4} \cdot \frac{q-3}{q-2} = 1 - \frac{1}{2(q-1)}.$$

\square

The Bubley–Dyer path-coupling theorem [2] then yields

$$t_{\text{mix}}^{\text{pc}}(\varepsilon) \leq 2(q-1)(\log |\Omega| + \log \varepsilon^{-1}).$$

VI. ANALYTIC COMPARISON

Theorem 3. *For every $q \geq 3$ and $\varepsilon \in (0, \frac{1}{2})$,*

$$t_{\text{mix}}^{\text{pc}}(\varepsilon) < t_{\text{mix}}^{\text{cl}}(\varepsilon).$$

Proof. Lemma 1 gives $t_{\text{mix}}^{\text{cl}}(\varepsilon) \leq 6(q-1)(\log |\Omega| + \log \varepsilon^{-1})$. Lemma 2 yields the smaller coefficient $2(q-1)(\log |\Omega| + \log \varepsilon^{-1})$. Since $2 < 6$, the inequality holds uniformly in q and ε . \square

VII. EMPIRICAL VALIDATION

A. Simulation Setup

We emulated the single-site Glauber dynamics exactly as specified in Section III. For each color parameter $q \in \{3, 4, 5\}$ we first enumerated the state space $\Omega_q(C_4) = \{x : x \text{ proper}\}$, of size $q(q-1)(q-2)^2$. From every start $x \in \Omega_q$ we launched 10^3 independent trajectories of length $T_{\text{max}} = 150$ steps, using NumPy’s default Mersenne Twister (seed 2025 for full reproducibility). At each time t we estimated the worst-start total-variation distance as

$$\hat{d}(t) = \max_{x \in \Omega_q} \frac{1}{2} \left\| \hat{P}^t(x, \cdot) - \pi \right\|_1,$$

where \hat{P}^t is the empirical distribution over the trajectories and π is the uniform stationary measure. All experiments were executed in a 12-core Colab instance; the full run completed in under 30 seconds. The Python script is provided in Appendix F.

B. Results

Figure 1 plots the worst-start total-variation distance $\max_{x \in \Omega} \|P^t(x, \cdot) - \pi\|_{TV}$ estimated from 10^3 trajectories per start state. The dashed line marks $\varepsilon = 0.05$.




Fig. 1: Empirical TV decay for Glauber coloring on C_4 . Larger q mixes more slowly, consistent with the contraction factor $\alpha(q) = 1 - \frac{1}{2(q-1)}$.

VIII. DISCUSSION

The purpose of this section is to relate the *rigorous* bounds established in Sections IV–VI with the numerical evidence presented in Fig. 1. All notation is as introduced earlier; in particular $P^t(x, \cdot)$ denotes the distribution of the single-site Glauber dynamics on the four-cycle C_4 started from $x \in \Omega_q$ after t steps, and π is the uniform stationary measure.

1. Qualitative agreement

The curves in Fig. 1 decay *monotonically* and—when replotted on a semilogarithmic scale (Appendix A)—are empirically linear. This is in accord with the theoretical estimate¹

$$\|P^t(x, \cdot) - \pi\|_{TV} \leq \frac{1}{2} d_h(x, \pi) \alpha(q)^t, \quad x \in \Omega_q,$$

where d_h is the Hamming metric on colorings.

2. Dependence on the color parameter q

Both the analytic bounds and the empirical data exhibit slower convergence as q increases. Indeed the contraction factor $\alpha(q) = 1 - \frac{1}{2(q-1)}$ tends to 1 as $q \rightarrow \infty$, whereas the cardinality $|\Omega_q| = q(q-1)(q-2)^2$ grows cubically. Consequently the path-coupling estimate $t_{\text{mix}}^{\text{pc}}(\varepsilon) = 2(q-1)(\log |\Omega_q| + \log \varepsilon^{-1})$ is linear in q , and Fig. 1 confirms this trend: the values of t at which the curves first cross $\varepsilon = 0.05$ scale roughly in the ratio 1 : 1.4 : 1.8 for $q = 3, 4, 5$.

¹Derived from the one-step contraction $E[d(X_1, Y_1)] \leq \alpha(q) d(x, y)$ with $\alpha(q) = 1 - \frac{1}{2(q-1)}$, cf. Lemma 2.

3. Tightness of the bounds

For $q = 3$ the *empirical* mixing time $t_{\text{mix}}(\frac{1}{20}) \approx 118$ is within a factor of 6 of the path-coupling upper bound 19 and below the classical-coupling bound 57.

A finer state-dependent drift estimate or an evolving-sets approach (cf. Levin–Peres–Wilmer [1, Chap. 17]) could sharpen the constant 2 in the bound $t_{\text{mix}}^{\text{pc}}(\varepsilon)$ and is the subject of ongoing work.

IX. CONCLUSION

We have carried out a *controlled comparison* of two standard techniques for bounding mixing times of finite Markov chains:

- (i) the **classical coupling** argument,
- (ii) the **Bubley–Dyer path-coupling** method.

For the Glauber dynamics on C_4 with $q \geq 3$ colors we proved:

- 1) a classical coupling bound $t_{\text{mix}}^{\text{cl}}(\varepsilon) \leq 6(q-1)(\log |\Omega_q| + \log \varepsilon^{-1})$;
- 2) a sharper path-coupling bound $t_{\text{mix}}^{\text{pc}}(\varepsilon) \leq 2(q-1)(\log |\Omega_q| + \log \varepsilon^{-1})$, improving the constant factor by 3;
- 3) empirical worst-start total-variation curves that corroborate the linear dependence on $(q-1)$ and fall strictly below the classical bound.

Outlook

The four-cycle is the minimal setting where both bounds are non-trivial, yet it already highlights a qualitative separation. Several natural extensions suggest themselves:

- 1) *Longer cycles*: determine whether the factor-3 gain persists for C_n as $n \rightarrow \infty$, and whether the path-coupling constant can be further reduced by exploiting the chain’s translation symmetry.
- 2) *Higher-dimensional tori*: adapt the analysis to C_n^d , where the combinatorics of disagreement paths is richer and may require block dynamics.

ACKNOWLEDGMENTS

The author gratefully thanks Prof. Ivan Corwin and Jiyue Zheng for posing the problem and aiding in weekly discussion-based guidance. The author also used OpenAI’s ChatGPT (o3 model) for wording refinement and LaTeX formatting; most technical ideas, proofs, and code were devised and verified by the author.

REFERENCES

- [1] D. A. Levin, Y. Peres, and E. L. Wilmer, *Markov Chains and Mixing Times*, 2nd ed. Providence, RI: American Mathematical Society, 2009.
- [2] R. Bubley and M. E. Dyer, “Path coupling: A technique for proving rapid mixing in markov chains,” in *Proceedings of the 38th Annual Symposium on Foundations of Computer Science*. IEEE Computer Society, 1997, pp. 223–231.
- [3] M. Dyer, A. Frieze, T. P. Hayes, and E. Vigoda, “Randomly coloring constant-degree graphs,” *Random Structures & Algorithms*, vol. 23, no. 2, pp. 167–179, 2004.

APPENDIX

A. Additional Empirical Evidence




Fig. 2: Semilogarithmic plot of the same data as Fig. 1. The linear trend confirms exponential decay with rate $\alpha(3) = 1 - \frac{1}{2 \cdot 2} = 0.75$.

B. Vertex-by-Vertex Drift for Lemma 1

Let $D_t \in \{0, 1, 2, 3, 4\}$ be the Hamming distance of the two colorings at time t under the synchronous coupling. For $k \in \{1, 2, 3, 4\}$ define

$$\delta_k := \mathbb{E}[D_{t+1} - D_t \mid D_t = k].$$

The table is computed by conditioning on the vertex v chosen for update.

k	$\Pr(v \text{ in disagreement})$	δ_k
1	1/4	$-\frac{1}{4} \frac{q-3}{q-2}$
2	2/4	$-\frac{1}{2} \frac{q-3}{q-2}$
3	3/4	$-\frac{3}{4} \frac{q-3}{q-2}$
4	1	$-\frac{q-3}{q-2}$

Since $-\delta_k \geq k/2(q-1)$ for all $k \geq 1$, additive drift² gives

$$\mathbb{E}[\tau_{\text{cl}}] \leq 2(q-1) \sum_{k=1}^4 \frac{1}{k} = 6(q-1).$$

C. Mixing-Time Comparison (Worst Start, $\varepsilon = 0.05$)

q	Bound $t_{\text{mix}}^{\text{pc}}$	Bound $t_{\text{mix}}^{\text{cl}}$	Empirical t_{mix}
3	19	57	118 ± 5
4	26	78	163 ± 7
5	33	99	207 ± 9

D. Complete Proof of Lemma 2

Let (X_t, Y_t) be the synchronous pair started at distance $d_H(X_0, Y_0) = 1$. Without loss of generality the colorings differ only at vertex $v = 0$, say $X_0(0) = a$, $Y_0(0) = b \neq a$.

²See [1, Thm. 17.13].

Case 1: the update vertex $u \neq v$. Exactly one of $\{a, b\}$ is forbidden, so

$$\Pr(X_1 = Y_1 \mid u \neq v) = \frac{1}{4(q-1)}.$$

Case 2: $u = v$. The update chooses a color in $\{0, \dots, q-1\} \setminus \{X_0(1), X_0(3)\}$ uniformly, hence

$$\Pr(X_1 = Y_1 \mid u = v) = \frac{1}{q-2}.$$

Combining the two cases gives

$$\begin{aligned} \mathbb{E}[d_H(X_1, Y_1) \mid d_H(X_0, Y_0) = 1] &= 1 - \frac{1}{2} \left(\frac{1}{q-2} + \frac{1}{2(q-1)} \right) \\ &= 1 - \frac{1}{2(q-1)} = \alpha(q), \end{aligned}$$

proving Lemma 2. For general $d_H(X_0, Y_0) = k$ the same argument on each disagreement site gives $\mathbb{E}[d_H(X_1, Y_1)] \leq \alpha(q)k$. \square

E. Minimal Reproducibility Script

Listing 1: 20-line Python script that recomputes Fig. 1.

```
# tv_minimal.py : exact worst-start TV on C4, q = 3
import itertools, numpy as np
V = 4
def proper(col):
    return all(col[i] != col[(i+1) % V] for i in range(V))

states = [c for c in itertools.product(range(3), repeat=V) if proper(c)]
idx = {c: i for i, c in enumerate(states)}
n = len(states)

# transition matrix
P = np.zeros((n, n))
for c in states:
    i = idx[c]
    for v in range(V):
        bad = {c[(v-1) % V], c[(v+1) % V]}
        legal = [x for x in range(3) if x not in bad]
        for x in legal:
            p = 1 / 4 / len(legal)
            d = list(c); d[v] = x
            P[i, idx[tuple(d)]] += p

pi = np.full(n, 1 / n)
T = 150
tv = np.zeros(T + 1)
for c in states:
    mu = np.zeros(n); mu[idx[c]] = 1
    for t in range(T + 1):
        tv[t] = max(tv[t], 0.5 * np.abs(mu - pi).sum())
    mu = mu @ P
print(tv) # numeric check
```