

# Chapter 5: Fourier Transform on $\mathbb{R}$

## Introduction & Motivation

(Iz Tecúm)

(Columbia University)

December 27, 2025

# High-level outline for today

- What problem does the Fourier transform solve?

# High-level outline for today

- What problem does the Fourier transform solve?
- Where does its kernel  $e^{-2\pi i x \xi}$  come from?

# High-level outline for today

- What problem does the Fourier transform solve?
- Where does its kernel  $e^{-2\pi i x \xi}$  come from?
- From Fourier series (discrete frequencies) to Fourier transform (continuous spectrum)

# High-level outline for today

- What problem does the Fourier transform solve?
- Where does its kernel  $e^{-2\pi i x \xi}$  come from?
- From Fourier series (discrete frequencies) to Fourier transform (continuous spectrum)
- Why inversion matters; why it's nontrivial

# High-level outline for today

- What problem does the Fourier transform solve?
- Where does its kernel  $e^{-2\pi i x \xi}$  come from?
- From Fourier series (discrete frequencies) to Fourier transform (continuous spectrum)
- Why inversion matters; why it's nontrivial
- Core properties you'll use repeatedly

# High-level outline for today

- What problem does the Fourier transform solve?
- Where does its kernel  $e^{-2\pi i x \xi}$  come from?
- From Fourier series (discrete frequencies) to Fourier transform (continuous spectrum)
- Why inversion matters; why it's nontrivial
- Core properties you'll use repeatedly
- Where this chapter sits: PDE, probability, signal processing

# Roadmap

- 1 Motivation
- 2 Core Properties & Examples
- 3 Fourier Inversion



# What problem are we solving?

**Goal:** Decompose a function  $f : \mathbb{R} \rightarrow \mathbb{C}$  into *pure oscillations* (frequencies).

## Two worlds

- **Time/space domain** (original  $x$ -variable):  $f(x)$

# What problem are we solving?

**Goal:** Decompose a function  $f : \mathbb{R} \rightarrow \mathbb{C}$  into *pure oscillations* (frequencies).

## Two worlds

- **Time/space domain** (original  $x$ -variable):  $f(x)$
- **Frequency domain** (frequency  $\xi$ ):  $\hat{f}(\xi)$

# What problem are we solving?

**Goal:** Decompose a function  $f : \mathbb{R} \rightarrow \mathbb{C}$  into *pure oscillations* (frequencies).

## Two worlds

- **Time/space domain** (original  $x$ -variable):  $f(x)$
- **Frequency domain** (frequency  $\xi$ ):  $\hat{f}(\xi)$

Why switch domains?

- Linear operations that are *hard* in time (e.g. convolution) become *easy* in frequency (pointwise products).

# What problem are we solving?

**Goal:** Decompose a function  $f : \mathbb{R} \rightarrow \mathbb{C}$  into *pure oscillations* (frequencies).

## Two worlds

- **Time/space domain** (original  $x$ -variable):  $f(x)$
- **Frequency domain** (frequency  $\xi$ ):  $\hat{f}(\xi)$

Why switch domains?

- Linear operations that are *hard* in time (e.g. convolution) become *easy* in frequency (pointwise products).
- Many PDEs and filters are *diagonal* in frequency.

# What problem are we solving?

**Goal:** Decompose a function  $f : \mathbb{R} \rightarrow \mathbb{C}$  into *pure oscillations* (frequencies).

## Two worlds

- **Time/space domain** (original  $x$ -variable):  $f(x)$
- **Frequency domain** (frequency  $\xi$ ):  $\hat{f}(\xi)$

Why switch domains?

- Linear operations that are *hard* in time (e.g. convolution) become *easy* in frequency (pointwise products).
- Many PDEs and filters are *diagonal* in frequency.
- Random variables: distribution  $\leftrightarrow$  characteristic function (Fourier transform).

Where does the kernel  $e^{-2\pi i x \xi}$  come from?

**Translation symmetry** on  $\mathbb{R}$ :  $(T_t f)(x) = f(x - t)$ .

Where does the kernel  $e^{-2\pi i x \xi}$  come from?

**Translation symmetry** on  $\mathbb{R}$ :  $(T_t f)(x) = f(x - t)$ .

**Eigenfunction requirement**

Seek functions  $\phi$  with  $T_t \phi = \lambda(t) \phi$  for all  $t \in \mathbb{R}$ :

$$\phi(x - t) = \lambda(t) \phi(x) \quad \forall x, t.$$

Where does the kernel  $e^{-2\pi i x \xi}$  come from?

**Translation symmetry** on  $\mathbb{R}$ :  $(T_t f)(x) = f(x - t)$ .

**Eigenfunction requirement**

Seek functions  $\phi$  with  $T_t \phi = \lambda(t) \phi$  for all  $t \in \mathbb{R}$ :

$$\phi(x - t) = \lambda(t) \phi(x) \quad \forall x, t.$$

Then  $\lambda(t + s) = \lambda(t)\lambda(s)$ , and with mild regularity  $\lambda(t) = e^{ct}$ .



# Where does the kernel $e^{-2\pi i x \xi}$ come from?

**Translation symmetry** on  $\mathbb{R}$ :  $(T_t f)(x) = f(x - t)$ .

## Eigenfunction requirement

Seek functions  $\phi$  with  $T_t \phi = \lambda(t) \phi$  for all  $t \in \mathbb{R}$ :

$$\phi(x - t) = \lambda(t) \phi(x) \quad \forall x, t.$$

Then  $\lambda(t + s) = \lambda(t)\lambda(s)$ , and with mild regularity  $\lambda(t) = e^{ct}$ .

- To keep  $|\lambda(t)| = 1$  (isometries in  $L^2$ ):  $c = 2\pi i \xi$ ,  $\xi \in \mathbb{R}$ .
- $\Rightarrow$  eigenfunctions:  $\phi_\xi(x) = e^{2\pi i \xi x}$ .

## Where does the kernel $e^{-2\pi i x \xi}$ come from?

**Translation symmetry** on  $\mathbb{R}$ :  $(T_t f)(x) = f(x - t)$ .

### Eigenfunction requirement

Seek functions  $\phi$  with  $T_t \phi = \lambda(t) \phi$  for all  $t \in \mathbb{R}$ :

$$\phi(x - t) = \lambda(t) \phi(x) \quad \forall x, t.$$

Then  $\lambda(t + s) = \lambda(t)\lambda(s)$ , and with mild regularity  $\lambda(t) = e^{ct}$ .

- To keep  $|\lambda(t)| = 1$  (isometries in  $L^2$ ):  $c = 2\pi i \xi$ ,  $\xi \in \mathbb{R}$ .
- $\Rightarrow$  eigenfunctions:  $\phi_\xi(x) = e^{2\pi i \xi x}$ .

**Therefore:** project  $f$  against these characters to read off “how much of frequency  $\xi$ ” it contains.

## From Fourier series to Fourier transform

- On the circle  $\mathbb{T}$ :  $f(x) \sim \sum_{n \in \mathbb{Z}} a_n e^{2\pi i n x}$ ,  $a_n = \int_0^1 f(x) e^{-2\pi i n x} dx$ .

## From Fourier series to Fourier transform

- On the circle  $\mathbb{T}$ :  $f(x) \sim \sum_{n \in \mathbb{Z}} a_n e^{2\pi i n x}$ ,  $a_n = \int_0^1 f(x) e^{-2\pi i n x} dx$ .
- On  $\mathbb{R}$ : frequencies become *continuous*.

# From Fourier series to Fourier transform

- On the circle  $\mathbb{T}$ :  $f(x) \sim \sum_{n \in \mathbb{Z}} a_n e^{2\pi i n x}$ ,  $a_n = \int_0^1 f(x) e^{-2\pi i n x} dx$ .
- On  $\mathbb{R}$ : frequencies become *continuous*.

Heuristic: let the period  $L \rightarrow \infty$ . Frequency grid  $n/L$  becomes continuous  $\xi$ , sums  $\rightarrow$  integrals.

# From Fourier series to Fourier transform

- On the circle  $\mathbb{T}$ :  $f(x) \sim \sum_{n \in \mathbb{Z}} a_n e^{2\pi i n x}$ ,  $a_n = \int_0^1 f(x) e^{-2\pi i n x} dx$ .
- On  $\mathbb{R}$ : frequencies become *continuous*.

Heuristic: let the period  $L \rightarrow \infty$ . Frequency grid  $n/L$  becomes continuous  $\xi$ , sums  $\rightarrow$  integrals.

## Definition (Fourier transform on $\mathbb{R}$ )

$$\widehat{f}(\xi) = \int_{-\infty}^{\infty} f(x) e^{-2\pi i x \xi} dx \quad (\text{at least for } f \in L^1(\mathbb{R})).$$

# From Fourier series to Fourier transform

- On the circle  $\mathbb{T}$ :  $f(x) \sim \sum_{n \in \mathbb{Z}} a_n e^{2\pi i n x}$ ,  $a_n = \int_0^1 f(x) e^{-2\pi i n x} dx$ .
- On  $\mathbb{R}$ : frequencies become *continuous*.

Heuristic: let the period  $L \rightarrow \infty$ . Frequency grid  $n/L$  becomes continuous  $\xi$ , sums  $\rightarrow$  integrals.

## Definition (Fourier transform on $\mathbb{R}$ )

$$\widehat{f}(\xi) = \int_{-\infty}^{\infty} f(x) e^{-2\pi i x \xi} dx \quad (\text{at least for } f \in L^1(\mathbb{R})).$$

Different texts use different  $2\pi$  conventions; here it's the Stein–Shakarchi normalization.

## First properties (why this change of variables is useful)

For nice  $f, g$  (e.g.  $L^1$  or Schwartz):

$$\text{Linearity: } \widehat{af + bg} = a\hat{f} + b\hat{g}.$$

$$\text{Translation: } \widehat{(T_t f)}(\xi) = e^{-2\pi i t \xi} \hat{f}(\xi).$$

$$\text{Scaling: } \widehat{(f(ax))}(\xi) = \frac{1}{|a|} \hat{f}\left(\frac{\xi}{a}\right).$$

$$\text{Convolution: } \widehat{(f * g)} = \hat{f} \cdot \hat{g}.$$

$$\text{Differentiation: } \widehat{(f')}(\xi) = (2\pi i \xi) \hat{f}(\xi).$$

$$\text{Multiplication by } x : \widehat{(xf(x))}(\xi) = \frac{1}{2\pi i} \frac{d}{d\xi} \hat{f}(\xi).$$



## First properties (why this change of variables is useful)

For nice  $f, g$  (e.g.  $L^1$  or Schwartz):

$$\text{Linearity: } \widehat{af + bg} = a\hat{f} + b\hat{g}.$$

$$\text{Translation: } \widehat{(T_t f)}(\xi) = e^{-2\pi i t \xi} \hat{f}(\xi).$$

$$\text{Scaling: } \widehat{(f(ax))}(\xi) = \frac{1}{|a|} \hat{f}\left(\frac{\xi}{a}\right).$$

$$\text{Convolution: } \widehat{(f * g)} = \hat{f} \cdot \hat{g}.$$

$$\text{Differentiation: } \widehat{(f')}(\xi) = (2\pi i \xi) \hat{f}(\xi).$$

$$\text{Multiplication by } x : \widehat{(xf(x))}(\xi) = \frac{1}{2\pi i} \frac{d}{d\xi} \hat{f}(\xi).$$

**Meta:** convolution  $\leftrightarrow$  multiplication; differentiation  $\leftrightarrow$  polynomial weights in  $\xi$ .

## Why inversion matters (and why it's delicate)

- We want  $\mathcal{F}^{-1}(\mathcal{F}f) = f$ : frequency analysis must be *reversible*.

# Why inversion matters (and why it's delicate)

- We want  $\mathcal{F}^{-1}(\mathcal{F}f) = f$ : frequency analysis must be *reversible*.
- Inversion theorem (informal): if  $f \in L^1(\mathbb{R})$  and  $\hat{f} \in L^1(\mathbb{R})$ , then

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i x \xi} d\xi \quad \text{for a.e. } x.$$

# Why inversion matters (and why it's delicate)

- We want  $\mathcal{F}^{-1}(\mathcal{F}f) = f$ : frequency analysis must be *reversible*.
- Inversion theorem (informal): if  $f \in L^1(\mathbb{R})$  and  $\hat{f} \in L^1(\mathbb{R})$ , then

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i x \xi} d\xi \quad \text{for a.e. } x.$$

- Delicate because absolute convergence and pointwise limits can fail without hypotheses.

# Why inversion matters (and why it's delicate)

- We want  $\mathcal{F}^{-1}(\mathcal{F}f) = f$ : frequency analysis must be *reversible*.
- Inversion theorem (informal): if  $f \in L^1(\mathbb{R})$  and  $\hat{f} \in L^1(\mathbb{R})$ , then

$$f(x) = \int_{-\infty}^{\infty} \hat{f}(\xi) e^{2\pi i x \xi} d\xi \quad \text{for a.e. } x.$$

- Delicate because absolute convergence and pointwise limits can fail without hypotheses.
- Strategy in the chapter: *prove inversion for very nice  $f$  (Schwartz), then extend by density.*

## Two running examples (intuition builders)

**Gaussian:**  $f(x) = e^{-\pi x^2} \Rightarrow \hat{f}(\xi) = e^{-\pi \xi^2}$  (self-dual).

## Two running examples (intuition builders)

**Gaussian:**  $f(x) = e^{-\pi x^2} \Rightarrow \hat{f}(\xi) = e^{-\pi \xi^2}$  (self-dual).

**Box/rect function:**  $f = \mathbf{1}_{[-1,1]} \Rightarrow$

$$\hat{f}(\xi) = \int_{-1}^1 e^{-2\pi i x \xi} dx = \frac{\sin(2\pi \xi)}{\pi \xi} \quad (\text{sinc}).$$

## Two running examples (intuition builders)

**Gaussian:**  $f(x) = e^{-\pi x^2} \Rightarrow \hat{f}(\xi) = e^{-\pi \xi^2}$  (self-dual).

**Box/rect function:**  $f = \mathbf{1}_{[-1,1]} \Rightarrow$

$$\hat{f}(\xi) = \int_{-1}^1 e^{-2\pi i x \xi} dx = \frac{\sin(2\pi \xi)}{\pi \xi} \quad (\text{sinc}).$$

### Lessons:

- Smooth  $\Rightarrow$  fast decay in frequency (Gaussian).
- Sharp edges  $\Rightarrow$  slowly decaying oscillations (sinc tails).



## Where does this chapter sit (big applications)

- **PDE / Heat equation:** Fourier turns  $\partial_t u = \Delta u$  into  $\partial_t \hat{u} = -4\pi^2 |\xi|^2 \hat{u}$  (decoupled ODEs).

## Where does this chapter sit (big applications)

- **PDE / Heat equation:** Fourier turns  $\partial_t u = \Delta u$  into  $\partial_t \hat{u} = -4\pi^2 |\xi|^2 \hat{u}$  (decoupled ODEs).
- **Probability:** characteristic function  $\varphi_X(\xi) = \mathbb{E}[e^{2\pi i \xi X}]$  is the Fourier transform of the law of  $X$ .

## Where does this chapter sit (big applications)

- **PDE / Heat equation:** Fourier turns  $\partial_t u = \Delta u$  into  $\partial_t \hat{u} = -4\pi^2 |\xi|^2 \hat{u}$  (decoupled ODEs).
- **Probability:** characteristic function  $\varphi_X(\xi) = \mathbb{E}[e^{2\pi i \xi X}]$  is the Fourier transform of the law of  $X$ .
- **Signals/filters:** convolution with a kernel  $\Leftrightarrow$  pointwise multiply by a transfer function.

## Where does this chapter sit (big applications)

- **PDE / Heat equation:** Fourier turns  $\partial_t u = \Delta u$  into  $\partial_t \hat{u} = -4\pi^2 |\xi|^2 \hat{u}$  (decoupled ODEs).
- **Probability:** characteristic function  $\varphi_X(\xi) = \mathbb{E}[e^{2\pi i \xi X}]$  is the Fourier transform of the law of  $X$ .
- **Signals/filters:** convolution with a kernel  $\Leftrightarrow$  pointwise multiply by a transfer function.
- **Uncertainty principles:** cannot be localized tightly in both  $x$  and  $\xi$  (Heisenberg-type bounds).

## What this chapter will / won't do

### Will do

- Define  $\mathcal{F}$  on  $\mathbb{R}$  with Stein–Shakarchi normalization.
- Prove inversion under reasonable  $L^1$  assumptions using approximation.
- Prepare ground for  $L^2$ : Plancherel/Parseval in the next chapter.

# What this chapter will / won't do

## Will do

- Define  $\mathcal{F}$  on  $\mathbb{R}$  with Stein–Shakarchi normalization.
- Prove inversion under reasonable  $L^1$  assumptions using approximation.
- Prepare ground for  $L^2$ : Plancherel/Parseval in the next chapter.

## Won't (yet)

- Full distribution theory and tempered distributions (later courses).
- Multidimensional transforms in generality (arrive later).
- Deep singular integral theory (needs more machinery).

## Learning roadmap for the chapter

- 1 Master the *definition* + core properties by rote (they drive every computation).

## Learning roadmap for the chapter

- 1 Master the *definition* + core properties by rote (they drive every computation).
- 2 Understand the *inversion proof strategy*: nice class  $\rightarrow$  approximation  $\rightarrow$  limit.



## Learning roadmap for the chapter

- ① Master the *definition* + core properties by rote (they drive every computation).
- ② Understand the *inversion proof strategy*: nice class  $\rightarrow$  approximation  $\rightarrow$  limit.
- ③ Drill the *two canon examples* (Gaussian, box  $\rightarrow$  sinc).

## Learning roadmap for the chapter

- 1 Master the *definition* + core properties by rote (they drive every computation).
- 2 Understand the *inversion proof strategy*: nice class  $\rightarrow$  approximation  $\rightarrow$  limit.
- 3 Drill the *two canon examples* (Gaussian, box  $\rightarrow$  sinc).
- 4 Keep applications in mind: PDE (heat), probability (characteristic functions), filtering.

## Micro summary (what to remember from this intro)

- Kernel  $e^{-2\pi i x \xi}$  is forced by translation symmetry (characters of  $(\mathbb{R}, +)$ ).

## Micro summary (what to remember from this intro)

- Kernel  $e^{-2\pi i x \xi}$  is forced by translation symmetry (characters of  $(\mathbb{R}, +)$ ).
- $\mathcal{F}$  transforms convolution to multiplication and derivatives to polynomials in  $\xi$ .

## Micro summary (what to remember from this intro)

- Kernel  $e^{-2\pi i x \xi}$  is forced by translation symmetry (characters of  $(\mathbb{R}, +)$ ).
- $\mathcal{F}$  transforms convolution to multiplication and derivatives to polynomials in  $\xi$ .
- Inversion is the central theorem; proofs proceed by approximation from very nice functions.

## Micro summary (what to remember from this intro)

- Kernel  $e^{-2\pi i x \xi}$  is forced by translation symmetry (characters of  $(\mathbb{R}, +)$ ).
- $\mathcal{F}$  transforms convolution to multiplication and derivatives to polynomials in  $\xi$ .
- Inversion is the central theorem; proofs proceed by approximation from very nice functions.
- This unlocks PDE, probability, and signal processing viewpoints.

# Roadmap

- 1 Motivation
- 2 Core Properties & Examples
- 3 Fourier Inversion

## Setup & domain

Fourier transform on  $\mathbb{R}$  (Stein–Shakarchi normalization)

For  $f \in L^1(\mathbb{R})$ ,

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx.$$



## Setup & domain

### Fourier transform on $\mathbb{R}$ (Stein–Shakarchi normalization)

For  $f \in L^1(\mathbb{R})$ ,

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx.$$

- **Minimal domain:**  $L^1(\mathbb{R})$  ensures the integral is absolutely convergent.

## Setup & domain

### Fourier transform on $\mathbb{R}$ (Stein–Shakarchi normalization)

For  $f \in L^1(\mathbb{R})$ ,

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx.$$

- **Minimal domain:**  $L^1(\mathbb{R})$  ensures the integral is absolutely convergent.
- **Nice class:** Schwartz  $\mathcal{S}(\mathbb{R})$  (smooth, rapidly decaying) is dense in  $L^p$  and stable under  $\mathcal{F}$ .

## Fourier transform on $\mathbb{R}$ (Stein–Shakarchi normalization)

For  $f \in L^1(\mathbb{R})$ ,

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx.$$

- **Minimal domain:**  $L^1(\mathbb{R})$  ensures the integral is absolutely convergent.
- **Nice class:** Schwartz  $\mathcal{S}(\mathbb{R})$  (smooth, rapidly decaying) is dense in  $L^p$  and stable under  $\mathcal{F}$ .
- **Conventions:** Different books place  $2\pi$  differently. We stick to Stein–Shakarchi.

## Linearity, conjugation, reflection

For  $f, g \in L^1$ ,  $a, b \in \mathbb{C}$ :

$$\widehat{af + bg} = a\hat{f} + b\hat{g}.$$

# Linearity, conjugation, reflection

For  $f, g \in L^1$ ,  $a, b \in \mathbb{C}$ :

$$\widehat{af + bg} = a\widehat{f} + b\widehat{g}.$$

If  $\overline{f}$  is the complex conjugate of  $f$ ,

$$\widehat{\overline{f}}(\xi) = \overline{\widehat{f}(-\xi)}.$$

# Linearity, conjugation, reflection

For  $f, g \in L^1$ ,  $a, b \in \mathbb{C}$ :

$$\widehat{af + bg} = a\widehat{f} + b\widehat{g}.$$

If  $\bar{f}$  is the complex conjugate of  $f$ ,

$$\widehat{\bar{f}}(\xi) = \overline{\widehat{f}(-\xi)}.$$

If  $f^\sharp(x) := f(-x)$  (time-reversal),

$$\widehat{f^\sharp}(\xi) = \widehat{f}(-\xi).$$

# Linearity, conjugation, reflection

For  $f, g \in L^1$ ,  $a, b \in \mathbb{C}$ :

$$\widehat{af + bg} = a\hat{f} + b\hat{g}.$$

If  $\bar{f}$  is the complex conjugate of  $f$ ,

$$\widehat{\bar{f}}(\xi) = \overline{\hat{f}(-\xi)}.$$

If  $f^\sharp(x) := f(-x)$  (time-reversal),

$$\hat{f}^\sharp(\xi) = \hat{f}(-\xi).$$

## Real/even/odd patterns:

- $f$  real  $\Rightarrow \hat{f}(-\xi) = \overline{\hat{f}(\xi)}$  (Hermitian symmetry).
- $f$  even real  $\Rightarrow \hat{f}$  real and even.
- $f$  odd real  $\Rightarrow \hat{f}$  purely imaginary and odd.

## Translation & modulation (prove it)

**Translation:**  $(T_t f)(x) := f(x - t)$ .

$$\widehat{(T_t f)}(\xi) = \int_{\mathbb{R}} f(x - t) e^{-2\pi i x \xi} dx \stackrel{u=x-t}{=} e^{-2\pi i t \xi} \int_{\mathbb{R}} f(u) e^{-2\pi i u \xi} du = e^{-2\pi i t \xi} \widehat{f}(\xi).$$



## Translation & modulation (prove it)

**Translation:**  $(T_t f)(x) := f(x - t)$ .

$$\widehat{(T_t f)}(\xi) = \int_{\mathbb{R}} f(x - t) e^{-2\pi i x \xi} dx \stackrel{u=x-t}{=} e^{-2\pi i t \xi} \int_{\mathbb{R}} f(u) e^{-2\pi i u \xi} du = e^{-2\pi i t \xi} \widehat{f}(\xi).$$

**Modulation:**  $M_{\xi_0} f(x) := e^{2\pi i \xi_0 x} f(x)$ .

$$\widehat{(M_{\xi_0} f)}(\xi) = \int f(x) e^{2\pi i \xi_0 x} e^{-2\pi i \xi x} dx = \widehat{f}(\xi - \xi_0).$$

**Moral:** shifts in time  $\leftrightarrow$  phase in frequency; tones in time  $\leftrightarrow$  shifts in frequency.

## Scaling & time reversal (derive)

For  $a \neq 0$ , define  $S_a f(x) := f(ax)$ .

$$\widehat{(S_a f)}(\xi) = \int f(ax) e^{-2\pi i x \xi} dx \stackrel{u=ax}{=} \frac{1}{|a|} \int f(u) e^{-2\pi i u (\xi/a)} du = \frac{1}{|a|} \widehat{f}\left(\frac{\xi}{a}\right).$$

## Scaling & time reversal (derive)

For  $a \neq 0$ , define  $S_a f(x) := f(ax)$ .

$$\widehat{(S_a f)}(\xi) = \int f(ax) e^{-2\pi i x \xi} dx \stackrel{u=ax}{=} \frac{1}{|a|} \int f(u) e^{-2\pi i u (\xi/a)} du = \frac{1}{|a|} \widehat{f}\left(\frac{\xi}{a}\right).$$

Time reversal  $f^\sharp(x) = f(-x)$  was:  $\widehat{f^\sharp}(\xi) = \widehat{f}(-\xi)$ . **Heuristic:** zoom in time  $\Rightarrow$  stretch out in frequency (and renormalize).

## Differentiation & multiplication (operational rules)

If  $f, f' \in L^1$ ,

$$\widehat{(f')}( \xi ) = \int f'(x) e^{-2\pi i x \xi} dx \stackrel{\text{IBP}}{=} \left[ f(x) e^{-2\pi i x \xi} \right]_{-\infty}^{\infty} + (2\pi i \xi) \int f(x) e^{-2\pi i x \xi} dx = (2\pi i \xi) \widehat{f}(\xi),$$

where the boundary term vanishes for  $f \in \mathcal{S}$ ; for  $L^1$  use approximation.

## Differentiation & multiplication (operational rules)

If  $f, f' \in L^1$ ,

$$\widehat{(f')}(\xi) = \int f'(x) e^{-2\pi i x \xi} dx \stackrel{\text{IBP}}{=} \left[ f(x) e^{-2\pi i x \xi} \right]_{-\infty}^{\infty} + (2\pi i \xi) \int f(x) e^{-2\pi i x \xi} dx = (2\pi i \xi) \widehat{f}(\xi),$$

where the boundary term vanishes for  $f \in \mathcal{S}$ ; for  $L^1$  use approximation.

$$\widehat{(xf(x))}(\xi) = \frac{1}{2\pi i} \frac{d}{d\xi} \widehat{f}(\xi) \quad (\text{differentiate under the integral}).$$

**Meta:** derivatives  $\leftrightarrow$  polynomial factors in  $\xi$ ; moments in time  $\leftrightarrow$  derivatives in frequency.

## Convolution theorem (key bridge)

$(f * g)(x) := \int_{\mathbb{R}} f(x-y)g(y) dy$  with  $f, g \in L^1$  gives  $f * g \in L^1$ .

$$\widehat{(f * g)}(\xi) = \int_{\mathbb{R}} \int_{\mathbb{R}} f(x-y)g(y) e^{-2\pi i x \xi} dy dx \stackrel{x-y=u}{=} \int g(y) e^{-2\pi i y \xi} dy \cdot \int f(u) e^{-2\pi i u \xi} du = \widehat{f}(\xi) \widehat{g}(\xi)$$

## Convolution theorem (key bridge)

$(f * g)(x) := \int_{\mathbb{R}} f(x-y)g(y) dy$  with  $f, g \in L^1$  gives  $f * g \in L^1$ .

$$\widehat{(f * g)}(\xi) = \int_{\mathbb{R}} \int_{\mathbb{R}} f(x-y)g(y) e^{-2\pi i x \xi} dy dx \stackrel{x-y=u}{=} \int g(y) e^{-2\pi i y \xi} dy \cdot \int f(u) e^{-2\pi i u \xi} du = \widehat{f}(\xi) \widehat{g}(\xi)$$

**Corollary:**  $f * g$  is a *smoothed* version of  $f$  when  $g$  is smooth/decaying; in frequency you just multiply by  $\widehat{g}$  (a low-pass filter).

## Riemann–Lebesgue lemma (decay)

### Lemma

If  $f \in L^1(\mathbb{R})$  then  $\hat{f}$  is bounded, uniformly continuous, and

$$\hat{f}(\xi) \rightarrow 0 \quad \text{as } |\xi| \rightarrow \infty.$$



# Riemann–Lebesgue lemma (decay)

## Lemma

If  $f \in L^1(\mathbb{R})$  then  $\widehat{f}$  is bounded, uniformly continuous, and

$$\widehat{f}(\xi) \rightarrow 0 \quad \text{as } |\xi| \rightarrow \infty.$$

## Sketch:

- Boundedness by  $\|\widehat{f}\|_\infty \leq \|f\|_1$  (triangle inequality).
- Approximate  $f$  by compactly supported continuous functions; oscillations cancel at large  $|\xi|$  (Dirichlet kernel trick).
- Uniform continuity from dominated convergence (difference quotient inside the integral).

**Intuition:** integrals against faster oscillations average out.

## Symmetry & uncertainty (preview)

- **Even/odd & real:** summarized earlier (Hermitian symmetry).

# Symmetry & uncertainty (preview)

- **Even/odd & real:** summarized earlier (Hermitian symmetry).
- **Time–frequency tradeoff (Heisenberg):**

$$\sigma_x(f) \sigma_\xi(\hat{f}) \geq \frac{1}{4\pi},$$

where  $\sigma$  denotes standard deviation (for  $L^2$ -normalized  $f$ ).

# Symmetry & uncertainty (preview)

- **Even/odd & real:** summarized earlier (Hermitian symmetry).
- **Time–frequency tradeoff (Heisenberg):**

$$\sigma_x(f) \sigma_\xi(\hat{f}) \geq \frac{1}{4\pi},$$

where  $\sigma$  denotes standard deviation (for  $L^2$ -normalized  $f$ ).

- Equality for Gaussians  $\Rightarrow$  optimally concentrated in both domains.

## Worked example I: Gaussian is self-dual

Let  $f(x) = e^{-\pi x^2}$ . Complete the square:

$$\widehat{f}(\xi) = \int_{\mathbb{R}} e^{-\pi x^2} e^{-2\pi i x \xi} dx = \int_{\mathbb{R}} e^{-\pi (x^2 + 2ix\xi)} dx = \int_{\mathbb{R}} e^{-\pi (x - i\xi)^2} e^{-\pi \xi^2} dx.$$

## Worked example I: Gaussian is self-dual

Let  $f(x) = e^{-\pi x^2}$ . Complete the square:

$$\hat{f}(\xi) = \int_{\mathbb{R}} e^{-\pi x^2} e^{-2\pi i x \xi} dx = \int_{\mathbb{R}} e^{-\pi (x^2 + 2ix\xi)} dx = \int_{\mathbb{R}} e^{-\pi (x-i\xi)^2} e^{-\pi \xi^2} dx.$$

Shift the contour (Schwarz class justification) or use standard Gaussian integral:

$$\int_{\mathbb{R}} e^{-\pi (x-i\xi)^2} dx = \int_{\mathbb{R}} e^{-\pi u^2} du = 1.$$

$$\Rightarrow \hat{f}(\xi) = e^{-\pi \xi^2}.$$

**Takeaway:** Gaussians minimize uncertainty and are fixed by  $\mathcal{F}$ .

## Worked example II: Box $\rightarrow$ sinc

$$f(x) = \mathbf{1}_{[-1,1]}(x).$$

$$\hat{f}(\xi) = \int_{-1}^1 e^{-2\pi i x \xi} dx = \left[ \frac{e^{-2\pi i x \xi}}{-2\pi i \xi} \right]_{-1}^1 = \frac{e^{-2\pi i \xi} - e^{2\pi i \xi}}{-2\pi i \xi} = \frac{\sin(2\pi \xi)}{\pi \xi}.$$

## Worked example II: Box $\rightarrow$ sinc

$$f(x) = \mathbf{1}_{[-1,1]}(x).$$

$$\hat{f}(\xi) = \int_{-1}^1 e^{-2\pi i x \xi} dx = \left[ \frac{e^{-2\pi i x \xi}}{-2\pi i \xi} \right]_{-1}^1 = \frac{e^{-2\pi i \xi} - e^{2\pi i \xi}}{-2\pi i \xi} = \frac{\sin(2\pi \xi)}{\pi \xi}.$$

### Observations:

- Decays like  $1/|\xi|$  (slow, due to discontinuities).
- Zeros at  $\xi \in \mathbb{Z} \setminus \{0\}$ .
- Sinc tails  $\Rightarrow$  Gibbs-type ripples on reconstruction near jumps.



## Worked example III: Triangle via convolution

Triangle  $\tau = \mathbf{1}_{[-1,1]} * \mathbf{1}_{[-1,1]}$ .

$$\widehat{\tau}(\xi) = \widehat{\mathbf{1}_{[-1,1]}}(\xi) \cdot \widehat{\mathbf{1}_{[-1,1]}}(\xi) = \left( \frac{\sin(2\pi\xi)}{\pi\xi} \right)^2.$$

## Worked example III: Triangle via convolution

Triangle  $\tau = \mathbf{1}_{[-1,1]} * \mathbf{1}_{[-1,1]}$ .

$$\widehat{\tau}(\xi) = \widehat{\mathbf{1}_{[-1,1]}}(\xi) \cdot \widehat{\mathbf{1}_{[-1,1]}}(\xi) = \left( \frac{\sin(2\pi\xi)}{\pi\xi} \right)^2.$$

**Moral:** Convolution in time smooths the edges  $\Rightarrow$  faster decay (here  $\sim 1/\xi^2$ ) in frequency.

## Common pitfalls & domain cautions

- Not every  $L^1$  function has  $\hat{f} \in L^1$ ; inversion needs care (will prove under extra hypotheses).

## Common pitfalls & domain cautions

- Not every  $L^1$  function has  $\hat{f} \in L^1$ ; inversion needs care (will prove under extra hypotheses).
- Functions with jumps decay slowly in frequency ( $1/\xi$  tails).

## Common pitfalls & domain cautions

- Not every  $L^1$  function has  $\hat{f} \in L^1$ ; inversion needs care (will prove under extra hypotheses).
- Functions with jumps decay slowly in frequency ( $1/\xi$  tails).
- Poles/impulses (e.g.  $\delta$ ) live in the world of distributions; we postpone this framework.

## Common pitfalls & domain cautions

- Not every  $L^1$  function has  $\hat{f} \in L^1$ ; inversion needs care (will prove under extra hypotheses).
- Functions with jumps decay slowly in frequency ( $1/\xi$  tails).
- Poles/impulses (e.g.  $\delta$ ) live in the world of distributions; we postpone this framework.
- Interchange of integrals (Fubini/Tonelli) requires integrability checks; we always justify via  $L^1$  or Schwartz approximation.

## Bridge to the next section

- We now have the toolkit: translation/modulation, scaling, convolution, differentiation rules.

## Bridge to the next section

- We now have the toolkit: translation/modulation, scaling, convolution, differentiation rules.
- Next: **Inversion theorem** — when and how  $\mathcal{F}^{-1}(\mathcal{F}f) = f$ .



## Bridge to the next section

- We now have the toolkit: translation/modulation, scaling, convolution, differentiation rules.
- Next: **Inversion theorem** — when and how  $\mathcal{F}^{-1}(\mathcal{F}f) = f$ .
- Strategy preview: prove for Schwartz, then extend to  $L^1$  by approximation (Gaussian mollifiers / dominated convergence).

# Roadmap

- 1 Motivation
- 2 Core Properties & Examples
- 3 Fourier Inversion

# The inversion problem

## Question

Given  $f : \mathbb{R} \rightarrow \mathbb{C}$ , can we *recover*  $f$  from its Fourier transform

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx ?$$

# The inversion problem

## Question

Given  $f : \mathbb{R} \rightarrow \mathbb{C}$ , can we *recover*  $f$  from its Fourier transform

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx ?$$

- Formal guess:  $f(x) \stackrel{?}{=} \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi.$

# The inversion problem

## Question

Given  $f : \mathbb{R} \rightarrow \mathbb{C}$ , can we *recover*  $f$  from its Fourier transform

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx ?$$

- Formal guess:  $f(x) \stackrel{?}{=} \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi$ .
- Issues: absolute convergence; interchanging integrals; pointwise vs. a.e. equality.

# The inversion problem

## Question

Given  $f : \mathbb{R} \rightarrow \mathbb{C}$ , can we *recover*  $f$  from its Fourier transform

$$\widehat{f}(\xi) = \int_{\mathbb{R}} f(x) e^{-2\pi i x \xi} dx ?$$

- Formal guess:  $f(x) \stackrel{?}{=} \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi$ .
- Issues: absolute convergence; interchanging integrals; pointwise vs. a.e. equality.
- Strategy (Chapter 5): prove for *very nice*  $f$  (Schwartz), extend by approximation.

## A standard sufficient condition

### Fourier inversion (one $L^1$ version)

If  $f \in L^1(\mathbb{R})$  and  $\widehat{f} \in L^1(\mathbb{R})$ , then

$$f(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi \quad \text{for a.e. } x \in \mathbb{R}.$$

# A standard sufficient condition

## Fourier inversion (one $L^1$ version)

If  $f \in L^1(\mathbb{R})$  and  $\widehat{f} \in L^1(\mathbb{R})$ , then

$$f(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi \quad \text{for a.e. } x \in \mathbb{R}.$$

### Notes:

- The a.e. qualifier is sharp in general.
- There are many variants (e.g. pointwise at continuity points).
- The  $L^2$ -theory (Plancherel) gives a clean inverse on  $L^2$ —next chapter.



# Approximate identities via Gaussians

## Base Gaussian

$g(x) := e^{-\pi x^2}$  satisfies  $\widehat{g}(\xi) = g(\xi)$ .

# Approximate identities via Gaussians

## Base Gaussian

$g(x) := e^{-\pi x^2}$  satisfies  $\widehat{g}(\xi) = g(\xi)$ .

## Scaled approximate identity

Define  $\phi_\varepsilon(x) := \frac{1}{\varepsilon} g\left(\frac{x}{\varepsilon}\right) = \frac{1}{\varepsilon} e^{-\pi(x/\varepsilon)^2}$ , so  $\int_{\mathbb{R}} \phi_\varepsilon = 1$  and  $\phi_\varepsilon \rightarrow \delta$  (weakly).

# Approximate identities via Gaussians

## Base Gaussian

$g(x) := e^{-\pi x^2}$  satisfies  $\widehat{g}(\xi) = g(\xi)$ .

## Scaled approximate identity

Define  $\phi_\varepsilon(x) := \frac{1}{\varepsilon} g\left(\frac{x}{\varepsilon}\right) = \frac{1}{\varepsilon} e^{-\pi(x/\varepsilon)^2}$ , so  $\int_{\mathbb{R}} \phi_\varepsilon = 1$  and  $\phi_\varepsilon \rightarrow \delta$  (weakly).

- **Fourier side:** by scaling,

$$\widehat{\phi_\varepsilon}(\xi) = \widehat{g}(\varepsilon\xi) = e^{-\pi\varepsilon^2\xi^2}.$$

- **Consequence:**  $f * \phi_\varepsilon \rightarrow f$  in  $L^1$  and pointwise at Lebesgue points.

## Key identity linking $f * \phi_\varepsilon$ and $\widehat{f}$

For  $f \in L^1(\mathbb{R})$  and  $\varepsilon > 0$ :

$$(f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} \widehat{\phi_\varepsilon}(\xi) d\xi = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi.$$

## Key identity linking $f * \phi_\varepsilon$ and $\widehat{f}$

For  $f \in L^1(\mathbb{R})$  and  $\varepsilon > 0$ :

$$(f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} \widehat{\phi_\varepsilon}(\xi) d\xi = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi.$$

**Why true?** For nice  $f$  (e.g. Schwartz),  $\widehat{f * \phi_\varepsilon} = \widehat{f} \cdot \widehat{\phi_\varepsilon}$ ; take inverse transform. For  $f \in L^1$ , use approximation by Schwartz and dominated convergence (details next).

## Proof skeleton: Schwartz case first

Let  $f \in \mathcal{S}(\mathbb{R})$ .

- ①  $\mathcal{S}$  is invariant under  $\mathcal{F}$  and convolution;  $\widehat{g} = g$  for  $g(x) = e^{-\pi x^2}$ .

## Proof skeleton: Schwartz case first

Let  $f \in \mathcal{S}(\mathbb{R})$ .

- ①  $\mathcal{S}$  is invariant under  $\mathcal{F}$  and convolution;  $\widehat{g} = g$  for  $g(x) = e^{-\pi x^2}$ .
- ② For  $\phi_\varepsilon(x) = \varepsilon^{-1}g(x/\varepsilon)$  we have  $\widehat{\phi}_\varepsilon(\xi) = e^{-\pi\varepsilon^2\xi^2}$ .

# Proof skeleton: Schwartz case first

Let  $f \in \mathcal{S}(\mathbb{R})$ .

- ①  $\mathcal{S}$  is invariant under  $\mathcal{F}$  and convolution;  $\widehat{g} = g$  for  $g(x) = e^{-\pi x^2}$ .
- ② For  $\phi_\varepsilon(x) = \varepsilon^{-1}g(x/\varepsilon)$  we have  $\widehat{\phi}_\varepsilon(\xi) = e^{-\pi\varepsilon^2\xi^2}$ .
- ③ Convolution theorem:  $\widehat{f * \phi_\varepsilon} = \widehat{f} \cdot \widehat{\phi_\varepsilon}$ .



# Proof skeleton: Schwartz case first

Let  $f \in \mathcal{S}(\mathbb{R})$ .

- ①  $\mathcal{S}$  is invariant under  $\mathcal{F}$  and convolution;  $\widehat{g} = g$  for  $g(x) = e^{-\pi x^2}$ .
- ② For  $\phi_\varepsilon(x) = \varepsilon^{-1}g(x/\varepsilon)$  we have  $\widehat{\phi_\varepsilon}(\xi) = e^{-\pi\varepsilon^2\xi^2}$ .
- ③ Convolution theorem:  $\widehat{f * \phi_\varepsilon} = \widehat{f} \cdot \widehat{\phi_\varepsilon}$ .
- ④ Invert (still in  $\mathcal{S}$ ):

$$(f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi\varepsilon^2\xi^2} d\xi.$$

## Proof skeleton: Schwartz case first

Let  $f \in \mathcal{S}(\mathbb{R})$ .

- ①  $\mathcal{S}$  is invariant under  $\mathcal{F}$  and convolution;  $\widehat{g} = g$  for  $g(x) = e^{-\pi x^2}$ .
- ② For  $\phi_\varepsilon(x) = \varepsilon^{-1}g(x/\varepsilon)$  we have  $\widehat{\phi_\varepsilon}(\xi) = e^{-\pi\varepsilon^2\xi^2}$ .
- ③ Convolution theorem:  $\widehat{f * \phi_\varepsilon} = \widehat{f} \cdot \widehat{\phi_\varepsilon}$ .
- ④ Invert (still in  $\mathcal{S}$ ):

$$(f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi\varepsilon^2\xi^2} d\xi.$$

- ⑤ As  $\varepsilon \rightarrow 0$ ,  $e^{-\pi\varepsilon^2\xi^2} \rightarrow 1$  pointwise and is  $\leq 1$ —use dominated convergence (since  $\widehat{f} \in \mathcal{S} \subset L^1$ ) to get

$$\lim_{\varepsilon \rightarrow 0} (f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi.$$

# Proof skeleton: Schwartz case first

Let  $f \in \mathcal{S}(\mathbb{R})$ .

①  $\mathcal{S}$  is invariant under  $\mathcal{F}$  and convolution;  $\widehat{g} = g$  for  $g(x) = e^{-\pi x^2}$ .

② For  $\phi_\varepsilon(x) = \varepsilon^{-1}g(x/\varepsilon)$  we have  $\widehat{\phi_\varepsilon}(\xi) = e^{-\pi\varepsilon^2\xi^2}$ .

③ Convolution theorem:  $\widehat{f * \phi_\varepsilon} = \widehat{f} \cdot \widehat{\phi_\varepsilon}$ .

④ Invert (still in  $\mathcal{S}$ ):

$$(f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi\varepsilon^2\xi^2} d\xi.$$

⑤ As  $\varepsilon \rightarrow 0$ ,  $e^{-\pi\varepsilon^2\xi^2} \rightarrow 1$  pointwise and is  $\leq 1$ —use dominated convergence (since  $\widehat{f} \in \mathcal{S} \subset L^1$ ) to get

$$\lim_{\varepsilon \rightarrow 0} (f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi.$$

⑥ But  $f * \phi_\varepsilon \rightarrow f$  uniformly (even in  $\mathcal{S}$ -topology), hence inversion holds for Schwartz  $f$ .

## Extension to $L^1$ with $\widehat{f} \in L^1$

Let  $f \in L^1(\mathbb{R})$  with  $\widehat{f} \in L^1(\mathbb{R})$ . For each  $\varepsilon > 0$ :

$$(f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi.$$

## Extension to $L^1$ with $\widehat{f} \in L^1$

Let  $f \in L^1(\mathbb{R})$  with  $\widehat{f} \in L^1(\mathbb{R})$ . For each  $\varepsilon > 0$ :

$$(f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi.$$

**Right-hand side:** dominated by  $|\widehat{f}(\xi)|$ , integrable by hypothesis; thus

$$\lim_{\varepsilon \rightarrow 0} \int \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi = \int \widehat{f}(\xi) e^{2\pi i x \xi} d\xi \quad (\text{DCT}).$$

## Extension to $L^1$ with $\widehat{f} \in L^1$

Let  $f \in L^1(\mathbb{R})$  with  $\widehat{f} \in L^1(\mathbb{R})$ . For each  $\varepsilon > 0$ :

$$(f * \phi_\varepsilon)(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi.$$

**Right-hand side:** dominated by  $|\widehat{f}(\xi)|$ , integrable by hypothesis; thus

$$\lim_{\varepsilon \rightarrow 0} \int \widehat{f}(\xi) e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi = \int \widehat{f}(\xi) e^{2\pi i x \xi} d\xi \quad (\text{DCT}).$$

**Left-hand side:**  $f * \phi_\varepsilon \rightarrow f$  in  $L^1$  and at *Lebesgue points* of  $f$ . Hence for a.e.  $x$ ,

$$f(x) = \int_{\mathbb{R}} \widehat{f}(\xi) e^{2\pi i x \xi} d\xi.$$

# Lebesgue points & pointwise recovery

## Lebesgue point

$x$  is a Lebesgue point of  $f \in L^1$  if

$$\lim_{r \downarrow 0} \frac{1}{2r} \int_{x-r}^{x+r} |f(y) - f(x)| dy = 0.$$

## Lebesgue point

$x$  is a Lebesgue point of  $f \in L^1$  if

$$\lim_{r \downarrow 0} \frac{1}{2r} \int_{x-r}^{x+r} |f(y) - f(x)| dy = 0.$$

- Almost every  $x$  is a Lebesgue point (Lebesgue differentiation theorem).
- For approximate identities  $(\phi_\varepsilon)$  with  $\int \phi_\varepsilon = 1$ , we have  $(f * \phi_\varepsilon)(x) \rightarrow f(x)$  at Lebesgue points.
- This is the engine behind the a.e. inversion conclusion.



## Technical justifications (what we used)

- **Tonelli/Fubini:** We swap integrals in convolution and transform definitions only when the integrand is absolutely integrable; else approximate via  $\mathcal{S}$ .

## Technical justifications (what we used)

- **Tonelli/Fubini:** We swap integrals in convolution and transform definitions only when the integrand is absolutely integrable; else approximate via  $\mathcal{S}$ .
- **Dominated convergence (DCT):**  $|\widehat{f}(\xi) e^{-\pi \varepsilon^2 \xi^2}| \leq |\widehat{f}(\xi)| \in L^1$ .

## Technical justifications (what we used)

- **Tonelli/Fubini:** We swap integrals in convolution and transform definitions only when the integrand is absolutely integrable; else approximate via  $\mathcal{S}$ .
- **Dominated convergence (DCT):**  $|\widehat{f}(\xi) e^{-\pi \varepsilon^2 \xi^2}| \leq |\widehat{f}(\xi)| \in L^1$ .
- **Approximate identity:**  $\phi_\varepsilon$  is positive, integrates to 1, and concentrates at 0; hence  $f * \phi_\varepsilon \rightarrow f$  in  $L^1$  and a.e.

## Technical justifications (what we used)

- **Tonelli/Fubini:** We swap integrals in convolution and transform definitions only when the integrand is absolutely integrable; else approximate via  $\mathcal{S}$ .
- **Dominated convergence (DCT):**  $|\widehat{f}(\xi) e^{-\pi \varepsilon^2 \xi^2}| \leq |\widehat{f}(\xi)| \in L^1$ .
- **Approximate identity:**  $\phi_\varepsilon$  is positive, integrates to 1, and concentrates at 0; hence  $f * \phi_\varepsilon \rightarrow f$  in  $L^1$  and a.e.
- **Schwartz density:**  $\mathcal{S}(\mathbb{R})$  is dense in  $L^1$  and stable under  $\mathcal{F}$ .

- If  $f \in L^1$  is continuous at  $x$ , the inversion formula often holds *pointwise at*  $x$  under milder hypotheses.

- If  $f \in L^1$  is continuous at  $x$ , the inversion formula often holds *pointwise at*  $x$  under milder hypotheses.
- If  $f, \widehat{f} \in L^1$ , we get inversion a.e. and *uniform* convergence of  $(f * \phi_\varepsilon)$  to  $f$  on continuity sets.

## Variants & remarks

- If  $f \in L^1$  is continuous at  $x$ , the inversion formula often holds *pointwise at*  $x$  under milder hypotheses.
- If  $f, \hat{f} \in L^1$ , we get inversion a.e. and *uniform* convergence of  $(f * \phi_\varepsilon)$  to  $f$  on continuity sets.
- $L^2$ -inversion (Plancherel) gives a unitary isomorphism on  $L^2$  without needing  $\hat{f} \in L^1$ —next chapter.

- If  $f \in L^1$  is continuous at  $x$ , the inversion formula often holds *pointwise at*  $x$  under milder hypotheses.
- If  $f, \hat{f} \in L^1$ , we get inversion a.e. and *uniform* convergence of  $(f * \phi_\varepsilon)$  to  $f$  on continuity sets.
- $L^2$ -inversion (Plancherel) gives a unitary isomorphism on  $L^2$  without needing  $\hat{f} \in L^1$ —next chapter.
- Edge behavior: discontinuities create oscillatory reconstruction (Gibbs-type phenomena).



## Worked example: inversion with Gaussian damping

Suppose  $f \in L^1$  and  $\hat{f} \in L^1$ .

$$f(x) = \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} \hat{f}(\xi) e^{2\pi i x \xi} \underbrace{e^{-\pi \varepsilon^2 \xi^2}}_{\text{damps high freq}} d\xi.$$

## Worked example: inversion with Gaussian damping

Suppose  $f \in L^1$  and  $\hat{f} \in L^1$ .

$$f(x) = \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}} \hat{f}(\xi) e^{2\pi i x \xi} \underbrace{e^{-\pi \varepsilon^2 \xi^2}}_{\text{damps high freq}} d\xi.$$

**Why useful?** Numerically stable: the Gaussian factor regularizes the integral; then send  $\varepsilon \downarrow 0$ .

- If  $f$  is smooth/decaying, the damping is harmless and accelerates convergence.
- If  $f$  has jumps, damping controls oscillations in the reconstruction.

## Quick check: box $\rightarrow$ sinc and back (heuristic)

For  $f = \mathbf{1}_{[-1,1]}$ ,  $\widehat{f}(\xi) = \frac{\sin(2\pi\xi)}{\pi\xi}$ .

$$f(x) \approx \int_{\mathbb{R}} \frac{\sin(2\pi\xi)}{\pi\xi} e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi \xrightarrow{\varepsilon \downarrow 0} f(x)$$

at all continuity points  $x \neq \pm 1$ . At  $x = \pm 1$  one gets midpoint limits.

## Quick check: box $\rightarrow$ sinc and back (heuristic)

For  $f = \mathbf{1}_{[-1,1]}$ ,  $\widehat{f}(\xi) = \frac{\sin(2\pi\xi)}{\pi\xi}$ .

$$f(x) \approx \int_{\mathbb{R}} \frac{\sin(2\pi\xi)}{\pi\xi} e^{2\pi i x \xi} e^{-\pi \varepsilon^2 \xi^2} d\xi \xrightarrow{\varepsilon \downarrow 0} f(x)$$

at all continuity points  $x \neq \pm 1$ . At  $x = \pm 1$  one gets midpoint limits. **Moral:** damping  $+\varepsilon \downarrow 0$  realizes inversion even for nonsmooth  $f$ .

## Summary: inversion section

- Proved inversion for  $\mathcal{S}$  via Gaussian approximate identity.

## Summary: inversion section

- Proved inversion for  $\mathcal{S}$  via Gaussian approximate identity.
- Extended to  $f \in L^1$  with  $\hat{f} \in L^1$  using DCT + Lebesgue points.

## Summary: inversion section

- Proved inversion for  $\mathcal{S}$  via Gaussian approximate identity.
- Extended to  $f \in L^1$  with  $\hat{f} \in L^1$  using DCT + Lebesgue points.
- Learned a practical reconstruction formula with Gaussian damping.

## Summary: inversion section

- Proved inversion for  $\mathcal{S}$  via Gaussian approximate identity.
- Extended to  $f \in L^1$  with  $\hat{f} \in L^1$  using DCT + Lebesgue points.
- Learned a practical reconstruction formula with Gaussian damping.
- Next:  $L^2$  theory (Plancherel), Hausdorff–Young, and applications.



# The Fourier Inversion Theorem

## Statement

If  $f \in L^1(\mathbb{R})$  and  $\widehat{f} \in L^1(\mathbb{R})$ , then for almost every  $x \in \mathbb{R}$ ,

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \widehat{f}(\xi) e^{ix\xi} d\xi.$$

# The Fourier Inversion Theorem

## Statement

If  $f \in L^1(\mathbb{R})$  and  $\hat{f} \in L^1(\mathbb{R})$ , then for almost every  $x \in \mathbb{R}$ ,

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}(\xi) e^{ix\xi} d\xi.$$

- The Fourier transform is **invertible**.
- Frequency and spatial domains are two sides of the same coin.
- This closes the circle:  $f \mapsto \hat{f} \mapsto f$ .

# Significance of Fourier Inversion

- **Theoretical:** Proves the transform is not just a computational trick, but a true dual representation.
- **Analysis:** Central in PDEs, signal processing, harmonic analysis.
- **Applications:** Image compression, MRI reconstruction, quantum mechanics, option pricing.

# Significance of Fourier Inversion

- **Theoretical:** Proves the transform is not just a computational trick, but a true dual representation.
- **Analysis:** Central in PDEs, signal processing, harmonic analysis.
- **Applications:** Image compression, MRI reconstruction, quantum mechanics, option pricing.

## Key Idea

The inversion formula shows that Fourier analysis is a *complete language* for functions.

## Further Directions (Not Covered Today)

- **Plancherel Theorem:** Extension of inversion to  $L^2(\mathbb{R})$ , establishing

$$\|f\|_{L^2} = \frac{1}{\sqrt{2\pi}} \|\hat{f}\|_{L^2}.$$

- **Hausdorff–Young Inequality:** Bounds  $\|\hat{f}\|_{L^q}$  in terms of  $\|f\|_{L^p}$ .
- These results expand Fourier analysis from  $L^1$  to the entire  $L^p$  scale.

## Further Directions (Not Covered Today)

- **Plancherel Theorem:** Extension of inversion to  $L^2(\mathbb{R})$ , establishing

$$\|f\|_{L^2} = \frac{1}{\sqrt{2\pi}} \|\widehat{f}\|_{L^2}.$$

- **Hausdorff–Young Inequality:** Bounds  $\|\widehat{f}\|_{L^q}$  in terms of  $\|f\|_{L^p}$ .
- These results expand Fourier analysis from  $L^1$  to the entire  $L^p$  scale.

$\implies$  The inversion theorem is just the *beginning* of the story.

## Plancherel: statement (lite)

**Theorem.**  $\mathcal{F}$  extends uniquely to a unitary map on  $L^2(\mathbb{R})$ :

$$\|f\|_{L^2} = \|\widehat{f}\|_{L^2}, \quad \langle f, g \rangle = \langle \widehat{f}, \widehat{g} \rangle.$$

**Why it matters:** energy conservation; clean inversion on  $L^2$  without assuming  $\widehat{f} \in L^1$ .

## Plancherel: proof sketch in 4 beats

- 1 Prove on Schwartz  $\mathcal{S}$ :  $\|\widehat{f}\|_2 = \|f\|_2$  (Gaussian integral + Fubini).
- 2 Density:  $\mathcal{S}$  is dense in  $L^2$ .
- 3 Extend by continuity: define  $\widehat{\cdot}$  on  $L^2$  as the  $L^2$ -limit.
- 4 Unitarity and inversion follow; Parseval  $\langle f, g \rangle = \langle \widehat{f}, \widehat{g} \rangle$ .



## Plancherel: quick example

If  $f(x) = e^{-\pi x^2}$ , then  $\widehat{f}(\xi) = e^{-\pi \xi^2}$  and

$$\int_{\mathbb{R}} |f|^2 = \int_{\mathbb{R}} e^{-2\pi x^2} dx = \int_{\mathbb{R}} |\widehat{f}|^2 \quad (\text{Gaussian is self-dual}).$$

**Takeaway:**  $L^2$  is the “natural home” of Fourier analysis.

# Hausdorff–Young (one-slide version)

**Theorem.** If  $f \in L^p(\mathbb{R})$  with  $1 \leq p \leq 2$  and  $\frac{1}{p} + \frac{1}{q} = 1$ , then  $\widehat{f} \in L^q(\mathbb{R})$  and

$$\|\widehat{f}\|_{L^q} \leq \|f\|_{L^p}.$$

**Idea.** Interpolate between the endpoints:  $L^1 \rightarrow L^\infty$  (trivial bound) and  $L^2 \rightarrow L^2$  (Plancherel).

**Use.** Controls  $\widehat{f}$  integrability; key in estimates PDE.

## Applications montage (pick 1 if time)

- Heat equation:  $\partial_t \hat{u} = -4\pi^2 |\xi|^2 \hat{u} \Rightarrow u = G_t * u_0$  (Gaussian kernel).
- Probability: characteristic functions  $\varphi_X(\xi) = \mathbb{E}[e^{2\pi i \xi X}]$ ; CLT via pointwise product limits.
- Signals: convolutional smoothing = low-pass multiplier  $\hat{g}$ ; de-noising; deconvolution caveats.