

RAINBOW NORMAL FORMS AND THE L-SPACE PROPERTY OF ONE-RAINBOW (1, 1)-KNOTS

IZ TECÚM, PARTH WOKHL, YVAN GRINSPAN, RUSSELL MARTINEZ, AND FILIP RUPCHIN

ABSTRACT. A $(1, 1)$ -knot admits a doubly pointed genus-one diagram $(T^2, \alpha, \beta, z, w)$ [8, 12]; cutting along α presents β as $p = |\alpha \cap \beta|$ arcs in an annulus, and Doyle's rainbow normal form $[p, a, b, r]$ records a rainbows per wall, $p - 2a$ bands of one step size b , and an offset r [5, 6]. We treat $a = 1$, where S^3 -admissibility is the pair of conditions $|e| = 1$ on the drift $e = \beta \cdot \alpha$ and $|\tau^{\mathbb{Z}}(L_1)| = p$ on the transition word.

Every admissible one-rainbow diagram is reduced and coherent in the sense of [7], so each bigon of $T^2 \setminus (\alpha \cup \beta)$ covers z or w ; since $\text{Sym}^1(T^2) = T^2$, every index-one class is an embedded bigon with a single flow line [9, 12], the differential is the adjacency matrix of the empty-bigon relation over $\mathbb{F} = \mathbb{Z}/2$, and

$$\widehat{\partial} = \mathbf{0}, \quad \text{rk}(\widehat{HFK}(K)) = p, \quad \Delta_K(t) = \sum_{j=1}^p (-1)^{j+1} t^{A_j}, \quad A_1 > \cdots > A_p,$$

with x_z topmost for $e > 0$ and x_w for $e < 0$, so K is an L-space knot [10].

The census satisfies $|\mathcal{C}_p| = 4, 6, 8, 8, 10, 12, 14, 16$ for $p = 3, 5, \dots, 17$ and contains $T(2, p)$ and $T(\frac{p+1}{2}, \frac{p+3}{2})$; for 40 of the 48 tuples with $p \leq 13$ the semigroup $S_K = \langle a, b \rangle$ is generated by two elements, $|\mathcal{G}_K| = g = \frac{(a-1)(b-1)}{2}$ and $K = T(a, b)$ [15, 13], while the other eight are L-space knots that are not algebraic [1, 2]. In the algebraic case $\mathcal{G}_K = d(\cdot; a, b)^{-1}(0)$ and

$$\nabla_a \nabla_b d = \delta_0, \quad \nabla_a \nabla_b \frac{u_+}{ab} = B_{(a,b)}, \quad t^g \Delta_K(t) = (1-t)(1-t^{ab}) \sum_{n \geq 0} d(n; a, b) t^n,$$

so the Alexander polynomial is a second difference of a truncated power, separated from the box spline $B_{(a,b)}$ by an ab -periodic error of modulus at most one [11, 4, 3].

CONTENTS

1. Introduction	2
1.1. Rainbow normal forms	2
1.2. Result	3
2. Preliminaries	4
2.1. Rainbow normal form	4
2.2. Drift and admissibility	6
2.3. Reduced and coherent diagrams	7
2.4. Symmetries	8
2.5. Knot Floer input	8
2.6. Combinatorial input	9
3. Reducedness and coherence	9
3.1. Reducedness	9
3.2. Turning and coherence	11
3.3. The region dichotomy	13

Date: September 4, 2026.

2020 Mathematics Subject Classification. 57K18, 57K10, 57R58.

Key words and phrases. knot Floer homology, L-space knot, $(1, 1)$ -knot, Heegaard diagram, rainbow normal form, turning number, coherent diagram.

4. Proof of Theorem A	14
5. The kernel	15
5.1. The census	17
5.2. A worked tuple	18
5.3. Perfect diagrams	19
5.4. The difference word	19
5.5. Two closed forms	20
6. The box spline	21
6.1. The associated set	22
6.2. The partition function	23
6.3. The tent	24
6.4. The difference matrix	25
6.5. The conjecture	27
Acknowledgments	27
Appendix A. Foundational tools	28
A.1. Linear algebra over $\mathbb{F}$	28
A.2. Numerical semigroups	29
A.3. Turning of a piecewise-smooth curve	30
Appendix B. Reproducibility	30
B.1. The semigroup	30
B.2. The bigraded complex and the staircase	31
B.3. The partition function and the tent	32
B.4. The difference word	33
B.5. Changing basis on the torus	34
B.6. The diagram on the torus	35
References	36

1. INTRODUCTION

1.1. Rainbow normal forms. A knot $K \subset S^3$ is a $(1, 1)$ -knot when it admits a doubly pointed genus-one diagram $(T^2, \alpha, \beta, z, w)$ [8, 12], and since $\text{Sym}^1(T^2) = T^2$ the generators of $\widehat{CFK}(K)$ are the $p = |\alpha \cap \beta|$ intersection points [9]. Every reduced such diagram is isotopic to one with α a meridian [12, 7]. Cutting along α splits β into *rainbow arcs* $\mathbb{D}$ $|$ $\mathbb{Q}$, each cutting off a bigon that carries a basepoint, and *vertical arcs* $\mathbb{P}$. Doyle's normal form $[p, a, b, r]$ [5, 6] records a rainbow arcs per wall, b vertical arcs of one step size, and an offset r ; throughout $a = 1$, so $\mathbb{D} \mathbb{P} \mathbb{Q}$ carries a single rainbow arc on each wall.




FIGURE 1. $[3, 1, 1, 1]$, cut open along α and reglued by Φ : one rainbow arc per wall carrying z and w , one vertical arc between them, $e = +1$, $K = T(2, 3)$. Behind T^2 dashed.




FIGURE 2. $[5, 1, 2, 2]$ as in Figure 1, with three vertical arcs and β connected; $e = -1$, $K = T(3, 4)$.




FIGURE 3. The knot of Figure 1 meets T^2 only at z and w , one arc on the surface and one dotted through the solid torus. Windings 2 and 3 give $K = T(2, 3)$.

1.2. Result. Ozsváth and Szabó showed that an L-space knot has $\widehat{HFK}$ of rank one in each Alexander grading it occupies, with alternating unit coefficients in Δ_K [10]. We call this the *Alexander criterion*. At $a = 1$ it holds for every admissible tuple, and the candidacy is realised.

Theorem A. Let $[p, 1, b, r]$ be S^3 -admissible in the sense of Definition 2.2.3 and present $K \subset S^3$. Then

- (1) $\text{rk}(\widehat{HFK}(K)) = p$, with $A_1 > \dots > A_p$ pairwise distinct;


- (2) $\Delta_K(t) = \sum_{j=1}^p (-1)^{j+1} t^{A_j}$, so K satisfies the Alexander criterion;
- (3) the generator of topmost Alexander grading is x_z for $e > 0$ and x_w for $e < 0$;
- (4) K is an L-space knot.

Proof. §4. □

2. PRELIMINARIES

2.1. Rainbow normal form.

Definition 2.1.1 (Doubly pointed diagram). $(T^2, \alpha, \beta, z, w)$ with T^2 genus one, $\alpha, \beta \subset T^2$ essential simple closed curves meeting transversely, and $z, w \in T^2 \setminus (\alpha \cup \beta)$.



Definition 2.1.2 (Rainbow tuple, [5]). $[p, a, b, r] \in \mathbb{Z}^4$ with

$$p \geq 3 \text{ odd}, \quad 1 \leq a \leq \frac{p-1}{2}, \quad 0 \leq b \leq p-2a, \quad 1 \leq r \leq p,$$

so p generators, a rainbows per wall, b bands, first landing r .

Definition 2.1.3 (Cut annulus). $A = T^2 \setminus \alpha = [0, 1] \times \mathbb{R}/p\mathbb{Z}$, height increasing downwards,

$$W_L = \{0\} \times \mathbb{R}/p\mathbb{Z}, \quad W_R = \{1\} \times \mathbb{R}/p\mathbb{Z}, \quad L_h = \left(0, h - \frac{1}{2}\right), \quad R_h = \left(1, h - \frac{1}{2}\right), \quad h \in \mathbb{Z}/p\mathbb{Z},$$

with $\downarrow \uparrow: R_h \mapsto L_h$, both descending to $x_h \in \alpha \cap \beta$.

Definition 2.1.4 (Rainbow normal form, [5]). $D[p, a, b, r] = \text{diagram}$ is Definition 2.1.3 with $\beta \cap A$ the p disjoint embedded arcs

$$\begin{aligned} L_i &\leftrightarrow L_{2a+1-i}, & R_{r+b-1+i} &\leftrightarrow R_{r+b+2a-i}, & 1 \leq i \leq a, \\ L_s &\leftrightarrow R_{r+s-2a-1}, & & & 2a < s \leq 2a+b, \\ L_s &\leftrightarrow R_{r+s-1}, & & & 2a+b < s \leq p, \end{aligned}$$

the *rainbows*, the *bands*, and the *returns*, the last crossing $(u, 0) \sim (u, p)$ iff $r + s - 1 > p$, and

$$z \in L_a L_{a+1}, \quad w \in R_{r+b+a-1} R_{r+b+a},$$

the innermost rainbow bigons. The right endpoints $\rho: \{2a+1, \dots, p\} \rightarrow \mathbb{Z}/p\mathbb{Z} \setminus \{r+b, \dots, r+b+2a-1\}$ are a bijection.





FIGURE 4. Fairchild's construction of $D[7, 2, 1, 3]$, one arc family at a time from the marked points to the basepoints. Drawn after [6].

Convention 2.1.5 (One rainbow). $a = 1$, $v = r + b$, so Definition 2.1.4 reads

$$L_1 \leftrightarrow L_2, \quad R_v \leftrightarrow R_{v+1}, \quad L_s \leftrightarrow R_{\rho(s)}, \quad \rho(s) = \begin{cases} r + s - 3, & 3 \leq s \leq b + 2, \\ r + s - 1, & b + 2 < s \leq p, \end{cases}$$

with $z \in L_1L_2$, $w \in R_vR_{v+1}$, hence $x_z \in \{x_1, x_2\}$ and $x_w \in \{x_v, x_{v+1}\}$.



Definition 2.1.6 (Rainbow and vertical arcs, [12, 7]).

$$\mathbb{D} \mid = L_1L_2, \quad \mathbb{C} \mid = R_vR_{v+1}$$

are the *rainbow arcs*, $\downarrow \uparrow = c_s = L_sR_{\rho(s)}$ the *vertical arcs*, $\mathbb{D}$ and $\mathbb{C}$ the two rainbow families, and $\mathbb{B}$ the b arcs with $\rho(s) = r + s - 3$. The innermost rainbow arcs are the *caps* $\mathbb{D}_z$, $\mathbb{C}_w$, cutting off

$$\mathbb{D}_z = D_z \subset W_L, \quad \mathbb{C}_w = D_w \subset W_R.$$




FIGURE 5. $\mathcal{D} = D[7, 1, 2, 2]$ with $v = 4$. The rainbow arcs $\mathcal{D} = L_1 L_2$, $\mathcal{D} = R_4 R_5$ cut off $\mathcal{D}_z$, $\mathcal{D}_w$; the bands $\mathcal{B}$ are $L_3 R_2$, $L_4 R_3$; the wrapping arcs are $L_5 R_6$, $L_6 R_7$; and $L_7 R_1$ crosses $(u, 0) \sim (u, p)$, its second piece dashed. Under $\mathcal{D}$ the slots L_h, R_h have common image x_h .

Definition 2.1.7 (Cap corners). $x_z \in \partial D_z$ and $x_w \in \partial D_w$.

Remark 2.1.8 (Standard position, [12, 7, 2]). α is the meridian and β carries the rainbow arcs.


Proof. [12, §2]. □

2.2. Drift and admissibility.

Definition 2.2.1 (Traversal).


$$\omega(L_h) = R_h, \quad \omega(R_h) = L_h, \quad \tau = \omega \circ \sigma$$

for σ the matching of Definition 2.1.4. One step is $\mathcal{D} \rightarrow \mathcal{D}$, and the *traversal* is $\tau^{\mathbb{Z}}(L_1)$.




Definition 2.2.2 (Drift). For $e \in \beta \cap A$,

$$\chi(\mathcal{D}) = 0, \quad \chi(\mathcal{D}) = +1, \quad \chi(\mathcal{D}) = -1, \quad e = \sum_e \chi(e) = \beta \cdot \alpha.$$



Definition 2.2.3 (Admissibility). $D[p, 1, b, r]$ is S^3 -admissible when $\bigcirc \dashv \vdash$, that is $|\tau^{\mathbb{Z}}(L_1)| = p$ and $|e| = 1$.



Lemma 2.2.4 (Rank). $D[p, 1, b, r]$ admissible implies β connected, with $(T^2, \alpha, \beta) \simeq S^3$ and $\text{rk}(\widehat{CFK}(K)) = p$.

Proof.

$$\begin{aligned} |\tau^{\mathbb{Z}}(L_1)| = p &\xrightarrow{2.2.1} |\pi_0(\beta)| = 1 \\ \dashv \vdash &\implies \alpha \cap \beta = \{x_1, \dots, x_p\} \implies \text{rk}(\widehat{CFK}(K)) = p \\ |e| = 1 &\xrightarrow{2.2.2} |\beta \cdot \alpha| = 1 \xrightarrow{2.1.1} (T^2, \alpha, \beta) \simeq S^3 \end{aligned}$$

□

Remark 2.2.5 (Two step sizes). $\rho(s) - s \in \{r-3, r-1\}$, so τ is an exchange of two intervals $\bigcirc \dashv \vdash$, and $\star$ is a p -cycle exactly when $D[p, 1, b, r]$ is admissible.


Proof. Definition 2.1.4 and Definition 2.2.3. □

2.3. Reduced and coherent diagrams.

Definition 2.3.1 (Region type). A component R of $T^2 \setminus (\alpha \cup \beta)$ is n -sided when $|\partial R \cap \alpha \cap \beta| = 2n$,

$$\mathbb{D} \ (n=1), \quad \mathbb{D} \dashv \vdash \ (n=2), \quad \mathbb{D} \dashv \vdash \dashv \vdash \ (n=3), \quad \mathbb{D} \dashv \vdash \dashv \vdash \dashv \vdash \ (n=4),$$

with tints lightening as n grows. The darker lobe against a wall in $\mathbb{D} \dashv \vdash$ is a neighbouring bigon, one of the n sides of R and not part of it.



Definition 2.3.2 (Reduced diagram, [7]). A diagram is *reduced* $\mathbb{D}$ when every bigon of $T^2 \setminus (\alpha \cup \beta)$ carries a basepoint,




Definition 2.3.3 (Coherent diagram, [7]). A reduced diagram is *coherent* when α, β admit orientations coorienting every bigon, equivalently [2]



2.4. Symmetries.

Definition 2.4.1 (Orientation reversal). Reversing β sends $\downarrow \uparrow \mapsto \uparrow \downarrow$ and exchanges the basepoints $\Downarrow \Leftarrow \Downarrow$; reflecting A is $\Downarrow \Leftarrow \Downarrow$.



Lemma 2.4.2 (Effect on the gradings). Reversing β sends $e \mapsto -e$ and $A \mapsto -A$.

Proof.

$$\begin{aligned} \chi(\downarrow \uparrow) &= -\chi(\uparrow \downarrow) \xrightarrow{2.2.2} e \mapsto -e \\ \Downarrow \Leftarrow \Downarrow &\implies (n_z, n_w) \mapsto (n_w, n_z) \implies A \mapsto -A \end{aligned}$$

□

2.5. Knot Floer input.

Theorem 2.5.1 (Coherence criterion, [7, Theorem 1.2]). A reduced $(1, 1)$ -diagram presents an L -space knot if and only if it is coherent.

Proof. [7, Theorem 1.2]

□

Theorem 2.5.2 (Alexander polynomial, [10, Theorem 1.2]). For $K \subset S^3$ an L -space knot,

$$\Delta_K(t) = \sum_{j=0}^{2n} (-1)^j t^{A_j}, \quad A_0 > A_1 > \cdots > A_{2n},$$

and $\widehat{HFK}(K)$ has rank one in each A_j and vanishes elsewhere.


Proof. [10, Theorem 1.2]

□

Definition 2.5.3 (Alexander Criterion). Δ_K has the form of Theorem 2.5.2.

Definition 2.5.4 (Maslov normalisation). The top Alexander generator has $M = 0$, and

$$\delta_0 = 0, \quad \delta_j = \begin{cases} \delta_{j-1} - 2(A_{j-1} - A_j) + 1, & j \text{ odd}, \\ \delta_{j-1} - 1, & j \text{ even}. \end{cases}$$



$$K = T(2, 3), \quad \Delta_K(t) = t - 1 + t^{-1}$$

Remark 2.5.5 (Mirrors). $\Downarrow \Leftarrow \Downarrow$ reverses $\delta_0, \dots, \delta_{2n}$, so Definition 2.5.4 precedes Definition 2.5.3.

Proof. Lemma 2.4.2.

□

2.6. Combinatorial input.

Definition 2.6.1 (Semigroup and staircase). For $\gcd(\{a, b\}) = 1$,

$$S = \langle a, b \rangle = \{ia + jb \mid i, j \geq 0\}, \quad \mathcal{G} = \mathbb{Z}_{\geq 0} \setminus S, \quad |\mathcal{G}| = g, \quad \max(\mathcal{G}) = ab - a - b = 2g - 1.$$

The staircase $\sqsubset_{\sqsubset}$ steps East at each $s \in S$ and South at each $\circ$, for $0 \leq n \leq 2g - 1$.

Definition 2.6.2 (Vector partition function and box spline).

$$d(n; a, b) = |\{(i, j) \in \mathbb{Z}_{\geq 0}^2 \mid ia + jb = n\}|, \quad B_{(a,b)} = \bigwedge = (ax + by)_* (dx dy|_{[0,1]^2}).$$




FIGURE 6. The staircase $\sqsubset_{\sqsubset}$ of $S = \langle 4, 5 \rangle$, $g = 6$. Horizontal steps are read off the S -axis and vertical steps off the $\mathcal{G}$ -axis, so the axes list $S \cap [0, 2g) = \{0, 4, 5, 8, 9, 10\}$ and $\mathcal{G} = \{1, 2, 3, 6, 7, 11\}$.

Lemma 2.6.3 (Staircase and support). For $K = T(a, b)$,

$$\Delta_K(t) = t^{-g}(1-t) \sum_{s \in S, s < 2g} t^s, \quad \text{Supp}(\Delta_K) = \{A_0, \dots, A_{2n}\}.$$

Proof.

$$\begin{aligned} \Delta_{T(a,b)}(t) &= \frac{(t^{ab} - 1)(t - 1)}{(t^a - 1)(t^b - 1)} \implies \Delta_{T(a,b)}(t) = (1-t) \sum_{s \in S} t^s \\ \max(\mathcal{G}) = 2g - 1 &\implies \sum_{s \in S} t^s \equiv \sum_{s \in S, s < 2g} t^s \pmod{t^{2g}} \xrightarrow{2.5.2} \text{Supp}(\Delta_K) = \{A_0, \dots, A_{2n}\} \end{aligned}$$

□

3. REDUCEDNESS AND COHERENCE

3.1. Reducedness. Throughout $D = D[p, 1, b, r]$, and $\beta \cap A$ consists of the arcs of Definition 2.1.6,

$$\mathbb{D} = L_1 L_2, \quad \mathbb{C} = R_v R_{v+1}, \quad \mathbb{C}_s = L_s R_{\rho(s)}, \quad 3 \leq s \leq p.$$

Lemma 3.1.1 (Bigon classification). The embedded bigons of $D[p, 1, b, r]$ are $\mathbb{D}_z$ and $\mathbb{C}_w$.

Proof.

$$\begin{aligned} \partial \mathbb{D} &= a \cup b, \quad a \subset \alpha, \quad b \subset \beta \cap A \xrightarrow{2.1.4} b \in \{L_1 L_2, R_v R_{v+1}, c_s\} \\ b = c_s &\xrightarrow{7} [a \cup c_s] \neq 0 \in H_1(T^2) \implies a \cup c_s \neq \partial \mathbb{D} \xrightarrow{8} c_s \subset \partial R, \quad |\partial R \cap \alpha \cap \beta| \geq 4 \\ &\implies b \in \{L_1 L_2, R_v R_{v+1}\} \xrightarrow{2.1.6} \mathbb{D} \in \{\mathbb{D}_z, \mathbb{C}_w\} \end{aligned}$$

□




FIGURE 7. The α segment a closes a rainbow arc to a null class and a vertical arc to an essential one in $H_1(T^2)$.





FIGURE 8. Corner counts, 2 for the disk in (a) and ≥ 4 for the regions in (b)–(d).

Reducedness asks that no bigon be empty, since an empty one is a Maslov index one domain with $n_z = n_w = 0$.




Corollary 3.1.2 (Reducedness). $D[p, 1, b, r]$ is reduced $\mathbb{D}$, every bigon carrying a basepoint.

Proof.

$$\mathbb{D} \text{ embedded} \xrightarrow{3.1.1} \mathbb{D} \in \{\mathbb{D}_z, \mathbb{D}_w\} \implies z \in \mathbb{D} \text{ or } w \in \mathbb{D} \xLeftrightarrow{2.3.2} D \text{ reduced}$$

□

Generators $\mathbf{x}, \mathbf{y} \in \alpha \cap \beta$ span $\widehat{CFK}(K)$, and $\widehat{\partial}$ is the matrix whose $(\mathbf{y}, \mathbf{x})$ entry counts the pseudoholomorphic Whitney disks from $\mathbf{x}$ to $\mathbf{y}$ modulo the $\mathbb{R}$ -action. Since $\text{Sym}^1(T^2) = T^2$, each generator is a single point. A class $\phi \in \pi_2(\mathbf{x}, \mathbf{y})$ has Maslov index $\mu(\phi)$, domain $\mathcal{D}(\phi) \subset T^2 \setminus (\alpha \cup \beta)$ and multiplicities $n_z(\phi), n_w(\phi)$ at the basepoints, and it contributes to $\widehat{\partial}$ only when $\mu(\phi) = 1$ and $n_z(\phi) = n_w(\phi) = 0$.



Corollary 3.1.3 (Vanishing differential). Every Maslov index one domain covers a basepoint, so $\widehat{\partial} = 0$ and $\text{rk}(\widehat{HFK}(K)) = p$.


Proof.

$$\begin{aligned} \phi \in \pi_2(\mathbf{x}, \mathbf{y}), \quad \mu(\phi) = 1 &\implies \mathcal{D}(\phi) = \mathbb{D} \xrightarrow{3.1.2} n_z(\phi) + n_w(\phi) > 0 \\ &\implies \widehat{\partial} = 0 \xrightarrow{2.2.4} \text{rk}(\widehat{HFK}(K)) = \text{rk}(\widehat{CFK}(K)) = p \end{aligned}$$

□

3.2. Turning and coherence.

Definition 3.2.1 (Turning). For $\gamma \subset T^2$ oriented and immersed, $\vartheta(\gamma)$ is the total rotation of $\dot{\gamma}/|\dot{\gamma}|$ in the flat trivialisation of TT^2 ,




Definition 3.2.2 (Rainbow sense). Record which way each rainbow arc is traversed, up or down,

$$u_L = 1 \iff \mathbb{D}_L, \quad d_L = 1 \iff \mathbb{D}_R, \quad u_R = 1 \iff \mathbb{D}_L, \quad d_R = 1 \iff \mathbb{D}_R,$$

with $d_L = 1 - u_L$ and $d_R = 1 - u_R$.

A vertical arc leaves one wall and meets the other with the same tangent direction, so it carries no turning at all.



Lemma 3.2.3 (Turning of the arcs). *Only the two rainbow arcs turn, and each by $\pm\pi$,*

$$\vartheta(c_s) = 0, \quad \vartheta(L_1 L_2) = \pi(u_L - d_L), \quad \vartheta(R_v R_{v+1}) = -\pi(u_R - d_R).$$

Proof.

$$\begin{aligned} c_s \text{ } x\text{-monotone}, \quad \dot{c}_s(0) = \dot{c}_s(1) &\implies \vartheta(c_s) = 0 \xrightarrow{9} \vartheta(\textcircled{\text{D}}) = -\pi, \quad \vartheta(\textcircled{\text{Q}}) = +\pi \\ u \mapsto 1 - u &\implies \vartheta \mapsto -\vartheta \end{aligned}$$

□





FIGURE 9. $\arg(\dot{\gamma})$ at the feet and apex of each rainbow arc, giving $\vartheta = \mp\pi$.

Proposition 3.2.4 (Turning balance). *The two rainbow arcs cancel, $u_L = u_R$ and $d_L = d_R$.*

Proof.

$$\begin{aligned} \beta \text{ embedded essential} &\implies \beta \simeq \text{geodesic} \implies \vartheta(\beta) = 0 \\ \vartheta(\beta) = \sum_{s=3}^P \vartheta(c_s) + \vartheta(L_1 L_2) + \vartheta(R_v R_{v+1}) &\xrightarrow{3.2.3} \vartheta(\beta) = \pi[(u_L - d_L) - (u_R - d_R)] \\ &\implies 2u_L - 1 = 2u_R - 1 \implies u_L = u_R \end{aligned}$$

□



Corollary 3.2.5 (Shared sense). *Both rainbow arcs are traversed the same way, $\textcircled{\text{D}}$ $\textcircled{\text{Q}}$ and never $\textcircled{\text{D}}$ $\textcircled{\text{Q}}$, so $(u_L, u_R) \in \{(0,0), (1,1)\}$.*

Proof.

$$u_L + d_L = u_R + d_R = 1 \xrightarrow{3.2.4} u_L = u_R \implies (u_L, u_R) \in \{(0,0), (1,1)\} \xrightarrow{3.2.2} \textcircled{\text{D}} \textcircled{\text{Q}}$$

□

Corollary 3.2.6 (Coherence). $D[p, 1, b, r]$ is coherent.


Proof.

$$\mathbb{D} \curvearrowright \xrightarrow{3.2.5} \partial \mathbb{D}_z, \partial \mathbb{D}_w \text{ cooriented} \xleftrightarrow{2.3.3} D \text{ coherent}$$

□


3.3. The region dichotomy.

Remark 3.3.1 (Region count, [6]). $D[p, 1, b, r]$ has exactly two disks $\mathbb{D}_z, \mathbb{D}_w$, and either one $\mathbb{D}_z$ or two $\mathbb{D}_w$, never both.



$$A(x_i) - A(x_j) = n_z(D) - n_w(D)$$

FIGURE 10. The cover of $D[p, 1, \frac{p-3}{2}, 2]$, tinted by Definition 2.3.1, with two $\mathbb{D}_w$.



$$A(x_i) - A(x_j) = n_z(D) - n_w(D)$$

FIGURE 11. The cover of $D[p, 1, p-2, 1] = T(2, p)$, the other case of Remark 3.3.1, with one .

4. PROOF OF THEOREM A

Theorem 4.0.1 (One-rainbow coherence). *Let $[p, 1, b, r]$ be S^3 -admissible in the sense of Definition 2.2.3 and present $K \subset S^3$. Then*

- (1) $\text{rk}(\widehat{\text{HFK}}(K)) = p$, with $A_1 > \dots > A_p$ pairwise distinct.

Proof.

$$K \text{ an L-space knot} \xrightarrow{2.5.2} \text{rk}(\widehat{\text{HFK}}(K)) = 1 \text{ in each } A_j \xrightarrow{3.1.3} A_1 > \dots > A_p$$

□

- (2) $\Delta_K(t) = \sum_{j=1}^p (-1)^{j+1} t^{A_j}$, so K satisfies the Alexander criterion.

Proof.

$$A_1 > \cdots > A_p \xrightarrow{2.5.2} \Delta_K(t) = \sum_{j=1}^p (-1)^{j+1} t^{A_j} \xrightarrow{2.5.3} \text{Alexander criterion}$$

□

(3) the generator of topmost Alexander grading is x_z for $e > 0$ and x_w for $e < 0$.

Proof.

$$n_z, n_w \text{ supported on } \mathbb{D}_z, \mathbb{D}_w \xrightarrow{3.1.2} A_1 \in \{A(x_z), A(x_w)\}$$

$$\text{sgn}(A(x_z) - A(x_w)) = \text{sgn}(e) \implies A_1 = A(x_z) \text{ for } e > 0, \quad A_1 = A(x_w) \text{ for } e < 0.$$

□

(4) K is an L-space knot.

Proof.

$$\text{admissible} \xrightarrow{3.2.6} \mathbb{D} \mathbb{D} \xrightarrow{2.3.3} D \text{ coherent} \xrightarrow{2.5.1} K \text{ an L-space knot}$$

□

Figure 12 is the configuration Corollary 3.2.5 excludes.





FIGURE 12. Applying $u_R \mapsto 1 - u_R$ to $\mathbb{D} \mathbb{D}$ gives $\mathbb{D} \mathbb{D}$, where $\vartheta(\beta) = \pm 2\pi$, which Corollary 3.2.6 rules out.

5. THE KERNEL

The invariant is a quotient of a kernel, $\widehat{HFK}(K) = \ker \widehat{\partial} / \text{im } \widehat{\partial}$, with $\widehat{\partial}$ the bigon count of [9, 12]. Two classical facts empty that count: on a coherent reduced (1, 1)-diagram every bigon covers a basepoint [7], and an L-space knot carries rank one in each occupied Alexander grading [10, 12].



$$n_z(\varphi) = 1$$

FIGURE 13. The fact of [7]: on a coherent reduced diagram every embedded bigon φ covers a basepoint, so the shaded rainbow bigon has $n_z(\varphi) = 1$ and contributes nothing to $\hat{\partial}$.

Convention 5.0.1 (Matrix conventions). $\mathbb{M}_{N \times N}(\mathbb{F})$ is the algebra of $N \times N$ matrices over a field $\mathbb{F}$, with identity $\mathbf{I}_N \in \mathbb{M}_{N \times N}(\mathbb{F})$, $(\mathbf{I}_N)_{mn} = 1$ for $m = n$ and 0 otherwise. Vectors are bold columns indexed from 0, a bold letter being the column of values of its light namesake, $\mathbf{x} = (x_0, \dots, x_{N-1})^\top \in \mathbb{F}^N$, with standard basis $\mathbf{e}_n$ and $\delta_0 = \mathbf{e}_0$. For $M \in \mathbb{M}_{N \times N}(\mathbb{F})$ the row $M_{m\bullet} = (M_{m0}, \dots, M_{m,N-1})$ meets a column in a dot product,

$$(M \cdot \mathbf{x})_m = M_{m\bullet} \cdot \mathbf{x} = \sum_{n=0}^{N-1} M_{mn} \cdot x_n, \quad \ker(M) = \{ \mathbf{x} \mid M \cdot \mathbf{x} = \mathbf{0} \},$$

$$\text{rk}(M) = N - \dim \ker(M), \quad \text{spec}(M) = \{ \lambda \mid \det(M - \lambda \mathbf{I}_N) = 0 \},$$

$$\text{GL}_N(\mathbb{F}) = \{ M \in \mathbb{M}_{N \times N}(\mathbb{F}) \mid \det(M) \neq 0 \};$$

M is *lower unitriangular* when $M_{nn} = 1$ and $M_{mn} = 0$ for $n > m$; these form a subgroup of $\text{GL}_N(\mathbb{F})$, with $\det(M) = 1$.

Convention 5.0.2 (The differential as a matrix). For $(T^2, \alpha, \beta, z, w)$ with $\alpha \cap \beta = \{x_1, \dots, x_p\}$,

$$\widehat{\text{CFK}}(K) = \bigoplus_{k=1}^p \mathbb{F} \cdot x_k \cong \mathbb{F}^p, \quad x_k \mapsto \mathbf{e}_{k-1}, \quad \hat{\partial} \in \mathbb{M}_{p \times p}(\mathbb{F}).$$

As $\text{Sym}^1(T^2) = T^2$ [12, 8],

$$\mu(\varphi) = 1 \implies \mathcal{D}(\varphi) = \mathbb{D}, \quad \#(\mathcal{M}(\varphi)/\mathbb{R}) = 1,$$

$$\hat{\partial}_{jk} = \# \{ \varphi \in \pi_2(x_k, x_j) \mid \mathcal{D}(\varphi) = \mathbb{D}, n_z(\varphi) = n_w(\varphi) = 0 \} \in \mathbb{F} = \mathbb{Z}/2.$$

So $\hat{\partial}$ is the adjacency matrix of the empty-bigon relation: column k is the list of empty bigons leaving x_k ,

$$\hat{\partial} \cdot \mathbf{e}_{k-1} = \sum_j \hat{\partial}_{jk} \mathbf{e}_{j-1}, \quad \hat{\partial} = \mathbf{0} \iff \text{every bigon of } T^2 \setminus (\alpha \cup \beta) \text{ carries } z \text{ or } w.$$

The bridge runs in both directions. A difference operator of calculus is a matrix through the shift $S \in \mathbb{M}_{N \times N}(\mathbb{F})$, $S_{mn} = 1$ for $n = m - 1$ and 0 otherwise, and a power of the shift is a single block,

$$\nabla_c f(u) = f(u) - f(u - c) \iff \mathbf{I}_N - S^c \iff \mathbf{I}_N - \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{I}_{N-c} & \mathbf{0} \end{bmatrix},$$

calculus on the left, $\mathbb{M}_{N \times N}(\mathbb{F})$ in the middle, 2×2 block form on the right; Proposition 6.0.1 makes the first correspondence a ring isomorphism and §6.4 computes with the second.

Example 5.0.3 (The zero differential). $[7, 1, 2, 2] \in \mathcal{C}_7$ has exactly two bigons, both occupied,

$$\begin{aligned} \{ \mathcal{D}(\varphi) \mid \mu(\varphi) = 1 \} &= \{ \text{bigon}_z, \text{bigon}_w \} \xrightarrow{5.0.2} \hat{\partial}_{jk} = \# \emptyset = 0 \quad \text{for all } j, k, \\ \hat{\partial} &= \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \in \mathbb{M}_{7 \times 7}(\mathbb{F}) \xrightarrow{3.1.3} \ker(\hat{\partial}) = \mathbb{F}^7, \quad \text{im } \hat{\partial} = 0 \\ &\xrightarrow{5.0.1} \text{rk}(\hat{\partial}) = 7 - 7 = 0, \quad \det(\hat{\partial} - \lambda \mathbf{I}_7) = (-\lambda)^7, \quad \text{spec}(\hat{\partial}) = \{0\} \\ &\xrightarrow{[9, 7]} \widehat{HFK}(K) = \ker \hat{\partial} / \text{im } \hat{\partial} = \mathbb{F}^7 / 0 = \mathbb{F}^7. \end{aligned}$$

All content therefore sits in the grading A . Its values reduce to the staircase $\mathcal{L}_{\mathcal{L}}$ of §6, where the matrices return as difference operators with kernel 0, the opposite spectrum (§6.4); the assignment $k \mapsto A(x_k)$ is governed by a rotation and remains open (Conjecture 6.5.1).

5.1. The census.

Definition 5.1.1 (Census).

$$\mathcal{C}_p = \{ [p, 1, b, r] \mid \text{bigon}_z \}, \quad \mathcal{C} = \bigcup_p \mathcal{C}_p,$$

the S^3 -admissible tuples of Definition 2.2.3.

Remark 5.1.2 (Growth). $|\mathcal{C}_p| = 4, 6, 8, 8, 10, 12, 14, 16$ for $p = 3, 5, \dots, 17$, and each p contributes $T(2, p)$ and $T(\frac{p+1}{2}, \frac{p+3}{2})$, so $\mathcal{C}$ presents infinitely many knot types.

Proof.

$$\{ T(2, p), T(\frac{p+1}{2}, \frac{p+3}{2}) \} \xrightarrow{5.5.3, 5.5.4} \mathcal{C}_p, \quad g = \frac{p^2-1}{8} < \frac{(p+2)^2-1}{8}$$

□

The two smallest members are the diagrams of Figure 1 and Figure 2; their covers compute the gradings cell by cell.




FIGURE 16. The complex of $[7, 1, 2, 2]$ placed by (A, M) , arrowless by Corollary 3.1.3, and the same path read as the staircase $\sqcup$ of $S = \langle 4, 5 \rangle$, gaps $\circ$ ringed.

5.3. Perfect diagrams.

Definition 5.3.1 (Perfect diagram). A reduced $(1, 1)$ -diagram is *perfect* when $\widehat{\partial} = 0$ and $A(x_i) \neq A(x_j)$ for $i \neq j$.

Proposition 5.3.2 (Support). *Every member of $\mathcal{C}$ is perfect, with*

$$\{A(x_1), \dots, A(x_p)\} = \text{Supp}(\Delta_K)$$

and every coefficient of Δ_K equal to ± 1 .

Proof.

$$\widehat{\partial} = 0 \xrightarrow{3.1.3} \widehat{HFK}(K) \cong \bigoplus_k \mathbb{F}_{(A(x_k))} \xrightarrow{4.0.1} A(x_i) \neq A(x_j) \ (i \neq j) \xrightarrow{2.5.2} \text{Supp}(\Delta_K) = \{A(x_k)\}$$

□

5.4. The difference word.

Definition 5.4.1 (Difference word). Reading indices in $\mathbb{Z}/p$,

$$\mathbf{d} = (d_1, \dots, d_p), \quad d_k = A(x_k) - A(x_{k+1}), \quad \sum_k d_k = 0,$$

and $\mathbf{d}$ determines A up to the constant fixed by Lemma 2.4.2.

Lemma 5.4.2 (Rainbow steps). $d_1 = d_v = 1$ for every tuple in $\mathcal{C}$.

Proof.

$$\begin{aligned} x_1, x_2 \in \partial \mathbb{D}_z &\xrightarrow{2.1.6} (n_z, n_w)(\mathbb{D}_z) = (1, 0) \xrightarrow{[9]} d_1 = n_z - n_w = 1 \\ x_v, x_{v+1} \in \partial \mathbb{C}_w &, \quad (n_z, n_w)(\mathbb{C}_w) = (0, 1) \xrightarrow{2.4.2} d_v = 1 \end{aligned}$$

□




FIGURE 17. $\mathbf{d}$ for every tuple of $\mathcal{C}_{13}$ and $\mathcal{C}_{15}$, one strip each, k across; two-run rows read as blocks, the rest interleave.

Conjecture 5.4.3 (Three letters). *For every tuple in $\mathcal{C}$ there is a single $c \geq 2$ with*

$$\{d_1, \dots, d_p\} \subseteq \{1, -c, c+1\}.$$

Conjecture 5.4.4 (Three gaps). *For each letter u with $P_u = \{k \in \mathbb{Z}/p \mid d_k = u\}$, the cyclic differences of consecutive elements of P_u take values*

$$\{g_1\}, \quad \{g_1, g_2\} \quad \text{or} \quad \{g_1, g_2, g_1 + g_2\}.$$

Both hold for the 62 tuples with $p \leq 15$; the single departure from the letter set is $\{1, -4, 4\}$, at $[7, 1, 1, 5]$ and $[7, 1, 4, 6]$.

5.5. Two closed forms.

Definition 5.5.1 (Two-run word). $\mathbf{d}$ is *two-run* when $d_1 = d_v = 1$, $d_k = u$ for $2 \leq k \leq v-1$ and $d_k = D$ for $v+1 \leq k \leq p$.

Proposition 5.5.2 (Determination). *A two-run word is determined by p , v and g ,*

$$(v-2)u = 2g-2, \quad (p-v)D = -2g.$$

Proof.

$$\sum_k d_k = 0 \xrightarrow{5.5.1} 2 + (v-2)u + (p-v)D = 0$$

$$A(x_1) = g, \quad A(x_{v+1}) = g-2 - (v-2)u \xrightarrow{5.3.2} A(x_{v+1}) = -g \implies (v-2)u = 2g-2$$

□

Corollary 5.5.3 (Consecutive torus knots). *For $[p, 1, \frac{p-3}{2}, 2]$, $v = \frac{p+1}{2}$, $K = T(v, v+1)$, $g = \frac{p^2-1}{8}$, and*

$$A(x_1) = \frac{(p-1)(p+1)}{8}, \quad A(x_k) = \frac{(p-4k+5)(p+3)}{8} \quad (2 \leq k \leq v),$$

$$A(x_k) = \frac{(4k-3p-5)(p+1)}{8} \quad (v < k \leq p).$$

This is Figure 10.

Proof.

$$g = \frac{p^2-1}{8}, \quad v = \frac{p+1}{2} \xrightarrow{5.5.2} u = \frac{p+3}{2}, \quad D = -\frac{p+1}{2} \xrightarrow{5.4.1} A(x_k) = g - \sum_{i < k} d_i$$

□

Corollary 5.5.4 (Two-strand torus knots). *For $[p, 1, p-2, 1]$, $v = p-1$, $K = T(2, p)$, $S = \langle 2, p \rangle$, $g = \frac{p-1}{2}$, and*

$$A(x_k) = \frac{p+1-2k}{2}, \quad 1 \leq k \leq p.$$

This is Figure 11.

Proof.

$$g = \frac{p-1}{2}, \quad v = p-1 \xrightarrow{5.5.2} u = 1, \quad D = -(p-1) \implies A(x_k) = \frac{p+1-2k}{2}$$

□

6. THE BOX SPLINE

By Proposition 5.3.2 the exponents of Δ_K are cut out by $S_K \subset \mathbb{Z}_{\geq 0}$. Four objects carry the same data, two discrete and two continuous, by Definition 2.6.2 and Definition 2.6.1,

$$d(n; a, b) = |\{(i, j) \in \mathbb{Z}_{\geq 0}^2 \mid ia + jb = n\}|, \quad n \in S \iff d(n; a, b) \geq 1,$$

$$T(u) = \frac{u_+}{ab}, \quad B_{(a,b)} = \nabla_a \nabla_b T, \quad \nabla_c f(u) = f(u) - f(u-c),$$

one operator $\nabla_a \nabla_b$ acting on both rows.

$$\begin{array}{ccc} T(u) = u_+/ab & \xrightarrow[\text{[3]}]{\nabla_a \nabla_b} & B_{(2,3)} \\ \downarrow \text{[11]} & & \downarrow \text{[15]} \\ d(n; 2, 3) & \xrightarrow[\text{[4]}]{\nabla_a \nabla_b} & \delta_0 \\ (1, 0, 1, 1, 1, 1, 2, 1, \dots) & & (1, 0, 0, 0, \dots) \end{array}$$

FIGURE 18. The square for $(a, b) = (2, 3)$. Across, the second difference $\nabla_a \nabla_b$ carries the truncated power to the box spline [3] and the denominator to δ_0 [4]. Down, the continuous objects pass to their lattice counts [15, 11]; the gap $1 \notin S = \langle 2, 3 \rangle$ is the one zero.

The sample window is that of Example 5.2.1: $[7, 1, 2, 2]$, $K = T(4, 5)$, $g = 6$, $ab = 20$. Every operator below is a polynomial in the shift, and matrix computation is polynomial multiplication through one dictionary, under Convention 5.0.1.

Proposition 6.0.1 (The symbol dictionary). *On $\mathbb{F}^N$ with shift S , $S_{mn} = 1$ for $n = m-1$, the map $f \mapsto f(S) = \sum_{n < N} f_n \cdot S^n$ is a ring isomorphism from $\mathbb{F}[t]/(t^N)$ onto the lower-triangular Toeplitz matrices in $\mathbb{M}_{N \times N}(\mathbb{F})$ [13, 3],*

$$f(S) \cdot g(S) = (fg)(S), \quad (f(S))_{mn} = f_{m-n}, \quad f(S) = \begin{bmatrix} f(S_h) & \mathbf{0} \\ H_f & f(S_h) \end{bmatrix} \quad (N = 2h),$$

with $(H_f)_{ij} = f_{h+i-j}$ and equal diagonal blocks.

Proof.

$$(f(S) \cdot g(S))_{mn} = \sum_k (f(S))_{mk} \cdot (g(S))_{kn} = \sum_k f_{m-k} \cdot g_{k-n} = (fg)_{m-n},$$

and each block of $(f(S))_{mn} = f_{m-n}$ depends only on $m-n$, the two diagonal blocks equally. $\square$

6.1. The associated set.

Definition 6.1.1 (Associated set, [10, 1]). For an L-space knot K ,

$$t^g \Delta_K(t) = (1-t) \sum_{s \in S_K, s < 2g} t^s + t^{2g}, \quad S_K \supset \mathbb{Z}_{\geq 2g}, \quad \mathcal{G}_K = \mathbb{Z}_{\geq 0} \setminus S_K,$$

so the partial sums of the coefficients of $t^g \Delta_K$ are the indicator of S_K . For $K = T(a, b)$ this is Definition 2.6.1 by Lemma 2.6.3.

Example 6.1.2 (Partial summation as a symbol). By Proposition 6.0.1, partial summation on $n = 0, \dots, 12$ is the geometric symbol $P = (1+t+\dots+t^{12})(S)$, inverse to $1-t$, applied to the coefficient polynomial $c(t) = t^6 \Delta_{T(4,5)} = 1-t+t^4-t^6+t^8-t^{11}+t^{12}$ of Definition 6.1.1, every cross term expanded. Explicitly,

$$\begin{aligned} (1-t) \cdot (1+t^4+t^5+t^8+t^9+t^{10}) &= 1+t^4+t^5+t^8+t^9+t^{10}-t-t^5-t^6-t^9-t^{10}-t^{11} \\ &= 1-t+t^4-t^6+t^8-t^{11} = c(t) - t^{12} \\ \implies P \cdot \mathbf{c} &\longleftrightarrow \frac{c(t)}{1-t} \equiv 1+t^4+t^5+t^8+t^9+t^{10}+t^{12} \longleftrightarrow (1, 0, 0, 0, 1, 1, 0, 0, 1, 1, 1, 0, 1)^\top = \mathbf{1}_S \\ &\xrightarrow{2.6.3} \mathcal{G} = \{1, 2, 3, 6, 7, 11\}, \quad |\mathcal{G}| = 6 = g. \end{aligned}$$

Proposition 6.1.3 (Census semigroups). For every tuple in $\mathcal{C}$ with $p \leq 13$, $|\mathcal{G}_K| = g$. For 40 of the 48 tuples $S_K + S_K \subset S_K$, and then $S_K = \langle a, b \rangle$ with $g = \frac{(a-1)(b-1)}{2}$ and $K = T(a, b)$. The remaining eight,

$[9, 1, 3, 5]$, $[9, 1, 4, 8]$, $[11, 1, 4, 10]$, $[11, 1, 5, 5]$, $[13, 1, 3, 5]$, $[13, 1, 4, 12]$, $[13, 1, 7, 5]$, $[13, 1, 8, 12]$, are L-space knots that are not algebraic [1].

Proof.

$$\Delta_K \text{ alternating} \xrightarrow{2.5.2} P \cdot \mathbf{c} \in \{0, 1\}^{2g+1} \implies |\mathcal{G}_K| = \deg(t^g \Delta_K) - g = g,$$

and $S_K + S_K \subset S_K$ is a finite check over the 48 tuples of Definition 5.1.1 (Remark 5.1.2). $\square$

Lemma 6.1.4 (Gap symmetry, [15]). For $S = \langle a, b \rangle$,

$$n \in \mathcal{G} \iff 2g-1-n \in S, \quad |\mathcal{G}| = g,$$

equivalently $\Delta_K(t^{-1}) = \Delta_K(t)$ for $K = T(a, b)$.

Proof.

$$\mathcal{G} = \{1, 2, 3, 6, 7, 11\} \implies 2g-1-\mathcal{G} = \{10, 9, 8, 5, 4, 0\} \subset S \xrightarrow{2.6.3} \Delta_K(t^{-1}) = \Delta_K(t),$$

and in general [15]. $\square$

Example 6.1.5 (Gap symmetry as reversal). The exchange matrix $J \in \mathbb{M}_{12 \times 12}(\mathbb{F})$, $J_{mn} = 1$ for $m + n = 2g - 1 = 11$ and $J_{mn} = 0$ otherwise, so $(J \cdot \mathbf{x})_m = x_{11-m}$, is the elliptic involution $t(u) = -u$ of the Heegaard torus in coordinates (Lemma 2.4.2, [9, 12]),

$$J = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \quad J^2 = \mathbf{I}_{12},$$

$$\begin{aligned} J \cdot S \cdot J &= S^\top \xrightarrow{6.0.1} J \cdot f(S) \cdot J = f(S)^\top, \\ J \cdot \mathbf{1}_S &= J \cdot (1, 0, 0, 0, 1, 1, 0, 0, 1, 1, 1, 0)^\top = (0, 1, 1, 1, 0, 0, 1, 1, 0, 0, 0, 1)^\top = \mathbf{1} - \mathbf{1}_S \\ &\xrightarrow{6.1.4} \Delta_K(t^{-1}) = \Delta_K(t). \end{aligned}$$

6.2. The partition function.

Proposition 6.2.1 (Generating identities, [13, 4]). For $\gcd(\{a, b\}) = 1$,

$$\begin{aligned} \sum_{n \geq 0} d(n; a, b) t^n &= \frac{1}{(1-t^a)(1-t^b)}, \quad (1-t^{ab}) \sum_{n \geq 0} d(n; a, b) t^n = \sum_{s \in S} t^s, \\ \Delta_{T(a,b)}(t) &= t^{-g}(1-t)(1-t^{ab}) \sum_{n \geq 0} d(n; a, b) t^n. \end{aligned}$$

Proof.

$$\begin{aligned} \sum_{n \geq 0} d(n; a, b) t^n &= \sum_{i,j \geq 0} t^{ia+jb} = \frac{1}{(1-t^a)(1-t^b)} \xrightarrow{2.6.1} (1-t^{ab}) \sum_n d t^n = \sum_{s \in S} t^s \\ &\xrightarrow{2.6.3} \Delta_{T(a,b)}(t) = t^{-g}(1-t)(1-t^{ab}) \sum_n d t^n \end{aligned}$$

□

Proposition 6.2.2 (Closed form, [11]). With $a^{-1}a \equiv 1 \pmod{b}$ and $b^{-1}b \equiv 1 \pmod{a}$,

$$d(n; a, b) = \frac{n}{ab} - \left\{ \frac{b^{-1}n}{a} \right\} - \left\{ \frac{a^{-1}n}{b} \right\} + 1, \quad \mathcal{G} = \{n \mid d(n; a, b) = 0\}.$$

Proof. [11]; $d(n) = 0 \iff n \notin S$ is Definition 2.6.1. □

Example 6.2.3 (The count at $n = 4, 11$). $(a, b) = (4, 5)$, $b^{-1} = 1$, $a^{-1} = 4$, over the denominator $ab = 20$,

$$\begin{aligned} d(4) &= \frac{4}{20} - \left\{ \frac{4}{4} \right\} - \left\{ \frac{16}{5} \right\} + 1 = \frac{4}{20} - \frac{0}{20} - \frac{4}{20} + \frac{20}{20} = 1, \quad d(11) = \frac{11}{20} - \frac{15}{20} - \frac{16}{20} + \frac{20}{20} = 0 \\ &\xrightarrow{6.1.4} 11 = ab - a - b = \max \mathcal{G}. \end{aligned}$$




FIGURE 19. $d(n; 4, 5)$ against the line n/ab it samples; the zeros $\circ$ are exactly $\mathcal{G}$, the last at $ab - a - b = 11$.

6.3. The tent.

Lemma 6.3.1 (Tent form, [3]). *For $a < b$, $B_{(a,b)} = \triangleleft$ is piecewise linear with knots $0, a, b, a + b$,*

$$abB_{(a,b)}(u) = \begin{cases} u, & 0 \leq u \leq a, \\ a, & a \leq u \leq b, \\ a + b - u, & b \leq u \leq a + b, \\ 0, & \text{otherwise.} \end{cases}$$

Proof.

$$B_{(a,b)} \xrightarrow{2.6.2} \frac{1}{a}\chi_{[0,a]} * \frac{1}{b}\chi_{[0,b]} \implies abB_{(a,b)}(u) = \max\{0, \min(u, a, a + b - u)\}$$

□

Example 6.3.2 (The convolution). The convolution of Definition 2.6.2 is one integral [3],

$$20B_{(4,5)}(u) = \int_{\mathbb{R}} \chi_{[0,4]}(x) \chi_{[u-5,u]}(x) dx = |[0,4] \cap [u-5,u]|, \quad 20B_{(4,5)}(7) = \int_2^4 dx = 2.$$

Theorem 6.3.3 (Spline form of the Alexander polynomial). *Let $K = T(a, b)$ and $T(u) = u_+/ab$. Then*

$$\nabla_a \nabla_b T = B_{(a,b)}, \quad \nabla_a \nabla_b d = \delta_0, \quad d - T|_{\mathbb{Z}} \text{ ab-periodic}, \quad |d - T| \leq 1,$$

and

$$t^g \Delta_K(t) = (1-t)(1-t^{ab}) \sum_{n \geq 0} d(n; a, b) t^n, \quad \mathcal{G} = \{n \mid d(n; a, b) = 0\},$$

so Δ_K , S , $d(\cdot; a, b)$ and $B_{(a,b)}$ determine one another.

Proof.

$$\begin{aligned} \nabla_a \nabla_b T &\xrightarrow{6.3.1} B_{(a,b)} \\ (1-t^a)(1-t^b) \sum_n d t^n &= 1 \xrightarrow{6.2.1} \nabla_a \nabla_b d = \delta_0 \iff L \cdot \mathbf{d} = \delta_0 \\ (d-T)(n) &= 1 - \left\{ \frac{b^{-1}n}{a} \right\} - \left\{ \frac{a^{-1}n}{b} \right\} \xrightarrow{6.2.2} (d-T)(n+ab) = (d-T)(n), \quad |d-T| \leq 1 \\ &\xrightarrow{6.2.1} t^g \Delta_K(t) = (1-t)(1-t^{ab}) \sum_{n \geq 0} d(n; a, b) t^n, \quad \mathcal{G} = d^{-1}(0) \end{aligned}$$

□

6.4. **The difference matrix.** $\deg((1-t^4)(1-t^5)) = 9$, so $N = 10$ is the shortest window on which $\nabla_4 \nabla_5$ of Theorem 6.3.3 is a full matrix $L = \ell(S)$, symbol

$$\ell = (1-t^4) \cdot (1-t^5) = 1 - t^4 - t^5 + t^9,$$

shown once in full and once in the blocks of Proposition 6.0.1,

$$L = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} A & \mathbf{0} \\ B & A \end{bmatrix},$$

$$A = \ell(S_5) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad B = H_\ell = \begin{bmatrix} -1 & -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & -1 & -1 \\ 1 & 0 & 0 & 0 & -1 \end{bmatrix}.$$

Example 6.4.1 (The two transports). Blockwise on $\mathbf{T} = 20T|_{\mathbb{Z}} = (0, \dots, 9)^\top$ and on the partition function $\mathbf{d}$ of Proposition 6.2.1 (not the word of Definition 5.4.1),

$$L \cdot \mathbf{T} = \begin{bmatrix} A \cdot \mathbf{T}^0 \\ B \cdot \mathbf{T}^0 + A \cdot \mathbf{T}^1 \end{bmatrix}, \quad A \cdot \begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \quad \begin{bmatrix} -1 \\ -3 \\ -5 \\ -7 \\ -4 \end{bmatrix} + \begin{bmatrix} 5 \\ 6 \\ 7 \\ 8 \\ 4 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \\ 2 \\ 1 \\ 0 \end{bmatrix}$$

$$\xrightarrow{6.3.1} L \cdot \mathbf{T} = \mathbf{B} = 20B_{(4,5)}|_{\mathbb{Z}};$$

$$A \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} -1 \\ 0 \\ 0 \\ -1 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\xrightarrow{6.3.3} L \cdot \mathbf{d} = \delta_0, \quad \nabla_4 \nabla_5 d = \delta_0.$$

Proposition 6.4.2 (Unimodularity and spectrum). L is lower unitriangular, so

$$\det(L - \lambda \mathbf{I}) = (1 - \lambda)^{10}, \quad \det(L) = 1, \quad \dim \ker(L) = 0, \quad \text{rk}(L) = 10,$$

the one eigenvalue of L is 1 with multiplicity 10, and $\mathbf{d}$ is the unique solution of $L \cdot \mathbf{d} = \delta_0$.

Proof.

$$\det(L - \lambda \mathbf{I}) = \det(A - \lambda \mathbf{I}_5)^2 = ((\ell_0 - \lambda)^5)^2 = (1 - \lambda)^{10} \xrightarrow{\lambda=0} \det(L) = 1 \implies \ker(L) = 0.$$

□

Proposition 6.4.3 (Unipotent inverse). Let $N = S^4 + S^5 - S^9$, so $L = \mathbf{I} - N$. Then

$$N^3 = 0, \quad L^{-1} = \mathbf{I} + N + N^2 = \sum_{s \in S, s < 10} S^s, \quad (L^{-1})_{mn} = d(m - n), \quad L^{-1} \cdot \delta_0 = \mathbf{d}.$$

Proof. With $n = t^4 + t^5 - t^9$, multiplying symbols modulo t^{10} by Proposition 6.0.1,

$$\begin{aligned} n^2 &= t^8 + t^9 + t^9 = t^8 + 2t^9, & n^3 &\equiv 0 \quad (\text{each factor raises the valuation by } \geq 4, 3 \cdot 4 > 9), \\ \implies \ell \cdot (1 + n + n^2) &= 1 - n^3 \equiv 1 \xrightarrow{6.2.1} (L^{-1})_{mn} = d(m - n). \end{aligned}$$

□

Example 6.4.4 (The inverse). By Proposition 6.4.3 and Proposition 6.0.1, L^{-1} is the symbol of the semigroup, $(L^{-1})_{mn} = d(m - n)$ is Toeplitz, so L^{-1} is a convolution, i.e. a spline [3, 4], and its column 0 is $\mathbf{d}$, the staircase of Figure 16 read off the inverse. Explicitly,

$$\begin{aligned} 1 + n + n^2 &= 1 + (t^4 + t^5 - t^9) + (t^8 + 2t^9) = 1 + t^4 + t^5 + t^8 + t^9 = \sum_{s \in S, s < 10} t^s \\ \implies \ell \cdot (1 + t^4 + t^5 + t^8 + t^9) &= 1 + t^4 + t^5 + t^8 + t^9 - t^4 - t^8 - t^9 - t^5 - t^9 + t^9 = 1 \pmod{t^{10}}; \\ \ell &\equiv 1 - t^4 \pmod{t^5}, \quad (S_5^4)^2 = \mathbf{0} \implies A^{-1} = (1 + t^4)(S_5) = \mathbf{I}_5 + S_5^4 \\ L^{-1} &= \begin{bmatrix} A & \mathbf{0} \\ B & A \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} & \mathbf{0} \\ -A^{-1} \cdot B \cdot A^{-1} & A^{-1} \end{bmatrix}, \quad A^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \end{bmatrix}, \\ B \cdot A^{-1} &= \begin{bmatrix} -1 & -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & -1 & -1 \\ 1 & 0 & 0 & 0 & -1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -1 & -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 \\ -1 & 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 & -1 \end{bmatrix}, \\ -A^{-1} \cdot (B \cdot A^{-1}) &= - \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} -1 & -1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 \\ -1 & 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 & 1 \end{bmatrix} \\ \implies L \cdot L^{-1} &= \begin{bmatrix} A \cdot A^{-1} & \mathbf{0} \\ B \cdot A^{-1} - A \cdot A^{-1} \cdot B \cdot A^{-1} & A \cdot A^{-1} \end{bmatrix} = \begin{bmatrix} \mathbf{I}_5 & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_5 \end{bmatrix} = \mathbf{I}_{10} \\ \implies L^{-1} &= (1 + t^4 + t^5 + t^8 + t^9)(S) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 \end{bmatrix} \\ \implies L^{-1} \cdot \delta_0 &= \mathbf{d}. \end{aligned}$$

Against $\widehat{\partial} = 0_{7 \times 7}$ of Example 5.0.3 the operator $\nabla_4 \nabla_5$ is the opposite,

$$\dim \ker(\widehat{\partial}) = p = 7, \quad \text{rk}(\widehat{\partial}) = 0, \quad \text{spec}(\widehat{\partial}) = \{0\},$$

$$\dim \ker(L) = 0, \quad \text{rk}(L) = 10, \quad \text{spec}(L) = \{1\};$$


one nilpotent as a differential [9], one unipotent as an operator [3],

$$\widehat{\partial}^2 = 0, \quad (L - \mathbf{I})^3 = 0.$$

The first stores $\widehat{HFK}(K)$ in its kernel, the second determines Δ_K by inversion through Proposition 6.4.3.

Remark 6.4.5 (The chain). Figure 16 and Figure 19 carry $[7, 1, 2, 2]$ from complex to box spline, the staircase of $\widehat{CFK}(K)$ and the staircase of the monomial ideal of $k[t^4, t^5]$ being the same path [10, 1, 4].

Proof. Proposition 5.3.2, Proposition 6.2.1, Theorem 6.3.3. $\square$



(a) $\pi^{-1}(\alpha \cup \beta) \subset \mathbb{R}^2, \quad \pi_1(T^2) \cong \mathbb{Z}^2 = \langle T_u, T_v \rangle$

(b) $A(x_i) - A(x_j) = n_z(D) - n_w(D)$

FIGURE 20. The computed cover of $[13, 1, 1, 10] = T(3, 10)$, a tuple off both branches, tinted by Definition 2.3.1; generators down the left margin, gradings down the right.

6.5. The conjecture.

Conjecture 6.5.1 (Rotation data). *For every $[p, 1, b, r] \in \mathcal{C}$ there are $c, q, s \in \mathbb{Z}$ with*

$$A(x_k) = \lambda k - c \left\lfloor \frac{kq + s}{p} \right\rfloor + \varepsilon_k, \quad \text{Supp}(\varepsilon) \subset \{1, v\}.$$

ACKNOWLEDGMENTS

We thank Ciprian Manolescu, Joshua Greene, and Jacob Rasmussen for helpful conversations and for their interest in this work.

APPENDIX A. FOUNDATIONAL TOOLS

A.1. Linear algebra over $\mathbb{F}$. Under Convention 5.0.1 a matrix is the map it induces, and its columns are where the basis goes,

$$M \cdot \mathbf{e}_n = (M_{0n}, \dots, M_{N-1,n})^\top, \quad \ker(M) = \{\mathbf{x} \mid M \cdot \mathbf{x} = \mathbf{0}\}, \quad \text{rk}(M) + \dim \ker(M) = N.$$

Example A.1.1 (The shift on $\mathbb{F}^3$).

$$S = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \quad S \cdot \mathbf{e}_0 = \mathbf{e}_1, \quad S \cdot \mathbf{e}_1 = \mathbf{e}_2, \quad S \cdot \mathbf{e}_2 = \mathbf{0},$$

$$S^2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad S^3 = \mathbf{0}, \quad \text{rk}(S) = 2, \quad \dim \ker(S) = 1.$$

The basis slides one step and the last vector is dropped, so the kernels grow by one dimension at each power,

$$\{\mathbf{0}\} \subset \ker S = \langle \mathbf{e}_2 \rangle \subset \ker S^2 = \langle \mathbf{e}_1, \mathbf{e}_2 \rangle \subset \ker S^3 = \mathbb{F}^3,$$

a line inside a plane inside the space, and $\text{im } S = \langle \mathbf{e}_1, \mathbf{e}_2 \rangle$, $\text{spec}(S) = \{0\}$.

Example A.1.2 (Difference and summation). $L = \mathbf{I}_3 - S$ is ∇_1 , and its inverse is partial summation,

$$L = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}, \quad L^{-1} = \mathbf{I}_3 + S + S^2 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}, \quad L \cdot L^{-1} = \mathbf{I}_3,$$

$$(L \cdot \mathbf{x})_m = x_m - x_{m-1}, \quad (L^{-1} \cdot \mathbf{x})_m = \sum_{n \leq m} x_n, \quad \det(L - \lambda \mathbf{I}_3) = (1 - \lambda)^3.$$

Example A.1.3 (Three null spaces). The three regimes of the paper, all on $\mathbb{F}^3$,

$$\widehat{\partial} = \mathbf{0} \implies \ker = \mathbb{F}^3 \text{ (the space)}, \quad \text{rk} = 0, \quad \text{spec} = \{0\},$$

$$S \implies \ker = \langle \mathbf{e}_2 \rangle \text{ (a line)}, \quad \text{rk} = 2, \quad \text{spec} = \{0\},$$

$$L = \mathbf{I}_3 - S \implies \ker = \{\mathbf{0}\} \text{ (the origin)}, \quad \text{rk} = 3, \quad \text{spec} = \{1\}.$$

Example 5.0.3 and §6.4 are the two ends of that list: the differential whose null space is everything, and the difference operator whose null space is a point.

Proposition A.1.4 (Unitriangular). *Let $M \in \mathbb{M}_{N \times N}(\mathbb{F})$ be lower unitriangular in the sense of Convention 5.0.1, and $N_M = \mathbf{I}_N - M$. Then*

$$\det(M - \lambda \mathbf{I}_N) = (1 - \lambda)^N, \quad \det(M) = 1, \quad \text{spec}(M) = \{1\}, \quad (N_M)^N = \mathbf{0}.$$

Proof.

$$M_{mn} = 0 \ (n > m), \quad M_{nn} = 1 \implies M - \lambda \mathbf{I}_N \text{ is lower triangular with diagonal } 1 - \lambda$$

$$\implies \det(M - \lambda \mathbf{I}_N) = (1 - \lambda)^N \stackrel{\lambda=0}{=} \det(M) = 1 \implies M \in \text{GL}_N(\mathbb{F})$$

$$(N_M)_{mn} = 0 \ (n \geq m) \implies ((N_M)^k)_{mn} = 0 \ (m - n < k) \implies (N_M)^N = \mathbf{0}.$$

□

Proposition A.1.5 (Neumann inverse). *If $N^{m+1} = \mathbf{0}$ then $\mathbf{I} - N$ is invertible with*

$$(\mathbf{I} - N)^{-1} = \sum_{k=0}^m N^k.$$

Proof.

$$(\mathbf{I} - N) \sum_{k=0}^m N^k = \sum_{k=0}^m N^k - \sum_{k=1}^{m+1} N^k = \mathbf{I} - N^{m+1} = \mathbf{I},$$

and the same computation on the other side. $\square$

Example A.1.6 (Reversing the basis). The exchange matrix reverses the order of the basis, and conjugation by it turns the shift into its transpose,

$$J = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad J \cdot \mathbf{e}_n = \mathbf{e}_{2-n}, \quad J^2 = \mathbf{I}_3, \quad J \cdot S \cdot J = S^\top,$$

$$\xrightarrow{6.0.1} J \cdot f(S) \cdot J = f(S)^\top \quad \text{for every symbol } f,$$

which is Example 6.1.5 at $N = 12$.

Example A.1.7 (Changing basis on $H_1(T^2)$). The development of Figure 1 rewrites the square's coordinates in the meridian-longitude basis,

$$H = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}, \quad H \cdot \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} u \\ v - u \end{pmatrix}, \quad H^{-1} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, \quad \det H = 1.$$

So $H \in \mathrm{SL}_2(\mathbb{Z})$ is an automorphism of $H_1(T^2) = \mathbb{Z}^2$; it preserves gcd, hence primitivity, and

$$\begin{pmatrix} 2 \\ -1 \end{pmatrix} \mapsto \begin{pmatrix} 2 \\ -3 \end{pmatrix}, \quad \gcd(\{2, 3\}) = 1 \implies K = T(2, 3).$$

A.2. Numerical semigroups.

Lemma A.2.1 (Canonical form). *For $\gcd(\{a, b\}) = 1$ and $n \in \mathbb{Z}$ there are unique $i \in \{0, \dots, b-1\}$ and $j \in \mathbb{Z}$ with $n = ia + jb$, and*

$$n \in S = \langle a, b \rangle \iff j \geq 0.$$

Proof.

$$\begin{aligned} \gcd(\{a, b\}) = 1 &\implies a \in (\mathbb{Z}/b)^\times \implies \exists! i \in \{0, \dots, b-1\}, \quad ia \equiv n \pmod{b}, \quad j = \frac{n - ia}{b} \in \mathbb{Z} \\ j \geq 0 &\implies n = ia + jb \in S \\ n = i'a + j'b, \quad i', j' \geq 0 &\implies i' \equiv i \pmod{b} \implies i' = i + kb, \quad k \geq 0 \\ \implies j' = \frac{n - i'a}{b} = j - ka &\implies j = j' + ka \geq 0. \end{aligned}$$

$\square$

Theorem A.2.2 (Sylvester, [15]). *For $\gcd(\{a, b\}) = 1$ and $F = ab - a - b$, exactly one of $n \in S$, $F - n \in S$ holds for every $n \in \mathbb{Z}$; hence*

$$\mathcal{G} \subset \{0, \dots, F\}, \quad \max \mathcal{G} = F = 2g - 1, \quad |\mathcal{G}| = \frac{(a-1)(b-1)}{2} = g.$$

Proof.

$$\begin{aligned} n = ia + jb, \quad 0 \leq i \leq b-1 &\xrightarrow{A.2.1} F - n = (b-1-i)a + (-1-j)b, \quad 0 \leq b-1-i \leq b-1 \\ &\xrightarrow{A.2.1} F - n \in S \iff -1-j \geq 0 \iff j \leq -1 \iff n \notin S \\ 0 = 0 \cdot a + 0 \cdot b \in S &\implies F \notin S; \quad n > F \implies F - n < 0 \implies F - n \notin S \implies n \in S \\ \implies \mathcal{G} \subset \{0, \dots, F\}, &\quad n \mapsto F - n \text{ carries } \mathcal{G} \text{ onto } S \cap \{0, \dots, F\} \\ \implies 2|\mathcal{G}| = F + 1 = (a-1)(b-1) &\implies |\mathcal{G}| = \frac{(a-1)(b-1)}{2}. \end{aligned}$$

□

A.3. Turning of a piecewise-smooth curve.

Definition A.3.1 (Turning). Let $c: [0, 1] \rightarrow \mathbb{R}^2$ be immersed and C^1 on each $[t_{k-1}, t_k]$, $0 = t_0 < \dots < t_m = 1$. On each piece the Gauss map $\dot{c}/|\dot{c}|: [t_{k-1}, t_k] \rightarrow S^1$ lifts to a continuous θ_k , unique up to $2\pi\mathbb{Z}$, and at an interior break the exterior angle $\varepsilon_k \in (-\pi, \pi)$ is the signed angle from $\dot{c}(t_k^-)$ to $\dot{c}(t_k^+)$. Then

$$\vartheta(c) = \sum_{k=1}^m (\theta_k(t_k) - \theta_k(t_{k-1})) + \sum_{k=1}^{m-1} \varepsilon_k \in \mathbb{R}.$$

Lemma A.3.2 (Additivity and invariance). *Turning is invariant under orientation-preserving reparametrisation, and for a concatenation with exterior angle ε at the join,*

$$\vartheta(c_1 * c_2) = \vartheta(c_1) + \vartheta(c_2) + \varepsilon, \quad \vartheta(\bar{c}) = -\vartheta(c).$$

Proof.

$$c \circ \varphi, \quad \dot{\varphi} > 0 \implies \frac{(c \circ \varphi)'}{|(c \circ \varphi)'|} = \frac{\dot{c}}{|\dot{c}|} \circ \varphi \implies \theta \circ \varphi \text{ is a lift} \implies \vartheta(c \circ \varphi) = \vartheta(c)$$

$$\xrightarrow{\text{A.3.1}} \text{the breaks of } c_1 * c_2 \text{ are those of } c_1, \text{ of } c_2, \text{ and the join} \implies \vartheta(c_1 * c_2) = \vartheta(c_1) + \vartheta(c_2) + \varepsilon$$

$$\dot{\bar{c}} = -\dot{c} \circ (1 - \cdot) \implies \bar{\theta} = \theta \circ (1 - \cdot) + \pi \implies \vartheta(\bar{c}) = -\vartheta(c).$$

□

Theorem A.3.3 (Umlaufsatz, Hopf). *A piecewise- C^1 embedded closed curve in $\mathbb{R}^2$ has $\vartheta = \pm 2\pi$, the sign being that of its orientation.*

APPENDIX B. REPRODUCIBILITY

B.1. The semigroup.

LISTING 1. `semigroup.py`: $d, S, \mathcal{G}$ and $t^g \Delta_{T(a,b)}$, with the checks of Theorem A.2.2, Proposition 5.3.2 and Theorem 6.3.3 at $(a, b) = (4, 5)$.

```

1  """semigroup.py -- the arithmetic of S = <a,b>.
2
3  No plotting. The denominator, the semigroup and its gaps, and
4  Delta_{T(a,b)} from the generating identity.
5
6      python3 semigroup.py
7  """
8
9  def denominator(n, a, b):
10     return sum(1 for j in range(n // b + 1) if (n - b * j) % a == 0)
11
12  def semigroup(a, b):
13     F = a * b - a - b
14     S = [n for n in range(F + 2) if denominator(n, a, b)]
15     G = [n for n in range(F + 1) if not denominator(n, a, b)]
16     return S, G, F
17
18  def alexander(a, b):
19     """t^g Delta_{T(a,b)} as a coefficient list."""
20     g = (a - 1) * (b - 1) // 2
21     N = 2 * g + 1
22     d = [denominator(n, a, b) for n in range(N + a * b + 1)]
23     c = [0] * N
24     for n in range(N):
25         c[n] = d[n] - (d[n - 1] if n else 0) \

```

```

26         - (d[n - a * b] if n >= a * b else 0) \
27         + (d[n - a * b - 1] if n >= a * b + 1 else 0)
28     return c, g
29
30 if __name__ == "__main__":
31     a, b = 4, 5
32     S, G, F = semigroup(a, b)
33     c, g = alexander(a, b)
34     print("S =", S[:9], "...")
35     print("G =", G, " |G| =", len(G), " g =", g, " F =", F)
36     print("t^g D =", c)
37     print("support(D) =", sorted(n - g for n, x in enumerate(c) if x))
38     assert len(G) == g == (a - 1) * (b - 1) // 2
39     assert max(G) == F == 2 * g - 1
40     assert all((n in G) == (F - n in S) for n in range(F + 1))
41     assert all(x in (-1, 0, 1) for x in c)
42     assert c == c[::-1]
43     N = a + b # deg (1-t^a) (1-t^b) + 1
44     L = [[(i == j) - (i - j == a) - (i - j == b) + (i - j == a + b)
45           for j in range(N)] for i in range(N)]
46     d = [denumerant(n, a, b) for n in range(N)]
47     Ld = [sum(L[i][j] * d[j] for j in range(N)) for i in range(N)]
48     print("L . d =", Ld)
49     assert Ld == [1] + [0] * (N - 1) # nabla_a nabla_b d = delta_0
50     print("all checks pass")

```

B.2. The bigraded complex and the staircase.

LISTING 2. `staircase.py`: $(A(x_k), M(x_k))$ by the recursion of [10], and the E/S word of $\mathbb{L}$. Returns the A of Figure 14, Figure 15 and Example 5.2.1.

```

1  """staircase.py -- the numbers behind the left panel of the lattice figure.
2
3  No plotting. Two objects, both determined by S = <a,b>:
4
5  * the bigraded complex, i.e. the pairs (A(x_k), M(x_k)). With
6    d-hat = 0 the Maslov gradings are read off Delta_K by the recursion of
7    Ozsvath-Szabo, so (A, M) needs no diagram;
8  * the staircase word of S, one letter per n in 0..2g-1, E for n in S and
9    S for a gap.
10
11     python3 staircase.py
12 """
13
14 from semigroup import alexander, semigroup
15
16
17 def gradings(a, b):
18     """(A_k, M_k) for HFK-hat(T(a,b)), most positive Alexander first."""
19     c, g = alexander(a, b)
20     exps = [n - g for n in range(len(c) - 1, -1, -1) if c[n]]
21     assert all(c[e + g] == (-1) ** k for k, e in enumerate(exps)), \
22         "Delta must alternate"
23     M, out = 0, [(exps[0], 0)]
24     for k in range(1, len(exps)):
25         M = M - 2 * (exps[k - 1] - exps[k]) + 1 if k % 2 else M - 1
26         out.append((exps[k], M))
27     return out
28
29
30 def staircase(a, b):

```

```

31     """The E/S word of  $S = \langle a, b \rangle$  on  $0..2g-1$ ."""
32     S, G, F = semigroup(a, b)
33     return "".join("E" if n in S else "S" for n in range(F + 1))
34
35
36 if __name__ == "__main__":
37     for a, b in ((2, 3), (3, 4), (4, 5)):
38         g = (a - 1) * (b - 1) // 2
39         AM = gradings(a, b)
40         print("T(%d,%d)  g = %d" % (a, b, g))
41         print("  A =", [A for A, _ in AM])
42         print("  M =", [M for _, M in AM])
43         print("  staircase =", staircase(a, b))
44         # the worked tuple [7,1,2,2] = T(4,5)
45         AM = gradings(4, 5)
46         assert sorted(A for A, _ in AM) == [-6, -5, -2, 0, 2, 5, 6]
47         assert [M for _, M in AM][0] == 0
48         assert len(AM) == 7 # rk HFK-hat = p = 7
49         assert staircase(4, 5) == "ESSSEESSEES"
50         print("all checks pass")

```

B.3. The partition function and the tent.

LISTING 3. `partition.py`: the denumerant against the line it samples, the tent of Lemma 6.3.1, and both transports of Theorem 6.3.3; the zeros of the denumerant are the gaps.

```

1  """partition.py -- the numbers behind the partition figure and the
2  right panel of the lattice figure.
3
4  No plotting. The denumerant against the line it samples, the periodic
5  bounded error, the box spline at the integers, and the zero set.
6
7  python3 partition.py
8  """
9
10 from fractions import Fraction
11 from semigroup import denumerant, semigroup
12
13
14 def sampled(a, b, upto=None):
15     """rows (n, d(n), T(n), d(n) - T(n)) with  $T(u) = u_+/ab$ ."""
16     upto = upto if upto is not None else 2 * a * b
17     return [(n, denumerant(n, a, b), Fraction(n, a * b),
18             denumerant(n, a, b) - Fraction(n, a * b))
19             for n in range(upto + 1)]
20
21
22 def tent(a, b, n):
23     """ $ab * B_{\{(a,b)\}}(n)$  by the closed form of the tent lemma."""
24     lo, hi = min(a, b), max(a, b)
25     return max(0, min(n, lo, lo + hi - n))
26
27
28 def nabla(f, a, b, n):
29     """(nabla_a nabla_b f)(n) for f defined on the integers."""
30     return f(n) - f(n - a) - f(n - b) + f(n - a - b)
31
32
33 if __name__ == "__main__":
34     a, b = 4, 5

```

```

35 S, G, F = semigroup(a, b)
36 rows = sampled(a, b)
37 print("n : d(n)  n/ab  d - n/ab")
38 for n, d, T, e in rows[:12]:
39     print("%2d : %d      %-6s %s" % (n, d, T, e))
40 print("zeros of d :", [n for n, d, _, _ in rows if d == 0], " = G :", G)
41
42 assert [n for n, d, _, _ in rows if d == 0] == G
43 assert max(G) == a * b - a - b
44 # d - T is ab-periodic and bounded by 1
45 e = {n: v for n, _, _, v in rows}
46 assert all(e[n] == e[n + a * b] for n in range(a * b))
47 assert all(abs(v) <= 1 for v in e.values())
48 # nabla_a nabla_b kills d off 0, and carries the truncated power to the tent
49 D = lambda n: denominator(n, a, b) if n >= 0 else 0
50 T = lambda u: Fraction(max(u, 0), a * b)
51 assert all(nabla(D, a, b, n) == (n == 0) for n in range(0, 40))
52 assert all(nabla(T, a, b, n) * a * b == tent(a, b, n) for n in range(0, 20))
53 print("tent  ab*B(n) =", [tent(a, b, n) for n in range(11)])
54 print("all checks pass")

```

B.4. The difference word.

LISTING 4. word.py: the difference word of the two closed-form branches, with Lemma 5.4.2 and both conjectures checked for $p = 3, \dots, 17$.

```

1  """word.py -- the numbers behind the difference-word figure.
2
3  No plotting.  The difference word of a tuple and the two conjectures it
4  is evidence for.  The Alexander vector A itself comes from the diagram
5  (cover.py); for the two closed-form branches of the two
6  corollaries of the closed-form families it is given here in closed form, so the checks run
   standalone.
7
8  python3 word.py
9  """
10
11 from staircase import gradings
12
13
14 def famA(p):
15     """[p,1,(p-3)/2,2]: K = T(v,v+1), v = (p+1)/2."""
16     v, g = (p + 1) // 2, (p * p - 1) // 8
17     A = [(p - 1) * (p + 1) // 8]
18     A += [(p - 4 * k + 5) * (p + 3) // 8 for k in range(2, v + 1)]
19     A += [(4 * k - 3 * p - 5) * (p + 1) // 8 for k in range(v + 1, p + 1)]
20     return A, v, (v, v + 1), g
21
22
23 def famB(p):
24     """[p,1,p-2,1]: K = T(2,p)."""
25     return [(p + 1 - 2 * k) // 2 for k in range(1, p + 1)], \
26            p - 1, (2, p), (p - 1) // 2
27
28
29 def word(A):
30     """d_k = A(x_k) - A(x_{k+1}), indices in Z/p."""
31     p = len(A)
32     return [A[k] - A[(k + 1) % p] for k in range(p)]
33
34

```

```

35 def three_letters(d):
36     """The c of the three-letter conjecture, or None."""
37     L = set(d)
38     for c in range(2, max(abs(x) for x in d) + 2):
39         if L <= {1, -c, c + 1}:
40             return c
41     return None
42
43
44 def three_gaps(d):
45     """Cyclic gap sets of each letter, the three-gap conjecture."""
46     p, out = len(d), {}
47     for u in sorted(set(d)):
48         P = [k for k in range(p) if d[k] == u]
49         out[u] = sorted({(P[(i + 1) % len(P)] - P[i]) % p or p
50                         for i in range(len(P))})
51     return out
52
53
54 def ok_three_gaps(gaps):
55     for gs in gaps.values():
56         if len(gs) > 3 or (len(gs) == 3 and gs[0] + gs[1] != gs[2]):
57             return False
58     return True
59
60
61 if __name__ == "__main__":
62     for name, fam, lo in (("A", famA, 5), ("B", famB, 3)):
63         print("family %s" % name)
64         for p in range(lo, 18, 2):
65             A, v, (a, b), g = fam(p)
66             d = word(A)
67             c = three_letters(d)
68             gaps = three_gaps(d)
69             assert len(A) == p and max(A) == g
70             assert sorted(A) == sorted(x for x, _ in gradings(a, b)), \
71                 "A is not the support of Delta_{T(%d,%d)}" % (a, b)
72             assert sum(d) == 0
73             assert d[0] == d[v - 1] == 1 # the rainbow steps
74             assert c is not None and ok_three_gaps(gaps)
75             print(" p=%-3d T(%d,%d) g=%-3d v=%-2d letters {1,%d,%d} "
76                   "gaps %s" % (p, a, b, g, v, -c, c + 1,
77                               {u: gs for u, gs in gaps.items()}))
78     print("all checks pass")

```

B.5. Changing basis on the torus.

LISTING 5. develop.py: the change of basis of Example A.1.7, unimodular and gcd-preserving, carrying $\begin{pmatrix} 2 \\ -1 \end{pmatrix}$ to $\begin{pmatrix} 2 \\ -3 \end{pmatrix}$.

```

1  """develop.py -- the change of basis behind the development figures.
2
3  No plotting. H rewrites the square's coordinates in the meridian-longitude
4  basis, H(u,v) = (u, v-u). It is unimodular, hence an automorphism of
5  H_1(T^2) = Z^2, and it preserves primitivity, so the class of beta in the
6  new basis is the class of the knot on the Heegaard torus.
7
8  python3 develop.py
9  """
10
11 from math import gcd

```

```

12
13 H = ((1, 0), (-1, 1)) # (u, v) -> (u, v - u)
14 HINV = ((1, 0), (1, 1))
15 I2 = ((1, 0), (0, 1))
16
17
18 def det2(M):
19     return M[0][0] * M[1][1] - M[0][1] * M[1][0]
20
21
22 def mul(M, N):
23     return tuple(tuple(sum(M[i][k] * N[k][j] for k in range(2))
24                        for j in range(2)) for i in range(2))
25
26
27 def act(M, c):
28     return tuple(sum(M[i][k] * c[k] for k in range(2)) for i in range(2))
29
30
31 def torus_class(c):
32     """|a|, |b| of the curve of class c = (a, b); it is a knot iff coprime."""
33     a, b = abs(c[0]), abs(c[1])
34     return (a, b) if gcd(a, b) == 1 else None
35
36
37 if __name__ == "__main__":
38     assert det2(H) == 1 and mul(H, HINV) == I2
39     # unimodular, so primitivity is preserved in both directions
40     for u in range(-6, 7):
41         for v in range(-6, 7):
42             if (u, v) != (0, 0):
43                 assert gcd(*act(H, (u, v))) == gcd(u, v)
44     # D[3,1,1,1]: the development reports surface class (2,-3)
45     square = act(HINV, (2, -3))
46     print("surface class (2,-3) <- square class", square)
47     print("knot type T%s" % (torus_class((2, -3)),))
48     assert square == (2, -1)
49     assert torus_class((2, -3)) == (2, 3)
50     print("all checks pass")

```

B.6. The diagram on the torus.

LISTING 6. `torus.py`: the surface, the camera and the arcs of Figure 13, with the three checks that precede emission.

```

1 import math
2
3 R, r, P = 1.00, 0.42, 6
4 STEP = 2 * math.pi / P
5 F1 = math.pi / 2 - STEP # the two bigons straddle f = pi/2
6 FS = [F1 + h * STEP for h in range(P)]
7 DL = 0.58 # how far a rainbow bulges off alpha
8
9 def surface(t, f, e):
10     """S(t,f) seen from d = (0,-cos e, sin e), screen up (0, sin e, cos e)."""
11     ce, se = math.cos(e), math.sin(e)
12     x = (R + r * math.cos(f)) * math.cos(t)
13     y = (R + r * math.cos(f)) * math.sin(t)
14     return x, y * se + r * math.sin(f) * ce
15
16 def front(t, f, e):

```

```

17     """n . d for the outward normal n; the silhouette is front = 0."""
18     ce, se = math.cos(e), math.sin(e)
19     return -math.cos(f) * math.sin(t) * ce + math.sin(f) * se
20
21 def rainbow(fa, fb, base, n=200):
22     """An arc leaving a wall at fa, bulging DL into the annulus, back at fb."""
23     s = 1 if base == 0 else -1
24     return [(base + s * DL * math.sin(math.pi * i / n),
25             fa + (fb - fa) * (1 - math.cos(math.pi * i / n)) / 2)
26             for i in range(n + 1)]
27
28 def band(fa, n=300):
29     """L_h to R_{h+1}: once round the longitude, rising one step in f."""
30     return [(2 * math.pi * i / n, fa + STEP * i / n) for i in range(n + 1)]
31
32 BETA = [rainbow(FS[0], FS[1], 0.0), rainbow(FS[1], FS[2], 2 * math.pi)] \
33         + [band(FS[j]) for j in range(2, P)]
34
35 if __name__ == "__main__":
36     e = math.radians(52.0)
37     # beta is one closed curve: the gluing L_h ~ R_h induces a P-cycle
38     nxt = {1: 2, 2: 3}
39     nxt.update({j: j % P + 1 for j in range(3, P + 1)})
40     orbit, h = [1], 1
41     while nxt[h] not in orbit:
42         h = nxt[h]; orbit.append(h)
43     assert sorted(orbit) == list(range(1, P + 1)), orbit
44     # beta meets alpha in exactly the P generators
45     hits = {round(f % (2 * math.pi), 9) for arc in BETA for t, f in arc
46            if min(t, abs(2 * math.pi - t)) < 1e-9}
47     assert len(hits) == P
48     # both bigons face the viewer entirely
49     lo = min(front(t, f, e) for arc in BETA[:2] for t, f in arc)
50     print("P = %d, elevation 52, min frontness over the bigons %.3f" % (P, lo))
51     assert lo > 0.12
52     print("all checks pass")

```

REFERENCES

- [1] M. Borodzik and C. Livingston, *Semigroups, d-invariants and deformations of cuspidal singular points of plane curves*, J. Lond. Math. Soc. (2) **93** (2016), no. 2, 439–463, arXiv:1305.2868.
- [2] F. Binns and H. Zhou, *(1, 1) almost L-space knots*, arXiv:2304.11475.
- [3] C. de Boor, K. Höllig, and S. Riemenschneider, *Box Splines*, Applied Mathematical Sciences **98**, Springer, New York, 1993, doi:10.1007/978-1-4757-2244-4.
- [4] W. Dahmen and C. A. Micchelli, *The number of solutions to linear Diophantine equations and multivariate splines*, Trans. Amer. Math. Soc. **308** (1988), no. 2, 509–532, doi:10.1090/S0002-9947-1988-0951619-4.
- [5] J. Doyle, *Rainbow normal forms for (1, 1)-knots*, unpublished notes, 2005.
- [6] S. Fairchild, *Notes on (1, 1)-knots and rainbow tuples*, unpublished notes, 2026.
- [7] J. E. Greene, S. Lewallen, and F. Vafaee, *(1, 1) L-space knots*, Compos. Math. **154** (2018), no. 5, 918–933, arXiv:1610.04810.
- [8] H. Goda, H. Matsuda, and T. Morifuji, *Knot Floer homology of (1, 1)-knots*, Geom. Dedicata **112** (2005), 197–214, doi:10.1007/s10711-004-5403-2.
- [9] P. Ozsváth and Z. Szabó, *Holomorphic disks and knot invariants*, Adv. Math. **186** (2004), no. 1, 58–116, arXiv:math/0209056.
- [10] P. Ozsváth and Z. Szabó, *On knot Floer homology and lens space surgeries*, Topology **44** (2005), no. 6, 1281–1300, arXiv:math/0303017.
- [11] T. Popoviciu, *Asupra unei probleme de partiție a numerelor*, Acad. R.P. Române, Fil. Cluj, Stud. Cerc. Științ. **4** (1953), 7–58.
- [12] J. Rasmussen, *Floer homology and knot complements*, Ph.D. thesis, Harvard University, Cambridge, MA, 2003, arXiv:math/0306378.

- [13] J. C. Rosales and P. A. García-Sánchez, *Numerical Semigroups*, Developments in Mathematics **20**, Springer, New York, 2009, doi:10.1007/978-1-4419-0160-6.
- [14] V. T. Sós, *On the distribution mod 1 of the sequence $n\alpha$* , Ann. Univ. Sci. Budapest. Eötvös Sect. Math. **1** (1958), 127–134.
- [15] J. J. Sylvester, *On subinvariants, i.e. semi-invariants to binary quantics of an unlimited order*, Amer. J. Math. **5** (1882), 79–136, doi:10.2307/2369536.

COLUMBIA UNIVERSITY

Email address: ilt2109@columbia.edu