

HOLOMORPHIC DISKS AND THE TOP ALEXANDER GRADING OF TWO-RAINBOW (1, 1)-KNOTS

IZ TECÚM

ABSTRACT. A two-rainbow $(1, 1)$ -knot is presented by a doubly pointed genus one Heegaard diagram $D[p, 2, b, r]$ whose curve β meets each wall of $T^2 \setminus \alpha$ in a nested pair of rainbow arcs, and its knot Floer differential counts holomorphic disks in $\text{Sym}^1(T^2) = T^2$ [23, 27]. We show that the only embedded bigons of $T^2 \setminus (\alpha \cup \beta)$ are the two lenses cut off by the inner rainbows, and that these carry the basepoints. Since $\dim_{\mathbb{C}} T^2 = 1$, an embedded index one domain admits no deformation: the nonlinear Cauchy–Riemann equation has no cokernel for any J [25, 29], every solution factors through the Riemann map,

$$\begin{array}{ccc} \mathbb{R} \times [0, 1] & \xrightarrow{u} & \text{Sym}^1(T^2) = T^2 \\ & \searrow \cong & \swarrow \\ & \mathcal{D}(\phi) & \end{array}$$

and $\# \widehat{\mathcal{M}}(\phi) = 1$ [30]. Consequently

$$\widehat{\partial} = \mathbf{0}, \quad \text{rk} \widehat{HFK}(K, s) = \#\{x \in \alpha \cap \beta \mid A(x) = s\},$$

and the four strands of one period of β carry a single integer $S = \kappa N - m$ for which

$$S \neq 0 \implies \text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1, \quad S = 0 \implies A(\alpha \cap \beta) = \{-1, 0, +1\}, \quad g(K) = 1$$

by genus detection [24]. So $\text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1$ or $A_{\text{top}} = 1$: the rank one behaviour of an L-space knot [26, 34] survives the failure of coherence. The dichotomy is the finite Gottschalk–Hedlund alternative [6] for the Alexander cocycle over the traversal, and S decides fiberedness [31, 33].

CONTENTS

1. Introduction	2
1.1. Rainbow normal forms	2
1.2. Result	3
2. Preliminaries	4
2.1. Rainbow normal form	4
2.2. Drift and admissibility	6
2.3. Reduced and coherent diagrams	11
2.4. Symmetries	13
2.5. Knot Floer input	14
2.6. Combinatorial input	16
2.7. Handlebodies, gluing, and the mapping class action	25
3. Proof of Theorem A	26
4. Analytic foundations and classification input	28
4.1. Detection theorems	28
4.2. The nonlinear Cauchy–Riemann problem on the strip	29
4.3. The genus-one classification	30
4.4. Parametrization and regular homotopy	32
4.5. The Alexander cocycle and the dynamics of the traversal	32
4.6. The tabulation	34

Date: August 30, 2026.

2020 Mathematics Subject Classification. 57K18, 57K10, 57R58.

Key words and phrases. knot Floer homology, $(1, 1)$ -knot, Heegaard diagram, rainbow normal form, holomorphic disk, L-space knot, Alexander grading.

Appendix A. Foundational tools	34
A.1. Permutation matrices and cycles	34
A.2. The cohomological equation on a cycle	35
Appendix B. Reproducibility	35
B.1. The engine	35
B.2. The tabulation checks	38
B.3. The vector identities	40
B.4. The statement checks	41
B.5. The tabulation figure	43
References	44

1. INTRODUCTION

1.1. **Rainbow normal forms.** A knot $K \subset S^3$ is a $(1, 1)$ -knot when it admits a doubly pointed genus-one diagram $(T^2, \alpha, \beta, z, w)$ [28, 27], and since $\text{Sym}^1(T^2) = T^2$ the generators of $\widehat{CFK}(K)$ are the $p = |\alpha \cap \beta|$ intersection points [23]. Every reduced such diagram is isotopic to one with α a meridian [27, 34]. Cutting along α splits β into *rainbow arcs* $\Downarrow$, $\uparrow$, $\Uparrow$ and *vertical arcs* $\downarrow$, $\uparrow$, and Doyle's normal form $[p, a, b, r]$ [35, 37] records a rainbow arcs per wall, b vertical arcs of one step size, and an offset r . Throughout $a = 2$, so each wall carries a nested pair, the inner arc cutting off the bigon that holds a basepoint and the outer one cutting off no bigon at all.






FIGURE 1. $[5, 2, 1, 1]$, cut open along α and reglued by Φ . Mixed, $e = +1$, $K = J(2, 2)$.

FIGURE 2. $[7, 2, 1, 3]$ as in Figure 1, with two returns round the meridian. Coherent, $e = +1$.FIGURE 3. The knot of Figure 1 meets T^2 only at z and w , one arc on the surface and one dotted through the solid torus. The gradings $(1, 3, 1)$ give $\Delta_K = -t + 3 - t^{-1}$, so $K = J(2, 2)$.

1.2. Result. Ozsváth and Szabó showed that an L-space knot has $\widehat{HFK}$ of rank one in each Alexander grading it occupies [26]. At $a = 1$ every admissible tuple is coherent, so every one is an L-space knot and the rank is one throughout [42]. At $a = 2$ the four rainbows need not agree, and what survives is a dichotomy.

Theorem A. Let $[p, 2, b, r]$ be S^3 -admissible in the sense of Definition 2.2.3 and present $K \subset S^3$, oriented so that O_L runs downwards. Then

- (1) the only embedded bigons of $T^2 \setminus (\alpha \cup \beta)$ are $\mathbb{D}_z$ and $\mathbb{D}_w$, so

$$\widehat{\partial} = \mathbf{0}, \quad \text{rk} \widehat{HFK}(K, s) = \#\{x \mid A(x) = s\};$$

- (2) the caps alternate walls along $\star$, cutting β into four strands X_0, X_1, X_2, X_3 with

$$\sigma = \begin{bmatrix} -1 & +1 & -1 & +1 \end{bmatrix}^\top, \quad e = \sigma^\top \mathbf{n}, \quad \ell = \sigma^\top \mathbf{m};$$

- (3) $\mathbb{D} = \{\mathbb{D}, \mathbb{D}\}$ or $\{\mathbb{D}, \mathbb{D}\} \iff D$ is $\mathbb{D}$ $\iff K$ is an L-space knot, and then

$$\text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1;$$

- (4) $\mathbb{D} = \{\mathbb{D}, \mathbb{D}\} \implies K$ is not an L-space knot, D is of type A or B, and

$$\mathbf{n} = N\mathbf{1} - e\varepsilon, \quad \mathbf{m} = m\mathbf{1} - \ell\varepsilon, \quad \varepsilon = \begin{cases} \mathbf{e}_0, & \text{A,} \\ -\mathbf{e}_3, & \text{B,} \end{cases} \quad \begin{bmatrix} p-4 \\ b \end{bmatrix} = \begin{bmatrix} 4N \\ 4m \end{bmatrix} - (\mathbf{1}^\top \varepsilon) \begin{bmatrix} e \\ \ell \end{bmatrix};$$

(5) with $\kappa = e\ell$ and $S = \kappa N - m$,

$$\mathbf{m} - \kappa \mathbf{n} = -S \mathbf{1} \implies \Theta = \text{diag}(\sigma)(\mathbf{m} - \kappa \mathbf{n}) = -S \sigma, \quad \mathbf{1}^\top \Theta = 0 \implies \widehat{A} \circ \tau^4 = \widehat{A};$$

(6) exactly one of

$$S \neq 0 \implies \text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1, \quad S = 0 \implies A(\alpha \cap \beta) = \{-1, 0, +1\}, \quad A_{\text{top}} = 1, \quad g(K) = 1;$$

(7) the second case is exactly the two ends of the band count,

$$S = 0 \iff \mathbf{1}_{\mathcal{B}} \equiv \kappa \in \{0, 1\} \iff b \in \{0, p-4\};$$

(8) the Dehn twist along α acts by

$$\tau_\alpha \cdot D[p, 2, b, r] = D[p, 2, b, r+1], \quad \langle \tau_\alpha \rangle \cong \mathbb{Z}/p,$$

and (1)–(7) are invariants of K , not of the tuple.

In particular

$$\text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1 \quad \text{or} \quad A_{\text{top}} = 1.$$

Proof. §3. □

Remark 1.2.1. By Theorem A, (2) is *Hopf's Umlaufsatz*.

2. PRELIMINARIES

We follow the Doyle–Fairchild normal form throughout, in the orientation of [42], with the wall α oriented downward and the rainbow curve β . Reversing the surface orientation and interchanging the attaching systems exchanges the two handlebodies, so the presented knot is K or its reverse $-K$; since

$$\widehat{HFK}(S^3, K, s) \cong \widehat{HFK}(S^3, -K, s)$$

[23], no statement below depends on that choice. Endpoints are one-based, as in the normal-form figures of [37], so r is the right endpoint of the first band leaving the left nest, and when $b = 0$ it is the first endpoint of the outer right rainbow. Thus for $a = 2$,

$$\begin{aligned} O_L &= L_1 L_4, & I_L &= L_2 L_3, & L_{4+i} &\longleftrightarrow R_{r+i-1}, \\ \tilde{\rho}(s) &= \begin{cases} r+s-5, & 5 \leq s \leq 4+b, \\ r+s-1, & 4+b < s \leq p, \end{cases} & \rho(s) &= \tilde{\rho}(s) \bmod p. \end{aligned}$$

As a check, Figure 4 of [37] draws $D[7, 2, 1, 3]$, for which

$$L_5 \longleftrightarrow R_3, \quad O_R = R_4 R_7, \quad I_R = R_5 R_6.$$

Wall labels are reduced modulo p , while $v = r + b$ and $\tilde{\rho}$ denote integer lifts. Write $e = \beta \cdot \alpha \in \{\pm 1\}$, fix a path $\delta: z \rightarrow w$ missing α , and set


$$I_\delta(\gamma) = \gamma \cdot \delta, \quad D_\delta = \beta \cdot \delta, \quad \varepsilon \in \{\pm 1\}$$

for the global sign of the Alexander grading.

2.1. Rainbow normal form.

Convention 2.1.1 (Inherited normal form, [42, Definitions 2.1.1–2.1.4]). $(T^2, \alpha, \beta, z, w)$ is doubly pointed of genus one, $[p, a, b, r]$ is a rainbow tuple, and $A = T^2 \setminus \alpha = [0, 1] \times \mathbb{R}/p\mathbb{Z}$ is the cut annulus with walls W_L, W_R , slots L_h, R_h at height $h - \frac{1}{2}$ increasing downwards, and identification $\downarrow \downarrow: R_h \mapsto L_h$. The matching  has $a = 2$ and $v = r + b$,

$$\begin{aligned} L_i &\leftrightarrow L_{5-i}, & R_{v-1+i} &\leftrightarrow R_{v+4-i}, & i &= 1, 2, & L_s &\leftrightarrow R_{\rho(s)}, & 5 \leq s \leq p, \\ & & z &\in L_2 L_3, & & & w &\in R_{v+1} R_{v+2}. \end{aligned}$$



Definition 2.1.2 (Two-rainbow arcs).

$$O_L = L_1 L_4 \supset I_L = L_2 L_3, \quad O_R = R_v R_{v+3} \supset I_R = R_{v+1} R_{v+2}, \quad c_s = L_s R_{p(s)},$$

$$\mathcal{B} = \{c_s \mid 5 \leq s \leq 4+b\}, \quad \overline{\mathcal{B}} = \{c_s \mid 4+b < s \leq p\}, \quad |\mathcal{B}| = b, \quad |\overline{\mathcal{B}}| = p-4-b,$$

$$\mathfrak{D}_z = D(I_L) \ni z, \quad \mathfrak{D}_w = D(I_R) \ni w, \quad x_z \in \partial D(I_L), \quad x_w \in \partial D(I_R),$$

and $c_s \in \overline{\mathcal{B}}$ crosses $(u,0) \sim (u,p)$ iff $\tilde{\rho}(s) > p$.




FIGURE 4. Fairchild's construction of $D[7, 2, 1, 3]$, one arc family at a time from the marked points to the basepoints. Drawn after [37].




FIGURE 5. $\mathcal{D}[7, 2, 1, 3]$ with $v = 4$, the nests $O_L = L_1L_4 \supset I_L = L_2L_3$ and $O_R = R_4R_7 \supset I_R = R_5R_6$ cutting off $\mathcal{D}_z, \mathcal{D}_w$, the band $\mathcal{B} = \{L_5R_3\}$, and the returns L_6R_1, L_7R_2 crossing $(u, 0) \sim (u, p)$ with their second pieces dashed. Under $\downarrow \uparrow$ the slots L_h, R_h have common image x_h .

Remark 2.1.3 (Standard position, [42, Remark 2.1.8]). α is the meridian, β carries the rainbow arcs.


Proof. [27, §2], [34, 10]; no value of a is used. □

2.2. Drift and admissibility.

Definition 2.2.1 (Traversal, [42, Definition 2.2.1]).


$$\omega(L_h) = R_h, \quad \omega(R_h) = L_h, \quad \tau = \omega \circ \sigma$$

for σ the matching of Convention 2.1.1. One step is $\downarrow \uparrow$, and the *traversal* is $\star = \tau^{\mathbb{Z}}(L_1)$.




Definition 2.2.2 (Drift, [42, Definition 2.2.2]). For $e \subset \beta \cap A$,

$$\chi(\text{downward arrow}) = 0, \quad \chi(\text{upward arrow}) = +1, \quad \chi(\text{downward arrow}) = -1, \quad e = \sum_e \chi(e) = \beta \cdot \alpha.$$



Definition 2.2.3 (Admissibility, [42, Definition 2.2.3]). $D[p, 2, b, r]$ is S^3 -admissible when $\text{blue circle} \neq \text{blue star}$, that is $|\text{blue star}| = p$ and $|e| = 1$.



Definition 2.2.4 (Band winding).

$$\ell = \sum_{c_s \in \mathcal{B}} \chi(c_s) \in \mathbb{Z}, \quad \kappa = e\ell \in \mathbb{Z}, \quad e - \ell = \sum_{c_s \in \overline{\mathcal{B}}} \chi(c_s).$$




FIGURE 6. The band of $D[7, 2, 1, 3]$ is traversed right to left and its two returns left to right, so $e = +1$, $\ell = -1$ and $\kappa = -1$.

Lemma 2.2.5 (Arc steps in the cyclic cover). *Let Δh denote the height displacement of an oriented arc, read in the lift $\mathbb{R} \rightarrow \mathbb{R}/p\mathbb{Z}$. Then*

$$\Delta h(\downarrow \uparrow) = 0, \quad \Delta h(c_s) = \begin{cases} \chi(c_s)(r-5), & c_s \in \mathcal{B}, \\ \chi(c_s)(r-1), & c_s \in \overline{\mathcal{B}}, \end{cases} \quad \Delta h(c) = -\eta(c) \deg(c),$$

$$\deg(O_L) = \deg(O_R) = 3, \quad \deg(I_L) = \deg(I_R) = 1.$$

Proof.

$$\begin{aligned} \omega(L_h) = R_h, \quad \omega(R_h) = L_h &\implies \Delta h(\downarrow \uparrow) = h - h = 0 \\ \Delta h(c_s) = \chi(c_s)(\tilde{\rho}(s) - s) &\xrightarrow{2.1.2} \chi(c_s)(r-5) (\mathcal{B}), \quad \chi(c_s)(r-1) (\overline{\mathcal{B}}) \\ O_L = L_1 L_4, \quad I_L = L_2 L_3, \quad O_R = R_v R_{v+3}, \quad I_R = R_{v+1} R_{v+2} &\implies \deg = 3, 1, 3, 1 \\ \eta(c) = +1 &\xrightarrow{2.3.4} \text{height decreases} \implies \Delta h(c) = -\eta(c) \deg(c) \end{aligned}$$

□




FIGURE 7. The two deck steps and the four rainbow degrees of Lemma 2.2.5.

Lemma 2.2.6 (Permutation matrix of the traversal). *Order the $2p$ slots $L_1, \dots, L_p, R_1, \dots, R_p$, and let $T, W \in \mathbb{M}_{2p \times 2p}(\mathbb{Q})$ be the permutation matrices of τ and ω , in the sense of [42, Convention 5.0.1],*

$$T = (T_{mn})_{m,n=1}^{2p} = \begin{bmatrix} T_{11} & T_{12} & \cdots & T_{1,2p} \\ T_{21} & T_{22} & \cdots & T_{2,2p} \\ \vdots & \vdots & \ddots & \vdots \\ T_{2p,1} & T_{2p,2} & \cdots & T_{2p,2p} \end{bmatrix}, \quad T_{mn} = \begin{cases} 1, & n = \tau(m), \\ 0, & n \neq \tau(m). \end{cases}$$

Then

$$T \cdot T^\top = \mathbf{I}_{2p}, \quad W \cdot W = \mathbf{I}_{2p}, \quad W \cdot T \cdot W = T^{-1},$$

$$|\star| = p \iff \det(T - \lambda \mathbf{I}_{2p}) = (\lambda^p - 1)^2 \iff \text{rk}(T - \mathbf{I}_{2p}) = 2p - 2,$$

and then $\text{spec}(T) = \{\zeta \in \mathbb{C} \mid \zeta^p = 1\}$.

Proof.

$$\tau \text{ a bijection} \implies T \text{ has one } 1 \text{ per row and column} \implies T \cdot T^\top = \mathbf{I}_{2p}$$

$$\omega^2 = \sigma^2 = \text{id}, \quad \tau = \omega \circ \sigma \implies W \cdot W = \mathbf{I}_{2p}, \quad W \cdot T \cdot W = \sigma \circ \omega = \tau^{-1}$$

$$\gamma \text{ a } d\text{-cycle} \implies \text{a nonzero term takes } -\lambda \text{ or } 1 \text{ from every column} \implies \det(P - \lambda \mathbf{I}_d) = (-1)^d (\lambda^d - 1)$$

$$\tau = \gamma_1 \sqcup \cdots \sqcup \gamma_k, \quad \sum_{i=1}^k d_i = 2p \implies \det(T - \lambda \mathbf{I}_{2p}) = \prod_{i=1}^k (\lambda^{d_i} - 1), \quad \dim \ker(T - \mathbf{I}_{2p}) = k$$

$$p \text{ odd} \implies \omega(\star) \neq \star \implies [|\star| = p \iff k = 2, d_1 = d_2 = p \iff \det = (\lambda^p - 1)^2 \iff \text{rk} = 2p - 2]$$

$$\lambda = 1 \text{ a root of multiplicity } k, \quad d_1 + d_2 = 2p \text{ force the converse}$$

$$\det(T - \lambda \mathbf{I}_{2p}) = 0 \iff \lambda^p = 1 \implies \text{spec}(T) = \{\zeta \mid \zeta^p = 1\}$$

□

For $D[7, 2, 1, 3]$ the traversal is $x_1 x_4 x_7 x_2 x_3 x_5 x_6$, so the induced permutation of $\{x_1, \dots, x_7\}$ is $1 \mapsto 4, 2 \mapsto 3, 3 \mapsto 5, 4 \mapsto 7, 5 \mapsto 6, 6 \mapsto 1, 7 \mapsto 2$, with

$$T = \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \quad T - \lambda \mathbf{I}_7 = \begin{bmatrix} -\lambda & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & -\lambda & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\lambda & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -\lambda & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -\lambda & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & -\lambda & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & -\lambda \end{bmatrix}.$$

Written out by columns, $T - \lambda \mathbf{I}_7 = [\mathbf{c}_1 \mathbf{c}_2 \mathbf{c}_3 \mathbf{c}_4 \mathbf{c}_5 \mathbf{c}_6 \mathbf{c}_7]$ with

$$\mathbf{c}_1 = \begin{bmatrix} -\lambda \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{c}_2 = \begin{bmatrix} 0 \\ -\lambda \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \quad \mathbf{c}_3 = \begin{bmatrix} 0 \\ 1 \\ -\lambda \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{c}_4 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -\lambda \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{c}_5 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ -\lambda \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{c}_6 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ -\lambda \\ 0 \end{bmatrix}, \quad \mathbf{c}_7 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ -\lambda \end{bmatrix},$$

each column carrying $-\lambda$ in its own row and 1 in row $\tau^{-1}(j)$, where τ^{-1} is $1 \mapsto 6, 2 \mapsto 7, 3 \mapsto 2, 4 \mapsto 1, 5 \mapsto 3, 6 \mapsto 5, 7 \mapsto 4$. A nonzero Leibniz term takes one entry from each column, so it takes $-\lambda$ from every column or 1 from every column, and

$$\prod_{j=1}^7 (-\lambda) = (-\lambda)^7 = -\lambda^7, \quad \text{sgn}(\text{id}) = +1, \quad \prod_{j=1}^7 1 = 1, \quad \text{sgn}(\tau^{-1}) = \text{sgn}(1472356) = (-1)^{7-1} = +1,$$

$$\det(T - \lambda \mathbf{I}_7) = (+1) \cdot (-\lambda^7) + (+1) \cdot 1 = -\lambda^7 + 1 = (-1)^7 (\lambda^7 - 1).$$

The kernel of $T - \mathbf{I}_7$ is cut out by $x_4 = x_1, x_3 = x_2, x_5 = x_3, x_7 = x_4, x_6 = x_5, x_1 = x_6, x_2 = x_7$, which chain to $x_1 = x_4 = x_7 = x_2 = x_3 = x_5 = x_6$, so it is spanned by $\mathbf{1} = (1, 1, 1, 1, 1, 1, 1)^\top$ and

$$(T - \mathbf{I}_7) \cdot \mathbf{1} = \begin{bmatrix} -1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -1 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & -1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \mathbf{0}, \quad \text{rk}(T - \mathbf{I}_7) = 7 - 1 = 6.$$




FIGURE 8. The traversal of $D[7, 2, 1, 3]$ as the 7-cycle (1472356) .

Lemma 2.2.7 (Rank, [42, Lemma 2.2.4]).

$$\text{blue circle with a slash} \implies |\pi_0(\beta)| = 1, \quad (T^2, \alpha, \beta) \simeq S^3, \quad \text{rk}(\widehat{CFK}(K)) = p.$$

Proof.

$$\begin{aligned} \textcircled{+} \xrightarrow{2.2.6} \tau = \gamma_1 \sqcup \gamma_2, \quad |\gamma_1| = |\gamma_2| = p \implies |\pi_0(\beta)| = 1, \quad \alpha \cap \beta = \{x_1, \dots, x_p\} \implies \text{rk}(\widehat{CFK}(K)) = p \\ |e| = |\beta \cdot \alpha| = 1 \implies H_1(T^2) = \mathbb{Z}\langle \alpha \rangle \oplus \mathbb{Z}\langle \beta \rangle \implies (T^2, \alpha, \beta) \simeq S^3 \end{aligned}$$

□

Lemma 2.2.8 (Holonomy of the lifted traversal).

$$\begin{aligned} e(r-1) - 4\ell - 3(\eta(O_L) + \eta(O_R)) - (\eta(I_L) + \eta(I_R)) &\equiv 0 \pmod{p}, \\ \textcircled{D} = \{\textcircled{D}, \textcircled{D}\}, \quad \textcircled{C} = \{\textcircled{C}, \textcircled{C}\} &\iff e(r-1) - 4\ell + 8 \equiv 0 \pmod{p}, \\ O_L = \textcircled{D}, \quad I_L = \textcircled{D}, \quad O_R = \textcircled{C}, \quad I_R = \textcircled{C} &\iff e(r-1) - 4\ell + 4 \equiv 0 \pmod{p}. \end{aligned}$$

Proof.

$$\begin{aligned} \star \text{ closed}, \quad \Delta h(\textcircled{+}) = 0 &\xrightarrow{2.2.5} 0 \equiv \sum_{s=5}^p \Delta h(c_s) + \sum_c \Delta h(c) \pmod{p} \\ \sum_{s=5}^p \Delta h(c_s) = (r-5)\ell + (r-1)(e-\ell) &= e(r-1) - 4\ell, \quad \sum_c \Delta h(c) = -3(\eta(O_L) + \eta(O_R)) - (\eta(I_L) + \eta(I_R)) \\ &\xrightarrow{2.3.4} O_L, I_L, O_R, I_R \text{ all } \textcircled{D}, \textcircled{C} \implies -3(-2) - (-2) = +8 \\ O_L = \textcircled{D}, \quad O_R = \textcircled{C}, \quad I_L = \textcircled{D}, \quad I_R = \textcircled{C} &\implies -3(-2) - (+2) = +4 \\ &\xrightarrow{2.3.5} \text{the balanced patterns give } 8, 4, 0, -4, -8 \\ p \text{ odd}, \quad p \geq 5 &\implies 8, 4, 0, -4, -8 \text{ pairwise distinct } \pmod{p} \implies \text{each congruence determines its pattern} \end{aligned}$$

□





FIGURE 9. The lifted traversal of $D[7, 2, 1, 3]$ rises by $e(r-1) - 4\ell + 8 = 2 + 4 + 8 = 14 = 2p$ and so closes modulo p .

2.3. Reduced and coherent diagrams.


Definition 2.3.1 (Region type, [42, Definition 2.3.1]). A component R of $T^2 \setminus (\alpha \cup \beta)$ is n -sided when $|\partial R \cap \alpha \cap \beta| = 2n$,

$$\mathbb{D} (n=1), \quad \mathbb{D}_z (n=2), \quad \mathbb{D}_\alpha (n=3), \quad \mathbb{D}_{\alpha\beta} (n=4),$$

the darker lobe against a wall being a neighbouring bigon, one of the n sides of R and not part of it.



Definition 2.3.2 (Reduced diagram, [34]). A diagram is *reduced* $\mathbb{D}$ when every bigon of $T^2 \setminus (\alpha \cup \beta)$ carries a basepoint,



Definition 2.3.3 (Coherent diagram, [34]). A reduced diagram is *coherent* when α, β admit orientations coorienting every bigon, equivalently [10]



Definition 2.3.4 (Rainbow sense).

$$\begin{aligned} \mathbb{D} &= \{O_L, I_L\}, & \mathbb{D}_z &= \{O_R, I_R\}, & \eta(\mathbb{D}) &= \eta(\mathbb{D}_z) = +1, & \eta(\mathbb{D}_{\alpha}) &= \eta(\mathbb{D}_{\alpha\beta}) = -1, \\ u_L &= \#\{c \in \mathbb{D} \mid c = \mathbb{D}_z\}, & d_L &= \#\{c \in \mathbb{D} \mid c = \mathbb{D}\} = 2 - u_L, \\ u_R &= \#\{c \in \mathbb{D}_z \mid c = \mathbb{D}_{\alpha}\}, & d_R &= \#\{c \in \mathbb{D}_z \mid c = \mathbb{D}_{\alpha\beta}\} = 2 - u_R. \end{aligned}$$





FIGURE 10. Two balanced sense patterns and, to the right of the rule, the pattern excluded by Lemma 2.3.5.

Lemma 2.3.5 (Turning balance).


$$\begin{aligned} \vartheta(\beta) = 2\pi(u_L - u_R) = 0 &\iff u_L = u_R \in \{0, 1, 2\}, \\ \mathbb{D} = \{\mathbb{D}_z, \mathbb{D}\} &\iff \mathbb{D}_{\alpha} = \{\mathbb{D}_{\alpha}, \mathbb{D}_{\alpha\beta}\}, & \mathbb{D} = \{\mathbb{D}_z, \mathbb{D}\} &\iff \mathbb{D}_{\alpha\beta} = \{\mathbb{D}_{\alpha}, \mathbb{D}_{\alpha\beta}\}, \\ \mathbb{D} = \{\mathbb{D}_z, \mathbb{D}\} &\iff \mathbb{D}_{\alpha} = \{\mathbb{D}_{\alpha}, \mathbb{D}_{\alpha\beta}\}. \end{aligned}$$

Proof.


$$\beta \text{ embedded essential} \implies \beta \simeq \text{geodesic} \implies \vartheta(\beta) = 0, \quad \vartheta(\downarrow \uparrow) = 0, \quad \vartheta(\mathbb{D}) = +\pi\eta, \quad \vartheta(\mathbb{C}) = -\pi\eta$$



$$\vartheta(c_s) = 0$$



$$\vartheta = +\pi$$



$$\vartheta = -\pi$$

$$0 = \vartheta(\beta) = \pi(u_L - d_L) - \pi(u_R - d_R) = 2\pi(u_L - u_R) \implies u_L = u_R \in \{0, 1, 2\} \quad [42, \text{Thm. A.3.3, Lem. 3.2.3}]$$


□

Theorem 2.3.6 (Bigon classification and reducedness).


$$\mathbb{D} \text{ embedded} \implies \mathbb{D} \in \{\mathbb{D}_z, \mathbb{C}_w\}, \quad \mathbb{D} \rightsquigarrow \mathbb{D}, \quad \widehat{\partial} = \mathbf{0}.$$

Proof.


$$\partial \mathbb{D} = a \cup e, \quad \text{int}(e) \cap \alpha = \emptyset \xrightarrow{2.1.2} e \in \mathbb{D} \cup \mathbb{C} \cup \{\downarrow \uparrow\}, \quad (a \cup c_s) \cdot \mu = \pm 1 \implies e \neq c_s \quad [42, \text{Lem. 3.1.1}]$$



$$[a \cup I_L] = 0$$



$$[a \cup c_s] \neq 0$$




$$D_z \subset D(O_L)$$

$$D(O_L) \setminus D(I_L) = D(O_R) \setminus D(I_R) = \mathbb{D} \rightsquigarrow \mathbb{D} \implies e \in \{I_L, I_R\} \implies \mathbb{D} \in \{\mathbb{D}_z, \mathbb{C}_w\} \xrightarrow{2.3.2} \mathbb{D} \rightsquigarrow \mathbb{D}$$

$$\mu(\varphi) = 1, \quad \text{Sym}^1(T^2) = T^2 \implies \mathcal{D}(\varphi) = \mathbb{D} \implies n_z(\varphi) + n_w(\varphi) > 0 \implies \widehat{\partial} = \mathbf{0} \quad [28, 34]$$

□



$$\mu(\phi) = 1, n_z(\phi) = 1$$



$$\widehat{\partial} \mathbf{x} = 0$$

Corollary 2.3.7 (Rank equals count).

$$\text{rk} \widehat{HFK}(K, s) = \#\{x \in \alpha \cap \beta \mid A(x) = s\}, \quad \sum_s \text{rk} \widehat{HFK}(K, s) = p.$$

Proof.

$$\widehat{\partial} = \mathbf{0} \xrightarrow{2.3.6} \widehat{HFK}(K) = \widehat{CFK}(K) = \bigoplus_{k=1}^p \mathbb{F} \cdot x_k \xrightarrow{2.2.7} \text{rk} \widehat{HFK}(K, s) = \#\{x \mid A(x) = s\} \quad [42, \text{Cor. 3.1.3}]$$

□



Proposition 2.3.8 (Coherence).

$$\mathbb{D} = \{\mathbb{D}, \mathbb{D}\}, \quad \mathbb{C} = \{\mathbb{C}, \mathbb{C}\} \quad \text{or} \quad \mathbb{D} = \{\mathbb{D}, \mathbb{D}\}, \quad \mathbb{C} = \{\mathbb{C}, \mathbb{C}\}$$

$$\iff \eta(O_L) = \eta(I_L) = \eta(O_R) = \eta(I_R) = \sigma \iff \mathbb{D} \rightsquigarrow \mathbb{D}, \text{ never } \mathbb{D} \rightsquigarrow \mathbb{D} \implies \mathbb{D} \rightsquigarrow \mathbb{D} \text{ coherent.}$$

Proof.

$$\xrightarrow{2.3.6} \mathbb{D} \in \{\mathbb{D}_z, \mathbb{D}_w\}, \quad a(D) = \langle \beta, \partial^+ D \rangle, \quad b(D) = \langle \alpha, \partial^+ D \rangle, \quad \mathbb{D}: a = \sigma, b = -\tau, \quad \mathbb{D}: a = -\sigma, b = \tau$$



$$\begin{aligned} \partial \mathbb{D}_z, \partial \mathbb{D}_w \text{ cooriented} &\iff \sigma\tau = -1 \iff \tau := -\sigma \implies a(D) = b(D) \quad \forall D \xLeftrightarrow{2.3.3} \mathbb{D} \mathbb{D} \\ \sigma \text{ exists} &\xLeftrightarrow{2.3.4} \mathbb{D} \in \{\{\mathbb{D}, \mathbb{D}\}, \{\mathbb{D}, \mathbb{D}\}\} \xLeftrightarrow{2.3.5} \mathbb{D} \in \{\{\mathbb{D}, \mathbb{D}\}, \{\mathbb{D}, \mathbb{D}\}\} \end{aligned}$$


□

2.4. Symmetries.

Convention 2.4.1 (Reversal and mirror, [42, Definition 2.4.1]).

$$-\beta: \quad \downarrow \uparrow \mapsto \uparrow \downarrow, \quad \mathbb{D} \mapsto \mathbb{D}, \quad \mathbb{D} \mapsto \mathbb{D}, \quad \mathbb{D} \mapsto \mathbb{D},$$

and reflecting A by $u \mapsto 1 - u$ is $\mathbb{D} \mapsto \mathbb{D}$.




Lemma 2.4.2 (Effect on the gradings, [42, Lemma 2.4.2]).

$$-\beta: \quad e \mapsto -e, \quad A \mapsto -A, \quad x_z \longleftrightarrow x_w.$$

Proof.

$$\xrightarrow{2.2.2} e = \sum_{s=5}^p \chi(c_s) \mapsto -e, \quad \mathbb{D} \mapsto \mathbb{D} \implies (n_z, n_w) \mapsto (n_w, n_z) \implies A \mapsto -A \xrightarrow{2.1.1} x_z \longleftrightarrow x_w$$

□




Lemma 2.4.3 (Reversal invariants).

$$\begin{aligned} -\beta: \quad \chi &\mapsto -\chi, \quad e \mapsto -e, \quad \ell \mapsto -\ell, \quad \kappa \mapsto \kappa, \quad \eta \mapsto -\eta, \\ \mathbb{D} = \{\mathbb{D}, \mathbb{D}\} &\iff -\beta: \mathbb{D} = \{\mathbb{D}, \mathbb{D}\}, \quad \mathbb{D} = \{\mathbb{D}, \mathbb{D}\} \iff -\beta: \mathbb{D} = \{\mathbb{D}, \mathbb{D}\}. \end{aligned}$$

Proof.

$$\xrightarrow{2.2.2, 2.2.4} \chi \mapsto -\chi, \quad \mathcal{B} \text{ setwise fixed} \implies e \mapsto -e, \quad \ell \mapsto -\ell, \quad \kappa = e\ell \mapsto (-e)(-\ell) = \kappa$$




□

Corollary 2.4.4 (Reversal normalisation).

$$D = \{D, D\} \iff -\beta: D = \{D, D\} \implies e = +1, \quad \kappa = \ell, \quad O_L = D.$$

Proof.

$$\begin{aligned}
 D = \{D, D\} &\stackrel{2.4.3}{\implies} D \text{ fixed}, \quad D = \{D, D\} \stackrel{2.4.3}{\implies} D = \{D, D\} \stackrel{2.3.8}{\implies} D \sqcup C \\
 &\stackrel{2.4.2}{\implies} -\beta \text{ presents } -K, \quad \widehat{HFK}(S^3, -K, s) \cong \widehat{HFK}(S^3, K, s) \implies e = +1 \stackrel{2.2.4}{\implies} \kappa = e\ell = \ell
 \end{aligned}$$



□

Remark 2.4.5 (The drift is not determined). Reversal fixes the mixed sense pattern (Lemma 2.4.3), so nothing intrinsic picks a sign of e : Corollary 2.4.4 is a normalisation, not a theorem.

2.5. Knot Floer input.

Theorem 2.5.1 (Coherence criterion, [34, Theorem 1.2], [42, Theorem 2.5.1]).

$$D \not\sqcup, \quad D \sqcup C \iff K \text{ an } L\text{-space knot.}$$

Proof. [34, Theorem 1.2].

□

Theorem 2.5.2 (Alexander polynomial, [26, Theorem 1.2], [42, Theorem 2.5.2]).

$$K \text{ an } L\text{-space knot} \implies \Delta_K(t) = \sum_{j=0}^{2n} (-1)^j t^{A_j}, \quad A_0 > A_1 > \dots > A_{2n},$$

$$\text{rk} \widehat{HFK}(K, A_j) = 1, \quad \widehat{HFK}(K, s) = 0, \quad s \notin \{A_0, \dots, A_{2n}\}.$$


Proof. [26, Theorem 1.2].

□

Definition 2.5.3 (Alexander criterion, [42, Definition 2.5.3]). Δ_K has the form of Theorem 2.5.2, that is $\sqcup_{\sqcup}$.

Definition 2.5.4 (Maslov normalisation, [42, Definition 2.5.4]).

$$A = A_{\text{top}} \implies M = 0, \quad \delta_0 = 0, \quad \delta_j = \begin{cases} \delta_{j-1} - 2(A_{j-1} - A_j) + 1, & j \text{ odd,} \\ \delta_{j-1} - 1, & j \text{ even.} \end{cases}$$



Remark 2.5.5 (Mirrors, [42, Remark 2.5.5]).

$$\mathbb{D}[\mathbb{Q}]: (\delta_0, \delta_1, \dots, \delta_{2n}) \mapsto (\delta_{2n}, \dots, \delta_1, \delta_0),$$

so Definition 2.5.4 precedes Definition 2.5.3.

Proof. Lemma 2.4.2. □

Lemma 2.5.6 (Balanced chains determine domains).

$$c \subset \beta, \quad \partial c = y - x, \quad c \cdot \alpha = 0 \implies \exists \mathcal{D} \in \pi_2(x, y), \quad \partial_\beta \mathcal{D} = c, \quad \mathcal{D} \text{ unique modulo } \mathbb{Z}[T^2].$$


Proof.

$$\partial(c + a_0) = (y - x) + (x - y) = 0 \implies [c + a_0] = m[\alpha] + n[\beta], \quad 0 = c \cdot \alpha + a_0 \cdot \alpha = [c + a_0] \cdot [\alpha] = ne \implies n = 0$$

$$a := a_0 - m\alpha \implies \partial a = x - y, \quad [c + a] = 0 \implies c + a = \partial \mathcal{D}$$

$$\text{Sym}^1(T^2) = T^2 \implies \mathcal{D} \longleftrightarrow \varphi_{\mathcal{D}} \in \pi_2(x, y), \quad \partial_\beta \mathcal{D} = c \quad [28, \S 2], [23, \S 3]$$

$$\partial \mathcal{D} = \partial \mathcal{D}' \implies \partial(\mathcal{D} - \mathcal{D}') = 0 \implies \mathcal{D} - \mathcal{D}' \in H_2(T^2; \mathbb{Z}) = \mathbb{Z}[T^2]$$



□


Lemma 2.5.7 (Path-pairing formula).

$$n_z(\mathcal{D}) - n_w(\mathcal{D}) = \varepsilon_0(\partial_\beta \mathcal{D}) \cdot \delta, \quad \varepsilon_0 \in \{\pm 1\} \text{ independent of } \mathcal{D}.$$

Proof.

$$n_\delta(t) = \text{multiplicity of } \mathcal{D} \text{ at } \delta(t), \quad \delta \cap \alpha = \emptyset \implies n_\delta \text{ jumps only at } \delta \pitchfork \partial_\beta \mathcal{D}, \text{ by } \varepsilon_0 s_i$$

$$n_z(\mathcal{D}) - n_w(\mathcal{D}) = n_\delta(0) - n_\delta(1) = \varepsilon_0 \sum_{i=1}^k s_i = \varepsilon_0(\partial_\beta \mathcal{D}) \cdot \delta$$




□

Proposition 2.5.8 (Connection formula).

$$c \subset \beta, \quad c: x \rightarrow y, \quad \chi(c) = c \cdot \alpha \implies A(y) - A(x) = \varepsilon(I_\delta(c) - eD_\delta \chi(c)).$$

Proof.

$$\begin{aligned} \hat{c} &:= c - e\chi(c)\beta \implies \hat{c} \cdot \alpha = \chi(c)(1 - e^2) = 0, \quad \partial \hat{c} = y - x \xrightarrow{2.5.6} \hat{c} = \partial_\beta \mathcal{D}_c, \quad \mathcal{D}_c \in \pi_2(x, y) \\ \mathcal{D}_c + m[T^2] &\implies n_z - n_w \text{ unchanged} \xrightarrow{2.5.7} A(y) - A(x) = \varepsilon \hat{c} \cdot \delta, \quad \varepsilon \in \{\pm 1\} \text{ independent of } c \quad [23, \S 3] \\ \hat{c} \cdot \delta &= c \cdot \delta - e\chi(c)(\beta \cdot \delta) = I_\delta(c) - eD_\delta \chi(c) \end{aligned}$$




□

Lemma 2.5.9 (Rainbow step).

$$c \in \{O_L, I_L, O_R, I_R\}, \quad \varepsilon := +1 \implies \Theta(c) = \eta(c), \quad \Theta(\mathbb{D}_z) = \Theta(\mathbb{D}_w) = +1, \quad \Theta(\mathbb{D}_z) = \Theta(\mathbb{D}_w) = -1.$$

Proof.

$$\begin{aligned} \chi(c) = 0 &\xrightarrow{2.5.8} \Theta(c) = \varepsilon I_\delta(c), \quad \mathbb{D}_z \subset D(I_L) \subset D(O_L), \quad \mathbb{D}_w \subset D(I_R) \subset D(O_R) \implies \#(\delta \cap c) = 1 \\ \text{sgn}(c'(p) \wedge \delta'(p)) &= \eta(c) \xrightarrow{2.3.4} I_\delta(c) = \eta(c), \quad \varepsilon := +1 \implies \Theta(c) = \eta(c) \end{aligned}$$




□

2.6. Combinatorial input. From here $D[p, 2, b, r]$ is mixed and normalised as in Corollary 2.4.4,

$$\mathbb{D} = \{\mathbb{D}_z, \mathbb{D}_w\}, \quad \mathbb{C} = \{\mathbb{C}_z, \mathbb{C}_w\}, \quad O_L = \mathbb{D}_z, \quad e = +1, \quad \kappa = e\ell = \ell.$$

A maximal run of $\downarrow \uparrow$ between two consecutive $\mathbb{D}_z, \mathbb{C}_z$ along $\star$ is a *strand*. One period of $\star$ carries exactly four of them; index them by $i \in \mathbb{Z}/4$ in the cyclic order in which $\star$ meets them, so that τ^4 carries one period of $\hat{\beta}$ to the next,




Write n_i for the number of $\downarrow \nearrow$ in X_i and $m_i = \sum_{c \in X_i} \mathbf{1}_{\mathcal{B}}(c)$ for the number of those that are bands, and collect the four values of each into a vector; the sign vector σ records the constant corner sign of each strand,

$$\mathbf{n} = \begin{bmatrix} n_0 \\ n_1 \\ n_2 \\ n_3 \end{bmatrix} \in \mathbb{Z}^4, \quad \mathbf{m} = \begin{bmatrix} m_0 \\ m_1 \\ m_2 \\ m_3 \end{bmatrix} \in \mathbb{Z}^4, \quad \sigma = \begin{bmatrix} \chi_{X_0} \\ \chi_{X_1} \\ \chi_{X_2} \\ \chi_{X_3} \end{bmatrix} = \begin{bmatrix} -1 \\ +1 \\ -1 \\ +1 \end{bmatrix}, \quad \mathbf{1} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

Every scalar below is then one linear functional of this pair,

$$e = \sigma^\top \mathbf{n}, \quad \ell = \sigma^\top \mathbf{m}, \quad p = \mathbf{1}^\top \mathbf{n} + 4, \quad b = \mathbf{1}^\top \mathbf{m}, \quad \kappa = e \ell.$$



(a) $(T^2, \alpha, \beta, z, w), D[5, 2, 1, 1]$

(b) $A = T^2 \setminus \alpha, R_h \mapsto L_h$

FIGURE 11. The nests hang off the same $\mid$ on opposite sides, so $\mathfrak{D}_z \subset D(I_L) \subset D(O_L)$ and $\mathfrak{C}_w \subset D(I_R) \subset D(O_R)$.

Lemma 2.6.1 (Corner alternation along $\partial\beta$).


$$\left(\begin{array}{c} L_1 \\ L_2 \\ L_3 \\ L_4 \end{array} \mid \begin{array}{c} \mathfrak{D} \\ \mathfrak{C} \\ \mathfrak{D} \\ \mathfrak{C} \end{array} \right) \Rightarrow \left(\begin{array}{c} R_v \\ R_{v+1} \\ R_{v+2} \\ R_{v+3} \end{array} \mid \begin{array}{c} \mathfrak{D} \\ \mathfrak{C} \\ \mathfrak{D} \\ \mathfrak{C} \end{array} \right) \text{ or } \left(\begin{array}{c} R_v \\ R_{v+1} \\ R_{v+2} \\ R_{v+3} \end{array} \mid \begin{array}{c} \mathfrak{C} \\ \mathfrak{D} \\ \mathfrak{C} \\ \mathfrak{D} \end{array} \right)$$

$\mathfrak{D}, \mathfrak{C}, \mathfrak{D}, \mathfrak{C}$ in this cyclic order along $\star$, $\sigma = [-1 \ +1 \ -1 \ +1]^\top$.

Proof.

$$O_L = \mathfrak{D} \xrightarrow{2,3,4} (\eta(O_L), \eta(I_L)) = (-1, +1), \quad \mathfrak{C} = \{\mathfrak{C}, \mathfrak{C}\} \xrightarrow{2,3,5} (\eta(O_R), \eta(I_R)) \in \{(-1, +1), (+1, -1)\}$$

$$\begin{aligned} \mathfrak{D} \text{ ends on } W_L \xrightarrow{\uparrow\uparrow} W_R, \quad \chi = -1 \text{ until } \mathfrak{C}, \quad \mathfrak{C} \text{ ends on } W_R \xrightarrow{\uparrow\uparrow} W_L, \quad \chi = +1 \text{ until } \mathfrak{D} \\ \Rightarrow \mathfrak{D}, \mathfrak{C}, \mathfrak{D}, \mathfrak{C} \text{ alternate along } \star, \quad \sigma = [-1 \ +1 \ -1 \ +1]^\top \end{aligned}$$



$$\sigma_i = (-1)^{i+1}$$

□

Lemma 2.6.2 (Alexander connecting arc).

$$\mathcal{B} = \{c_s \mid 5 \leq s \leq 4+b\}, \quad \overline{\mathcal{B}} = \{c_s \mid 5+b \leq s \leq p\}, \quad \tilde{\rho}(\mathcal{B}) = \{r, r+1, \dots, r+b-1\},$$

$$\exists \delta \subset \left(\left(\cdot \right) \right), \quad \delta = \left(\begin{array}{c} \text{Diagram} \end{array} \right) : z \rightarrow w \text{ embedded}, \quad \delta \cap \alpha = \emptyset,$$


$$\delta \cap \beta = \{I_L, O_L\} \cup \mathcal{B} \cup \{O_R, I_R\}, \quad \#(\delta \cap c_s) = \mathbf{1}_{\mathcal{B}}(c_s), \quad 5 \leq s \leq p.$$

Proof.

$$\tilde{\rho}(s) = r + s - 5 \ (\mathcal{B}), \quad \tilde{\rho}(s) = r + s - 1 \ (\overline{\mathcal{B}}) \implies \tilde{\rho} \text{ strictly increasing} \implies A \setminus \bigcup_{s=5}^p c_s = \bigsqcup_{j=1}^{p-4} F_j$$

$$z \in \mathbb{D}_z \subset F_L, \quad w \in \mathbb{D}_w \subset F_R, \quad \tilde{\rho}(c_{4+b}) = v-1 \implies \text{sep}(F_L, F_R) = \mathcal{B}$$

$$\delta = \eta_z * \gamma^\vee * \eta_w \text{ embedded}, \quad \delta \cap \alpha = \emptyset, \quad \delta \cap \beta = \{I_L, O_L\} \cup \mathcal{B} \cup \{O_R, I_R\}$$



$$\#(\delta \cap c_s) = \mathbf{1}_{\mathcal{B}}(c_s)$$

□


Corollary 2.6.3 (Alexander increments of the elementary domains). *Each bracket is Θ of the arc it draws,*

$$\left(\begin{array}{c} \text{arc} \\ \text{---} \end{array} \right) = \eta(c), \quad \left(\begin{array}{c} \text{arc} \\ \text{---} \end{array} \right) = \chi(c_s)(1 - \kappa), \quad \left(\begin{array}{c} \text{arc} \\ \text{---} \end{array} \right) = -\chi(c_s) \kappa.$$

Proof.

$$\begin{aligned} & \xrightarrow{2.6.2, 2.5.9} D_\delta = \beta \cdot \delta = \underbrace{(-1) + (+1) + (-1) + (+1)}_{\eta \text{ of the four caps}} + \sum_{c_s \in \mathcal{B}} \chi(c_s) = \ell \\ & \xrightarrow{2.5.8} \Theta(c_s) = \chi(c_s) \mathbf{1}_{\mathcal{B}}(c_s) - e \ell \chi(c_s) = \chi(c_s) (\mathbf{1}_{\mathcal{B}}(c_s) - \kappa), \quad \chi(c) = 0 \implies \Theta(c) = \eta(c) \end{aligned}$$

□



(a) $\pi^{-1}(\alpha \cup \beta) \subset \mathbb{R}^2, \quad \pi_1(T^2) \cong \mathbb{Z}^2 = \langle T_u, T_v \rangle$

(b) $A(x_i) - A(x_j) = n_z(\mathcal{D}) - n_w(\mathcal{D})$

FIGURE 12. The cover of $D[7, 2, 3, 1]$, type A, the arrow tinting each complementary face F by $\#(\partial F \cap \alpha)$ as in Definition 2.3.1, so the two $\mathbb{D}$ per fundamental domain take the darkest wash and $A = (1, 0, -1, 0, 1, 0, -1)$.





FIGURE 13. The cover of $D[9, 2, 3, 2]$, type B, read the same way through Definition 2.3.1, giving $A = (1, 2, 1, 0, 0, -1, -2, -1, 0)$.

Lemma 2.6.4 (Lifted strands in the cyclic cover). *Let $\pi: \widehat{T^2} \rightarrow T^2$ unroll α , with $\widehat{X}$ the lift of a strand X oriented in $+x$ and τ the deck translation by one cell.*

- (1) $\widehat{T^2} = \mathbb{R} \times \mathbb{R}/p\mathbb{Z}$, $\widehat{\alpha} = \mathbb{Z} \times \mathbb{R}/p\mathbb{Z}$, $C_k = [k, k+1] \times \mathbb{R}/p\mathbb{Z}$;
- (2) $a_i^k = \widehat{R}_{v+i} \in \widehat{\alpha} \cap \partial N_R^k$, $b_j^k = \widehat{L}_{j+1} \in \widehat{\alpha} \cap \partial N_L^k$, $\tau a_i^k = a_i^{k+1}$, $\tau b_j^k = b_j^{k+1}$;
- (3) $\widehat{X} \cap \widehat{Y} = \emptyset$ ($X \neq Y$) $\iff \beta$ embedded: $\left(\begin{array}{c} \text{diagram of two crossing strands} \end{array} \right) = \emptyset$;
- (4) $v_i = n_i + 1$: $\left(\begin{array}{c} \text{diagram of a strand crossing a vertical line} \end{array} \right)$.

Proof.

$$\begin{aligned} \pi^{-1}(\alpha) &= \mathbb{Z} \times \mathbb{R}/p\mathbb{Z}, & \tau(C_k, N_L^k, N_R^k) &= (C_{k+1}, N_L^{k+1}, N_R^{k+1}) \implies (1), (2) \\ \beta \text{ embedded} &\iff \widehat{\beta} \text{ embedded} \implies (3) \\ N_R^k &\xrightarrow{\uparrow\downarrow} C_{k+1} \xrightarrow{c^{(1)}} \dots \xrightarrow{c^{(n_i)}} C_{k+n_i+1} \xrightarrow{\uparrow\downarrow} N_L^{k+n_i+1} \implies v_i = n_i + 1 \implies (4) \end{aligned}$$



□


Lemma 2.6.5 (Shift rigidity of the strand matching).

$$\left(\begin{array}{c} \text{strand matching} \end{array} \right) \implies F(4k+i) = 4(k+v_i) + \pi(i) = 4k+i+c \iff 4v_i + \pi(i) - i = c \quad \forall i, \quad \left(\begin{array}{c} \text{crossing} \end{array} \right) = \emptyset.$$

Proof.

$$m < n, F(m) > F(n) \implies \text{interleaved ends} \xrightarrow{2.6.4} \widehat{X}_m \cap \widehat{X}_n \neq \emptyset \implies \perp, \quad F\tau = \tau F \implies F(n+4) = F(n) + 4$$

$$F(i+1) - F(i) \geq 1, \quad \sum_{i=0}^3 (F(i+1) - F(i)) = (F(0) + 4) - F(0) = 4 \implies F(n) = n + c \iff 4v_i + \pi(i) - i = c$$



□

Lemma 2.6.6 (The two mixed diagram types).


$$\left(\begin{array}{c} \text{Diagram A} \end{array} \right) \text{ or } \left(\begin{array}{c} \text{Diagram B} \end{array} \right), \quad \left(\begin{array}{c} \text{Diagram C} \end{array} \right) = \emptyset.$$

Proof.

$$\xrightarrow{2.6.5} \pi(i) - i \equiv c \pmod{4} \quad \forall i, \quad (U) \xrightarrow{2.6.1} (\pi(i) - i) \in \{(2, 0, -2, 0), (0, 2, 0, -2)\} \implies \nexists c \implies (U) = \emptyset$$

$$\pi_A = (3, 0, 1, 2) \implies v_1 = v_2 = v_3 = v_0 + 1, \quad \pi_B = (1, 2, 3, 0) \implies v_0 = v_1 = v_2 = v_3 - 1, \quad \xrightarrow{2.6.4} n_i = v_i - 1$$

$$\xrightarrow{2.5.9} (\eta(O_L), \eta(O_R), \eta(I_L), \eta(I_R)) = (-1, -1, +1, +1) \text{ in both,} \quad A: O_L, O_R, I_L, I_R, \quad B: O_L, I_R, I_L, O_R$$



□


Lemma 2.6.7 (Product domains between consecutive strands).

$$v_{j-1} = v_j = v_{j+1} \implies \mathcal{D}_{j-1,j}, \mathcal{D}_{j,j+1} = \left(\begin{array}{c} \text{strand matching} \end{array} \right) \text{ product domains,} \quad \alpha \cap \beta \cap \text{int } \mathcal{D}_{j\pm 1,j} = \emptyset,$$

$$(X_{j-1}, X_j, X_{j+1}) = (\text{rev}(X_j) - 1, X_j, \text{rev}(X_j) + 1), \quad m_{j-1} = m_j = m_{j+1}, \quad j = \begin{cases} 2, & \text{A,} \\ 1, & \text{B.} \end{cases}$$

Proof.

$$\begin{aligned} \xRightarrow{2.6.6} v_{j-1} = v_j = v_{j+1} &\implies \partial \mathcal{D}_{j\pm 1, j} \subset \widehat{X} \cup \widehat{\alpha}, \quad x \in \alpha \cap \beta \cap \text{int } \mathcal{D} \implies N_R \hookrightarrow \text{int } \mathcal{D} \implies \perp \\ W_L^k \cap (X_{j-1} \cup X_j \cup X_{j+1}) &= \{t_k - 1, t_k, t_k + 1\} \subset \text{one branch of } \tilde{\rho} \implies \tilde{\rho}(t_k + \varepsilon) = \tilde{\rho}(t_k) + \varepsilon \\ \mathbf{1}_{\mathcal{B}} \circ \tilde{\rho} \text{ constant on each branch} &\implies m_{j-1} = m_j = m_{j+1} \end{aligned}$$



$$\tilde{\rho}(t_k + \varepsilon) = \tilde{\rho}(t_k) + \varepsilon$$

□

Lemma 2.6.8 (Conjugate strand counts).

$$\text{A} = \left(\begin{array}{c} \text{Diagram A} \end{array} \right) \implies \mathbf{n} = N\mathbf{1} - e\mathbf{e}_0, \quad \mathbf{m} = m\mathbf{1} - \ell\mathbf{e}_0,$$


$$\text{B} = \left(\begin{array}{c} \text{Diagram B} \end{array} \right) \implies \mathbf{n} = N\mathbf{1} + e\mathbf{e}_3, \quad \mathbf{m} = m\mathbf{1} + \ell\mathbf{e}_3.$$

$$\mathbf{n} = N\mathbf{1} - e\varepsilon, \quad \mathbf{m} = m\mathbf{1} - \ell\varepsilon, \quad \varepsilon = \begin{cases} \mathbf{e}_0, & \text{A,} \\ -\mathbf{e}_3, & \text{B,} \end{cases} \quad \sigma^\top \varepsilon = -1, \quad \mathbf{1}^\top \varepsilon = \pm 1.$$

Proof.

$$\xRightarrow{2.6.1} e = \sum_{s=5}^p \chi(c_s) = \sigma^\top \mathbf{n}, \quad \ell = \sigma^\top \mathbf{m}, \quad \xRightarrow{2.6.6, 2.6.7} \mathbf{n} - N\mathbf{1}, \mathbf{m} - m\mathbf{1} \text{ supported on one strand}$$

$$\begin{aligned} \text{A: } e &= \sigma^\top (N\mathbf{1} + (n_0 - N)\mathbf{e}_0) = N - n_0 \implies \mathbf{n} = N\mathbf{1} - e\mathbf{e}_0, & \text{B: } e &= n_3 - N \implies \mathbf{n} = N\mathbf{1} + e\mathbf{e}_3 \\ \text{likewise } \mathbf{m} &\text{ with } \ell, & \sigma^\top \mathbf{e}_0 &= -1, \quad \sigma^\top \mathbf{e}_3 = +1 \implies \sigma^\top \varepsilon = -1, \quad \mathbf{1}^\top \varepsilon = \pm 1 \end{aligned}$$



$$\mathbf{n} = N\mathbf{1} - e\varepsilon, \quad \mathbf{m} = m\mathbf{1} - \ell\varepsilon$$


□

Corollary 2.6.9 (Monotonicity of A along a strand). *Lift A to $\hat{A} = A \circ \pi$ on $\hat{\alpha} \cap \hat{\beta}$, so that $d\hat{A} := \hat{A}(y) - \hat{A}(x) = \Theta(c)$ across each arc $\hat{c}: x \rightarrow y$ of $\hat{\beta}$. Then*

$$\begin{aligned} d\hat{A}|_{\hat{X}_i} &= \sigma_i(\mathbf{1}_{\mathcal{B}} - \kappa) \in \sigma_i \cdot \{-\kappa, 1 - \kappa\}, & \sigma_i \cdot d\hat{A}|_{\hat{X}_i} \leq 0 &\iff \kappa \geq 1, \\ \kappa \geq 1 &\implies \hat{A}|_{X_0, X_2} = \left(\text{blue strand with slope } \kappa \right), & \hat{A}|_{X_1, X_3} &= \left(\text{blue strand with slope } 1 - \kappa \right), \\ \kappa \leq 0 &\implies \hat{A}|_{X_0, X_2} = \left(\text{blue strand with slope } 1 - \kappa \right), & \hat{A}|_{X_1, X_3} &= \left(\text{blue strand with slope } \kappa \right). \end{aligned}$$

Proof.

$$\stackrel{2.6.3, 2.6.1}{\implies} \sigma_i \cdot \Theta(c_s) = \mathbf{1}_{\mathcal{B}}(c_s) - \kappa \in \{-\kappa, 1 - \kappa\}, \quad \kappa \geq 1 \implies \leq 0, \quad \kappa \leq 0 \implies \geq 0$$




□

Corollary 2.6.10 (Total Alexander increment of a strand).

$$\begin{aligned} \begin{bmatrix} p-4 \\ b \end{bmatrix} &= \begin{bmatrix} \mathbf{1}^\top \mathbf{n} \\ \mathbf{1}^\top \mathbf{m} \end{bmatrix} = \begin{bmatrix} 4N \\ 4m \end{bmatrix} - (\mathbf{1}^\top \varepsilon) \begin{bmatrix} e \\ \ell \end{bmatrix}, & \mathbf{1}^\top \varepsilon &= \begin{cases} +1, & A, \\ -1, & B, \end{cases} \quad S = \kappa N - m \\ \mathbf{m} - \kappa \mathbf{n} = -S \mathbf{1} &\implies \Theta = \text{diag}(\sigma)(\mathbf{m} - \kappa \mathbf{n}) = -S \sigma = S \begin{bmatrix} +1 \\ -1 \\ +1 \\ -1 \end{bmatrix}, & \mathbf{1}^\top \Theta &= 0, \\ \sum_{\hat{X}_0} d\hat{A} = \sum_{\hat{X}_2} d\hat{A} &= \left(\text{blue strand with slope } \kappa \right) = S, & \sum_{\hat{X}_1} d\hat{A} = \sum_{\hat{X}_3} d\hat{A} &= \left(\text{blue strand with slope } 1 - \kappa \right) = -S. \end{aligned}$$

Proof.

$$\begin{aligned} \mathbf{1}^\top \mathbf{n} &= \sum_{s=5}^p 1 = p-4, \quad \mathbf{1}^\top \mathbf{m} = b \stackrel{2.6.8}{\implies} \begin{bmatrix} p-4 \\ b \end{bmatrix} = \begin{bmatrix} 4N \\ 4m \end{bmatrix} - (\mathbf{1}^\top \varepsilon) \begin{bmatrix} e \\ \ell \end{bmatrix} \\ \kappa e = e^2 \ell = \ell &\implies \mathbf{m} - \kappa \mathbf{n} = (m - \kappa N) \mathbf{1} + (\kappa e - \ell) \varepsilon = -S \mathbf{1} \\ \Theta(X_i) = \sigma_i(m_i - \kappa n_i) &\implies \Theta = \text{diag}(\sigma)(\mathbf{m} - \kappa \mathbf{n}) = -S \sigma, \quad \mathbf{1}^\top \Theta = -S \mathbf{1}^\top \sigma = 0 \end{aligned}$$



□

Lemma 2.6.11 (The Alexander profile of $\partial\beta$). *Order the eight generators of one period of $\hat{\beta}$ along $\star$ and let $\hat{\Theta} \in \mathbb{Z}^8$ be the vector of successive increments of $\hat{A}$. Then*


$$\hat{\Theta} = [-1 \quad S \quad -1 \quad -S \quad +1 \quad S \quad +1 \quad -S]^\top, \quad \sum_k \hat{\Theta}_k = 0 \implies \hat{A} \circ \tau^4 = \hat{A},$$

and the graph of $\widehat{A}$ over one period, wall steps in the wall colour and strand steps in the rainbow colour, is

$$\begin{aligned} A: & \left(\begin{array}{c} \text{graph of } \widehat{A} \text{ over one period} \end{array} \right), & B: & \left(\begin{array}{c} \text{graph of } \widehat{A} \text{ over one period} \end{array} \right), \\ A: & \max \widehat{A} = S \text{ at one point}, & B: & \max \widehat{A} = S + 1 \text{ at one point} \quad (S \geq 1). \end{aligned}$$

Proof.

$$\begin{aligned} & \xrightarrow{2.5.9, 2.6.10} \text{cap steps } \eta(c), \quad \text{strand totals } \pm S, \quad \xrightarrow{2.6.6} \eta_A = (-1, -1, +1, +1), \quad \eta_B = (-1, +1, +1, -1) \\ & \widehat{\Theta}_A = [-1 \ S \ -1 \ -S \ +1 \ S \ +1 \ -S]^\top, \quad A: 0, -1, S-1, S-2, -2, -1, S-1, S, 0 \\ & \widehat{\Theta}_B = [-1 \ S \ +1 \ -S \ +1 \ S \ -1 \ -S]^\top, \quad B: 0, -1, S-1, S, 0, 1, S+1, S, 0, \quad \sum_k \widehat{\Theta}_k = 0 \end{aligned}$$




□

Lemma 2.6.12 (Simplicity of the top Alexander grading).

$$S \neq 0 \implies A(\star) = \left(\begin{array}{c} \text{graph of } \widehat{A} \text{ over one period} \end{array} \right) \implies \text{rk} \widehat{HFK}(K, A_{\text{top}}) = \#\{x \mid A(x) = A_{\text{top}}\} = 1.$$

Proof.

$$\begin{aligned} & \xrightarrow{2.6.9} |\kappa| \geq 2 \implies 0 \notin \{\kappa, \kappa - 1\} \implies \widehat{A} \text{ strictly monotone on every strand} \xrightarrow{2.6.11} A_{\text{top}} \text{ attained once} \\ & \kappa = 1 \implies \ell = 1 \xrightarrow{2.2.8} r = 1 \implies \overline{\mathcal{B}} = \emptyset \implies b = p - 4, \quad \mathbf{m} = \mathbf{n}, \quad S = 0 \quad (\text{vacuous}) \\ & \kappa = 0, S = -m \neq 0 \xrightarrow{2.2.8} r = p - 3, \quad \tilde{\rho}(s) \equiv s - 8(\mathcal{B}), \quad \tilde{\rho}(s) \equiv s - 4(\overline{\mathcal{B}}), \quad m \geq 1, \ell = 0 \implies b \geq 2 \\ & \xrightarrow{2.6.11} \rho(s_*) \in \{1, 2\}, \quad c_* \in \overline{\mathcal{B}} \implies s_* \in \{5, 6\} \subset \mathcal{B} \implies \perp \implies c_* \in \mathcal{B}, \quad |\Theta(c_*)| = 1 \xrightarrow{A} \text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1 \end{aligned}$$



□

FIGURE 14. The splitting $S^3 = V_\alpha \cup_{T^2} V_\beta$, the cut annulus $A = T^2 \setminus \alpha$ carrying the gluing $\uparrow \downarrow$, and the resulting K .

Remark 2.6.13 (What §2.6 computes). Every statement above is phrased in $\mathbf{n}, \mathbf{m}, \sigma$, which are read off one period of $\widehat{\beta}$; isotopy of β in T^2 , handle slides over α and stabilisation act on $D[p, 2, b, r]$ without changing K , so the chain

$$\left(\begin{array}{c} \text{graph of } \widehat{A} \text{ over one period} \end{array} \right) \xrightarrow{2.6.8} [\mathbf{n} \ \mathbf{m}] = [N\mathbf{1} \ m\mathbf{1}] - \varepsilon \cdot [e \ \ell] \xrightarrow{2.6.10} \Theta = -S\sigma \xrightarrow{2.6.12} \text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1$$

is an invariant of K and not of the tuple that presented it. In particular $S \neq 0$ forces the top group to be one dimensional, so A_{top} is detected by a single generator and

$$S \neq 0 \implies \text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1, \quad S = 0 \implies \max A - \min A = 2.$$

2.7. Handlebodies, gluing, and the mapping class action. Behind the splitting of Figure 14 sits one handle decomposition,

$$V = h^0 \cup_{\varphi_1} h^1 \cong S^1 \times D^2, \quad \varphi_1 : S^0 \times D^2 \hookrightarrow \partial h^0, \quad D = \{0\} \times D^2, \quad \partial D = \alpha;$$

compressing disks of a solid torus are isotopic, so all information sits in φ , that is in (α, β, z, w) , with $||[\alpha] \cdot [\beta]|| = |e|$ as in Definition 2.2.3.

Lemma 2.7.1 (The wall twists r). *Let τ_α be the Dehn twist along α , acting on φ by regluing A after one further turn. Then*

$$\tilde{\rho} \xrightarrow{\tau_\alpha} \tilde{\rho} + 1 \iff \tau_\alpha \cdot D[p, 2, b, r] = D[p, 2, b, r+1], \quad \langle \tau_\alpha \rangle \cong \mathbb{Z}/p,$$

$$\left(\begin{array}{c} L_s \\ \text{blue curve} \\ R_{\tilde{\rho}(s)} \end{array} \right) \xrightarrow{\tau_\alpha} \left(\begin{array}{c} L_s \\ \text{blue curve} \\ R_{\tilde{\rho}(s)+1} \end{array} \right).$$

Proof.

$$\begin{aligned} \tilde{\rho}(s) - r \text{ and } v - r = b \text{ are independent of } r &\implies r \mapsto r+1 \implies \tilde{\rho} \mapsto \tilde{\rho} + 1, \quad \{R_v, \dots, R_{v+3}\} \mapsto \{R_{v+1}, \dots, R_{v+4}\} \\ &\implies \text{the right wall rotates by one foot} = \tau_\alpha \cdot \varphi, \quad \tilde{\rho} + p \equiv \tilde{\rho} \implies \tau_\alpha^p = \text{id}, \quad \langle \tau_\alpha \rangle \cong \mathbb{Z}/p \end{aligned}$$

□




FIGURE 15. $\text{MCG}(T^2) \cong \text{SL}(2, \mathbb{Z})$ acting on $H_1(T^2) = \mathbb{Z}\mu \oplus \mathbb{Z}\lambda$ with $\mu = [\alpha]$, by $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and its inverse.

Neither condition of Definition 2.2.3 is τ_α -invariant, so a $\mathbb{Z}/p$ orbit leaves the admissible locus, and all three failure modes occur inside one orbit.




FIGURE 16. The $\langle \tau_\alpha \rangle$ -orbit of $D[7, 2, 3, 1]$: the sign of e flips, then $|e| \neq 1$, then β disconnects.

Faded panels are not S^3 -admissible.

Remark 2.7.2 (Reversal and mirror). The reversal of Convention 2.4.1 is $-I \in \text{SL}(2, \mathbb{Z})$, so $\tau_\alpha, -I, \mathbb{D}, \mathbb{C}$ generate the visible action on tuples, and Corollary 2.4.4 picks $e = +1$ out of each $\pm I$ pair.

What §2.6 counts is holomorphic: the $\mathbb{D}$ of Corollary 2.6.3 and the product domain of Lemma 2.6.7 are the only embedded index one shapes, each carrying exactly one strip of Definition 4.2.1; see Proposition 4.2.4 and Example 4.2.5.

Example 2.7.3 (Three diagrams and their vectors). From the traversal, via Lemma 2.6.8,

$$\begin{aligned}
 [5, 2, 0, 2]: \quad \mathbf{n} &= \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \quad \mathbf{m} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad e = \sigma^\top \mathbf{n} = 1, \quad \ell = 0, \quad \kappa = 0, \quad \Theta = \mathbf{0}, \quad S = 0, \\
 [7, 2, 3, 1]: \quad \mathbf{n} &= \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix} = 1 \cdot \mathbf{1} - 1 \cdot \mathbf{e}_0, \quad \mathbf{m} = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad \kappa = 1, \quad \mathbf{m} - \kappa \mathbf{n} = \mathbf{0}, \quad S = 0, \\
 [13, 2, 6, 2]: \quad \mathbf{n} &= \begin{bmatrix} 2 \\ 2 \\ 2 \\ 3 \end{bmatrix} = 2 \cdot \mathbf{1} + 1 \cdot \mathbf{e}_3, \quad \mathbf{m} = \begin{bmatrix} 2 \\ 2 \\ 2 \\ 0 \end{bmatrix} = 2 \cdot \mathbf{1} - 2 \cdot \mathbf{e}_3, \quad \kappa = -2, \\
 \mathbf{m} - \kappa \mathbf{n} &= \begin{bmatrix} 2 \\ 2 \\ 2 \\ 0 \end{bmatrix} + 2 \cdot \begin{bmatrix} 2 \\ 2 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 6 \\ 6 \\ 6 \end{bmatrix} = -S \mathbf{1}, \quad \Theta = \text{diag}(\sigma) \begin{bmatrix} 6 \\ 6 \\ 6 \\ 6 \end{bmatrix} = \begin{bmatrix} -6 \\ +6 \\ -6 \\ +6 \end{bmatrix}, \quad S = -6.
 \end{aligned}$$

$D[7, 2, 3, 1]$ is the sharp case of Lemma 2.6.12: $S = 0$ and A_{top} is attained twice, so $S \neq 0$ cannot be dropped.

3. PROOF OF THEOREM A

Theorem 3.0.1 (Two-rainbow coherence, [37]). *Let $[p, 2, b, r]$ be S^3 -admissible in the sense of Definition 2.2.3 and present $K \subset S^3$, oriented so that O_L runs downwards. Then*

(1) *the only embedded bigons of $T^2 \setminus (\alpha \cup \beta)$ are $\mathbb{D}_z$ and $\mathbb{D}_w$, so*

$$\widehat{\partial} = \mathbf{0}, \quad \text{rk} \widehat{HFK}(K, s) = \#\{x \mid A(x) = s\};$$

(2) *the caps alternate walls along $\star$, cutting β into four strands X_0, X_1, X_2, X_3 with*

$$\sigma = [-1 \quad +1 \quad -1 \quad +1]^\top, \quad e = \sigma^\top \mathbf{n}, \quad \ell = \sigma^\top \mathbf{m};$$

(3) $\mathbb{D} = \{\mathbb{D}_z, \mathbb{D}_w\}$ or $\{\mathbb{D}_z, \mathbb{D}_w\} \iff D$ is $\mathbb{D}$ $\iff K$ is an L -space knot, and then

$$\text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1;$$

(4) $\mathbb{D} = \{\mathbb{D}_z, \mathbb{D}_w\} \implies K$ is not an L -space knot, D is of type A or B, and

$$\mathbf{n} = N \mathbf{1} - e \varepsilon, \quad \mathbf{m} = m \mathbf{1} - \ell \varepsilon, \quad \varepsilon = \begin{cases} \mathbf{e}_0, & \text{A,} \\ -\mathbf{e}_3, & \text{B,} \end{cases} \quad \begin{bmatrix} p-4 \\ b \end{bmatrix} = \begin{bmatrix} 4N \\ 4m \end{bmatrix} - (\mathbf{1}^\top \varepsilon) \begin{bmatrix} e \\ \ell \end{bmatrix};$$

(5) *with $\kappa = e \ell$ and $S = \kappa N - m$,*

$$\mathbf{m} - \kappa \mathbf{n} = -S \mathbf{1} \implies \Theta = \text{diag}(\sigma)(\mathbf{m} - \kappa \mathbf{n}) = -S \sigma, \quad \mathbf{1}^\top \Theta = 0 \implies \widehat{A} \circ \tau^4 = \widehat{A};$$

(6) *exactly one of*

$$S \neq 0 \implies \text{rk} \widehat{HFK}(K, A_{\text{top}}) = 1, \quad S = 0 \implies A(\alpha \cap \beta) = \{-1, 0, +1\}, \quad A_{\text{top}} = 1, \quad g(K) = 1;$$

(7) *the second case is exactly the two ends of the band count,*

$$S = 0 \iff \mathbf{1}_{\mathcal{B}} \equiv \kappa \in \{0, 1\} \iff b \in \{0, p-4\};$$

(8) *the Dehn twist along α acts by*

$$\tau_\alpha \cdot D[p, 2, b, r] = D[p, 2, b, r+1], \quad \langle \tau_\alpha \rangle \cong \mathbb{Z}/p,$$

and (1)–(7) are invariants of K , not of the tuple.

In particular

$$\widehat{\text{rkHFK}}(K, A_{\text{top}}) = 1 \quad \text{or} \quad A_{\text{top}} = 1.$$




FIGURE 17. The four cases of Theorem 3.0.1 at $p = 9$, each drawn on the annulus $T^2 \setminus \alpha$ whose two red edges are glued by $\downarrow \downarrow$, with the sign word $(\eta(O_L), \eta(O_R), \eta(I_L), \eta(I_R))$ printed beneath and the number beside a foot equal to $A(x_h)$.

Proof. (1),

$$\text{admissible} \xrightarrow{2.3.6} \mathbb{D} \in \{\mathbb{D}_z, \mathbb{D}_w\}, \quad \mathbb{D} \xrightarrow{\text{blue}} \mathbb{D} \xrightarrow{\text{red}} \mathbb{D}, \quad \widehat{\partial} = \mathbf{0} \implies \widehat{\text{rkHFK}}(K, s) = \#\{x \mid A(x) = s\}$$

(2),

$$\mathbb{D}, \mathbb{D} \text{ nested} \xrightarrow{2.6.1} \mathbb{D}, \mathbb{D} \text{ alternate} \implies \sigma = [-1 \quad +1 \quad -1 \quad +1]^\top$$

(3),

$$\mathbb{D} = \{\mathbb{D}, \mathbb{D}\}, \{\mathbb{D}, \mathbb{D}\} \xrightarrow{2.3.8} \mathbb{D} \xrightarrow{2.5.1} K \text{ an L-space knot} \xrightarrow{2.5.2} \widehat{\text{rkHFK}}(K, A_{\text{top}}) = 1$$

(4),

$$\mathbb{D} = \{\mathbb{D}, \mathbb{D}\} \xrightarrow{2.3.8} \neg \mathbb{D} \xrightarrow{2.5.1} K \text{ not an L-space knot}, \quad \xrightarrow{2.6.6} \text{A or B}$$

$$\xrightarrow{2.6.8} \mathbf{n} = N\mathbf{1} - e\varepsilon, \quad \mathbf{m} = m\mathbf{1} - \ell\varepsilon \xrightarrow{2.6.10} \begin{bmatrix} p-4 \\ b \end{bmatrix} = \begin{bmatrix} \mathbf{1}^\top \mathbf{n} \\ \mathbf{1}^\top \mathbf{m} \end{bmatrix}$$

(5),

$$\kappa e = e^2 \ell = \ell \xrightarrow{2.6.10} \mathbf{m} - \kappa \mathbf{n} = -S\mathbf{1}, \quad \Theta = -S\sigma, \quad \mathbf{1}^\top \Theta = -S\mathbf{1}^\top \sigma = 0 \xrightarrow{2.6.11} \widehat{A} \circ \tau^4 = \widehat{A}$$

(6),

$$S \neq 0 \xrightarrow{2.6.12} \widehat{\text{rkHFK}}(K, A_{\text{top}}) = 1, \quad S = 0 \xrightarrow{2.6.10, 2.6.9} d\widehat{A}|_{X_i} = 0 \xrightarrow{2.6.11} A(\alpha \cap \beta) = \{-1, 0, +1\}$$

$$\implies A_{\text{top}} = 1, \quad \widehat{\text{HFK}}(K, A_{\text{top}}) \neq 0 \implies g(K) = \max\{s \mid \widehat{\text{HFK}}(K, s) \neq 0\} = 1$$

by genus detection [24]: a one signed sum vanishes only termwise, and the three strand levels are centred by $A = -A \circ \text{conj.}$ (7),

$$d\widehat{A}|_{X_i} = \sigma_i(\mathbf{1}_{\mathcal{B}} - \kappa) \implies \left[S = 0 \iff \mathbf{1}_{\mathcal{B}} \equiv \kappa \in \{0, 1\} \iff \mathbf{m} \in \{\mathbf{0}, \mathbf{n}\} \iff b \in \{0, p-4\} \right]$$

(8),

$$\xrightarrow{2.7.1} \tau_\alpha \cdot D[p, 2, b, r] = D[p, 2, b, r+1], \quad \tau_\alpha^p = \text{id} \xrightarrow{2.6.13} \text{invariants of } K$$

The last line is (3) and (6) together. □




FIGURE 18. The groups $\widehat{\text{rkHFK}}(K, s)$ of the four diagrams of Figure 17 on one pair of axes, the columns and the marks at $s = \pm A_{\text{top}}$ in red, so that the gaps of the $\bowtie$ staircase, the single top of $S \neq 0$ and the three levels of $S = 0$ are read side by side.

Remark 3.0.2 (The two ends of the band count). By 3.0.1,

$$\Delta_K(t) = \pm \left[(N+1)(t+t^{-1}) - (2N+2+\varepsilon) \right], \quad \varepsilon = \begin{cases} +1, & \text{B,} \\ -1, & \text{A,} \end{cases} \quad N = \left\lfloor \frac{p-3}{4} \right\rfloor.$$

4. ANALYTIC FOUNDATIONS AND CLASSIFICATION INPUT

4.1. Detection theorems.

Theorem 4.1.1 (Euler characteristic, [23, Equation (1)]).

$$\sum_{s \in \mathbb{Z}} \chi(\widehat{\text{HFK}}(K, s)) \cdot t^s = \Delta_K(t).$$

Proof. [23, Equation (1)]; see also [27]. □

Theorem 4.1.2 (Genus detection and symmetry, [24, Theorem 1.2]).

$$g(K) = \max\{s \in \mathbb{Z} \mid \widehat{\text{HFK}}(K, s) \neq 0\}, \quad \widehat{\text{HFK}}_M(K, s) \cong \widehat{\text{HFK}}_{M-2s}(K, -s).$$

Proof. [24, Theorem 1.2] and [23, §3]. □

Theorem 4.1.3 (Fiberedness detection, [31, Theorem 1.1], [33]).

$$K \text{ fibered} \iff \widehat{\text{rkHFK}}(K, g(K)) = 1 \implies \Delta_K \text{ monic}.$$

Proof. [31, Theorem 1.1], the genus one case in [33]; monicity by Theorem 4.1.1 at $s = g(K)$. □

Theorem 4.1.4 (Alternating knots, [21, Theorem 1.3], [22, Theorem 1.4]).

$$K \text{ alternating}, \quad \Delta_K(t) = \sum_s a_s \cdot t^s \implies \widehat{\text{rkHFK}}_M(K, s) = \begin{cases} |a_s|, & M = s + \frac{\sigma(K)}{2}, \\ 0, & \text{otherwise,} \end{cases} \quad \tau(K) = -\frac{\sigma(K)}{2}.$$

Proof. [21, Theorem 1.3], [22, Theorem 1.4]. □

Example 4.1.5 (Detection at $D[9, 2, 3, 2]$). From the second panel of Figure 17, via Corollary 2.3.7, Theorem 4.1.1, Theorem 4.1.2 and Theorem 4.1.3,

$$\begin{aligned} \widehat{\text{rkHFK}}(K, *) &= \left(\begin{array}{c} \text{blue bars at } -2, -1, 0, 1, 2 \\ \text{red triangles at } -2, 2 \end{array} \right) = (1, 2, 3, 2, 1), \quad \sum_s \text{rk} = 9 = p, \\ \sum_{s \in \mathbb{Z}} \chi(\widehat{\text{HFK}}(K, s)) \cdot t^s &= -t^2 + 2t - 1 + 2t^{-1} - t^{-2} = \Delta_K(t), \quad \Delta_K(1) = 1, \\ g(K) = 2 = A_{\text{top}}, \quad \widehat{\text{rkHFK}}(K, 2) &= 1 \implies K \text{ fibered}, \quad \Delta_K \text{ monic.} \end{aligned}$$

At $s = 0$ three generators cancel to $\chi = -1$: Theorem 4.1.1 is strictly weaker than Corollary 2.3.7.

Example 4.1.6 (The thin case $D[5, 2, 1, 1]$). For the diagram of Figure 11, by Theorem 4.1.4 with $\sigma(K) = 0$,

$$\Delta_K(t) = -t + 3 - t^{-1}, \quad (\text{rk} \widehat{HFK}(K, s))_{s=1,0,-1} = (1, 3, 1) = (|a_1|, |a_0|, |a_{-1}|),$$

$$\widehat{HFK}_M(K, s) \neq 0 \implies M = s + \frac{\sigma(K)}{2} = s, \quad \tau(K) = -\frac{\sigma(K)}{2} = 0:$$

this is 4_1 , the first entry of Remark 3.0.2, revisited in Example 4.3.3.

4.2. The nonlinear Cauchy–Riemann problem on the strip. Every count in §2.6 is a count of solutions of a nonlinear elliptic boundary value problem with totally real boundary conditions on the strip $\mathbb{R} \times [0, 1]$. The problem is stated in Definition 4.2.1, linearised in Theorem 4.2.2, compactified in Theorem 4.2.3, and solved for the two domains of Corollary 2.6.3 and Lemma 2.6.7 in Proposition 4.2.4.


Definition 4.2.1 (Floer strip). For $x, y \in \alpha \cap \beta$, $\phi \in \pi_2(x, y)$ and J an almost complex structure on $\text{Sym}^1(T^2) = T^2$,

$$\begin{aligned} \mathcal{M}_J(\phi) &= \left\{ u: \mathbb{R} \times [0, 1] \rightarrow T^2 \mid \partial_s u + J(u) \partial_t u = 0, \ u(\mathbb{R}, 0) \subset \beta, \ u(\mathbb{R}, 1) \subset \alpha, \ [u] = \phi \right\}, \\ \lim_{s \rightarrow -\infty} u(s, \cdot) &= x, \quad \lim_{s \rightarrow +\infty} u(s, \cdot) = y, \quad |u(s, t) - x| = O(e^{-\delta|s|}), \quad \delta > 0. \end{aligned}$$

The equation is the $\bar{\partial}$ -operator of the complex structure J , and α, β are totally real in $\text{Sym}^1(T^2) = T^2$, so u is smooth up to the boundary and the energy is the area of the domain,

$$E(u) = \frac{1}{2} \int_{\mathbb{R} \times [0, 1]} |du|^2 ds dt = \int_{\mathbb{R} \times [0, 1]} u^* \omega = \text{Area}(\mathcal{D}(\phi)) = \sum_{R \in \pi_0(T^2 \setminus (\alpha \cup \beta))} n_R(\phi) \cdot \text{Area}(R),$$

so $E(u) \geq 0$ forces $n_R(\phi) \geq 0$ for every region: a solution exists only for a nonnegative domain,



Theorem 4.2.2 (Fredholm index and transversality, [23, §3], [29, §4]). For $p > 2$ and $\delta > 0$ smaller than the smallest angle at x, y ,

$$D_u \bar{\partial}_J: W_\delta^{1,p}(u^* TT^2; \alpha, \beta) \longrightarrow L_\delta^p(\Omega^{0,1} \otimes u^* TT^2), \quad D_u \bar{\partial}_J \xi = \nabla_s \xi + J(u) \nabla_t \xi + (\nabla_\xi J) \partial_t u$$


is Fredholm, and Riemann–Roch computes its index as the Maslov index, a combinatorial quantity in the domain [29, Corollary 4.10],

$$\text{ind } D_u \bar{\partial}_J = \mu(\phi) = e(\mathcal{D}(\phi)) + n_x(\mathcal{D}(\phi)) + n_y(\mathcal{D}(\phi)), \quad e(\mathcal{D}) = \chi(\mathcal{D}) - \frac{1}{4}(\# \text{corners}),$$

n_x, n_y being the averages of the four local multiplicities at x and at y . The universal moduli space is cut out transversally, so by Sard–Smale a generic J makes every $\mathcal{M}_J(\phi)$ a smooth manifold with

$$\dim \mathcal{M}_J(\phi) = \mu(\phi), \quad \mathbb{R} \curvearrowright \mathcal{M}_J(\phi) \text{ freely by } (\sigma \cdot u)(s, t) = u(s - \sigma, t), \quad \widehat{\mathcal{M}}(\phi) = \mathcal{M}_J(\phi) / \mathbb{R};$$

at $\mu(\phi) = 2$,



the flow lines being the orbits of the translation.

Proof. [23, §3], [29, §4]; the linear theory as in [25, Appendix C]. □

Theorem 4.2.3 (Gromov compactness, [23, §3], [29, §7]).

$$\begin{aligned}
 E(u) = \text{Area}(\mathcal{D}(\phi)) < \infty &\text{ depends only on } \phi \implies \text{a priori bound on } \|du\|_{L^2}, \\
 \pi_2(T^2) = 0, \quad n_z(\phi) = 0 &\implies \text{no sphere and no boundary bubbling}, \\
 \mu(\phi) = 1 &\implies \widehat{\mathcal{M}}(\phi) \text{ a compact 0-manifold}, \\
 \mu(\phi) = 2 \implies \partial \widehat{\mathcal{M}}(\phi) = \bigsqcup_{\substack{\phi_1 * \phi_2 = \phi \\ \mu(\phi_1) = \mu(\phi_2) = 1}} \widehat{\mathcal{M}}(\phi_1) \times \widehat{\mathcal{M}}(\phi_2) &\implies \widehat{\partial}^2 = \mathbf{0},
 \end{aligned}$$

the ends of the quotient being the broken strips,

$$\partial \left(\text{wavy line} \right) = \left(\text{B-shape} \right) \sqcup \left(\text{P-shape} \right).$$

Proof. [23, §3], [29, §7]; compactness as in [25, Chapter 4]. $\square$

Proposition 4.2.4 (Automatic regularity of the two shapes).

$$\mathcal{D}(\phi) = \left(\text{B-shape} \right) \text{ or } \left(\text{P-shape} \right) \text{ embedded} \implies \# \widehat{\mathcal{M}}(\phi) = 1 \text{ for every } J.$$

Proof.

$\mathcal{D}(\phi)$ embedded $\implies \mathcal{M}_J(\phi) = \{ \text{biholomorphisms } \mathbb{R} \times [0, 1] \rightarrow \mathcal{D}(\phi) \}$ by the Riemann mapping theorem

$$\dim_{\mathbb{C}} T^2 = 1 \implies \text{coker } D_u \bar{\partial}_J = 0 \quad \forall J \implies \# \widehat{\mathcal{M}}(\phi) = \#(\text{Aut}(\mathbb{R} \times [0, 1]) / \mathbb{R}) = 1 \quad [28, §2], [30, §3]$$

The two shapes are exactly Corollary 2.6.3 and Lemma 2.6.7, so every count in §2.6 is a count of solutions of Definition 4.2.1. $\square$

Example 4.2.5 (The index of the two shapes). A disk with $2n$ convex corners has $e = \chi - \frac{2n}{4} = 1 - \frac{n}{2}$, and a generator at which $\mathcal{D}(\phi)$ occupies k of its four quadrants contributes $\frac{k}{4}$; Theorem 4.2.2 then reads

$$\begin{aligned}
 \left(\text{B-shape} \right) : \quad \mu &= \underbrace{1 - \frac{2}{4}}_e + \underbrace{\frac{1}{4}}_{n_x} + \underbrace{\frac{1}{4}}_{n_y} = \frac{1}{2} + \frac{1}{2} = 1, \\
 \left(\text{P-shape} \right) : \quad \mu &= \underbrace{1 - \frac{4}{4}}_e + \underbrace{\frac{2}{4}}_{n_x} + \underbrace{\frac{2}{4}}_{n_y} = 0 + 1 = 1.
 \end{aligned}$$

On the diagram of Figure 11 the bigon $\mathfrak{B}_z$ has $n_z = 1$, so it is excluded from $\widehat{\partial}$ by Definition 4.2.1; every other embedded bigon is excluded by Theorem 2.3.6, and $\widehat{\partial} = \mathbf{0}$ as in Corollary 2.3.7.

4.3. The genus-one classification.

Theorem 4.3.1 (Schubert classification, [12], [13, Chapter 12]). For p, p' odd, $\gcd(p, q) = \gcd(p', q') = 1$, $0 < q < p$, $0 < q' < p'$,

$$\mathfrak{b}(p, q) \cong \mathfrak{b}(p', q') \iff p = p', \quad q' \equiv q^{\pm 1} \pmod{p}, \quad \overline{\mathfrak{b}(p, q)} \cong \mathfrak{b}(p, p - q),$$

and every $\mathfrak{b}(p, q)$ is a $(1, 1)$ -knot.

Proof. [12], [13, Chapter 12]; the $(1, 1)$ property in [28, §1]. $\square$

Theorem 4.3.2 (Genus one two-bridge knots, [13, Prop. 12.25], [20, §1]). K is genus one and two-bridge $\iff K \cong J(2m, 2\epsilon n)$ with $m, n > 0$, $\epsilon = \pm 1$, and then

$$\begin{aligned} J(2m, 2\epsilon n) &= \mathbf{b}(4mn - \epsilon, 4mn - 2n - \epsilon), \quad V = \begin{bmatrix} m & 1 \\ 0 & \epsilon n \end{bmatrix}, \\ \det(V - t \cdot V^\top) &= \det \begin{bmatrix} m(1-t) & 1 \\ -t & \epsilon n(1-t) \end{bmatrix} = \epsilon mn(1-t)^2 + t = \epsilon mnt^2 + (1 - 2\epsilon mn)t + \epsilon mn, \\ \Delta_K(t) &= t^{-1} \cdot \det(V - t \cdot V^\top) = \epsilon mn(t + t^{-1}) + 1 - 2\epsilon mn, \quad \sigma(K) = 1 + \epsilon, \\ (\widehat{\text{rkHFK}}(K, 1), \widehat{\text{rkHFK}}(K, 0), \widehat{\text{rkHFK}}(K, -1)) &= (mn, 2mn - \epsilon, mn). \end{aligned}$$

Proof. [13, Prop. 12.25], [20, §1], [19]; two-bridge knots are alternating [13, Chapter 12], so the ranks are $|a_s|$ by Theorem 4.1.4. $\square$

Example 4.3.3 (Two tabulated rows as double twist knots). At $(m, n, \epsilon) = (1, 1, -1)$, Theorem 4.3.2 gives

$$\begin{aligned} J(2, -2) &= \mathbf{b}(4 + 1, 4 - 2 + 1) = \mathbf{b}(5, 3), \quad V = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}, \quad V - t \cdot V^\top = \begin{bmatrix} 1-t & 1 \\ -t & t-1 \end{bmatrix}, \\ \det(V - t \cdot V^\top) &= (1-t)(t-1) - 1 \cdot (-t) = -t^2 + 3t - 1 \implies \Delta_K(t) = -t + 3 - t^{-1}, \quad \sigma(K) = 0, \\ (\widehat{\text{rkHFK}}(K, 1), \widehat{\text{rkHFK}}(K, 0), \widehat{\text{rkHFK}}(K, -1)) &= (mn, 2mn - \epsilon, mn) = (1, 3, 1), \end{aligned}$$

which is $D[5, 2, 1, 1]$ of Figure 11 and Example 4.1.6. At $(m, n, \epsilon) = (1, 2, -1)$,

$$J(2, -4) = \mathbf{b}(8 + 1, 8 - 4 + 1) = \mathbf{b}(9, 5), \quad \Delta_K(t) = -2t + 5 - 2t^{-1}, \quad (\widehat{\text{rkHFK}}) = (2, 5, 2),$$

which is both $S = 0$ rows at $p = 9$ – the last two panels of Figure 17 and of Figure 18. These are 4_1 and 6_1 of Remark 3.0.2; Theorem 4.3.7 and Theorem 4.3.4 show they exhaust the possibilities there.

Theorem 4.3.4 (Hatcher–Thurston, [14]).

$$K \text{ two-bridge} \implies S^3 \setminus \text{int } \nu(K) \text{ atoroidal} \implies K \text{ not a nontrivial satellite.}$$

Proof. [14]. $\square$

Theorem 4.3.5 (Morimoto–Sakuma, [15]). *The satellite (1,1)-knots are exactly the Morimoto–Sakuma family; each has tunnel number one and a nontrivial torus knot companion.*

Proof. [15]. $\square$

Theorem 4.3.6 (Satellite genus, [11]). *For $K = P(C)$ with pattern $P \subset V$, winding number w and companion C ,*

$$g(K) = |w| \cdot g(C) + g_V(P), \quad g(K) = 1, \quad g(C) \geq 1, \quad P \not\subset \text{core}(V) \implies w = 0.$$

Proof.

$$\begin{aligned} 1 &= |w| \cdot g(C) + g_V(P), \quad g(C) \geq 1 \implies (|w|, g_V(P)) \in \{(1, 0), (0, 1)\} \\ |w| = 1, \quad g_V(P) = 0 &\implies \exists \text{ annulus } A \subset V, \partial A = P \sqcup \lambda \implies P \simeq \text{core}(V) \implies \perp \implies w = 0 \end{aligned}$$

$\square$

Theorem 4.3.7 (Matsuda, [16]).

$$K \text{ genus one } (1, 1) \iff K \cong J(2m, 2n), \quad mn \neq 0, \quad \text{or} \quad K \in \mathcal{MS}_1,$$

where $\mathcal{MS}_1$ is the genus one Morimoto–Sakuma family.

Proof. [16], [15], with Theorem 4.3.2 and Theorem 4.3.5; see also [36, §1] for the (1,1) satellites and [40, 41] for the two-bridge side. $\square$

Corollary 4.3.8 (Where $S = 0$ lands).

$$S = 0 \xrightarrow{3.0.1} A_{\text{top}} = 1 \xrightarrow{4.1.2} g(K) = 1 \xrightarrow{4.3.7} K \cong J(2m, 2n) \text{ or } K \in \mathcal{MS}_1,$$

and in the second case Theorem 4.3.6 forces $w(P) = 0$.

Proof. Theorem 3.0.1 (6), Theorem 4.1.2, Theorem 4.3.7, Theorem 4.3.6. $\square$

Theorem 4.3.9 (Whitehead doubles, [32, Theorems 1.1–1.2]). $\widehat{HFK}(D_{\pm}(C, t))$ is determined by t and the filtered chain homotopy type of $\widehat{CFK}(C)$, and

$$\tau(D_{\pm}(C, t)) = \begin{cases} 1, & t < 2\tau(C), \\ 0, & t \geq 2\tau(C); \end{cases} \quad \{D_{\pm}(U, t)\}_{t \in \mathbb{Z}} = \{\text{twist knots}\} \cup \{U\}.$$

Proof. [32, Theorems 1.1–1.2]. □

4.4. Parametrization and regular homotopy.

Theorem 4.4.1 (Winding numbers on surfaces, [1], [3]). For S compact oriented, $v \in \Gamma(TS)$ nowhere zero, and regular immersions $\gamma_0, \gamma_1 : S^1 \looparrowright S$,

$$\begin{aligned} \gamma_0 \simeq_{\text{reg}} \gamma_1 &\iff [\gamma_0]_{\text{free}} = [\gamma_1]_{\text{free}} \text{ and } \text{wind}_v(\gamma_0) = \text{wind}_v(\gamma_1), \\ S = \mathbb{R}^2 &\implies [\gamma_0 \simeq_{\text{reg}} \gamma_1 \iff \text{rot}(\gamma_0) = \text{rot}(\gamma_1)]. \end{aligned}$$

Proof. [1] for the plane, [3] in general; see [5, §1.2]. □

This is the analytic fact behind Lemma 2.3.5: an embedded essential $\beta \subset T^2$ is regularly homotopic to a geodesic, so $\vartheta(\beta) = 0$ and the cap senses balance, which is what forces $\sigma = [-1 \quad +1 \quad -1 \quad +1]^{\top}$ in Lemma 2.6.1 and hence $e = \sigma^{\top} \mathbf{n}$ in Lemma 2.6.8. See also [2], [4].

Theorem 4.4.2 (Mapping class parametrization, [18]).

$$\text{Mod}(T^2, \{z, w\}) \twoheadrightarrow \{(1, 1)\text{-knots}\} / \cong, \quad \varphi \longmapsto K_{\varphi}, \quad D_{\varphi} = (T^2, \alpha, \varphi(\beta), z, w).$$

Proof. [18]. □

Under this surjection the orbit of Lemma 2.7.1 is the image of $\langle \tau_{\alpha} \rangle \cong \mathbb{Z}/p$ in $\text{MCG}(T^2) \cong \text{SL}(2, \mathbb{Z})$ of Figure 15, and Figure 16 is the corresponding walk through the fibres; the admissible locus is not τ_{α} -invariant, so the walk leaves it after finitely many steps. Compare [17, 38, 39] for other parametrizations of the same set.

Lemma 4.4.3 (Finite disk bound, [37, Prop. 3.1]). For every tuple $D[p, a, b, r]$ the set of embedded Whitney disks is finite, and

$$\#(\mathcal{D} \cap \tilde{\beta}) \leq \left\lfloor \frac{p}{2} \right\rfloor.$$

Proof. [35], [37, Prop. 3.1]. □

4.5. The Alexander cocycle and the dynamics of the traversal.

Definition 4.5.1 (Return map, cocycle). On $\alpha \cap \beta = \{x_1, \dots, x_p\} \cong \mathbb{Z}/p$, ordered by wall height,

$$F = \omega \circ \sigma : \mathbb{Z}/p \longrightarrow \mathbb{Z}/p, \quad \Theta(h) := \Theta(c_h) = \begin{cases} \eta(c_h), & c_h \text{ a cap,} \\ \chi(c_h)(\mathbf{1}_{\mathcal{D}}(c_h) - \kappa), & c_h \text{ a crossing arc,} \end{cases}$$

where c_h is the outgoing arc of $\star$ at x_h (Definition 2.2.1, Corollary 2.6.3, Lemma 2.5.9).

Lemma 4.5.2 (Eight branches).

$$F(h) - h \in \{\pm 1, \pm 3\} \cup \{\pm(r-5), \pm(r-1)\} \pmod{p}, \quad \text{blue cycle} \iff F \text{ a } p\text{-cycle},$$

$$F(D[9, 2, 3, 2]) = \left(\begin{array}{c} \text{Diagram with 9 vertices } x_1 \dots x_9 \text{ and directed edges} \end{array} \right) = (147632589),$$

cap steps in the wall colour, crossing steps in the rainbow colour.

Proof.

$$\xrightarrow{2.2.5} F(h) - h = \Delta h(c_h) \in \{\pm 1, \pm 3\} \cup \{\pm(r-5), \pm(r-1)\}, \quad F \text{ a } p\text{-cycle} \iff |\star| = p \xleftrightarrow{2.2.7} \text{diagram}$$

□

Theorem 4.5.3 (Transfer function, [6]). *For admissible $D[p, 2, b, r]$,*

$$\sum_{h \in \mathbb{Z}/p} \Theta(h) = 0, \quad A \circ F - A = \Theta, \quad \varphi \circ F - \varphi = \Theta \implies \varphi = A + \text{const},$$

the constant fixed by $A = -A \circ \text{conj}$; the orbit of the skew product $\tilde{F}(h, k) = (F(h), k + \Theta(h))$ through $(h, A(x_h))$ is the graph of A ,

$$\left(\text{graph of } A \right) = \text{graph } A \text{ of } D[9, 2, 3, 2], \quad \text{one period, closing to } A(x_1) = 1.$$

Proof.

$$\xrightarrow{2.6.10} \sum_h \Theta(h) = \mathbf{1}^\top \Theta + \sum_{c \text{ cap}} \eta(c) = 0 + 0 = 0, \quad \xrightarrow{2.5.8} A(F(h)) - A(h) = \Theta(c_h) = \Theta(h)$$

$$(\varphi - A) \circ F = \varphi - A \xrightarrow{4.5.2} \varphi - A \text{ constant on the single } F\text{-orbit}$$

For minimal systems, solvability $\iff$ bounded Birkhoff sums is [6]; on a finite orbit the obstruction is exactly the mean. □

Theorem 4.5.4 (Oscillation equals drift). *For mixed $D[p, 2, b, r]$, normalised as in Corollary 2.4.4,*

$$\begin{aligned} \max A - \min A &= |S| + 2, \quad A_{\text{top}} = \frac{1}{2}|S| + 1, \quad S \in 2\mathbb{Z}, \\ g(K) &= \frac{1}{2}|\kappa N - m| + 1, \quad \widehat{\text{rkHFK}}(K, g(K)) = \begin{cases} 1, & S \neq 0, \\ N + 1, & S = 0. \end{cases} \end{aligned}$$

Proof.

$$\begin{aligned} &\xrightarrow{2.6.9} \text{extrema of } A \text{ at the eight cap nodes} \xrightarrow{2.6.11} \max A - \min A = |S| + 2 \quad \forall S \in \mathbb{Z}, \text{ both types} \\ A = -A \circ \text{conj} &\implies A_{\text{top}} = \frac{1}{2}(\max A - \min A) = \frac{1}{2}|S| + 1 \in \mathbb{Z} \implies S \in 2\mathbb{Z} \xrightarrow{2.3.6, 4.1.2} g(K) = A_{\text{top}} \\ S \neq 0 &\xrightarrow{2.6.12} \widehat{\text{rkHFK}}(K, A_{\text{top}}) = 1, \quad S = 0 \xrightarrow{2.6.10, 2.6.9} A \text{ constant on strands} \\ &\implies \{A = A_{\text{top}}\} = \text{one strand and its two cap feet} \xrightarrow{2.6.8} \widehat{\text{rkHFK}}(K, A_{\text{top}}) = N + 1 \end{aligned}$$

□

Corollary 4.5.5 (Fibredness).

$$K \text{ fibred} \iff \widehat{\text{rkHFK}}(K, g(K)) = 1 \iff S \neq 0 \text{ or } N = 0,$$

and then Δ_K is monic; the unique fibred member with $S = 0$ is 4_1 at $p = 4N + 5 = 5$.

Proof. Theorem 4.1.3, Theorem 4.5.4; $N = 0$ forces type B and $p = 5$ by Corollary 2.6.10; monicity is Theorem 4.1.1 at $s = g(K)$. □

4.6. The tabulation.

Definition 4.6.1 (Tabulation).

$$\mathcal{T}_p = \{D[p, 2, b, r] \mid \text{diagram}\}, \quad \mathcal{T} = \bigcup_p \mathcal{T}_p,$$

the S^3 -admissible tuples of Definition 2.2.3.

Remark 4.6.2 (Growth). $|\mathcal{T}_p| = 8, 6, 12, 12, 18, 16, 22$ for $p = 5, 7, \dots, 17$, ninety four in all, split forty eight $\mathbb{D}$ $\mathbb{Q}$ against forty six mixed; each p contributes the mixed row $D[p, 2, 0, p-3]$, so $\mathcal{T}$ presents infinitely many knot types.

Proof.

$$D[p, 2, 0, p-3] \in \mathcal{T}_p \xrightarrow{3.0.2} \Delta_K = \pm \left[(N+1)(t+t^{-1}) - (2N+2+\varepsilon) \right], \quad N = \left\lfloor \frac{p-3}{4} \right\rfloor \text{ unbounded}$$

□

p, b, r	type	ℓ	κ	N	m	S	A_{top}	$\#A_{\text{top}}$	span
5, 0, 2	B	0	0	0	0	0	1	1	2
5, 1, 1	B	1	1	0	0	0	1	1	2
7, 0, 4	A	0	0	1	0	0	1	2	2
7, 3, 1	A	1	1	1	1	0	1	2	2
9, 0, 6	B	0	0	1	0	0	1	2	2
9, 3, 2	B	-1	-1	1	1	-2	2	1	4
9, 5, 1	B	1	1	1	1	0	1	2	2
11, 0, 8	A	0	0	2	0	0	1	3	2
11, 1, 4	A	-1	-1	2	0	-2	2	1	4
11, 7, 1	A	1	1	2	2	0	1	3	2
13, 0, 10	B	0	0	2	0	0	1	3	2
13, 6, 2	B	-2	-2	2	2	-6	4	1	8
13, 8, 10	B	0	0	2	2	-2	2	1	4
13, 9, 1	B	1	1	2	2	0	1	3	2
15, 0, 12	A	0	0	3	0	0	1	4	2
15, 2, 4	A	-2	-2	3	0	-6	4	1	8
15, 8, 12	A	0	0	3	2	-2	2	1	4
15, 11, 1	A	1	1	3	3	0	1	4	2
17, 0, 14	B	0	0	3	0	0	1	4	2
17, 3, 10	B	-1	-1	3	1	-4	3	1	6
17, 8, 14	B	0	0	3	2	-2	2	1	4
17, 9, 2	B	-3	-3	3	3	-12	7	1	14
17, 13, 1	B	1	1	3	3	0	1	4	2

FIGURE 19. The normalised tabulation: the mixed tuples with $e = +1$ and $p \leq 17$, with type, ℓ , κ , N , m , $S = \kappa N - m$, A_{top} , its multiplicity, and $\text{span} A$; the identities $A_{\text{top}} = \frac{1}{2}|S| + 1$ and $\text{span} A = |S| + 2$ of Theorem 4.5.4 hold in every row.

APPENDIX A. FOUNDATIONAL TOOLS

A.1. Permutation matrices and cycles. A permutation π of $\{1, \dots, d\}$ acts by $P_\pi \cdot \mathbf{e}_n = \mathbf{e}_{\pi(n)}$, one 1 per row and per column, and

$$P_\pi \cdot P_\tau = P_{\pi\tau}, \quad P_\pi^{-1} = P_\pi^\top, \quad \det(P_\pi - \lambda \mathbf{I}_d) = \prod_{i=1}^k (-1)^{d_i} (\lambda^{d_i} - 1), \quad \dim \ker(P_\pi - \mathbf{I}_d) = k,$$

for π a disjoint union of k cycles of lengths $d_1, \dots, d_k$: a nonzero Leibniz term takes $-\lambda$ or 1 from every column of a cycle, and a vector fixed by P_π is constant on each cycle.

Example A.1.1 (The 4-cycle).

$$\pi = (1234), \quad P_\pi = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \quad \det(P_\pi - \lambda \mathbf{I}_4) = \lambda^4 - 1, \quad \ker(P_\pi - \mathbf{I}_4) = \langle \mathbf{1} \rangle.$$

Lemma 2.2.6 applies this with $d = 2p$ and $k = 2$, $d_1 = d_2 = p$, and Lemma 4.5.2 reads the same cycle as the return map F .

A.2. The cohomological equation on a cycle. For F a d -cycle on a finite set X and $\theta : X \rightarrow \mathbb{Z}$, the *cohomological equation* is $\varphi \circ F - \varphi = \theta$, and

$$\exists \varphi \iff \sum_{x \in X} \theta(x) = 0, \quad \varphi(F^k x_0) = \varphi(x_0) + \sum_{j=0}^{k-1} \theta(F^j x_0), \quad \varphi \text{ unique up to a constant,}$$

existence by telescoping, uniqueness because an F -invariant function is constant on the single orbit. The *oscillation* $\max \varphi - \min \varphi$ is the deviation of the partial sums, and does not depend on the choice of constant.

Example A.2.1 (On $\mathbb{Z}/4$).

$$\theta = (2, -1, -2, 1), \quad \sum \theta = 0, \quad \varphi = (0, 2, 1, -1) + c, \quad \max \varphi - \min \varphi = 3.$$

§4.5 applies this with F the traversal, $\theta = \Theta$ and $\varphi = A$; [6] is the compact minimal analogue, where existence is equivalent to bounded Birkhoff sums.

APPENDIX B. REPRODUCIBILITY

B.1. The engine.

LISTING 1. `tworainbow.py`: the normal form, the traversal, drift, admissibility, geometry and gradings of $D[p, 2, b, r]$; every number quoted in the paper is computed here at render time.

```

1 import math
2
3 A = 2
4
5
6 def mod(h, p):
7     return (h - 1) % p + 1
8
9
10 def matching(p, b, r, a=A):
11     arcs = []
12     for i in range(1, a + 1):
13         arcs.append(("L", mod(i, p)), ("L", mod(2 * a + 1 - i, p)))
14         arcs.append(("R", mod(r + b - 1 + i, p)),
15                     ("R", mod(r + b + 2 * a - i, p)))
16     for s in range(2 * a + 1, p + 1):
17         rho = r + s - 2 * a - 1 if s <= 2 * a + b else r + s - 1
18         arcs.append(("L", mod(s, p)), ("R", mod(rho, p)))
19     return arcs
20
21
22 def traverse(p, b, r, a=A):
23     arcs = matching(p, b, r, a)
24     ends = [e for arc in arcs for e in arc]
25     if len(set(ends)) != 2 * p:
26         return None
27     nxt = {}
28     for u, v in arcs:
29         nxt[u], nxt[v] = v, u
30     walk, cur = [], ("L", 1)
31     for _ in range(p):
32         head = nxt[cur]
33         walk.append((frozenset((cur, head)), cur, head))
34         side, h = head
35         cur = ("R" if side == "L" else "L", h)
36     if cur != ("L", 1) or len(walk) != p:
37         return None
38     return walk
39
40
41 def invariants(p, b, r, a=A):
42     walk = traverse(p, b, r, a)
43     if walk is None:
44         return None
45     v = r + b
46     true = {}
47     for i in range(1, a + 1):

```

```

48     true[("L", mod(i, p))] = i
49     true[("L", mod(2 * a + 1 - i, p))] = 2 * a + 1 - i
50     true[("R", mod(v - 1 + i, p))] = v - 1 + i
51     true[("R", mod(v + 2 * a - i, p))] = v + 2 * a - i
52
53     drift = 0
54     caps = []
55     blocks, cur = [], []
56     for k, (_, tail, head) in enumerate(walk):
57         if tail[0] == head[0]:
58
59             sense = 1 if true[head] < true[tail] else -1
60             caps.append((k, tail[0], sense))
61             blocks.append(cur)
62             cur = []
63         else:
64             chi = 1 if tail[0] == "L" else -1
65             drift += chi
66             cur.append((k, chi, tail, head))
67     if cur:
68         if blocks:
69             blocks[0] = cur + blocks[0]
70         else:
71             blocks = [cur]
72     if abs(drift) != 1:
73         return None
74
75     uL = sum(1 for _, wall, s in caps if wall == "L" and s == 1)
76     uR = sum(1 for _, wall, s in caps if wall == "R" and s == 1)
77     dL, dR = a - uL, a - uR
78     assert uL - dL == uR - dR, (p, b, r, uL, uR)
79     return {"p": p, "b": b, "r": r, "drift": drift, "caps": caps,
80            "uL": uL, "uR": uR, "blocks": blocks,
81            "word": "".join("+" if s == 1 else "-" for _, _, s in caps),
82            "sides": "".join(w for _, w, _ in caps)}
83
84
85 def tabulate(pmax=17, a=A):
86     out = []
87     for p in range(3, pmax + 1, 2):
88         for b in range(0, p - 2 * a + 1):
89             for r in range(1, p + 1):
90                 inv = invariants(p, b, r, a)
91                 if inv is not None:
92                     out.append(inv)
93     return out
94
95
96 if __name__ == "__main__":
97     C = tabulate()
98     print(f"admissible [p,2,b,r] with p <= 17 : {len(C)}")
99     by_p = {}
100     for t in C:
101         by_p.setdefault(t["p"], []).append(t)
102     print(" by p:", {p: len(v) for p, v in sorted(by_p.items())})
103
104     bad = [t for t in C if t["uL"] != t["uR"]]
105     print(f"turning balance u_L = u_R : {len(C) - len(bad)}/{len(C)} "
106           f"f{failures {len(bad)}}")
107
108     vals = sorted({(t["uL"], t["uR"]) for t in C})
109     print(" (u_L,u_R) observed:", vals)
110
111     sides = sorted({t["sides"] for t in C})
112     print(" cap side words:", sides)
113
114     coh = [t for t in C if t["uL"] in (0, 2)]
115     mix = [t for t in C if t["uL"] == 1]
116     print(f"coherent {len(coh)} mixed {len(mix)}")
117     print(" coherent cap words:", sorted({t['word'] for t in coh}))
118     print(" mixed cap words: ", sorted({t['word'] for t in mix}))
119
120     nblk = sorted({len(t["blocks"]) for t in C})
121     print(" blocks per diagram:", nblk)
122
123
124 D_OUT, D_IN = 0.215, 0.095
125 U0, U1 = 0.14, 0.86
126

```

```

127
128 def _smooth(x):
129     x = min(1.0, max(0.0, x))
130     return x * x * (3 - 2 * x)
131
132
133 def geometry(p, b, r, a=A):
134     v = r + b
135     deps = [D_OUT - (D_OUT - D_IN) * (i - 1) / max(a - 1, 1)
136             for i in range(1, a + 1)]
137     out = {}
138
139     def rain(wall, h1, h2, dep, n=60):
140         ym, amp = 0.5 * (h1 + h2) - 0.5, 0.5 * (h2 - h1)
141         pts = [(dep * math.sin(math.pi * i / n) if wall == "L"
142                else 1.0 - dep * math.sin(math.pi * i / n),
143                ym - amp * math.cos(math.pi * i / n)) for i in range(n + 1)]
144         key = frozenset(((wall, mod(h1, p)), (wall, mod(h2, p))))
145         out[key] = ((wall, mod(h1, p)), pts)
146
147     for i in range(1, a + 1):
148         rain("L", i, 2 * a + 1 - i, deps[i - 1])
149         rain("R", v - 1 + i, v + 2 * a - i, deps[i - 1])
150     for s in range(2 * a + 1, p + 1):
151         rho = r + s - 2 * a - 1 if s <= 2 * a + b else r + s - 1
152         y0, y1 = s - 0.5, rho - 0.5
153         pts = [(i / 120, y0 + (y1 - y0) * _smooth((i / 120 - U0) / (U1 - U0)))
154                for i in range(121)]
155         key = frozenset(((("L", mod(s, p)), ("R", mod(rho, p))))
156         out[key] = (("L", mod(s, p)), pts)
157     zq = (0.42 * D_IN, float(a))
158     wq = (1.0 - 0.42 * D_IN, (v + a - 1 - 0.5) % p + 0.5)
159     return out, zq, wq
160
161
162 def _cross(a1, a2, b1, b2):
163     def cr(o, x, y):
164         return (x[0] - o[0]) * (y[1] - o[1]) - (x[1] - o[1]) * (y[0] - o[0])
165     d1, d2 = cr(b1, b2, a1), cr(b1, b2, a2)
166     d3, d4 = cr(a1, a2, b1), cr(a1, a2, b2)
167     if (d1 > 0) != (d2 > 0) and (d3 > 0) != (d4 > 0):
168         return 1 if cr(a1, a2, b2) > 0 else -1
169     return 0
170
171
172 def _pair(curve, delta, p):
173     tot = 0
174     for k in (-2, -1, 0, 1, 2):
175         for i in range(len(curve) - 1):
176             c1 = (curve[i][0], curve[i][1] + k * p)
177             c2 = (curve[i + 1][0], curve[i + 1][1] + k * p)
178             for j in range(len(delta) - 1):
179                 tot += _cross(c1, c2, delta[j], delta[j + 1])
180     return tot
181
182
183 _EPS = None
184
185
186 def gradings(p, b, r, a=A):
187     walk = traverse(p, b, r, a)
188     if walk is None:
189         return None
190     geo, zq, wq = geometry(p, b, r, a)
191     drift = sum(1 if t[0] == "L" else -1 for _, t, h in walk if t[0] != h[0])
192     delta = [(zq[0] + 0.0031, zq[1] + 0.0173), (0.5, 0.5 * (zq[1] + wq[1]) + 0.037),
193             (wq[0] - 0.0031, wq[1] + 0.0119)]
194
195     steps = []
196     for key, tail, head in walk:
197         start, pts = geo[key]
198         if start != tail:
199             pts = pts[:-1]
200             chi = 0 if tail[0] == head[0] else (1 if tail[0] == "L" else -1)
201             steps.append((tail, head, chi, _pair(pts, delta, p)))
202     beta_d = sum(s[3] for s in steps)
203
204     A_ = {}
205     cur = 0.0

```

```

206 for tail, head, chi, ixn in steps:
207     if tail[1] not in A_:
208         A_[tail[1]] = cur
209         cur += _EPS * (ixn - drift * chi * beta_d)
210     assert abs(cur) < 1e-9, (p, b, r, cur)
211     vals = A_
212     shift = 0.5 * (max(vals.values()) + min(vals.values()))
213     vals = {h: v - shift for h, v in vals.items()}
214     ints = {h: round(v) for h, v in vals.items()}
215     assert all(abs(vals[h] - ints[h]) < 1e-6 for h in ints), (p, b, r)
216     assert sorted(ints.values()) == sorted(-x for x in ints.values()), \
217         (p, b, r, sorted(ints.values()))
218     sgn = {}
219     for _, head, _, _ in steps:
220         sgn[head[1]] = 1 if head[0] == "R" else -1
221     dpoly = {}
222     for h in ints:
223         dpoly[ints[h]] = dpoly.get(ints[h], 0) + sgn[h]
224     if sum(dpoly.values()) < 0:
225         dpoly = {k: -v for k, v in dpoly.items()}
226     assert abs(sum(dpoly.values())) == 1, (p, b, r, dpoly)
227     return {"A": ints, "drift": drift, "delta": dpoly}
228
229
230 def _calibrate():
231     global _EPS
232     for eps in (1, -1):
233         _EPS = eps
234         ok = True
235         for (p, b, r) in [(3, 1, 1), (5, 2, 2)]:
236             g = gradings(p, b, r, a=1)
237             assert g is not None, (p, b, r)
238             top = max(g["A"], key=lambda h: g["A"][h])
239             v = r + b
240             zc = {1, 2}
241             wc = {mod(v, p), mod(v + 1, p)}
242             want = zc if g["drift"] > 0 else wc
243             if top not in want:
244                 ok = False
245                 break
246         if ok:
247             return
248         raise AssertionError("calibration failed")
249
250
251 _calibrate()

```

B.2. The tabulation checks.

LISTING 2. `chk26.py`: the four-strand matching over the tabulation – cap wall word, χ per block, π by type (Lemma 2.6.6), the shift constant of Lemma 2.6.5, the paired-return identities of Lemma 2.6.8, block totals and the gate of Lemma 2.6.12.

```

1 from collections import Counter
2 import tworainbow as T
3
4 PI_A = (3, 0, 1, 2)
5 PI_B = (1, 2, 3, 0)
6
7
8 def analyse(p, b, r):
9     walk = T.traverse(p, b, r)
10    if walk is None:
11        return None
12    v = r + b
13    true = {}
14    for i in (1, 2):
15        true[("L", T.mod(i, p))] = i
16        true[("L", T.mod(5 - i, p))] = 5 - i
17        true[("R", T.mod(v - 1 + i, p))] = v - 1 + i
18        true[("R", T.mod(v + 4 - i, p))] = v + 4 - i
19
20    def name(t, h):
21        hs = {t[1], h[1]}
22        if t[0] == "L":

```

```

23     return "OL" if hs == {T.mod(1, p), T.mod(4, p)} else "IL"
24     return "OR" if hs == {T.mod(v, p), T.mod(v + 3, p)} else "IR"
25
26 k0 = [k for k, (_, t, h) in enumerate(walk)
27       if t[0] == h[0] and name(t, h) == "OL"][0]
28 w = walk[k0:] + walk[:k0]
29
30 caps, blocks, cur = [], [], []
31 for _, t, h in w:
32     if t[0] == h[0]:
33         caps.append((name(t, h), t, h, 1 if true[h] < true[t] else -1))
34         blocks.append(cur)
35         cur = []
36     else:
37         s = None
38         L = t if t[0] == "L" else h
39         for ss in range(5, p + 1):
40             if T.mod(ss, p) == L[1]:
41                 rho = r + ss - 5 if ss <= 4 + b else r + ss - 1
42                 R = h if t[0] == "L" else t
43                 if T.mod(rho, p) == R[1]:
44                     s = ss
45                     break
46         cur.append((1 if t[0] == "L" else -1, s is not None and s <= 4 + b))
47     blocks = blocks[1:] + [cur]
48     if len(blocks) != 4 or len(caps) != 4:
49         return None
50     e = sum(c for B in blocks for c, _ in B)
51     if abs(e) != 1:
52         return None
53     ell = sum(c for B in blocks for c, _ in B if ib)
54
55     order = "A" if caps[1][0] == "OR" else "B"
56     du = "D" if caps[1] if order == "A" else 3[3] == -1 else "U"
57     walls = "".join("L" if n in ("OL", "IL") else "R" for n, _, _ in caps)
58     word = "".join("+" if s == 1 else "-" for _, _, s in caps)
59
60
61 def aidx(pt):
62     return (true[pt] - v) % p
63
64 def bidx(pt):
65     return true[pt] - 1
66
67 pi, nu = {}, {}
68 names = ["P1", "Q1", "P2", "Q2"]
69 for j, B in enumerate(blocks):
70
71     c0, c1 = caps[j], caps[(j + 1) % 4]
72     if names[j][0] == "P":
73         bpt, apt = c0[2], c1[1]
74     else:
75         apt, bpt = c0[2], c1[1]
76     i = aidx(apt)
77     pi[i] = bidx(bpt)
78     nu[i] = len(B) + 1
79     if sorted(pi) != [0, 1, 2, 3] or sorted(pi.values()) != [0, 1, 2, 3]:
80         return dict(bad="pi not a bijection", p=p, b=b, r=r, pi=pi)
81     cs = {4 * nu[i] + pi[i] - i for i in range(4)}
82
83     n = {names[j]: len(B) for j, B in enumerate(blocks)}
84     m = {names[j]: sum(1 for _, ib in B if ib) for j, B in enumerate(blocks)}
85     return dict(p=p, b=b, r=r, e=e, ell=ell, order=order, du=du,
86               walls=walls, word=word, pi=tuple(pi[i] for i in range(4)),
87               nu=tuple(nu[i] for i in range(4)), cs=cs, n=n, m=m,
88               chis=tuple(tuple(sorted({c for c, _ in B})) for B in blocks))
89
90
91 rows = [analyse(p, b, r) for p in range(5, 18, 2)
92         for b in range(0, p - 3) for r in range(1, p + 1)]
93 rows = [x for x in rows if x and "bad" not in x]
94 mix = [x for x in rows if x["word"].count("+") == 2 and x["word"] != "++--"
95        and x["word"][0] == "-" and set(x["word"]) == {"+", "-"}
96        and x["word"] in ("--+", "-+-")]
97 coh = [x for x in rows if x["word"] in ("----", "++++")]
98
99 print(f"total {len(rows)}   mixed {len(mix)}   coherent {len(coh)}")
100 print("(1) cap wall word:", Counter(x["walls"] for x in mix))
101 print("(2) chi per block:", Counter(x["chis"] for x in mix))

```

```

102 print("(3) order/DU split:", Counter((x["order"], x["du"]) for x in mix))
103 print("    pi by order: ",
104       Counter((x["order"], x["pi"]) for x in mix))
105 badpi = [x for x in mix
106          if x["pi"] != (PI_A if x["order"] == "A" else PI_B)]
107 print("    pi failures:", len(badpi))
108 print("(4) 4 nu_i + pi(i) - i constant:",
109       sum(1 for x in mix if len(x["cs"]) == 1), "/", len(mix))
110 print("    c mod 4 by order:",
111       Counter((x["order"], next(iter(x["cs"]))) % 4) for x in mix))
112
113 bad = []
114 for x in mix:
115     n, m, e, l, o = x["n"], x["m"], x["e"], x["ell"], x["order"]
116     if o == "A":
117         N, M = n["Q2"], m["Q2"]
118         ok = (n["Q2"] == n["P2"] == n["Q1"] and m["Q2"] == m["P2"] == m["Q1"]
119              and n["P1"] - n["P2"] == -e and m["P1"] - m["P2"] == -l
120              and x["p"] == 4 * N + 4 - e and x["b"] == 4 * M - l)
121     else:
122         N, M = n["P2"], m["P2"]
123         ok = (n["P2"] == n["Q1"] == n["P1"] and m["P2"] == m["Q1"] == m["P1"]
124              and n["Q1"] - n["Q2"] == -e and m["Q1"] - m["Q2"] == -l
125              and x["p"] == 4 * N + 4 + e and x["b"] == 4 * M + l)
126     if not ok:
127         bad.append(x)
128 print("(5) paired / block-total identities, failures:", len(bad))
129
130 blk = []
131 for x in mix:
132     k = x["e"] * x["ell"]
133     N = x["n"]["Q2"] if x["order"] == "A" else x["n"]["P2"]
134     M = x["m"]["Q2"] if x["order"] == "A" else x["m"]["P2"]
135     S = k * N - M
136     Th = {X: (1 if X.startswith("Q") else -1) * (x["m"][X] - k * x["n"][X])
137           for X in ("P1", "Q1", "P2", "Q2")}
138     if not (Th["P1"] == Th["P2"] == S and Th["Q1"] == Th["Q2"] == -S):
139         blk.append((x, S, Th))
140 print("    block totals Theta(P_i)=S, Theta(Q_i)=-S, failures:", len(blk))
141
142 gate = []
143 for x in mix:
144     k = x["e"] * x["ell"]
145     N = x["n"]["Q2"] if x["order"] == "A" else x["n"]["P2"]
146     M = x["m"]["Q2"] if x["order"] == "A" else x["m"]["P2"]
147     S = k * N - M
148     g = T.gradings(x["p"], x["b"], x["r"])(["A"])
149     top = max(g.values())
150     mult = sum(1 for u in g.values() if u == top)
151     span = max(g.values()) - min(g.values())
152     if S != 0 and mult != 1:
153         gate.append(("S!=0 but multiple tops", x["p"], x["b"], x["r"], S, mult))
154     if S == 0 and span != 2:
155         gate.append(("S=0 but span", x["p"], x["b"], x["r"], span))
156 print("(6) gate lemma + span, failures:", len(gate), gate[:3])
157 print("    kappa values:", Counter(x["e"] * x["ell"] for x in mix))
158 print("    S values:", dict(sorted(Counter(
159     (x["e"] * x["ell"]) * (x["n"]["Q2"] if x["order"] == "A" else x["n"]["P2"])
160     - (x["m"]["Q2"] if x["order"] == "A" else x["m"]["P2"])
161     for x in mix).items()))))

```

B.3. The vector identities.

LISTING 3. chkvec2.py: $\mathbf{n} = N\mathbf{1} - e\boldsymbol{\varepsilon}$, $\mathbf{m} = m\mathbf{1} - \ell\boldsymbol{\varepsilon}$, $\mathbf{m} - \kappa\mathbf{n} = -S\mathbf{1}$ and $\Theta = -S\boldsymbol{\sigma}$
(Lemma 2.6.8, Corollary 2.6.10) on all forty six mixed tuples.

```

1 import chk26 as C
2
3 SIG = (-1, 1, -1, 1)
4 NAMES = ("P1", "Q1", "P2", "Q2")
5 dot = lambda u, v: sum(a * b for a, b in zip(u, v))
6
7 fail = {k: 0 for k in ("functional", "eps", "shift", "theta", "onetheta")}
8 seen = 0
9 for x in C.mix:
10     seen += 1

```



```

11 n = tuple(x["n"][k] for k in NAMES)
12 m = tuple(x["m"][k] for k in NAMES)
13 e, ell, o = x["e"], x["ell"], x["order"]
14 kap = e * ell
15
16 if not (dot(SIG, n) == e and dot(SIG, m) == ell
17       and sum(n) + 4 == x["p"] and sum(m) == x["b"]):
18     fail["functional"] += 1
19
20 eps = (1, 0, 0, 0) if o == "A" else (0, 0, 0, -1)
21 N = n[1] if o == "A" else n[0]
22 M = m[1] if o == "A" else m[0]
23 if not (dot(SIG, eps) == -1 and abs(sum(eps)) == 1):
24     fail["eps"] += 1
25 if any(n[i] != N - e * eps[i] for i in range(4)) or \
26     any(m[i] != M - ell * eps[i] for i in range(4)):
27     fail["shift"] += 1
28
29 S = kap * N - M
30 if any(m[i] - kap * n[i] != -S for i in range(4)):
31     fail["theta"] += 1
32 Th = tuple(SIG[i] * (m[i] - kap * n[i]) for i in range(4))
33 if Th != tuple(-S * s for s in SIG) or sum(Th) != 0:
34     fail["onetheta"] += 1
35
36 print(f"mixed tuples checked: {seen}")
37 for k, v in fail.items():
38     print(f" {k:<12} failures: {v}")

```

B.4. The statement checks.

LISTING 4. `chk27.py`: checks (1)–(9) – Figure 19, the vectors of Example 2.7.3, span and parity, the extreme rows of Remark 3.0.2, $\tau_\alpha: r \mapsto r+1$ (Lemma 2.7.1), the orbit of Figure 16, the dichotomy (Theorem 3.0.1 (6)–(7)), the four drawn cases of Figure 17, and §4.5.

```

1 import re
2 import chk26 as C
3 import tworainbow as T
4
5 NAMES = ("P1", "Q1", "P2", "Q2")
6
7
8 def scalars(x):
9     o, e, ell = x["order"], x["e"], x["ell"]
10    kappa = e * ell
11    N = x["n"]["Q2"] if o == "A" else x["n"]["P2"]
12    m = x["m"]["Q2"] if o == "A" else x["m"]["P2"]
13    g = T.gradings(x["p"], x["b"], x["r"])
14    A = g["A"]
15    top = max(A.values())
16    return dict(order=o, e=e, ell=ell, kappa=kappa, N=N, m=m,
17              S=kappa * N - m, top=top,
18              cnt=sum(1 for v in A.values() if v == top),
19              span=top - min(A.values()), delta=g["delta"])
20
21
22 def find(p, b, r):
23     return next(z for z in C.mix if (z["p"], z["b"], z["r"]) == (p, b, r))
24
25
26 import tab2 as V
27 rows = V.rows()
28 bad = 0
29 want = sorted((x["p"], x["b"], x["r"]) for x in C.mix if x["e"] == 1)
30 if sorted((q["p"], q["b"], q["r"]) for q in rows) != want:
31     bad += 1
32     print(" row set mismatch")
33 for q in rows:
34     d = scalars(find(q["p"], q["b"], q["r"]))
35     got = (d["order"], d["ell"], d["kappa"], d["N"], d["m"], d["S"],
36           d["top"], d["cnt"], d["span"])
37     drawn = (q["order"], q["ell"], q["kappa"], q["N"], q["m"], q["S"],
38             q["top"], q["cnt"], q["span"])
39     if d["e"] != 1 or got != drawn:
40         bad += 1

```

```

41     print(" row mismatch", (q["p"], q["b"], q["r"]), drawn, got)
42 print(f"(1) fig_tab2: {len(rows)} rows, mismatches {bad}")
43
44
45 bad = 0
46 for tgt, vn, vm in (((5, 0, 2), [0, 0, 0, 1], [0, 0, 0, 0]),
47                    ((7, 3, 1), [0, 1, 1, 1], [0, 1, 1, 1]),
48                    ((13, 6, 2), [2, 2, 2, 3], [2, 2, 2, 0])):
49     x = find(*tgt)
50     if [x["n"][k] for k in NAMES] != vn or [x["m"][k] for k in NAMES] != vm:
51         bad += 1
52     print(" vector mismatch", tgt)
53 print(f"(2) Example 2.7.1 vectors: mismatches {bad}")
54
55
56 f1 = f2 = f3 = f4 = 0
57 for x in C.mix:
58     d = scalars(x)
59     if d["S"] % 2:
60         f1 += 1
61     if d["top"] != abs(d["S"]) // 2 + 1:
62         f2 += 1
63     if d["span"] != 2 * d["top"]:
64         f3 += 1
65     if d["S"] == 0 and d["cnt"] != d["N"] + 1:
66         f4 += 1
67 print(f"(3) tabulation identities over {len(C.mix)} mixed tuples: "
68       f"S even {f1}, A_top {f2}, span {f3}, rank at S=0 {f4}")
69
70
71 bad, seen = 0, 0
72 for x in C.mix:
73     p, b = x["p"], x["b"]
74     if x["e"] != 1 or b not in (0, p - 4):
75         continue
76     seen += 1
77     d = scalars(x)
78     N, eps = d["N"], (1 if d["order"] == "B" else -1)
79     want = {1: N + 1, -1: N + 1, 0: -(2 * N + 2 + eps)}
80     flip = {k: -v for k, v in want.items()}
81     if d["delta"] not in (want, flip) or N != (p - 3) // 4:
82         bad += 1
83     print(" extreme row mismatch", (p, b, x["r"]), d["delta"], want)
84 print(f"(4) Remark 2.7.3 on {seen} extreme rows: mismatches {bad}")
85
86
87 bad, seen = 0, 0
88 for p in range(5, 18, 2):
89     for b in range(0, p - 3):
90         for r in range(1, p + 1):
91             seen += 1
92             for s in range(5, p + 1):
93                 r1 = (r % p) + 1
94                 a0 = r + s - 5 if s <= 4 + b else r + s - 1
95                 a1 = r1 + s - 5 if s <= 4 + b else r1 + s - 1
96                 if T.mod(a1, p) != T.mod(a0 + 1, p):
97                     bad += 1
98 print(f"(5) Lemma 2.7.1, tau_alpha: r -> r+1 on {seen} tuples, mismatches {bad}")
99
100
101 row = []
102 for r in range(1, 6):
103     walk = T.traverse(7, 3, r)
104     if walk is None:
105         row.append((r, "disconnected"))
106         continue
107     e = sum(1 if t[0] == "L" else -1 for _, t, h in walk if t[0] != h[0])
108     row.append((r, f"e={e:d}"))
109 print(f"(6) orbit of D[7,2,3,1] under tau_alpha:",
110       ", ".join(f"r={r}: {v}" for r, v in row))
111
112
113 f = dict(branch=0, values=0, top=0, band=0, iff_b=0)
114 for x in C.mix:
115     d = scalars(x)
116     p, b, S, k = x["p"], x["b"], d["S"], d["kappa"]
117     g = T.gradings(p, b, x["r"])
118     vals = set(g["A"].values())
119     if S != 0 and d["cnt"] != 1:

```

```

120     f["branch"] += 1
121     if S == 0 and vals != {-1, 0, 1}:
122         f["values"] += 1
123     if S == 0 and d["top"] != 1:
124         f["top"] += 1
125     if S == 0 and k not in (0, 1):
126         f["band"] += 1
127     if (S == 0) != (b in (0, p - 4)):
128         f["iff_b"] += 1
129     print("(7) dichotomy clauses (6)-(7) over 46 mixed tuples: "
130           + ", ".join(f"{k} {v}" for k, v in f.items()))
131
132
133     import main2 as M
134     f8 = 0
135     want = [("coherent", None, 5, 1), ("mixed", -2, 2, 1),
136            ("mixed", 0, 1, 2), ("mixed", 0, 1, 2)]
137     for (p, a, b, r), (kind, S, top, mult) in zip(M.CASES, want):
138         d = M.data(p, b, r)
139         got = ("coherent" if d["coh"] else "mixed", d["S"], d["top"], d["mult"])
140         if got != (kind, S, top, mult):
141             f8 += 1
142             print(" figure case mismatch", (p, b, r), got, (kind, S, top, mult))
143         if not d["coh"] and (d["S"] == 0) != (b in (0, p - 4)):
144             f8 += 1
145     print(f"(8) Figures 20-21, four drawn cases: mismatches {f8}")
146
147
148     f9 = dict(steps=0, span=0, parity=0, toprank=0, fibered=0)
149     for x in C.mix:
150         p, b, r = x["p"], x["b"], x["r"]
151         d = scalars(x)
152         walk = T.traverse(p, b, r)
153         seq = [t[1] for _, t, h in walk]
154         F = {seq[k]: seq[(k + 1) % p] for k in range(p)}
155         allowed = set()
156         for v in (1, 3, (r - 5) % p, (r - 1) % p):
157             allowed |= {v % p, (-v) % p}
158         if any((F[h] - h) % p not in allowed for h in seq):
159             f9["steps"] += 1
160         A = T.gradings(p, b, r)["A"]
161         top, mult = max(A.values()), 0
162         mult = sum(1 for v in A.values() if v == top)
163         if max(A.values()) - min(A.values()) != abs(d["S"]) + 2:
164             f9["span"] += 1
165         if d["S"] % 2:
166             f9["parity"] += 1
167         if mult != (1 if d["S"] != 0 else d["N"] + 1):
168             f9["toprank"] += 1
169         if (mult == 1) != (d["S"] != 0 or d["N"] == 0):
170             f9["fibered"] += 1
171     print("(9) Lemma 4.5.2 / Theorem 4.5.4 / Corollary 4.5.5 over 46 tuples: "
172           + ", ".join(f"{k} {v}" for k, v in f9.items()))

```

B.5. The tabulation figure.

LISTING 5. tab2.py: Figure 19, every cell read from chk26.py and tworainbow.py.

```

1 import numpy as np
2 import matplotlib
3 matplotlib.use("Agg")
4 import matplotlib.pyplot as plt
5
6 import tworainbow as T
7
8 plt.rcParams.update({"font.family": "serif", "mathtext.fontset": "cm",
9                     "pdf.fonttype": 42, "savefig.dpi": 300})
10
11 INK, RULE, SOFT = "#000000", "#000000", "#F2F2F2"
12
13 HEAD = [r"$p, b, r$", r"type", r"$\ell$", r"$\kappa$", r"$N$", r"$m$",
14         r"$S$", r"$A_{\mathrm{top}}$", r"$\mathrm{\#}A_{\mathrm{top}}$",
15         r"span"]
16
17
18 def rows():

```

```

19 import chk26 as C
20 out = []
21 for x in sorted(C.mix, key=lambda z: (z["p"], z["b"], z["r"])):
22     if x["e"] != 1:
23         continue
24     o, ell = x["order"], x["ell"]
25     N = x["n"]["Q2"] if o == "A" else x["n"]["P2"]
26     m = x["m"]["Q2"] if o == "A" else x["m"]["P2"]
27     A = T.gradings(x["p"], x["b"], x["r"])["A"]
28     top = max(A.values())
29     S = ell * N - m
30     assert S % 2 == 0
31     assert top == abs(S) // 2 + 1
32     assert top - min(A.values()) == abs(S) + 2
33     out.append(dict(p=x["p"], b=x["b"], r=x["r"], order=o, ell=ell,
34                   kappa=ell, N=N, m=m, S=S, top=top,
35                   cnt=sum(1 for v in A.values() if v == top),
36                   span=top - min(A.values())))
37 return out
38
39
40 def cells(d):
41     return [rf"${d['p']},{d['b']},{d['r']}$",
42            rf"${\mathrm{{{d['order']}}}}$",
43            rf"${d['ell']}$", rf"${d['kappa']}$", rf"${d['N']}$",
44            rf"${d['m']}$", rf"${d['S']}$", rf"${d['top']}$",
45            rf"${d['cnt']}$", rf"${d['span']}$"]
46
47
48 def main(out="fig_tab2"):
49     R = rows()
50     n = len(R)
51     w = np.array([2.05, 1.00, 0.82, 0.90, 0.78, 0.80, 0.88, 1.05, 1.10,
52                  0.95])
53     edge = np.concatenate([[0.0], np.cumsum(w)])
54     mid = 0.5 * (edge[:-1] + edge[1:])
55     W = edge[-1]
56     H = n + 2.6
57
58     fig, ax = plt.subplots(figsize=(0.685 * W, 0.148 * H), dpi=300)
59     fs = 6.4
60
61     for k in range(n):
62         if k % 2:
63             ax.add_patch(plt.Rectangle((0.0, k + 0.5), W, 1.0, color=SOFT,
64                                       lw=0, zorder=0))
65     for x, h in zip(mid, HEAD):
66         ax.text(x, -0.25, h, fontsize=fs, ha="center", va="center",
67               color=INK, zorder=3)
68     for k, d in enumerate(R):
69         for x, c in zip(mid, cells(d)):
70             ax.text(x, k + 1.0, c, fontsize=fs, ha="center", va="center",
71                   color=INK, zorder=3)
72
73     for y, lw in ((-0.95, 0.9), (0.35, 0.55), (n + 0.55, 0.9)):
74         ax.plot([0.0, W], [y, y], color=RULE, lw=lw, zorder=4,
75               solid_capstyle="butt")
76
77     ax.set_xlim(-0.10, W + 0.10)
78     ax.set_ylim(n + 1.05, -1.45)
79     ax.axis("off")
80     for ext in ("png", "pdf"):
81         fig.savefig(f"{out}.{ext}", bbox_inches="tight", pad_inches=0.05,
82                 facecolor="white")
83     plt.close(fig)
84     print(f"wrote {out}: {n} rows, all identities asserted")
85     return R
86
87
88 if __name__ == "__main__":
89     main()

```

REFERENCES

- [1] Hassler Whitney, *On regular closed curves in the plane*, *Compositio Mathematica* **4** (1937), 276–284. Numdam.
- [2] Bruce L. Reinhart, *The winding number on two manifolds*, *Annales de l’Institut Fourier* **10** (1960), 271–283. doi:10.5802/aif.100.

- [3] David R. J. Chillingworth, *Winding numbers on surfaces, I*, Mathematische Annalen **196** (1972), 218–249. doi:10.1007/BF01428050.
- [4] David R. J. Chillingworth, *Winding numbers on surfaces, II*, Mathematische Annalen **199** (1972), 131–153. doi:10.1007/BF01431419.
- [5] Benson Farb and Dan Margalit, *A primer on mapping class groups*, Princeton Mathematical Series **49**, Princeton University Press, Princeton, NJ, 2012. doi:10.1515/9781400839049.
- [6] Walter H. Gottschalk and Gustav A. Hedlund, *Topological Dynamics*, AMS Colloquium Publications **36**, American Mathematical Society, Providence, RI, 1955. doi:10.1090/coll/036.
- [7] Michael Keane, *Interval exchange transformations*, Mathematische Zeitschrift **141** (1975), 25–31. doi:10.1007/BF01236981.
- [8] Isaac P. Cornfeld, Sergei V. Fomin, and Yakov G. Sinai, *Ergodic Theory*, Grundlehren der mathematischen Wissenschaften **245**, Springer, New York, 1982. doi:10.1007/978-1-4615-6927-5.
- [9] Anton Zorich, *Deviation for interval exchange transformations*, Ergodic Theory and Dynamical Systems **17** (1997), no. 6, 1477–1499. doi:10.1017/S0143385797086215.
- [10] F. Binns and H. Zhou, $(1, 1)$ almost L -space knots, arXiv:2304.11475.
- [11] Horst Schubert, *Knoten und Vollringe*, Acta Mathematica **90** (1953), 131–286. doi:10.1007/BF02392437.
- [12] Horst Schubert, *Knoten mit zwei Brücken*, Mathematische Zeitschrift **65** (1956), 133–170. doi:10.1007/BF01473875.
- [13] Gerhard Burde and Heiner Zieschang, *Knots*, second revised and extended edition, de Gruyter Studies in Mathematics **5**, Walter de Gruyter, Berlin, 2003. doi:10.1515/9783110198034.
- [14] Allen Hatcher and William Thurston, *Incompressible surfaces in 2-bridge knot complements*, Inventiones Mathematicae **79** (1985), no. 2, 225–246. doi:10.1007/BF01388971.
- [15] Kanji Morimoto and Makoto Sakuma, *On unknotting tunnels for knots*, Mathematische Annalen **289** (1991), 143–167. doi:10.1007/BF01446565.
- [16] Hiroshi Matsuda, *Genus one knots which admit $(1, 1)$ -decompositions*, Proceedings of the American Mathematical Society **130** (2002), no. 7, 2155–2163. doi:10.1090/S0002-9939-01-06314-6.
- [17] Doo Ho Choi and Ki Hyoung Ko, *Parameterizations of 1-bridge torus knots*, Journal of Knot Theory and Its Ramifications **12** (2003), no. 4, 463–491. doi:10.1142/S0218216503002445; arXiv:math/0112102.
- [18] Alessia Cattabriga and Michele Mulazzani, *$(1, 1)$ -knots via the mapping class group of the twice punctured torus*, Advances in Geometry **4** (2004), no. 2, 263–277. doi:10.1515/adv.2004.016; arXiv:math/0205138.
- [19] Anh T. Tran, *Nonabelian representations and signatures of double twist knots*, Journal of Knot Theory and Its Ramifications **25** (2016), no. 3, Art. 1640013, 9 pp. doi:10.1142/S0218216516400137; arXiv:1507.05241.
- [20] Anh T. Tran, *Twisted Alexander polynomials of genus one two-bridge knots*, Kodai Mathematical Journal **41** (2018), no. 1, 86–97. doi:10.2996/kmj/1521424825; arXiv:1506.05035.
- [21] Peter Ozsváth and Zoltán Szabó, *Heegaard Floer homology and alternating knots*, Geometry & Topology **7** (2003), 225–254. doi:10.2140/gt.2003.7.225; arXiv:math/0209149.
- [22] Peter Ozsváth and Zoltán Szabó, *Knot Floer homology and the four-ball genus*, Geometry & Topology **7** (2003), 615–639. doi:10.2140/gt.2003.7.615; arXiv:math/0301026.
- [23] Peter Ozsváth and Zoltán Szabó, *Holomorphic disks and knot invariants*, Advances in Mathematics **186** (2004), no. 1, 58–116. doi:10.1016/j.aim.2003.05.001; arXiv:math/0209056.
- [24] Peter Ozsváth and Zoltán Szabó, *Holomorphic disks and genus bounds*, Geometry & Topology **8** (2004), 311–334. doi:10.2140/gt.2004.8.311; arXiv:math/0311496.
- [25] Dusa McDuff and Dietmar Salamon, *J-holomorphic Curves and Symplectic Topology*, AMS Colloquium Publications **52**, American Mathematical Society, Providence, RI, 2004. doi:10.1090/coll/052.
- [26] Peter Ozsváth and Zoltán Szabó, *On knot Floer homology and lens space surgeries*, Topology **44** (2005), no. 6, 1281–1300. doi:10.1016/j.top.2005.05.001; arXiv:math/0303017.
- [27] Jacob A. Rasmussen, *Floer homology and knot complements*, Ph.D. thesis, Harvard University, Cambridge, MA, 2003. arXiv:math/0306378.
- [28] Hiroshi Goda, Hiroshi Matsuda, and Takayuki Morifuji, *Knot Floer homology of $(1, 1)$ -knots*, Geometriae Dedicata **112** (2005), 197–214. doi:10.1007/s10711-004-5403-2; arXiv:math/0311084.
- [29] Robert Lipshitz, *A cylindrical reformulation of Heegaard Floer homology*, Geometry & Topology **10** (2006), 955–1097. doi:10.2140/gt.2006.10.955; arXiv:math/0502404.
- [30] Sucharit Sarkar and Jiajun Wang, *An algorithm for computing some Heegaard Floer homologies*, Annals of Mathematics **171** (2010), no. 2, 1213–1236. doi:10.4007/annals.2010.171.1213; arXiv:math/0607777.
- [31] Yi Ni, *Knot Floer homology detects fibred knots*, Inventiones Mathematicae **170** (2007), no. 3, 577–608; erratum, *ibid.* **177** (2009), no. 1, 235–238. doi:10.1007/s00222-007-0075-9; arXiv:math/0607156; erratum DOI; erratum arXiv:0808.0940.
- [32] Matthew Hedden, *Knot Floer homology of Whitehead doubles*, Geometry & Topology **11** (2007), no. 4, 2277–2338. doi:10.2140/gt.2007.11.2277; arXiv:math/0606094.
- [33] Paolo Ghiggini, *Knot Floer homology detects genus-one fibred knots*, American Journal of Mathematics **130** (2008), no. 5, 1151–1169. doi:10.1353/ajm.0.0016; arXiv:math/0603445.
- [34] Joshua Evan Greene, Sam Lewallen, and Faramarz Vafaee, *$(1, 1)$ L -space knots*, Compositio Mathematica **154** (2018), no. 5, 918–933. doi:10.1112/S0010437X17007989; arXiv:1610.04810.
- [35] Gabriel Doyle, *Calculating the knot Floer homology of $(1, 1)$ knots*, senior thesis, Princeton University, Princeton, NJ, 2005. Princeton DataSpace.
- [36] Philip J. P. Ordning, *On knot Floer homology of satellite $(1, 1)$ knots*, Ph.D. thesis, Columbia University, New York, 2006. Thesis PDF.
- [37] M. Fairchild, $\widehat{\text{HFK}}$ of $(1, 1)$ -knots, unpublished notes, PCMI, 2026.
- [38] Alessia Cattabriga and Michele Mulazzani, *Representations of $(1, 1)$ -knots*, Fund. Math. **188** (2005), 45–57. arXiv:math/0501234.

- [39] Philip J. P. Ording, *Constructing doubly-pointed Heegaard diagrams compatible with $(1, 1)$ knots*, J. Knot Theory Ramifications **22** (2013), no. 11, Art. 1350071. arXiv:1110.5675.
- [40] Luigi Grasselli and Michele Mulazzani, *Genus one 1-bridge knots and Dunwoody manifolds*, Forum Math. **13** (2001), no. 3, 379–397. doi:10.1515/form.2001.014.
- [41] Hyeran Cho, Sang Youl Lee, and Hyun-Jong Song, *Derivation of Schubert normal forms of 2-bridge knots from $(1, 1)$ -diagrams*, J. Knot Theory Ramifications **28** (2019), no. 8, Art. 1950048. doi:10.1142/S0218216519500482.
- [42] I. Tecúm, *Rainbow normal forms and the L-space property of one-rainbow $(1, 1)$ -knots*, preprint, 2026.

COLUMBIA UNIVERSITY

Email address: `ilt2109@columbia.edu`