

Homework 4 - Topics in Topology

5 April 2023

Please return Thursday April 13 2023 (extended deadline)

1. Fill in the gaps in the proof of Lemma 5 from Lecture 13.
2. In this exercise we revisit the formula for the product of the algebra \mathbb{D} constructed from the pair of Hopf algebras \mathbb{O} and \mathbb{U} . Recall the product formula was

$$u^p o^q = \sum_{i,j,k,\ell} \overline{\langle o^q, \tilde{u}^i \tilde{u}^j u^\ell \rangle} \langle o^i o^k o^\ell, u^p \rangle o^j u^k$$

- (a) Prove that for all $x \in \mathbb{O}, y \in \mathbb{U}$ we have $\overline{\langle x, y \rangle} = \langle x, \iota_{\mathbb{U}}^{-1}(y) \rangle$.
- (b) Show that $\tilde{u}^i \tilde{u}^j u^\ell = \iota_{\mathbb{U}}(\iota_{\mathbb{U}}^{-1}(u^\ell) u^j u^i)$ so $\overline{\langle o^q, \tilde{u}^i \tilde{u}^j u^\ell \rangle} = \langle o^q, \iota_{\mathbb{U}}^{-1}(u^\ell) u^j u^i \rangle$.
- (c) If we write $(\Delta_{\mathbb{U}})_{123}^1(y) = \sum y_1' y_2'' y_3'''$ then show that (using $o^q = x$ and $u^p = y$) $\langle o^i o^k o^\ell, u^p \rangle = \langle o^i, y' \rangle \langle o^k, y'' \rangle \langle o^\ell, y''' \rangle$ and likewise prove that $(\Delta_{\mathbb{O}})_{123}^1(x) = \sum x_1' x_2'' x_3'''$ implies

$$\langle o^q, \iota_{\mathbb{U}}^{-1}(u^\ell) u^j u^i \rangle = \langle \iota_{\mathbb{O}}^{-1}(x'), u^\ell \rangle \langle x'', u^j \rangle \langle x''', u^i \rangle$$

- (d) We extend the pairing $\langle \rangle$ to unordered tensor products by setting $P_{ij} : \mathbb{O}^{\otimes\{i\}} \times \mathbb{U}^{\otimes\{j\}} \rightarrow k$ to be the pairing applied to tensor factors i and j . Combine this notation with your the results from the previous parts to show that

$$\sum_{i,j,k,\ell} \overline{\langle o^q, \tilde{u}^i \tilde{u}^j u^\ell \rangle} \langle o^i o^k o^\ell, u^p \rangle o^j u^k = m_1^{\bar{2},2} P_{3,1} P_{1,3} (\iota_{\mathbb{O}}^{-1})_{\bar{1}} (\Delta_{\mathbb{O}})_{123}^1(o^q) (\Delta_{\mathbb{U}})_{123}^1(u^p)$$

3. Prove that the element $\alpha_1 = m_1^{123}(X_{31}C_2)$ in \mathbb{D} has the following properties (hint: sketch the corresponding tangles!):
 - (a) α is central: $x\alpha = \alpha x$ for all $x \in \mathbb{D}$.
 - (b) $\iota(\alpha) = \iota^{-1}(\alpha) = \alpha^{\pm 1}$, (you decide if it is + or -)
 - (c) $\Delta_{12}^1(\alpha) = m_1^{135} m_2^{246} (\alpha_5 \alpha_6 X_{12} X_{43})$
4. Suppose K is a 0-framed, rotation 0, one strand XC -diagram for a (long knot). Define $K^{(2)}$ to be a long knot diagram (it has one strand!) with 0 rotation, 0 writhe that passes through K twice. Write an explicit formula for $Z_{\mathbb{D}}(K^{(2)})$ in terms of $Z_{\mathbb{D}}(K)$ and Hopf algebra operations and elements of (tensor powers of) \mathbb{D} .