

LECTURE 8: THE ALEXANDER POLYNOMIAL REVISITED

In the last lecture, we proved that for a certain endomorphism XC -algebra $A = \text{End}(V)$ over the field $\mathbb{Q}(q)$, the invariant Z_A restricted to XC -links produced a polynomial invariant that satisfied certain skein relation, and after a process called deframing and some normalisation, we obtained the Jones polynomial as a instance of the invariant Z_A .

Today we will start showing that the Alexander polynomial that we studied in Lecture 3 can also be recovered from another endomorphism XC -algebra using a very similar scheme, although there is an important difference on how the Alexander polynomial will come out. After that, we would like to give some context about how these XC -algebra structures on $\text{End}(V)$ arise in nature.

1. THE ALEXANDER POLYNOMIAL REVISITED

Recall from Lecture 3 that the Alexander polynomial is a link invariant


$$\Delta : \mathcal{L} \longrightarrow \mathbb{Z}[t^{1/2}, t^{-1/2}]$$

that is uniquely determined by the following two properties:

- (1) $\Delta_{\text{unknot}}(t) = 1$,
- (2) the following *skein relation* holds:

$$\Delta_{L_+}(t) - \Delta_{L_-}(t) = (t^{-1/2} - t^{1/2})\Delta_{L_0}(t),$$

where L_+, L_-, L_0 denote three links that are identical except in a neighbourhood of some point where they look like below,



Let V be a 2-dimensional vector space over the field $\mathbb{Q}(q)$ with a fixed basis (e_1, e_2) , and throughout this section let $A := \text{End}(V) \cong \mathcal{M}_2(\mathbb{Q}(q))$.

Proposition 1. *The elements*

$$\begin{aligned} X_{ij} := & \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} q^{-2} & 0 \\ 0 & q^{-1} \end{pmatrix}_j + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_i \begin{pmatrix} q & 0 \\ 0 & -q^2 \end{pmatrix}_j \\ & + (q^{-2} - q^2) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}_j \in A^{\otimes\{i,j\}} \end{aligned}$$

and

$$C_s := \begin{pmatrix} q^2 & 0 \\ 0 & -q^2 \end{pmatrix}_s \in A^{\otimes\{s\}}$$

define an XC -algebra structure on A .

Proof. We claim that the inverses of these elements are given by

$$\begin{aligned} X_{ij}^- := & \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} q^2 & 0 \\ 0 & q \end{pmatrix}_j + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_i \begin{pmatrix} q^{-1} & 0 \\ 0 & -q^{-2} \end{pmatrix}_j \\ & + (q^2 - q^{-2}) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}_j \in A^{\otimes\{i,j\}} \end{aligned}$$

and

$$C_s^- := \begin{pmatrix} q^{-2} & 0 \\ 0 & -q^{-2} \end{pmatrix}_s \in A^{\otimes\{s\}}.$$

The proof of this and the rest of the axioms follow from some simple computations as in the previous lecture and they are left to the reader. \square

There is a key different with respect to the XC -algebra structure on A that returned the Jones polynomial:

Example 2. The unknot (with writhe 0) as a XC -diagram is simply $O = \check{m}_t^{s,s}(\check{C}_s)$. Now we obtain

$$Z_A(O) = m_t^{s,s}(C_s) = \text{tr} \begin{pmatrix} q^2 & 0 \\ 0 & -q^2 \end{pmatrix} = q^2 - q^2 = 0.$$

For the closed 0-writhe right-handed XC -trefoil 3_1 that was depicted in Lecture 5, a longer but elementary computation shows that $Z_A(3_1) = 0$. So no Alexander here!

This is no incidental: by proceeding as we did for the Jones polynomial during last lecture, we simply get a trivial invariant. This will be part of the homework.

To circumvent this problem, we proceed as follows: let D be a XC -diagram of a link in \mathbb{R}^3 and cut one closed component open by removing a copy of \check{C}^\pm and pull from the endpoints of the just-created open strand so that its endpoints lie in the uppermost and lowermost part, to obtain a XC -tangle diagram \dot{D}^s whose only open strand is labelled s . Note that $\check{m}_v^{u,u}\check{m}_u^{s,t}(\dot{D}^s\check{C}_t^\pm) = D$

Proposition 3. For a XC -diagram \dot{D}^s as above we have

$$Z_A(\dot{D}^s) = \lambda \cdot \text{Id}_s \in A^{\otimes \{s\}}$$

for some $\lambda \in \mathbb{Q}[q, q^{-1}] \subset \mathbb{Q}(q)$.

Proof. Homework. □

So we are forced to obtain a link invariant out of Z_A in a slightly different way, as follows: let D and \dot{D}^s be as before. Then put

$$Z_A^\circ(D) := \lambda = \frac{1}{2} m_t^{s,s}(Z_A(\dot{D}^s)) \in \mathbb{Q}[q, q^{-1}] \subset \mathbb{Q}(q)$$

where λ is the scalar of Proposition 3.

Lemma 4. The previous value is a well-defined invariant of D , that is, it is independent of the choice of strand we cut open.

Proof. Let us write $D = \check{m}_q^{t,t}\check{m}_p^{s,s}\check{m}_t^{t,v}\check{m}_s^{s,u}(E\check{C}_u\check{C}_v^-)$ for some XC -diagram E with two open strands s, t whose endpoints lie in the uppermost and lowermost part (draw a diagram for this!). A similar argument to that one must use in Proposition 3 ensures that

$$Z_A(E) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_s \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}_t + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_s \begin{pmatrix} \mu_1 & 0 \\ 0 & \mu_2 \end{pmatrix}_t + \begin{pmatrix} 0 & \gamma_1 \\ 0 & 0 \end{pmatrix}_s \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}_t + \begin{pmatrix} 0 & 0 \\ \gamma_2 & 0 \end{pmatrix}_s \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}_t$$

for some scalars $\lambda_i, \mu_i, \gamma_i \in \mathbb{Q}(q)$. Now, applying Z_A to the equality of XC -tangle diagrams

$$\check{m}_2^{2,t}\check{m}_2^{2,3}\check{m}_1^{1,s}\check{m}_1^{1,4}(\check{X}_{1,2}\check{X}_{3,4}E) = \check{m}_2^{2,3}\check{m}_2^{t,2}\check{m}_1^{1,4}\check{m}_1^{s,1}(\check{X}_{1,2}\check{X}_{3,4}E)$$

(draw a picture of this!) and comparing coefficients yields

$$\gamma_2 = q^2\gamma_1, \quad \mu_1 - \lambda_2 = (q - q^{-3})\gamma_2. \quad (1)$$

Likewise, applying Z_A to the equality of XC -diagrams

$$\check{m}_3^{3,6}\check{m}_2^{t,4}\check{m}_1^{s,5}(\check{X}_{4,3}\check{X}_{5,6}E) = \check{m}_3^{3,6}\check{m}_2^{4,t}\check{m}_1^{5,s}(\check{X}_{4,3}\check{X}_{5,6}E)$$

and again comparing coefficients yields

$$\lambda_1 - \lambda_2 = q\gamma_2 = \mu_1 - \mu_2, \quad \lambda_1 - \mu_1 = q^{-1}\gamma_1 = \lambda_2 - \mu_2. \quad (2)$$

Now, note that

$$\dot{D}^s = \check{m}_q^{t,t}\check{m}_t^{t,v}(E\check{C}_v^-), \quad \dot{D}^t = \check{m}_p^{s,s}\check{m}_s^{s,u}(E\check{C}_u).$$

Therefore we have

$$c = m_q^{t,t}m_t^{t,v}(Z_A(E)C_v^-) = q^{-2} \begin{pmatrix} \lambda_1 - \lambda_2 & 0 \\ 0 & \mu_1 - \mu_2 \end{pmatrix}_s = q^{-1} \begin{pmatrix} \gamma_2 & 0 \\ 0 & \gamma_2 \end{pmatrix}_s = q\gamma_1 \text{Id}_s$$

and similarly

$$Z_A(\dot{D}^t) = m_p^{s,s}m_s^{s,u}(Z_A(E)C_u) = q^2 \begin{pmatrix} \lambda_1 - \mu_1 & 0 \\ 0 & \lambda_2 - \mu_2 \end{pmatrix}_t = q\gamma_1 \text{Id}_s,$$

where we have used (1) and (2). Therefore $Z_A(\mathring{D}^s) = Z_A(\mathring{D}^t)$ and we conclude. \square

Theorem 5. *The invariant Z_A° satisfies the following properties:*

- (1) $Z_A^\circ(\text{unknot}) = 1$,
- (2) *Let $L, L_\alpha, L_{\alpha^-}$ be XC -diagrams of links in \mathbb{R}^3 that are identical except in a neighbourhood of a point where L looks like $\check{1}$, L_α like the positive kink $\check{\alpha}$ and L_{α^-} like the negative kink $\check{\alpha}^-$. Then*

$$Z_A^\circ(L_\alpha) = Z_A^\circ(L) = Z_A^\circ(L_{\alpha^-}).$$

- (3) *Let L_+, L_-, L_0 denote three XC -diagrams of links in \mathbb{R}^3 that are identical except in a neighbourhood of some point where they look like \check{X}, \check{X}^- and $\check{1}\check{1}$, respectively. Then the following skein relation holds:*

$$Z_A^\circ(L_+) - Z_A^\circ(L_-) = (q^{-2} - q^2)Z_A^\circ(L_0).$$

Proof. (1) is trivial, (2) follows from the equalities

$$Z_A(\check{\alpha}_i) = Z_A(\check{\alpha}_i^-) = Z_A(\check{1}_i) = \text{Id}_i,$$

which are readily verified, and (3) follows from an argument similar to the one given in the last lecture using the fundamental equality (easily verified)

$$X_{ij} - X_{ji}^{-1} = (q^{-2} - q^2)P_{ij},$$

where

$$P_{ij} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_j + \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}_j + \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}_i \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}_j + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_i \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_j$$

as in Lecture 7. \square

In particular, (2) in the previous theorem says that Z_A° is an invariant of links in \mathbb{R}^3 as it is invariant under the classical $(\Omega 1)$ move, so we do not need to deframe it.

Using the last theorem, the following becomes elementary.

Theorem 6. *Let D be a XC -diagram of a link L in \mathbb{R}^3 . Then*

$$\Delta_L(t) = Z_A^\circ(D)|_{q^2=t^{1/2}}$$

2. PRODUCING NEW XC -ALGEBRAS

What we would like to do next is to give a context to the XC -algebra structures on $\text{End}(V) \cong \mathcal{M}_2(\mathbb{Q}(q))$ that recover the Jones and Alexander polynomial of links via the invariant Z_A . This algebra (with either of the two XC structures, of course) is four-dimensional and then computations are rather elementary (although some can be involved). Now for an arbitrary XC -algebra A , the computation of $Z_A(D)$ can be potentially difficult, e.g. if A is infinite-dimensional and non-commutative.

How can we deal with this? One possibility is to focus not on the value $Z_A(K)$ but instead on the value $f(Z_A(K)) \in B$ where $f : A \rightarrow B$ is a certain map, and then studying an alternative way of describing $(f \circ Z_A)(K)$ that does not require to compute the value $Z_A(K)$ first. Since Z_A is a knot invariant, so is $f \circ Z_A$.

In fact, we will focus on a very particular case of this where B is another algebra and $f : A \rightarrow B$ is an algebra map. The following observation is key:

Lemma 7. *Let A be an XC -algebra over a field k with preferred elements $X \in A \otimes A$ and $C \in A$, and let B be a k -algebra.*

Any k -algebra homomorphism $f : A \rightarrow B$ induces an XC -algebra structure on B where the preferred elements are given by

$$X' := (f \otimes f)(X) \quad , \quad C' := f(C).$$

Proof. This follows directly from the fact that f is an algebra homomorphism. \square

Therefore, if A is large and difficult to understand, we can try to pass the XC -algebra structure to another algebra B instead that is easier to handle. The price to pay then is that the universal invariant $Z_B(K)$ produced with the new XC -algebra B is potentially weaker. What is sure is that the universal invariant with respect to B is determined by that with respect to A .

Definition 8. Let A, B be XC -algebras preferred elements X, X' and C, C' , respectively. An XC -algebra homomorphism is an algebra map $f : A \rightarrow B$ such that $X' = (f \otimes f)(X)$ and $C' = f(C)$.

Corollary 9. Let A, B be XC -algebras and let $f : A \rightarrow B$ be a XC -algebra map. Then for XC -tangle T , the value $Z_B(T)$ is determined by $Z_A(T)$,

$$Z_B(T) = f^{\otimes \mathcal{L}(T)}(Z_A(T)).$$

The main example when we want to apply this is when $B = \text{End}(V)$. In particular, we will see that the XC -algebra structures on $\text{End}(V)$ that recover the Jones and Alexander polynomials are transferred from XC -algebra structures on a more complicated algebra (that we have already seen!).

3. REPRESENTATIONS

In this section, we will let A be simply an algebra over some field k .

Definition 10. A *representation* of A or a (left) A -module is a pair (V, ρ) where V is k -vector space and ρ is a k -algebra homomorphism

$$\rho : A \rightarrow \text{End}(V),$$

where $\text{End}(V)$ is viewed as a k -algebra with multiplication given by composition, $f \cdot g := f \circ g$, and unit given by the identity map Id_V .

In other words, a representation of A is a k -vector space V together with a bilinear map

$$A \times V \rightarrow V$$

such that

$$(ab) \cdot v = a \cdot (b \cdot v) \quad , \quad 1 \cdot v = v$$

for all $a, b \in A, v \in V$.

Exercise 11. Show that the two phrasings in the definition above are indeed the same via

$$a \cdot v = \rho(a)(v).$$

We will freely use both notations interchangeably.

Examples 12. (1) For any k -algebra A , then $V = 0$ gives a representation, called the *zero representation*.

(2) If $A = k$, then a k -representation is simply a k -vector space.

(3) If $V = A$, then multiplication in the algebra defines a representation, $\rho(a)b := ab$, called the *regular representation* of A .

(4) Let $k := \mathbb{Q}$ and $A := \mathbb{Q}[\mathbb{Z}/2]$ the rational group algebra of $\mathbb{Z}/2$. If V is any \mathbb{Q} -vector space with a choice of basis (e_1, e_2) , then

$$\rho(0) := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad , \quad \rho(1) := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

extended \mathbb{Q} -linearly (that is, $\rho(\lambda \cdot 0 + \mu \cdot 1) := \lambda \rho(0) + \mu \rho(1)$) defines a representation. Indeed, $0 = 1 \cdot 0^1$ is the trivial element of the group algebra so it must be mapped to the identity of V . Since

$$\rho(0) = \rho(1 + 1) = \rho(1)\rho(1) = \rho(1)^2,$$

then the condition for ρ to be a representation is that the matrix defined by $\rho(1)$ squares to the identity, and it is indeed the case.

¹Watch out! In the expression $1 \cdot 0 \in \mathbb{Q}[\mathbb{Z}/2]$, we have $1 \in \mathbb{Q}$ and $0 \in \mathbb{Z}/2$.

- (5) Let us now consider $k = \mathbb{Z}/2$ (this is a field of characteristic 2) and $A := k[\mathbb{Z}/2]$ the group algebra of the group $\mathbb{Z}/2$. Then

$$\rho(0) := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad , \quad \rho(1) := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

defines a representation of A , because $\rho(1)$ squares to the identity mod 2.

Exercise 13. Show that the representation from Examples 12.(5) above is precisely the regular representation of $k[\mathbb{Z}/2]$, where $k = \mathbb{Z}/2$.

Definition 14. Let V be a representation of A . A *subrepresentation* of V is a k -linear subspace $W \subset V$ such that $\rho(a)w \in W$ for all $w \in W$, so that W is a representation of A on its own.

Observe that for any representation V of A , both 0 and V are always subrepresentations.

Definition 15. A representation $V \neq 0$ of A is *irreducible* (or *simple*) if its only subrepresentations are 0 and V .

Example 16. The representation $V = \text{span}_{\mathbb{Q}}(e_1, e_2)$ from Examples 12.(4) is not irreducible because both $W := \mathbb{Q}e_1 \subset V$ and $W' := \mathbb{Q}e_2 \subset V$ are one-dimensional subrepresentations.

The reason we are studying representation theory of algebras is the following.

Corollary 17. Let A be a XC -algebra, let (V, ρ) be a representation of A and endow $B := \text{End}(V)$ with the XC -algebra structure inherited from A . Then for any XC -tangle T we have

$$Z_B(T) = \rho^{\otimes \mathcal{L}(T)}(Z_A(T)).$$

4. THE $U(\mathfrak{sl}_2)$ EXAMPLE

Recall from Lecture 4 that $U(\mathfrak{sl}_2)$ is the \mathbb{C} -algebra generated by the elements X, Y, H and relations

$$[H, X] = 2X \quad , \quad [H, Y] = -2Y \quad , \quad [X, Y] = H$$

where $[u, v] := uv - vu$.

It follows from Lecture 4 that giving a representation

$$\rho : U(\mathfrak{sl}_2) \longrightarrow \text{End}(V)$$

on a vector space V (where we fix a basis) consists on giving matrices $\rho(X), \rho(Y), \rho(H)$ such that

$$[\rho(H), \rho(X)] = 2\rho(X) \quad , \quad [\rho(H), \rho(Y)] = -2\rho(Y) \quad , \quad [\rho(X), \rho(Y)] = \rho(H)$$

where again $[u, v] := uv - vu$.

Proposition 18. For $n \geq 1$, the matrices

$$\rho_n(X) = \begin{pmatrix} 0 & n-1 & 0 & \cdots & 0 \\ 0 & 0 & n-2 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \ddots & \ddots & 1 \\ 0 & 0 & \cdots & 0 & 0 \end{pmatrix} \quad , \quad \rho_n(Y) = \begin{pmatrix} 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \\ 0 & 2 & \ddots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & n-1 & 0 \end{pmatrix} ,$$

$$\rho_n(H) = \begin{pmatrix} n-1 & 0 & \cdots & 0 & 0 \\ 0 & n-3 & \cdots & 0 & 0 \\ \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & -n+3 & 0 \\ 0 & 0 & \cdots & 0 & -n+1 \end{pmatrix}$$

define a representation of $U(\mathfrak{sl}_2)$ on an n -dimensional complex vector space V_n .

Proof. It is left to the reader to check that the above matrix relations indeed hold. \square

Proposition 19. Each of the representations V_n above is irreducible for $n \geq 1$.

Proof. Let (e_1, \dots, e_n) be the fixed basis for V_n . First we make the following claims:

Claim 1: Every finite-dimensional representation W of $U(\mathfrak{sl}_2)$ has a vector $w \in W$ such that w is an eigenvector of H and $w \in \ker X$.

Claim 2: Any vector $w \in V_n$ satisfying the properties of Claim 1 is proportional to e_1 .

To see how the proposition follows from the claims, let $V' \subset V_n$ be a subrepresentation. Let $v' \in V'$ satisfy the above-mentioned properties for $W = V'$. Then this element viewed in $W = V$ also satisfies the above properties. Since $e_1 \in V_n$ satisfies these properties, we must have that v' and e_1 are proportional, thus $e_1 \in V'$. But this means that $e_2 = Ye_1$, $e_3 = \frac{1}{2}Y^2e_1$, etc all belong to V' as well, so $V' = V$.

Let us now prove the Claim 1. Since W is finite-dimensional and we are working over \mathbb{C} , the operator H has some eigenvector w , $Hw = \lambda w$. If $w \in \ker X$, we are done; else consider the sequence of vectors $(X^n w)_n$. It is easy to see from the relations defining $U(\mathfrak{sl}_2)$ that

$$H(X^n w) = X^n(H + 2n)w = (\lambda + 2n)(X^n w),$$

so we get a sequence of eigenvectors of H with different eigenvalues. This implies that there must be some n such that $X^n w \neq 0$ and $X^{n+1}w = 0$, as W is finite-dimensional. In this case $X^n w$ is the desired element.

To see Claim 2, such a w must be a scalar multiple of some e_i as it is an eigenvector of H , and in particular of e_1 as none of the others belong to $\ker X$. \square

Remark 20. In fact, it is not hard to show (although we will not do it in this course) that

- (1) These are the only irreducible representations (up to isomorphism²) of $U(\mathfrak{sl}_2)$,
- (2) Every finite-dimensional representation of $U(\mathfrak{sl}_2)$ is isomorphic to a direct sum of the irreducible representations V_n .

This gives a full description of all finite-dimensional representations of $U(\mathfrak{sl}_2)$.

5. THE $U_q(\mathfrak{sl}_2)$ EXAMPLE

Now recall that $U_q(\mathfrak{sl}_2)$ is the $\mathbb{C}(q)$ -algebra generated by X, Y, K, K^{-1} subject to the relations

$$KK^{-1} = 1 = K^{-1}K \quad , \quad KX = q^2XK \quad , \quad KY = q^{-2}YK \quad , \quad [X, Y] = \frac{K - K^{-1}}{q - q^{-1}}.$$

For any integer n , let us put

$$[n]_q := \frac{q^n - q^{-1}}{q - q^{-1}} = q^{n-1} + q^{n-3} + \dots + q^{-n+3} + q^{-n+1}.$$

Proposition 21. For $n \geq 1$ and $\varepsilon = \pm 1$, the matrices

$$\begin{aligned} \rho_{\varepsilon,n}(X) &= \varepsilon \begin{pmatrix} 0 & [n-1]_q & 0 & \dots & 0 \\ 0 & 0 & [n-2]_q & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \ddots & \ddots & [1]_q \\ 0 & 0 & \dots & 0 & 0 \end{pmatrix}, \quad \rho_{\varepsilon,n}(Y) = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 \\ [1]_q & 0 & \dots & 0 & 0 \\ 0 & [2]_q & \ddots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & [n-1]_q & 0 \end{pmatrix}, \\ \rho_{\varepsilon,n}(K) &= \varepsilon \begin{pmatrix} q^{n-1} & 0 & \dots & 0 & 0 \\ 0 & q^{n-3} & \dots & 0 & 0 \\ \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & q^{-n+3} & 0 \\ 0 & 0 & \dots & 0 & q^{-n+1} \end{pmatrix} \end{aligned}$$

define an n -dimensional representation of $U_q(\mathfrak{sl}_2)$, that we denote $V_{\varepsilon,n}$.

²An isomorphism between two representations V_1, V_2 of an algebra A is a linear isomorphism $\varphi : V_1 \rightarrow V_2$ that preserves the action of A , that is $\varphi(a \cdot v) = a \cdot \varphi(v)$ for any $v \in V_1$

Proof. This is again left to the reader to check. The last relation amounts to showing that

$$[n - 2i + 1]_q = [n - i]_q[i]_q - [i - 1]_q[n - i + 1]_q$$

for which the equalities

$$[-n]_q = -[n]_q \quad , \quad [n + m]_q = q^n[m]_q + q^{-m}[n]_q$$

are very helpful. \square

Remark 22. As in the $U(\mathfrak{sl}_2)$ case, one can show the following:

- (1) Each of these representations $V_{\varepsilon, n}$ is irreducible,
- (2) These are the only irreducible representations (up to isomorphism) of $U_q(\mathfrak{sl}_2)$,
- (3) Every finite-dimensional representation of $U_q(\mathfrak{sl}_2)$ is isomorphic to a direct sum of the irreducible representations $V_{\varepsilon, n}$.

This gives a full description of all finite-dimensional representations of $U_q(\mathfrak{sl}_2)$.

As we hinted before, our interest in studying the representation theory of $U_q(\mathfrak{sl}_2)$ was motivated by the possibility of transferring an XC -algebra structure on $U_q(\mathfrak{sl}_2)$ to $\text{End}(V)$. This XC -structure indeed exists.

Theorem 23 (\triangle). *The algebra $U_q(\mathfrak{sl}_2)$ has an XC -algebra structure determined by the elements*

$$X_{ij} = q^{H_i H_j / 2} \sum_{n=0}^{\infty} q^{n(n-1)/2} \frac{(q - q^{-1})^{2n}}{[n]_q!} X_i^n Y_j^n \quad , \quad C_i := K_i$$

where $K = q^H$ and $[n]_q! := [n]_q[n-1]_q \cdots [1]_q$.

Warning 24. The reader should notice that, strictly speaking, the previous statement is not correct, as the expression for X is not a well-defined element in $U_q(\mathfrak{sl}_2)$ (it is given by an infinite sum and $q^{H_i H_j / 2}$ is not defined). However it is possible to make it correct by replacing $U_q(\mathfrak{sl}_2)$ by a larger algebra $U_h(\mathfrak{sl}_2)$ where the element X does make sense. The definition of this larger algebra $U_h(\mathfrak{sl}_2)$ requires some technical considerations that we will not cover in this lecture course, and so we will content ourselves with this ill-defined version.

Now let $V := V_{1,2}$ be the 2-dimensional representation of $U_q(\mathfrak{sl}_2)$ for the choice $\varepsilon = +1$, and put $\rho = \rho_{1,2}$. Explicitly,

$$\rho(X) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad , \quad \rho(Y) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad , \quad \rho(K) = \begin{pmatrix} q & 0 \\ 0 & q^{-1} \end{pmatrix}.$$

Theorem 25. *The XC -algebra structure on $\text{End}(V)$ induced by the algebra map*

$$\rho : U_q(\mathfrak{sl}_2) \longrightarrow \text{End}(V)$$

is precisely the XC -algebra structure of Proposition 12 from Lecture 7 that recovered the Jones polynomial (after a change of variables $q' := q^2$).

Proof sketch. We recover the element C for $\text{End}(V)$ directly from $\rho(K)$ (after replacing q by q^2). For the element X , using the extension algebra one can show that

$$\rho^{\otimes\{i,j\}}(q^{H_i H_j / 2}) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix}_j + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_i \begin{pmatrix} q^{-1/2} & 0 \\ 0 & q^{1/2} \end{pmatrix}_j$$

Then we simply note that $\rho(X)^2 = 0 = \rho(Y)^2$, so after applying $\rho^{\otimes\{i,j\}}$ to X_{ij} the sum becomes finite, in fact it only has two summands,

$$\begin{aligned} \rho^{\otimes\{i,j\}}(X_{ij}) &= \left[\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix}_j + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_i \begin{pmatrix} q^{-1/2} & 0 \\ 0 & q^{1/2} \end{pmatrix}_j \right] \\ &\quad \times \left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}_i \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}_j + (q - q^{-1}) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}_j \right] \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} q^{1/2} & 0 \\ 0 & q^{-1/2} \end{pmatrix}_j + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}_i \begin{pmatrix} q^{-1/2} & 0 \\ 0 & q^{1/2} \end{pmatrix}_j \end{aligned}$$

$$+ (q^{1/2} - q^{-3/2}) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}_i \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}_j$$

so we obtain the same elements X and C as in Lecture 7 after replacing q by q^2 . \square

Remark 26. Using the rest of irreducible representations $V_{1,n}$ of $U_q(\mathfrak{sl}_2)$, one similarly obtains an XC -algebra structure on $\text{End}(V_{1,n})$ that similarly leads to a collection J_n of link polynomial invariants called the *coloured Jones polynomials*, where $J = J_2$.

It is possible to show, but we will not do it here, that the XC -algebra structure on $\text{End}(V)$ from Proposition 1 that recovers the Alexander polynomial also arises from a 2-dimensional representation of $U_\xi(\mathfrak{sl}_2)$ where $q = \xi = e^{\pi i/2}$ is a root of unity.