

The Category Zoo

Rough, unfinished notes! Use at your own risk.

Jorge Becerra

July 13, 2026

Contents

1	Linear Categories	4
1.1	Semiadditive category	4
1.2	Additive category	4
1.3	Preaddivitive category	5
1.4	$\mathbb{k}$ -linear category or $\mathbb{k}$ -category	5
1.5	Additive envelope	5
1.6	Karoubi envelope	5
1.7	Cauchy completion	5
1.8	Simple objects (non-abelian version)	6
1.9	Semisimple category (non-abelian version)	6
1.10	Generators	6
2	Miscellaneous	7
2.1	Action groupoid	7
2.2	Action of a group on a category	7
2.3	Homotopy fixed point for an action on a category	8
3	Limits and Colimits	8
3.1	Kernel (via equalizer/pullback)	8
3.2	Cokernel (via Coequalizer)	9
3.3	Left/Right Exact Functors	9
3.4	RAPL / LAPC	9
3.5	Linear functor + source semisimple imply exact	9
3.6	Pullbacks from Products and Equalizers	9
3.7	Pushouts from Coproducts and Coequalizers	10
3.8	Ends and coends	11
3.9	Homotopy fibre of a functor	13
3.10	Homotopy pullback	14
3.11	Yoneda lemma	14
3.12	Bimodules and the hom	16
3.13	coYoneda lemma	18
4	Monoidal Categories and Related Structures	20
4.1	Monoidal category	20
4.2	Left rigid category	20

4.3	Right rigid category	20
4.4	Rigid or autonomous category	20
4.5	Pivotal or sovereign category	21
4.6	Spherical category (classical definition)	21
4.7	Braided category	21
4.8	Ribbon or tortile category (take 1)	22
4.9	Balanced category	22
4.10	Braided pivotal = balanced rigid	22
4.11	Ribbon or tortile category (take 2)	22
4.12	Monoidal $\mathbb{k}$ -linear category	22
4.13	Pre-fusion category (non-abelian version)	23
4.14	Fusion category (non-abelian version)	23
4.15	Modular category (non-abelian, fusion version)	23
4.16	Drinfeld center	23
4.17	Müger center	24
4.18	Equivalent Characterizations of Modular Categories	24
4.19	Grothendieck-Verdier category	25
4.20	Pivotal Grothendieck-Verdier category	25
4.21	Braided Grothendieck-Verdier category	25
4.22	Ribbon Grothendieck-Verdier category	25
5	Abelian Categories and Related Structures	26
5.1	Kernels and cokernels	26
5.2	Monos and epis	26
5.3	Abelian category	26
5.4	Linear + object semisimple imply abelian	27
5.5	The Hom bifunctor is left exact in both variables	28
5.6	The functor $- \otimes_R M$ is right exact	30
5.7	Flat module	30
5.8	Right exact functors that coincide on projective objects coincide everywhere	30
5.9	Monoidal abelian category	32
5.10	Monoidal abelian $\mathbb{k}$ -linear category	32
5.11	Projective and injective objects	32
5.12	Direct sums and summands of projective and injective objects	33
5.13	Projective cover and injective hull	34
5.14	Socle and top	35
5.15	Semisimple category	35
5.16	Semisimplicity vs. Projectivity in Abelian Categories	36
5.17	Locally finite abelian $\mathbb{k}$ -linear category	37
5.18	Finite abelian $\mathbb{k}$ -linear category	37
5.19	A finite abelian category has a projective generator	38
5.20	Left/right exactness, adjointness and (dual) representability	38
5.21	Deligne product	39
5.22	Kelly product	39
5.23	(Multi)tensor, fusion and ring categories	40
5.24	Indecomposable projective objects in a finite tensor category	41
5.25	Nakayama functors: classical setting	42
5.26	The Nakayama permutation	43

5.27	The Nakayama automorphism	44
5.28	Comparison: quasi-Frobenius vs Frobenius algebras	45
5.29	Nakayama functors: Morita-independent version	46
5.30	The canonical coend Δ	47
5.31	The distinguished invertible object (take 1)	47
5.32	Unimodular finite tensor category	47
5.33	The distinguished invertible object (take 2)	48
5.34	Self-duality of Δ in a unimodular ftc	49
5.35	Spherical finite tensor category à la Douglas-Schommer-Pries-Snyder	49
5.36	Eilenberg-Watts theorem	50
5.37	The canonical Hopf algebra in a braided finite tensor category	50
5.38	Non-degenerate braided finite tensor category	52
5.39	Weakly-factorisable braided finite tensor category	52
5.40	Factorisable braided finite tensor category	52
5.41	Modular category (non-ss)	53
5.42	Hopf algebra terminology and relation with categorical ones	53
6	Module categories	54
6.1	Definition	54
6.2	$\mathcal{C}$ -module functor	55
6.3	Module category over a multitensor category	55
6.4	Finite module categories	56
6.5	Balanced Deligne product	56
6.6	Relative Serre functors	56
6.7	Pivotal module categories	56
6.8	Finite categories	56
6.9	Finite tensor categories	57
6.10	Finite module categories	57
6.11	Internal homs	57
6.12	Deligne product	58
6.13	Balanced Deligne product	59
6.14	The distinguished invertible element	60
6.15	stuff	61
6.16	Relative Serre functors	61
6.17	Pivotal module categories	63
6.18	Bimodules and the hom	64
7	Cyclic and modular operads	65
7.1	Classical operads and their algebras	65
7.2	Classical cyclic operads and their algebras	67
7.3	Classical modular operads	67
7.4	Ordinary, cyclic and modular operads à la Costello-Müller-Woike	68
7.5	The correspondence between groupoids and 1-homotopy types	70
7.6	The (framed) little disc operad E_n and fE_n	71
7.7	Disc algebras vs. framed E_n -algebras	72
7.8	The cyclic and modular endomorphism operad	74
7.9	Classifications of cyclic and modular algebras	75
8	Factorisation homology	76

8.1	Definition	76
8.2	Properties	76
8.3	Quantum structure sheaf or distinguished object	76
8.4	Moduli algebra	77
8.5	Skein algebra (categorical version))	77
9	Factorization homology	77
9.1	A brief review of factorization homology	77
9.2	The $\mathcal{A}^{\boxtimes n}$ -module structure	79
10	Modified traces	80
10.1	Twisted traces	80
10.2	Module-compatibility of twisted traces	80
10.3	Uniqueness	81
10.4	Left/right modified traces	81
10.5	Two-sided modified traces	83
11	Modular functors	83
11.1	Bakalov-Kirillov's modular functors	83
11.2	Open modular functors	85

These notes have been partially written copying snippets from the ChatGPT/Gemini AIs. Use at your own risk!

The author would like to thank Lukas Woike for explaining and clarifying to him many of the ideas appearing here.

1 Linear Categories

1.1 Semiadditive category

A category with a zero object (i.e., both initial and terminal object; and therefore zero morphisms) and with finite products and coproducts in which the canonical map

$$\coprod_i X_i \rightarrow \prod_i X_i$$

is an isomorphism—hence finite biproducts exist, and these are typically called direct sums $\oplus$.

1.2 Additive category

Any preadditive category is naturally enriched over commutative monoids: given maps $f, g : X \rightarrow Y$, define $f + g$ as the composite:

$$X \rightarrow X \times X \cong X \amalg X \rightarrow Y \times Y \cong Y \amalg Y \rightarrow Y$$

where the first map is the diagonal map (id, id) , the second map is $(f, 0) \amalg (0, g)$, and the third one is $\text{id} \amalg \text{id}$.

If such an enrichment extends to abelian groups, that is, if every map f has an inverse $-f$ for the previously defined sum, then the category is said to be an additive category.

Alternatively, additive means **Ab**-enriched plus finite products and coproducts exist (and because of the enrichment they are automatically isomorphic). Typically, the biproduct is represented by the direct sum sign $\oplus$.

A functor between additive categories is additive if it preserves the finite biproducts, i.e., $F(0) = 0$ and $F(X \oplus Y) \cong F(X) \oplus F(Y)$.

1.3 Preadditive category

A category enriched over abelian groups.

1.4 $\mathbb{k}$ -linear category or $\mathbb{k}$ -category

A category enriched over $\mathbb{k}$ -modules, where $\mathbb{k}$ is a commutative ring with unit. A linear category might *not* be additive (finite biproducts might not exist).

A functor between $\mathbb{k}$ -linear categories is linear if $F_{X,Y}$ between the hom's is a linear map. Linear functors are automatically additive, i.e., they preserve the biproducts.

1.5 Additive envelope

As mentioned before, a linear category might not have all finite biproducts. But it always embeds into a category that has them. The additive envelope $\mathbf{Add}(\mathcal{C})$ of a linear category $\mathcal{C}$ has objects formal direct sums $\bigoplus_i X_i$ of objects of $\mathcal{C}$ and arrows given by matrices of arrows in $\mathcal{C}$, with composition modelled by matrix multiplication. There is a canonical functor $\mathcal{C} \hookrightarrow \mathbf{Add}(\mathcal{C})$, and it is determined by the universal property that if $\mathcal{D}$ has all finite biproducts, then there is a bijection

$$\mathbf{LinFun}(\mathcal{C}, \mathcal{D}) \cong \mathbf{LinFun}(\mathbf{Add}(\mathcal{C}), \mathcal{D})$$

1.6 Karoubi envelope

Remember that an idempotent in a category is an endomorphism $f : X \rightarrow X$ of a certain object such that $f \circ f = f$. A splitting for an idempotent f is a pair of arrows $r : Y \rightarrow X$ and $s : Y \rightarrow X$ such that $s \circ r = f$ and $r \circ s = \text{id}_Y$. Splittings for idempotents are unique up to unique isomorphism.

A category $\mathcal{C}$ is idempotent complete or Karoubian if all idempotents split. A linear category might have idempotents that do not split. But it always embeds into a category where they do. The Karoubian envelope or idempotent completion $\mathbf{Kar}(\mathcal{C})$ of a linear category $\mathcal{C}$ has objects idempotents $f : X \rightarrow X$ (more precisely pairs (X, f)).

An arrow between two idempotents (X, f) and (Y, g) is an arrow $h : X \rightarrow Y$ such that $h = g \circ h \circ f$, with composition taken from $\mathcal{C}$. There is a canonical functor $\mathcal{C} \hookrightarrow \mathbf{Kar}(\mathcal{C})$, taking X to (X, id) , and it is determined by the universal property that if $\mathcal{D}$ is idempotent complete, then there is a bijection

$$\mathbf{LinFun}(\mathcal{C}, \mathcal{D}) \cong \mathbf{LinFun}(\mathbf{Kar}(\mathcal{C}), \mathcal{D})$$

1.7 Cauchy completion

A linear category is Cauchy complete if it has all finite biproducts and it is idempotent complete. The Cauchy completion of $\mathcal{C}$ is $\mathbf{Cauchy}(\mathcal{C}) := \mathbf{Kar}(\mathbf{Add}(\mathcal{C}))$ (warning: this is *not* equivalent to $\mathbf{Add}(\mathbf{Kar}(\mathcal{C}))$).

Every linear abelian category is Cauchy complete; the converse is not true.

1.8 Simple objects (non-abelian version)

If X is an object of a $\mathbb{k}$ -linear category, then $\text{End}(X)$ is naturally a $\mathbb{k}$ -algebra with unit the identity. It is not hard to see that the following items are equivalent:

- (i) the map $\mathbb{k} \rightarrow \text{End}_{\mathcal{C}}(X)$, $k \mapsto k \cdot \text{id}$ is an isomorphism of $\mathbb{k}$ -modules;
- (ii) the map $\mathbb{k} \rightarrow \text{End}_{\mathcal{C}}(X)$, $k \mapsto k \cdot \text{id}$ is an isomorphism of $\mathbb{k}$ -algebras;
- (iii) the $\mathbb{k}$ -algebra $\text{End}_{\mathcal{C}}(X)$ is isomorphic to $\mathbb{k}$;
- (iv) the $\mathbb{k}$ -module $\text{End}_{\mathcal{C}}(X)$ is free of rank 1.

If an object X satisfies any of these equivalent descriptions, we say that X is simple.

1.9 Semisimple category (non-abelian version)

Suppose $\mathcal{C}$ is a Cauchy complete linear category whose isomorphism classes of objects form a set. Then the following conditions are equivalent:

- (S) There is a set S and an equivalence of categories $\mathcal{C} \cong \text{Vec}_{\text{fd}}(S)$ (the category of S -graded vector spaces and grading-preserving maps).
- (End) For every object $X \in \mathcal{C}$, the endomorphism space $\text{End}(X)$ is a finite-dimensional complex semisimple algebra, i.e., a multimatrix algebra.
- (Obj) Every $X \in \mathcal{C}$ is isomorphic to a finite direct sum of simple objects $X = \bigoplus_{i=1}^n X_i$, where each pair X_i, X_j are either isomorphic or distinct.
- (Müger) There is a set of pairwise distinct simple objects $\{X_s\}_{s \in S}$ such that for any $A, B \in \mathcal{C}$, the composition map

$$\bigoplus_{s \in S} \mathcal{C}(A \rightarrow X_s) \otimes_{\mathbb{k}} \mathcal{C}(X_s \rightarrow B) \xrightarrow{\circ} \mathcal{C}(A \rightarrow B)$$

is an isomorphism. (The direct sum is taken in the category of finite-dimensional vector spaces.)

Being *finitely* semisimple means that S is a finite set, i.e., there are finitely many isomorphism classes of simple objects.

Fact: A Cauchy complete linear category is semisimple if and only if it is abelian and all short exact sequences split.

More important fact: Linearity and object semisimplicity already imply abelianness! See Section 5.4.

1.10 Generators

Let $\mathcal{C}$ be an additive category. An object $G \in \mathcal{C}$ is called a generator if any object of $\mathcal{C}$ is a quotient of some direct sum of copies of G , that is, if there is an epimorphism $G^{\oplus n} \twoheadrightarrow X$.

Lemma. G is a generator if and only if $\text{Hom}(G, -)$ is faithful.

Proof. Let $\mathcal{C}$ be an additive category and $G \in \mathcal{C}$.

Step 1: If G is a generator, then $\text{Hom}(G, -)$ is faithful.

Suppose G is a generator. By definition, for every $X \in \mathcal{C}$, there exists an epimorphism

$$f : G^{\oplus I} \twoheadrightarrow X$$

for some index set I .

Let $g : X \rightarrow Y$ be a morphism such that $\text{Hom}(G, g) = 0$. We want to show $g = 0$.

Consider the composition $g \circ f : G^{\oplus I} \rightarrow Y$. Since $\text{Hom}(G, g) = 0$, it follows that $g \circ f = 0$. But f is an epimorphism, so $g = 0$. Therefore, $\text{Hom}(G, -)$ is faithful.

Step 2: If $\text{Hom}(G, -)$ is faithful, then G is a generator.

Suppose $\text{Hom}(G, -)$ is faithful. Let $X \in \mathcal{C}$ be arbitrary. Consider the morphism

$$c : \bigoplus_{f \in \text{Hom}(G, X)} G \longrightarrow X$$

which sends the f -th copy of G via f to X . We claim that c is an epimorphism.

Indeed, suppose $h : X \rightarrow Y$ satisfies $h \circ c = 0$. Then for every $f \in \text{Hom}(G, X)$, we have $h \circ f = 0$. By the faithfulness of $\text{Hom}(G, -)$, this implies $h = 0$. Hence c is an epimorphism, and X is a quotient of a sum of copies of G . $\square$

2 Miscellaneous

2.1 Action groupoid

If G is a group and X is a G -set, the *action groupoid* $X//G$ is the groupoid (=cat with all arrows isos) with objects the elements of X , and an arrow $x \rightarrow y$ is a $g \in G$ such that $gx = y$. Composition = multiplication on G .

This specialises to the usual way of viewing a group G as a groupoid with one object and arrows G . This is then $*//G$.

2.2 Action of a group on a category

Recall that if X is a set or a topological space, a G -set or a G -space is the same as a functor $X : *//G \rightarrow \text{Set} / \text{Top}$.

Now let us view $*//G$ as a 2-category with only identity 2-cells, and let $\mathbf{Cat}$ be the 2-category of categories, functors and natural transformations. Then a (non-strict/homotopy) action of G on a category $\mathcal{C}$ is a pseudofunctor $\mathcal{C} : *//G \rightarrow \mathbf{Cat}$.

Explicitly, this is the data of: for every $g \in G$, an equivalence $\varphi_g : \mathcal{C} \xrightarrow{\sim} \mathcal{C}$, together with the data of natural isos $\varphi_g \varphi_h \cong \varphi_{gh}$, subject to the associativity condition that the isomorphism $\varphi_g \varphi_h \varphi_k \cong \varphi_g \varphi_{hk} \cong \varphi_{ghk}$ equals the iso $\varphi_g \varphi_h \varphi_k \cong \varphi_{gh} \varphi_k \cong \varphi_{ghk}$.

Alternative point of view: regard G as a discrete strict monoidal category with monoidal product equals the product of G . Recall that $\text{End}(\mathcal{C})$ is a monoidal category with monoidal product composition. Then an action of G on $\mathcal{C}$ amounts to the data of a strong monoidal functor $G \rightarrow \text{End}(\mathcal{C})$.

For an object $X \in \mathcal{C}$ we write $g.X := \varphi_g(X)$.

2.3 Homotopy fixed point for an action on a category

Let $\mathcal{C}$ be a G -category. A *homotopy fixed point* is the data of an object $X \in \mathcal{C}$ together with isomorphisms $\alpha_g : g.X \xrightarrow{\cong} X$ satisfying

$$\begin{array}{ccccc} h.g.X & \xrightarrow{h\alpha_g} & h.X & \xrightarrow{\alpha_h} & X \\ \downarrow \cong & & \nearrow \alpha_{hg} & & \\ (hg).X & & & & \end{array}$$

The category $\mathcal{C}^{hG}$ of homotopy fixed points has objects htpy fixed points and arrows maps $f : X \rightarrow Y$ in $\mathcal{C}$ such that

$$\begin{array}{ccc} g.X & \xrightarrow{\alpha_g} & X \\ \downarrow g.f & & \downarrow f \\ g.Y & \xrightarrow{\alpha'_g} & Y \end{array}$$

3 Limits and Colimits

3.1 Kernel (via equalizer/pullback)

Let $\mathcal{C}$ be a category with a zero object 0 (an object that is both initial and terminal) and suppose equalizers exist in $\mathcal{C}$. For a morphism $f : X \rightarrow Y$, the zero morphism $0_{XY} : X \rightarrow Y$ is the unique factorization of $X \rightarrow 0 \rightarrow Y$.

The **kernel** of f can be defined as the equalizer of the pair of parallel morphisms f and 0_{XY} :

$$\ker(f) \xrightarrow{k} X \rightrightarrows_{0_{XY}}^f Y.$$

That is, $k : \ker(f) \rightarrow X$ is an object and morphism such that $f \circ k = 0_{XY} \circ k (= 0_{\ker(f), Y})$ and it is universal with this property (any other $g : Z \rightarrow X$ with $f \circ g = 0_{XY} \circ g$ factors uniquely through k).

Since 0 is initial, the morphism $0 \rightarrow X$ exists uniquely. Since 0 is terminal, the morphism $X \rightarrow 0$ exists uniquely. Their composite $X \rightarrow 0 \rightarrow Y$ is the zero morphism 0_{XY} .

Alternatively, in a category $\mathcal{C}$ with a zero object 0 (which is both initial and terminal) and pullbacks, the **kernel** of a morphism $f : X \rightarrow Y$ can be defined using a pullback.

Specifically, the kernel $k : \ker(f) \rightarrow X$ is the pullback of f along the unique zero morphism $0_Y : 0 \rightarrow Y$. The pullback diagram is:

$$\begin{array}{ccc} \ker(f) & \longrightarrow & 0 \\ k \downarrow & & \downarrow 0_Y \\ X & \xrightarrow{f} & Y \end{array}$$

The map k automatically satisfies $f \circ k = 0$, and the universal property of the pullback ensures that k is universal among all maps into X whose composition with f is zero.

3.2 Cokernel (via Coequalizer)

Dually, let $\mathcal{C}$ be a category with a zero object 0 and suppose coequalizers exist. For a morphism $f : X \rightarrow Y$, consider the zero morphism $0_{XY} : X \rightarrow Y$.

The **cokernel** of f can be defined as the coequalizer of the pair f and 0_{XY} :

$$X \rightrightarrows_{0_{XY}}^f Y \xrightarrow{q} \text{coker}(f).$$

That is, $q : Y \rightarrow \text{coker}(f)$ is an object and morphism such that $q \circ f = q \circ 0_{XY} (= 0_{Y, \text{coker}(f)})$ and it is universal with this property (any other $g : Y \rightarrow Z$ with $g \circ f = g \circ 0_{XY}$ factors uniquely through q).

3.3 Left/Right Exact Functors

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor between categories where certain limits or colimits exist.

The functor F is **left exact** if it preserves all finite limits that exist in $\mathcal{C}$. This means that if L is the limit of a finite diagram $D : J \rightarrow \mathcal{C}$ (where J is a finite category), then $F(L)$ is the limit of the diagram $F \circ D : J \rightarrow \mathcal{D}$. Common examples include preserving terminal objects, products, and equalizers (and thus pullbacks and kernels if they exist).

The functor F is **right exact** if it preserves all finite colimits that exist in $\mathcal{C}$. This means that if C is the colimit of a finite diagram $D : J \rightarrow \mathcal{C}$, then $F(C)$ is the colimit of the diagram $F \circ D : J \rightarrow \mathcal{D}$. Common examples include preserving initial objects, coproducts, and coequalizers (and thus pushouts and cokernels if they exist).

A functor is **exact** if it is both left exact and right exact (preserves all finite limits and finite colimits).

In the context of abelian categories, these definitions are equivalent to the definitions involving preservation of kernels/cokernels or exact sequences, assuming the functor is also additive.

3.4 RAPL / LAPC

Right Adjoints Preserve Limits, in particular right adjoints are left exact.

Left Adjoints Preserve Colimits, in particular left adjoints are right exact.

For finite abelian categories (see later), it is actually an iff.

Perhaps a better way to remember this:

Left adjoints preserve **L**imits IS WRONG.

3.5 Linear functor + source semisimple imply exact

If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a linear (additive) functor between linear abelian categories and $\mathcal{C}$ is semisimple, then F is exact.

This follows as in $\mathcal{C}$ short exact sequences split and then by additivity so do the images under F .

In particular any linear functor $\text{vect} \rightarrow \mathcal{A}$ is exact.

3.6 Pullbacks from Products and Equalizers

The existence of pullbacks in a category $\mathcal{C}$ is closely related to the existence of binary products and equalizers.

Specifically, if a category $\mathcal{C}$ has all binary products and all equalizers, then it has all pullbacks.

To construct the pullback of a diagram $X \xrightarrow{f} Z \xleftarrow{g} Y$:

1. Form the binary product $X \times Y$ with its projection morphisms $p_X : X \times Y \rightarrow X$ and $p_Y : X \times Y \rightarrow Y$.
2. Consider the two morphisms from the product to Z : $f \circ p_X : X \times Y \rightarrow Z$ and $g \circ p_Y : X \times Y \rightarrow Z$.
3. Take the equalizer $e : P \rightarrow X \times Y$ of this pair:

$$P \xrightarrow{e} X \times Y \rightrightarrows_{g \circ p_Y}^{f \circ p_X} Z.$$

The resulting object P together with the morphisms $p_1 = p_X \circ e : P \rightarrow X$ and $p_2 = p_Y \circ e : P \rightarrow Y$ forms the pullback square:

$$\begin{array}{ccc} P & \xrightarrow{p_2} & Y \\ p_1 \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

Conversely, if a category has all pullbacks and a terminal object (which acts as the empty product), it has all finite products. If it has pullbacks and products, it also has equalizers (an equalizer of $h, k : A \rightarrow B$ is the pullback of $(h, k) : A \rightarrow B \times B$ along the diagonal $\Delta_B : B \rightarrow B \times B$).

Therefore, for a category $\mathcal{C}$ the following are equivalent:

- $\mathcal{C}$ has all finite limits,
- $\mathcal{C}$ has finite products and equalizers,
- $\mathcal{C}$ has pullbacks and a terminal object

In any of these equivalent cases we say that $\mathcal{C}$ is *finitely complete*.

Of course, as it is well known, $\mathcal{C}$ is complete (ie it has all small limits) iff $\mathcal{C}$ has all products and equalizers.

3.7 Pushouts from Coproducts and Coequalizers

Dually, the existence of pushouts is related to the existence of binary coproducts and coequalizers.

If a category $\mathcal{C}$ has all binary coproducts and all coequalizers, then it has all pushouts. To construct the pushout of a diagram $X \xleftarrow{f} Z \xrightarrow{g} Y$:

1. Form the binary coproduct $X + Y$ (often written $X \amalg Y$) with its injection morphisms $i_X : X \rightarrow X + Y$ and $i_Y : Y \rightarrow X + Y$.
2. Consider the two morphisms from Z to the coproduct: $i_X \circ f : Z \rightarrow X + Y$ and $i_Y \circ g : Z \rightarrow X + Y$.
3. Take the coequalizer $q : X + Y \rightarrow P$ of this pair:

$$Z \rightrightarrows_{i_Y \circ g}^{i_X \circ f} X + Y \xrightarrow{q} P.$$

The resulting object P together with the morphisms $q_1 = q \circ i_X : X \rightarrow P$ and $q_2 = q \circ i_Y : Y \rightarrow P$ forms the pushout square:

$$\begin{array}{ccc} Z & \xrightarrow{f} & X \\ g \downarrow & & \downarrow q_1 \\ Y & \xrightarrow{q_2} & P \end{array}$$

Conversely, if a category has all pushouts and an initial object (empty coproduct), it has all finite coproducts. If it has pushouts and coproducts, it also has coequalizers (constructed dually to equalizers using pushouts and codiagonals).

Therefore, for a category $\mathcal{C}$ the following are equivalent:

- $\mathcal{C}$ has all finite colimits,
- $\mathcal{C}$ has finite coproducts and coequalizers,
- $\mathcal{C}$ has pushouts and an initial object.

In any of these equivalent cases we say that $\mathcal{C}$ is *finitely cocomplete*.

Of course, as it is well known, $\mathcal{C}$ is cocomplete (ie it has all small colimits) iff $\mathcal{C}$ has all coproducts and coequalizers.

3.8 Ends and coends

These concepts are fundamental tools for working with functors $T : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{D}$. Ends and coends provide universal objects capturing information “along the diagonal” $T(c, c)$, defined via dinatural transformations.

Dinatural Transformations (cf. T-V B.1.1)

Let $\mathcal{C}$ be a category and $T : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

- A **dinatural transformation from an object D of $\mathcal{D}$ to T** is a family of morphisms

$$\alpha = \{\alpha_c : D \rightarrow T(c, c)\}_{c \in \text{Ob}(\mathcal{C})}$$

such that for every morphism $f : c \rightarrow d$ in $\mathcal{C}$, the following diagram commutes:

$$\begin{array}{ccc} D & \xrightarrow{\alpha_c} & T(c, c) \\ \alpha_d \downarrow & & \downarrow T(1_c, f) \\ T(d, d) & \xleftarrow{T(f, 1_d)} & T(d, c) \end{array}$$

The commutativity condition is $T(1_c, f) \circ \alpha_c = T(f, 1_d) \circ \alpha_d$. This family is denoted $\alpha : D \div T$ (or $\alpha : \Delta_D \div T$).

- A **dinatural transformation from T to an object D of $\mathcal{D}$** is a family of morphisms

$$\beta = \{\beta_c : T(c, c) \rightarrow D\}_{c \in \text{Ob}(\mathcal{C})}$$

such that for every morphism $f : c \rightarrow d$ in $\mathcal{C}$, the following diagram commutes:

$$\begin{array}{ccc} T(d, c) & \xrightarrow{T(1_d, f)} & T(d, d) \\ T(f, 1_c) \downarrow & & \downarrow \beta_d \\ T(c, c) & \xrightarrow{\beta_c} & D \end{array}$$

The commutativity condition is $\beta_d \circ T(1_d, f) = \beta_c \circ T(f, 1_c)$. This family is denoted $\beta : T \div D$ (or $\beta : T \div \Delta_D$).

(Note: The general definition of a dinatural transformation between two functors $S, T : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{D}$ requires a hexagonal diagram, as shown in T-V B.1.1. The square diagrams above are special cases for constant source/target functors.)

Coends (T-V B.1.3)

Let $T : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{D}$ be a functor. The **coend** of T , denoted $\int^c T(c, c)$, is an object of $\mathcal{D}$ equipped with a dinatural transformation

$$\omega' : T \rightrightarrows \int^c T(c, c)$$

satisfying the universal property:

For any object $x \in \mathcal{D}$ and any dinatural transformation $\beta : T \rightrightarrows x$, there exists a **unique** morphism

$$h : \int^c T(c, c) \rightarrow x$$

such that $\beta_c = h \circ \omega'_c$ for all c :

$$\begin{array}{ccc} T(c, c) & \xrightarrow{\omega'_c} & \int^c T(c, c) \\ \beta_c \downarrow & & \downarrow h \\ x & \xlongequal{\quad} & x \end{array}$$

The coend is the universal recipient for dinatural maps *from* T . If $\mathcal{D}$ has the necessary colimits, it can be constructed as a coequalizer:

$$\int^c T(c, c) = \text{coeq} \left(\coprod_{f:c \rightarrow d} T(c, d) \rightrightarrows \coprod_c T(c, c) \right)$$

where the parallel arrows are induced by the morphisms $T(1_c, f)$ and $T(f, 1_d)$ into the respective summands.

- **Example (T-V B.1.3, Ex 1):** Tensor product

$$F \otimes_{\mathcal{C}} G = \int^c (F(c) \times G(c))$$

for $F : \mathcal{C}^{op} \rightarrow \mathbf{Set}$ and $G : \mathcal{C} \rightarrow \mathbf{Set}$.

Ends (T-V B.1.2)

Let $T : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{D}$ be a functor. The **end** of T , denoted $\int_c T(c, c)$, is an object of $\mathcal{D}$ equipped with a dinatural transformation

$$\omega : \int_c T(c, c) \rightrightarrows T$$

satisfying the universal property:

For any object $x \in \mathcal{D}$ and any dinatural transformation $\alpha : x \rightrightarrows T$, there exists a **unique** morphism

$$h : x \rightarrow \int_c T(c, c)$$

such that $\alpha_c = \omega_c \circ h$ for all c :

$$\begin{array}{ccc} x & \xrightarrow{h} & \int_c T(c, c) \\ \parallel & & \downarrow \omega_c \\ x & \xrightarrow{\alpha_c} & T(c, c) \end{array}$$

The end is the universal source for dinatural maps to T . If $\mathcal{D}$ has the necessary limits, it can be constructed as an equalizer:

$$\int_c T(c, c) = \text{eq} \left(\prod_c T(c, c) \rightrightarrows \prod_{f:c \rightarrow d} T(d, c) \right)$$

where the parallel arrows arise from the dinaturality condition applied to the projections $\pi_c : \prod_b T(b, b) \rightarrow T(c, c)$. The first map's f -component is $T(f, 1_c) \circ \pi_c$, and the second map's f -component is $T(1_d, f) \circ \pi_d$.

- **Example (T-V B.1.2, Ex 1):** Natural transformations

$$\text{Nat}(F, G) = \int_c \text{Hom}_{\mathcal{D}}(Fc, Gc).$$

Properties (cf. Turaev–Virelizier Appendix B)

- **Hom-Set Isomorphisms (T-V B.1.4):** Mapping into an end or out of a coend corresponds to taking the end of the relevant Hom-functor in **Set**:

$$\text{Hom}_{\mathcal{D}}(x, \int_c T(c, c)) \cong \int_c \text{Hom}_{\mathcal{D}}(x, T(c, c))$$

$$\text{Hom}_{\mathcal{D}}(\int_c T(c, c), x) \cong \int_c \text{Hom}_{\mathcal{D}}(T(c, c), x).$$

- **Fubini Theorem (T-V B.1.5):** Under suitable conditions, the order of integration can be swapped:

$$\begin{aligned} \int_{c \in \mathcal{C}} \int_{d \in \mathcal{D}} T(c, d, c, d) &\cong \int_{d \in \mathcal{D}} \int_{c \in \mathcal{C}} T(c, d, c, d), \\ \int_{c \in \mathcal{C}} \int_{d \in \mathcal{D}} T(c, d, c, d) &\cong \int_{d \in \mathcal{D}} \int_{c \in \mathcal{C}} T(c, d, c, d). \end{aligned}$$

- **Other properties** include preservation by certain functors and relations under adjunctions (T-V B.1.6).

3.9 Homotopy fibre of a functor

Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a functor, and let $b \in \mathcal{B}$. The homotopy fibre $\text{hofib}_b(F)$ of F at b is defined as follows: an **object** of the homotopy fiber $\text{hofib}_b(F)$ is a pair (a, ϕ) where:

- a is an object in $\mathcal{A}$.
- ϕ is a morphism in $\mathcal{B}$ of the form $\phi : F(a) \rightarrow b$.

The morphism ϕ serves as a "bridge" or "path" demonstrating how the image of a is related to the target object b .

A **morphism** in $\text{hofib}_b(F)$ from an object (a, ϕ) to an object (a', ϕ') is a morphism $h : a \rightarrow a'$ in the category $\mathcal{A}$ such that the following diagram commutes in $\mathcal{B}$:

$$\begin{array}{ccc} F(a) & \xrightarrow{F(h)} & F(a') \\ & \searrow \phi & \swarrow \phi' \\ & b & \end{array}$$

The commutativity condition is explicitly: $\phi' \circ F(h) = \phi$.

Forgetting ϕ we get a functor $\text{hofib}_b(F) \rightarrow \mathcal{A}$.

3.10 Homotopy pullback

Consider a diagram of categories and functors:

$$\mathcal{A} \xrightarrow{F} \mathcal{B} \xleftarrow{G} \mathcal{C}$$

The **homotopy pullback**, denoted $\mathcal{A} \times_{\mathcal{B}}^h \mathcal{C}$, is a new category constructed as follows.

An **object** of the homotopy pullback category $\mathcal{A} \times_{\mathcal{B}}^h \mathcal{C}$ is a triplet (a, c, ϕ) where:

- a is an object in $\mathcal{A}$.
- c is an object in $\mathcal{C}$.
- ϕ is an isomorphism in $\mathcal{B}$ of the form $\phi : F(a) \rightarrow G(c)$.

A **morphism** in $\mathcal{A} \times_{\mathcal{B}}^h \mathcal{C}$ from an object (a, c, ϕ) to an object (a', c', ϕ') is a pair of morphisms (h_A, h_C) where:

- $h_A : a \rightarrow a'$ is a morphism in $\mathcal{A}$.
- $h_C : c \rightarrow c'$ is a morphism in $\mathcal{C}$.

This pair must satisfy the condition that the following diagram commutes in the category $\mathcal{B}$:

$$\begin{array}{ccc} F(a) & \xrightarrow{F(h_A)} & F(a') \\ \phi \downarrow & & \downarrow \phi' \\ G(c) & \xrightarrow{G(h_C)} & G(c') \end{array}$$

The commutativity condition is explicitly: $\phi' \circ F(h_A) = G(h_C) \circ \phi$.

Note we get a square

$$\begin{array}{ccc} \mathcal{A} \times_{\mathcal{B}}^h \mathcal{C} & \longrightarrow & \mathcal{C} \\ \downarrow & & \downarrow \\ \mathcal{A} & \longrightarrow & \mathcal{B} \end{array}$$

which is commutative up to natural isomorphism, ie the two resulting functors $\mathcal{A} \times_{\mathcal{B}}^h \mathcal{C} \rightarrow \mathcal{B}$ are naturally isomorphic.

The homotopy fibre $\text{hofib}_b(F)$ of a functor $F : \mathcal{A} \rightarrow \mathcal{B}$ is nothing but the homotopy pullback of the cospan $\mathcal{A} \xrightarrow{F} \mathcal{B} \xleftarrow{b} *$.

3.11 Yoneda lemma

Setup. Let $\mathcal{C}$ be a locally small category, $A \in \mathcal{C}$, and $F : \mathcal{C} \rightarrow \mathbf{Set}$ a functor. Write $\mathcal{C}(A, -)$ for the covariant representable functor $\text{Hom}_{\mathcal{C}}(A, -)$.

Classical statement. There is a bijection, natural in both A and F ,

$$\text{Nat}(\mathcal{C}(A, -), F) \cong FA, \tag{1}$$

given by $\eta \mapsto \eta_A(\text{id}_A)$, with inverse sending $x \in FA$ to the natural transformation with components $\eta_X(f) = (Ff)(x)$ for $f \in \mathcal{C}(A, X)$.

Natural transformations as an end

Recall that for functors $F, G : \mathcal{C} \rightarrow \mathcal{D}$, the set of natural transformations can be computed as an *end*:

$$\text{Nat}(F, G) \cong \int_{X \in \mathcal{C}} \mathcal{D}(FX, GX). \quad (2)$$

Here the end is the limit of the diagram sending $X \mapsto \mathcal{D}(FX, GX)$ and $f : X \rightarrow Y$ to the two maps

$$\mathcal{D}(FX, GX) \xrightarrow{(Gf)_*} \mathcal{D}(FX, GY) \xleftarrow{(Ff)^*} \mathcal{D}(FY, GY), \quad (3)$$

i.e. a family $(\varphi_X)_X$ lies in the end iff $Gf \circ \varphi_X = \varphi_Y \circ Ff$ for all f — exactly naturality.

The “ninja” Yoneda lemma

Applying this to $\mathcal{C}(A, -)$ and F gives the end-form of Yoneda:

$$FA \cong \int_{X \in \mathcal{C}} \mathbf{Set}(\mathcal{C}(A, X), FX). \quad (4)$$

Proof. By definition,

$$\int_X \mathbf{Set}(\mathcal{C}(A, X), FX) = \text{Nat}(\mathcal{C}(A, -), F), \quad (5)$$

so it suffices to reprove the classical statement, but it is instructive to see how the *wedge condition itself* forces everything. A wedge is a family

$$\varphi_X : \mathcal{C}(A, X) \longrightarrow FX, \quad X \in \mathcal{C}, \quad (6)$$

such that for every $f : X \rightarrow Y$ the square

$$\begin{array}{ccc} \mathcal{C}(A, X) & \xrightarrow{\varphi_X} & FX \\ f \circ - \downarrow & & \downarrow Ff \\ \mathcal{C}(A, Y) & \xrightarrow{\varphi_Y} & FY \end{array}$$

commutes. Take $X = A$, $g \in \mathcal{C}(A, X)$ arbitrary, and apply the square to $f = g : A \rightarrow X$ evaluated at id_A :

$$\varphi_X(g) = \varphi_X(g \circ \text{id}_A) = (Fg)(\varphi_A(\text{id}_A)). \quad (7)$$

So every component φ_X is completely determined by the single value $x := \varphi_A(\text{id}_A) \in FA$, via

$$\varphi_X(g) = (Fg)(x), \quad g \in \mathcal{C}(A, X). \quad (8)$$

Conversely, given any $x \in FA$, this formula defines a wedge (functoriality of F gives exactly the required commuting squares). This exhibits a bijection

$$\int_X \mathbf{Set}(\mathcal{C}(A, X), FX) \xrightarrow{\cong} FA, \quad \varphi \longmapsto \varphi_A(\text{id}_A), \quad (9)$$

with the evaluation map $x \mapsto (g \mapsto (Fg)(x))$ as inverse. ■ □

Remarks

- This proof is the “universal element” argument in disguise: the wedge condition, applied at $X = A$ with $g = \text{id}_A$, is nothing but the statement that $\mathcal{C}(A, -)$ is the **free/representing** functor on the “generic element” $\text{id}_A \in \mathcal{C}(A, A)$, and $x = \varphi_A(\text{id}_A)$ is where that generic element must go.
- The contravariant version, for $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$, reads

$$FA \cong \int_{X \in \mathcal{C}} \mathbf{Set}(\mathcal{C}(X, A), FX). \quad (10)$$

- The end formula is the one that generalizes cleanly to **enriched** category theory: for a $\mathcal{V}$ -category $\mathcal{C}$ and $F : \mathcal{C} \rightarrow \mathcal{V}$,

$$FA \cong \int_{X \in \mathcal{C}} [\mathcal{C}(A, X), FX], \quad (11)$$

where $[-, -]$ is the internal hom of $\mathcal{V}$ — this is the version used e.g. for Vect-enriched or Cat-enriched Yoneda, and it is the natural language for the **coend/density formula**

$$F \cong \int^{X \in \mathcal{C}} \mathcal{C}(X, -) \cdot FX \quad (12)$$

(every functor is a colimit of representables), which is Yoneda’s *coend* dual and the starting point for Kan extension formulas via the **Fubini/ninja Yoneda** manipulations (co-Yoneda reduction).

3.12 Bimodules and the hom

Setup. Let k be a field and A, B finite-dimensional k -algebras. Let M, M' be left A -modules and N, N' right B -modules, all finite-dimensional over k . As before, $M \otimes_k N$ and $M' \otimes_k N'$ carry natural A - B -bimodule structures

$$a \cdot (m \otimes n) \cdot b = (a \cdot m) \otimes (n \cdot b). \quad (13)$$

For an A - B -bimodule P , let $\text{Hom}_B(N, P)$ denote the space of right B -module maps $N \rightarrow P$ (forgetting the A -action on P). This carries a left A -module structure via

$$(a \cdot \varphi)(n) := a \cdot \varphi(n), \quad \varphi \in \text{Hom}_B(N, P), \quad a \in A, \quad n \in N, \quad (14)$$

which is well-defined precisely because the A - and B -actions on P commute:

$$(a \cdot \varphi)(nb) = a \cdot \varphi(nb) = a \cdot (\varphi(n) \cdot b) = (a \cdot \varphi(n)) \cdot b = (a \cdot \varphi)(n) \cdot b, \quad (15)$$

so $a \cdot \varphi$ is again B -linear.

Statement. There is an isomorphism, natural in M, M', N, N' ,

$$\boxed{\text{Hom}_{A\text{-mod-}B}(M \otimes_k N, M' \otimes_k N') \cong \text{Hom}_A(M, \text{Hom}_B(N, M' \otimes_k N'))}. \quad (16)$$

Proof

Proof. Start from the classical tensor–hom adjunction over k ,

$$\Xi : \text{Hom}_k(M \otimes_k N, M' \otimes_k N') \xrightarrow{\cong} \text{Hom}_k(M, \text{Hom}_k(N, M' \otimes_k N')), \quad \Xi(\theta)(m)(n) = \theta(m \otimes n). \quad (17)$$

It suffices to show Ξ restricts to a bijection between the two subspaces of bimodule/module-compatible maps.

θ is a morphism of A - B -bimodules if and only if, for all $a \in A, b \in B, m \in M, n \in N$,

$$\theta((a \cdot m) \otimes (n \cdot b)) = a \cdot \theta(m \otimes n) \cdot b. \quad (18)$$

Writing $\hat{\theta} := \Xi(\theta)$, i.e. $\hat{\theta}(m)(n) = \theta(m \otimes n)$, this condition reads

$$\hat{\theta}(a \cdot m)(n \cdot b) = a \cdot \hat{\theta}(m)(n) \cdot b \quad \text{for all } a, b, m, n. \quad (*)$$

Setting $a = 1$ in $(*)$: $\hat{\theta}(m)(n \cdot b) = \hat{\theta}(m)(n) \cdot b$ for all m, n, b , i.e. $\hat{\theta}(m) \in \text{Hom}_B(N, M' \otimes_k N')$ for every $m \in M$.

Setting $b = 1$ in $(*)$, given the above: $\hat{\theta}(a \cdot m)(n) = a \cdot \hat{\theta}(m)(n)$ for all m, n , i.e. $\hat{\theta}(a \cdot m) = a \cdot \hat{\theta}(m)$ in $\text{Hom}_B(N, M' \otimes_k N')$, using the A -module structure defined above. This says exactly that $\hat{\theta} : M \rightarrow \text{Hom}_B(N, M' \otimes_k N')$ is A -linear.

Conversely, if $\hat{\theta}(m)$ is B -linear for every m and $\hat{\theta}$ itself is A -linear into $\text{Hom}_B(N, M' \otimes_k N')$, then combining the two displayed identities recovers $(*)$ for all a, b .

Hence θ is a bimodule map if and only if $\hat{\theta}$ lies in $\text{Hom}_A(M, \text{Hom}_B(N, M' \otimes_k N'))$, so Ξ restricts to the desired isomorphism

$$\text{Hom}_{A\text{-mod-}B}(M \otimes_k N, M' \otimes_k N') \xrightarrow{\cong} \text{Hom}_A(M, \text{Hom}_B(N, M' \otimes_k N')), \quad \theta \longmapsto \hat{\theta}. \quad \blacksquare$$

□

Remarks

- This is a genuine adjunction: the functor $(-) \otimes_k N : A\text{-mod} \rightarrow A\text{-mod-}B$ is left adjoint to $\text{Hom}_B(N, -) : A\text{-mod-}B \rightarrow A\text{-mod}$, with unit and counit induced from the classical tensor–hom adjunction, restricted along the fully-faithful inclusion of bimodule maps described above. No finite-dimensionality is needed for this statement (unlike the pure tensor-factorization formula $\text{Hom}_A(M, M') \otimes_k \text{Hom}_B(N, N')$ from before, which does use it).
- If N is finite-dimensional, evaluation/coevaluation gives $\text{Hom}_B(N, P) \cong (N^\vee \otimes_k P)^B$ (right- B -invariants for the diagonal action, using the induced left B -action on $N^\vee$), so the formula can be rewritten as

$$\text{Hom}_{A\text{-mod-}B}(M \otimes_k N, M' \otimes_k N') \cong \text{Hom}_A(M, (N^\vee \otimes_k M' \otimes_k N')^B), \quad (19)$$

the same evaluation-morphism circle of ideas that appears in the canonical coend

$$\mathbb{F} = \int^X X^\vee \otimes X \quad (20)$$

for a finite tensor category: the coend $\mathbb{F}$ is built from exactly this kind of evaluation pairing between an object and its dual, and the adjunction above is its bimodule-level shadow.

- If A, B are algebras in a finite tensor category $\mathcal{A}$, then the isomorphism reads

$$\boxed{\text{Hom}_{A\text{-mod-}B}(M \otimes N, M' \otimes N') \cong \text{Hom}_A(M, \underline{\text{Hom}}_B(N, M' \otimes N'))}. \quad (21)$$

3.13 coYoneda lemma

Setup. Let $\mathcal{C}$ be a locally small category and $F : \mathcal{C} \rightarrow \mathbf{Set}$ a functor. Write $\mathcal{C}(X, -)$ for the covariant representable functor at $X \in \mathcal{C}$. The co-Yoneda lemma expresses F itself as a colimit of representables — it is the formal statement that “every functor is built out of its representable pieces.”

Statement. There is an isomorphism, natural in F ,

$$F \cong \int^{X \in \mathcal{C}} \mathcal{C}(X, -) \cdot FX, \quad (22)$$

where $\int^X$ denotes the *coend* and $\mathcal{C}(X, -) \cdot FX$ is the copower (a coproduct of $|FX|$ copies of the functor $\mathcal{C}(X, -)$):

$$\mathcal{C}(X, -) \cdot FX = \coprod_{x \in FX} \mathcal{C}(X, -). \quad (23)$$

If $F : \mathcal{C}^{op} \rightarrow \mathbf{Set}$ then the formula reads

$$F \cong \int^{X \in \mathcal{C}} \mathcal{C}(-, X) \cdot FX, \quad (24)$$

Evaluated at an object $A \in \mathcal{C}$, this reads

$$FA \cong \int^{X \in \mathcal{C}} \mathcal{C}(X, A) \cdot FX. \quad (25)$$

If the categories are $\mathbb{k}$ -linear, then this reads

$$FA \cong \int^{X \in \mathcal{C}} \mathcal{C}(X, A) \otimes FX \quad (26)$$

or in the contravariant case

$$FA \cong \int^{X \in \mathcal{C}} \mathcal{C}(A, X) \otimes FX. \quad (27)$$

And in particular for $F = \mathcal{C}(-, B)^*$ we get

$$\mathcal{C}(A, B)^* \cong \int^{X \in \mathcal{C}} \mathcal{C}(X, A) \otimes \mathcal{C}(X, B)^*. \quad (28)$$

or similarly taking the contravariant version

$$\boxed{\mathcal{C}(A, B)^* \cong \int^{X \in \mathcal{C}} \mathcal{C}(B, X) \otimes \mathcal{C}(A, X)^*}. \quad (29)$$

Coends as colimits Dually to the end description of naturality, the coend $\int^X \mathcal{C}(X, A) \times FX$ is the *coequalizer*

$$\coprod_{f: X \rightarrow Y} \mathcal{C}(Y, A) \times FX \xrightarrow[\text{id} \times Ff]{f^* \times \text{id}} \coprod_{X \in \mathcal{C}} \mathcal{C}(X, A) \times FX \longrightarrow \int^X \mathcal{C}(X, A) \times FX$$

That is, elements are pairs (g, x) with $g : X \rightarrow A$, $x \in FX$, modulo the relation

$$(g \circ f, x) \sim (g, (Ff)(x)), \quad \text{for } f : X \rightarrow Y, g : Y \rightarrow A, x \in FX. \quad (30)$$

We write $[g, x]$ for the equivalence class of (g, x) .

Proof

Proof. Define a map

$$\Phi : \int^X \mathcal{C}(X, A) \times FX \longrightarrow FA, \quad \Phi[g, x] = (Fg)(x). \quad (31)$$

This is well-defined on equivalence classes: for $f : X \rightarrow Y$,

$$\Phi[g \circ f, x] = (F(g \circ f))(x) = (Fg)((Ff)(x)) = \Phi[g, (Ff)(x)], \quad (32)$$

using functoriality of F , which is exactly the coequalizer relation.

Conversely, define

$$\Psi : FA \longrightarrow \int^X \mathcal{C}(X, A) \times FX, \quad \Psi(a) = [\text{id}_A, a]. \quad (33)$$

$$\Phi \circ \Psi = \text{id}: \Phi\Psi(a) = \Phi[\text{id}_A, a] = (F\text{id}_A)(a) = a.$$

$$\Psi \circ \Phi = \text{id}: \text{for any representative } (g, x) \text{ with } g : X \rightarrow A,$$

$$\Psi\Phi[g, x] = \Psi((Fg)(x)) = [\text{id}_A, (Fg)(x)] = [\text{id}_A \circ g, x] = [g, x], \quad (34)$$

using the coequalizer relation with $f = g$. Hence Φ and Ψ are mutually inverse bijections, natural in A and F . ■ □

Remarks

- Every element of $\int^X \mathcal{C}(X, A) \times FX$ can be represented (non-uniquely) as $[g, x]$ with $g : X \rightarrow A$, $x \in FX$; the coend relation says exactly that “pushing f along g ” and “pushing f along F ” give the same class. The isomorphism above shows the *canonical* representative of $a \in FA$ is $[\text{id}_A, a]$.
- This is the coend/co-Yoneda dual of the end form of Yoneda,

$$FA \cong \int_{X \in \mathcal{C}} \mathbf{Set}(\mathcal{C}(A, X), FX), \quad (35)$$

and the two are related by the general **Yoneda–density** philosophy: F is simultaneously a limit of the diagram of hom-sets *into* it (end form) and a colimit of representables glued along F ’s values (coend form).

- The enriched version, for $F : \mathcal{C} \rightarrow \mathcal{V}$ with $\mathcal{C}$ a $\mathcal{V}$ -category and $\mathcal{V}$ cocomplete closed symmetric monoidal, reads

$$F \cong \int^{X \in \mathcal{C}} \mathcal{C}(X, -) \otimes FX, \quad (36)$$

with $\otimes$ the tensoring of $\mathcal{V}$ -objects over $\mathcal{V}$ -functors. This is the formula underlying the **density theorem** (every presheaf is a colimit of representables) and is the standard tool for **Fubini/ninja Yoneda** manipulations, e.g. rewriting coends of Hom functors such as

$$\int^{X \in \mathcal{C}} \mathcal{C}(X, A) \otimes G(X) \cong G(A) \quad (37)$$

for any $G : \mathcal{C} \rightarrow \mathcal{V}$ — the “co-Yoneda reduction” used constantly when simplifying coend expressions (e.g. in the canonical coend algebra $F = \int^X X^\vee \otimes X$ for a finite tensor category).

4 Monoidal Categories and Related Structures

4.1 Monoidal category

A tuple $(\mathcal{C}, \otimes, \mathbf{1}, a, l, r)$ of a category $\mathcal{C}$ with a bifunctor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, an object $\mathbf{1}$ and natural isomorphisms

$$a_{X,Y,Z} : (X \otimes Y) \otimes Z \rightarrow X \otimes (Y \otimes Z), \quad l_X : \mathbf{1} \otimes X \rightarrow X, \quad r_X : X \otimes \mathbf{1} \rightarrow X$$

satisfying the so-called pentagon equation and the triangle equation.

4.2 Left rigid category

A monoidal category in which every object admits a left dual, that is, for every object X there exists a pair $({}^\vee X, \text{ev}_X)$ where ${}^\vee X$ is another object and a left evaluation

$$\text{ev}_X : {}^\vee X \otimes X \rightarrow \mathbf{1}$$

is a non-degenerate pairing. This means that there exists another map

$$\text{coev}_X : \mathbf{1} \rightarrow X \otimes {}^\vee X$$

called the left coevaluation satisfying

$$(\text{Id} \otimes \text{ev}_X)(\text{coev}_X \otimes \text{Id}) = \text{Id}, \quad (\text{ev}_X \otimes \text{Id})(\text{Id} \otimes \text{coev}_X) = \text{Id}.$$

Left duals are unique up to unique isomorphism.

4.3 Right rigid category

A monoidal category in which every object admits a right dual, that is, for every object X there exists a pair $(X^\vee, \widetilde{\text{ev}}_X)$ where $X^\vee$ is another object and a right evaluation

$$\widetilde{\text{ev}}_X : X \otimes X^\vee \rightarrow \mathbf{1}$$

is a non-degenerate pairing. This means that there exists another map

$$\widetilde{\text{coev}}_X : \mathbf{1} \rightarrow X^\vee \otimes X$$

called the right coevaluation satisfying

$$(\text{Id} \otimes \widetilde{\text{ev}}_X)(\widetilde{\text{coev}}_X \otimes \text{Id}) = \text{Id}, \quad (\widetilde{\text{ev}}_X \otimes \text{Id})(\text{Id} \otimes \widetilde{\text{coev}}_X) = \text{Id}.$$

Right duals are unique up to unique isomorphism.

4.4 Rigid or autonomous category

A monoidal category which is both left and right rigid.

Note that there are natural isomorphisms

$$({}^\vee(-))^\vee \xrightarrow{\cong} \text{Id} \xleftarrow{\cong} {}^\vee((-)^\vee)$$

4.5 Pivotal or sovereign category

A pivotal structure on a rigid category is the data of a monoidal natural isomorphism $\text{Id} \xrightarrow{\cong} (-)^{\vee\vee}$. Alternatively, this is the same as a monoidal natural isomorphism ${}^\vee(-) \xrightarrow{\cong} (-)^\vee$.

If we start with a right rigid category, then a pivotal structure is also a monoidal natural isomorphism $\text{Id} \xrightarrow{\cong} (-)^{\vee\vee}$, and it defines a left rigid structure with ${}^\vee X := X^\vee$ (in that case we put X^*).

By the uniqueness of duals, both approaches are equivalent. The key thing is that "rigid category" means a category that admits right and left rigidity; this is different from a category with distinguished duality, where the exact duality is fixed.

4.6 Spherical category (classical definition)

If $\mathcal{C}$ is a pivotal category, we have the notions of left and right traces of an endomorphism $f : X \rightarrow X$, defined as follows:

$$\text{tr}_l(f) := \text{ev}_X(\text{Id}_{X^*} \otimes f) \widetilde{\text{coev}}_X : \mathbf{1} \rightarrow \mathbf{1}, \quad \text{tr}_r(f) := \widetilde{\text{ev}}_X(f \otimes \text{Id}_{X^*}) \text{coev}_X : \mathbf{1} \rightarrow \mathbf{1}.$$

We also have the notions of left and right dimensions of an object X :

$$\dim_l(X) := \text{ev}_X \widetilde{\text{coev}}_X, \quad \dim_r(X) := \widetilde{\text{ev}}_X \text{coev}_X.$$

A spherical category is a pivotal category whose left and right traces coincide (and therefore also the left and right dimensions).

4.7 Braided category

A monoidal category endowed with a natural isomorphism

$$\tau_{X,Y} : X \otimes Y \rightarrow Y \otimes X$$

which is $\otimes$ -multiplicative in the sense that

$$\tau_{X,Y \otimes Z} = (\text{id}_Y \otimes \tau_{X,Z})(\tau_{X,Y} \otimes \text{id}_Z), \quad \tau_{X \otimes Y,Z} = (\tau_{X,Z} \otimes \text{id}_Y)(\text{id}_X \otimes \tau_{Y,Z})$$

The case of a braided rigid category If $\mathcal{C}$ is a braided rigid category, then for every object $X \in \mathcal{C}$ there is a natural isomorphism $X \xrightarrow{\cong} X^{\vee\vee}$, which is the "twist" in the graphical language – it is an isomorphism by the Whitney trick in the graphical language. This is the composite

$$X \xrightarrow{\text{coev}_X \vee \otimes \text{id}} X \otimes X^{\vee\vee} \otimes X^\vee \xrightarrow{\text{id} \otimes \tau_{X^{\vee\vee}, X^\vee}} X \otimes X^\vee \otimes X^{\vee\vee} \xrightarrow{\widetilde{\text{ev}}_X \otimes \text{id}} X^{\vee\vee} \quad (38)$$

This is called the **Drinfeld isomorphism**. See also Shibata-Shimizu §6.4.

This isomorphism is NOT canonical; in general there are infinitely many isomorphisms $X \cong X^{\vee\vee}$. Furthermore, this natural isomorphism is NOT a monoidal natural transformation and therefore does NOT define a pivotal structure.

4.8 Ribbon or tortile category (take 1)

Let $\mathcal{C}$ be a braided pivotal category. The left twist is defined as the following family of morphisms (in fact natural isomorphisms):

$$\theta_X^l := (\text{ev}_X \otimes \text{id}_X)(\text{id}_{X^*} \otimes \tau_{X,X})(\widetilde{\text{coev}}_X \otimes \text{id}_X) : X \rightarrow X$$

whereas the right twist is

$$\theta_X^r := (\text{id}_X \otimes \widetilde{\text{ev}}_X)(\tau_{X,X} \otimes \text{id}_{X^*})(\text{id}_X \otimes \text{coev}_X) : X \rightarrow X.$$

A ribbon category is a braided pivotal category in which the left and right twists coincide, $\theta = \theta^l = \theta^r$. It automatically satisfies that the twist is self-dual in the sense that $(\theta_X)^* = \theta_{X^*}$.

4.9 Balanced category

A braided category together with a *twist*, that is, a natural isomorphism $\theta : X \rightarrow X$, called the twist, satisfying

$$\theta_{X \otimes Y} = \tau_{Y,X} \tau_{X,Y} (\theta_X \otimes \theta_Y).$$

From this condition it automatically follows that $\theta_1 = \text{id}_1$, see Kassel Lemma XIV.3.3.

4.10 Braided pivotal = balanced rigid

(Deligne, see [43, Prop. 2.11]), also (Selinger, Lemma 4.20)

Let $\mathcal{C}$ be a braided rigid category. Then giving a twist θ on $\mathcal{C}$ (making $\mathcal{C}$ into a balanced rigid category) is equivalent to giving a pivotal structure $\text{Id} \xrightarrow{\cong} (-)^{\vee\vee}$ (making $\mathcal{C}$ into a braided pivotal category).

The correspondence goes as follows: given a twist θ , a pivotal structure is given by the composite

$$X \xrightarrow{\theta_X} X \longrightarrow X^{\vee\vee}$$

where the second arrow is (38). Conversely, given a pivotal structure $\omega : \text{Id} \xrightarrow{\cong} (-)^{\vee\vee}$, a twist is obtained as the composite

$$X \xrightarrow{\omega_X} X^{\vee\vee} \longrightarrow X$$

where the second arrow is the inverse of (38).

4.11 Ribbon or tortile category (take 2)

A balanced left rigid category in which the twist and the rigid structure are compatible in the sense that

$${}^\vee(\theta_X) = \theta_{\vee X}.$$

4.12 Monoidal $\mathbb{k}$ -linear category

A monoidal category which is also $\mathbb{k}$ -linear and whose monoidal product is $\mathbb{k}$ -bilinear.

4.13 Pre-fusion category (non-abelian version)

A pre-fusion $\mathbb{k}$ -category is a monoidal $\mathbb{k}$ -linear category $\mathcal{C}$ with the property that there exists a collection of simple objects $(V_i)_{i \in I}$ indexed by some set I satisfying that:

- (a) the monoidal unit $\mathbf{1}$ belongs to the family (and it is typically indexed as $V_0 = \mathbf{1}$);
- (b) $\text{Hom}_{\mathcal{C}}(V_i, V_j) = 0$ if $i \neq j$;
- (c) every object of $\mathcal{C}$ is a direct sum of finitely many objects from the indexed collection $(V_i)_{i \in I}$.

It readily follows that every simple object in $\mathcal{C}$ is isomorphic to exactly one object from $(V_i)_{i \in I}$. Also, (b) and (c) imply that the hom-sets are finite-free $\mathbb{k}$ -modules and that for any pair of objects X, Y :

$$\text{Hom}_{\mathcal{C}}(X, Y) \cong \bigoplus_{i \in I} \text{Hom}_{\mathcal{C}}(X, V_i) \otimes_{\mathbb{k}} \text{Hom}_{\mathcal{C}}(V_i, Y).$$

4.14 Fusion category (non-abelian version)

A pre-fusion rigid linear category in which the index set I is finite.

The dimension of a pivotal fusion linear category is given by:

$$\dim(\mathcal{C}) := \sum_{i \in I} \dim_l(V_i) \dim_r(V_i),$$

and if the category happens to be spherical then this formula becomes:

$$\dim(\mathcal{C}) = \sum_{i \in I} (\dim_l(V_i))^2.$$

4.15 Modular category (non-abelian, fusion version)

A ribbon fusion $\mathbb{k}$ -linear category with the property that, if:

$$s_{i,j} := \text{tr}(\tau_{V_j, V_i} \tau_{V_i, V_j}) \in \text{End}_{\mathcal{C}}(\mathbf{1}) = \mathbb{k}$$

(where τ denotes the braiding), then the S-matrix $S = (s_{ij})_{i,j \in I}$ is invertible over $\mathbb{k}$.

4.16 Drinfeld center

Let $(\mathcal{C}, \otimes, \mathbf{1})$ be a monoidal category. The **Drinfeld center** $\mathcal{Z}(\mathcal{C})$ is defined as follows:

Objects: Pairs (Z, β_Z) , where $Z \in \mathcal{C}$ and $\beta_Z = \{\beta_{Z,X} : Z \otimes X \xrightarrow{\cong} X \otimes Z\}_{X \in \text{Ob}(\mathcal{C})}$ is a family of natural isomorphisms (called the *half-braiding*) satisfying:

- Naturality: for any $f : X \rightarrow Y$, the diagram

$$\begin{array}{ccc} Z \otimes X & \xrightarrow{\beta_{Z,X}} & X \otimes Z \\ \text{id}_Z \otimes f \downarrow & & \downarrow f \otimes \text{id}_Z \\ Z \otimes Y & \xrightarrow{\beta_{Z,Y}} & Y \otimes Z \end{array}$$

commutes.

- Coherence: $\beta_{Z, X \otimes Y} = (\text{id}_X \otimes \beta_{Z,Y}) \circ (\beta_{Z,X} \otimes \text{id}_Y)$;
- Unit compatibility: $\beta_{Z, \mathbf{1}} = \text{id}_Z$.

Morphisms: A morphism $f : (Z, \beta_Z) \rightarrow (W, \beta_W)$ is a morphism $f : Z \rightarrow W$ in $\mathcal{C}$ such that:

$$(\text{id}_X \otimes f) \circ \beta_{Z,X} = \beta_{W,X} \circ (f \otimes \text{id}_X)$$

for all $X \in \mathcal{C}$, i.e.,

$$\begin{array}{ccc} Z \otimes X & \xrightarrow{\beta_{Z,X}} & X \otimes Z \\ f \otimes \text{id}_X \downarrow & & \downarrow \text{id}_X \otimes f \\ W \otimes X & \xrightarrow{\beta_{W,X}} & X \otimes W \end{array}$$

Monoidal structure: $\mathcal{Z}(\mathcal{C})$ is a braided monoidal category with:

- Tensor product: $(Z, \beta_Z) \otimes (W, \beta_W) := (Z \otimes W, \beta_{Z \otimes W})$, where:

$$\beta_{Z \otimes W, X} = (\beta_{Z,X} \otimes \text{id}_W) \circ (\text{id}_Z \otimes \beta_{W,X})$$

- Unit: $(1, \beta_1)$, where $\beta_{1,X} = \text{id}_X$;
- Braiding: $c_{(Z, \beta_Z), (W, \beta_W)} = \beta_{Z,W}$.

4.17 Müger center

Let $(\mathcal{C}, \otimes, 1, c)$ be a braided monoidal category. The **Müger center** $\mathcal{Z}_2(\mathcal{C})$ is the full subcategory of $\mathcal{C}$ whose objects Z satisfy:

$$c_{X,Z} \circ c_{Z,X} = \text{id}_{Z \otimes X} \quad \text{for all } X \in \mathcal{C}.$$

$\mathcal{Z}_2(\mathcal{C})$ inherits the monoidal structure and is a symmetric monoidal category.

4.18 Equivalent Characterizations of Modular Categories

Theorem: Let $\mathcal{C}$ be a braided fusion category over an algebraically closed field $\mathbb{k}$. Let $\{X_i\}_{i \in I}$ be a complete set of representatives of isomorphism classes of simple objects with $X_0 = \mathbf{1}$. Let $c_{X,Y}$ denote the braiding and $\bar{\mathcal{C}}$ the same category with inverse braiding.

The following are equivalent:

- (i) **Trivial Müger center:** $\mathcal{Z}_2(\mathcal{C}) \cong \text{Vec}_{\mathbb{k}}$;
- (ii) **Invertible S-matrix:** The matrix $S = (S_{ij})$, with entries

$$S_{ij} = \text{Tr}(c_{X_j, X_i} \circ c_{X_i, X_j})$$

is invertible over $\mathbb{k}$;

- (iii) **Drinfeld center equivalence:** $\mathcal{Z}(\mathcal{C}) \cong \mathcal{C} \boxtimes \bar{\mathcal{C}}$.

Remarks:

- The pair (S, T) , with $T_{ii} = \theta_i$, is called the *modular data*.
- These equivalences are due to Müger.
- Condition (i) expresses maximal non-degeneracy.
- Condition (iii) shows that the center is determined by the braiding and its inverse.

4.19 Grothendieck-Verdier category

(Following Müller-Woike)

A *Grothendieck-Verdier category* is a monoidal category $\mathcal{C}$ together with an object $K \in \mathcal{C}$ such that $\mathcal{C}(- \otimes Y, K)$ is representable for every $Y \in \mathcal{C}$ and such that the functor $D : \mathcal{C}^{op} \rightarrow \mathcal{C}$ sending Y to a representing object DY for $\mathcal{C}(- \otimes Y, K)$ is an equivalence. The object K is referred to as *dualizing object*. The functor D is referred to as *duality functor*.

In more detail, the functor D is defined by *choosing* an object $DY \in \mathcal{C}$ and a natural isomorphism $\mathcal{C}(- \otimes Y, K) \simeq \mathcal{C}(-, DY)$ for every $Y \in \mathcal{C}$. The assignment $Y \mapsto DY$ extends to a functor by the Yoneda Lemma. Note that representability of $\mathcal{C}(- \otimes Y, K)$ is a property; the pair of the choice of DY as representing object and the isomorphism $\mathcal{C}(- \otimes Y, K) \simeq \mathcal{C}(-, DY)$, however, is only unique up to canonical isomorphism. While D is only essentially unique in the sense just explained, the requirement that D is an equivalence does not depend on the choice involved in the definition of D .

Since D is an equivalence we have

$$\mathcal{C}(X \otimes Y, K) \cong \mathcal{C}(X, DY) \cong \mathcal{C}(Y, D^{-1}X)$$

so $\mathcal{C}^{rev}$ is also a GV-category with D^{-1} . Note: $D(\mathbf{1}) \cong K$ as $- \otimes \mathbf{1} \cong (-) = id$, and similarly $D^{-1}(\mathbf{1}) \cong K$. Hence we have canonical isomorphisms

$$D^2(\mathbf{1}) \cong DK \cong DD^{-1}\mathbf{1} \cong \mathbf{1}$$

and

$$K \cong D\mathbf{1} \cong D^2D^{-1}\mathbf{1} \cong D^2K.$$

Example: Every (right) rigid monoidal category is an example of a Grothendieck-Verdier category with $D = (-)^\vee$ and $K = \mathbf{1}$.

4.20 Pivotal Grothendieck-Verdier category

A pivotal structure on a GV-category is a monoidal natural isomorphism $\omega : id \xRightarrow{\cong} D^2$ such that $\omega_K : K \rightarrow D^2$ agrees with the canonical isomorphism from above.

4.21 Braided Grothendieck-Verdier category

A braided category equipped with a GV duality is called a braided Grothendieck-Verdier category.

Analogous to Section 4.10, we have (BD, Prop 7.1):

If $\mathcal{C}$ is a braided GV-category, then there is a bijection between

- pivotal structures on $\mathcal{C}$
- twists satisfying $\theta_K = id_K$.

Hence pivotal braided GV = balanced GV with $\theta_K = id_K$.

4.22 Ribbon Grothendieck-Verdier category

This is a pivotal braided GV-category (= balanced GV-category with $\theta_K = id_K$) with the additional compatibility condition that $D\theta_X = \theta_{DX}$.

5 Abelian Categories and Related Structures

5.1 Kernels and cokernels

Let $\mathcal{C}$ be a category with a zero object, and let $f : X \rightarrow Y$ be a morphism. A *kernel* of f is a pair $(\ker(f), i : \ker(f) \rightarrow X)$ such that $f \circ i = 0$, and it is universal with respect to this property: if $g : Z \rightarrow X$ satisfies $f \circ g = 0$, then there exists a unique $g' : Z \rightarrow \ker(f)$ such that $i \circ g' = g$:

$$\begin{array}{ccccc} & & Z & & \\ & \swarrow \exists! g' & \downarrow g & & \\ \ker(f) & \xrightarrow{i} & X & \xrightarrow{f} & Y \end{array}$$

A *cokernel* of f is a pair $(\operatorname{coker}(f), \pi : Y \rightarrow \operatorname{coker}(f))$ such that $\pi \circ f = 0$, and it is universal with respect to this property: if $g : Y \rightarrow Z$ satisfies $g \circ f = 0$, then there exists a unique $g' : \operatorname{coker}(f) \rightarrow Z$ such that $g' \circ \pi = g$:

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{\pi} & \operatorname{coker}(f) \\ & & \downarrow g & \swarrow \exists! g' & \\ & & Z & & \end{array}$$

A kernel of a cokernel is called the *image* of f , denoted $\operatorname{im}(f) \rightarrow Y$. A cokernel of a kernel is called the *coimage*, denoted $X \rightarrow \operatorname{coim}(f)$. The universal properties guarantee uniqueness up to unique isomorphism.

5.2 Monos and epis

If $\mathcal{C}$ is an arbitrary category, a morphism $i : X \rightarrow Y$ is a *monomorphism* (or *monic*) if $i \circ f = i \circ g$ for morphisms $f, g : Z \rightarrow X$ implies $f = g$. Dually, a morphism $\pi : X \rightarrow Y$ is an *epimorphism* (or *epi*) if $f \circ \pi = g \circ \pi$ for morphisms $f, g : Y \rightarrow Z$ implies $f = g$.

If $\mathcal{C}$ is additive, then:

- $i : X \rightarrow Y$ is monic if and only if $i \circ f = 0$ implies $f = 0$;
- $\pi : X \rightarrow Y$ is epi if and only if $f \circ \pi = 0$ implies $f = 0$.

Another characterisation (still assuming additivity):

- $i : X \rightarrow Y$ is monic if and only if the unique morphism $0 \rightarrow X$ is a kernel of i ;
- $\pi : X \rightarrow Y$ is epi if and only if the unique morphism $Y \rightarrow 0$ is a cokernel of π .

5.3 Abelian category

If $\mathcal{C}$ is an additive category, one can easily see that all kernels are monic and all cokernels are epi. The converse might not hold. When it does, then we talk of an abelian category.

More precisely, an additive category $\mathcal{C}$ is *abelian* if:

- every morphism has a kernel and a cokernel,
- every monomorphism is the kernel of its cokernel,
- every epimorphism is the cokernel of its kernel.

If $F : \mathcal{A} \rightarrow \mathcal{B}$ is a functor between abelian categories, then:

- F is *left exact* if it is additive and preserves kernels: $F(\ker(f)) \cong \ker(Ff)$;
- F is *right exact* if it is additive and preserves cokernels: $F(\operatorname{coker}(f)) \cong \operatorname{coker}(Ff)$;
- F is *exact* if it is both left and right exact.

Exactness can also be expressed in terms of short exact sequences, where these are defined in abelian categories using kernels and images.

5.4 Linear + object semisimple imply abelian

This proof uses the key consequence that in a $\mathbb{k}$ -linear semisimple category, every monomorphism and every epimorphism splits.

Assumptions

- $\mathcal{C}$ is $\mathbb{k}$ -linear with finite biproducts $(\oplus)$.
- $\mathcal{C}$ is semisimple (every object is finite $\oplus$ of simples S_i , Schur's Lemma holds).
- **Key Consequence:** Every mono splits; every epi splits.

Goal: Show $\mathcal{C}$ is Abelian

(A) **Zero Object:** Exists (empty $\oplus$).

(B) **Finite Biproducts:** Exists (by assumption).

(C) **Existence of Kernels and Cokernels:**

- **Kernel of $f : X \rightarrow Y$:**
 1. Decompose $X \cong \bigoplus_{i \in I} S_i$. Let $J = \{i \in I \mid f \circ \operatorname{incl}_{S_i} = 0\}$.
 2. Define $K = \bigoplus_{j \in J} S_j$ and let $k : K \rightarrow X$ be the canonical inclusion.
 3. Check $f \circ k = 0$: True by definition of J .
 4. Check Universality: Let $g : Z \rightarrow X$ satisfy $f \circ g = 0$. Since $k : K \rightarrow X$ is a monomorphism (inclusion of summands), it splits by the Key Consequence. Let $p_K : X \rightarrow K$ be a projection such that $p_K \circ k = \operatorname{id}_K$. Define $h = p_K \circ g : Z \rightarrow K$. We must show $k \circ h = g$. Let $X \cong K \oplus K'$ where $k' : K' \rightarrow X$. Any map $g : Z \rightarrow X$ corresponds to a pair $(g_K, g_{K'}) = (p_K \circ g, p_{K'} \circ g)$.

We have

$$f \circ g = f \circ (k \circ g_K + k' \circ g_{K'}) = (f \circ k) \circ g_K + (f \circ k') \circ g_{K'} = 0 + (f \circ k') \circ g_{K'} = 0.$$

By construction of K , f restricted to K' (via $f \circ k'$) is non-zero on the simple components of K' . This forces $g_{K'} = 0$.

So g corresponds to $(g_K, 0)$, which means $g = k \circ g_K$. Taking $h = g_K = p_K \circ g$ gives $k \circ h = g$. Uniqueness of h follows because k is mono.

5. Kernels exist.

• **Cokernel of $f : X \rightarrow Y$:**

1. Let $I \subseteq Y$ be the subobject generated by the images of all simple summands of X under f . (Technically, $I = \sum \text{Im}(f \circ \text{incl}_{S_i})$.) Since $\mathcal{C}$ is semisimple, the subobject I is a direct summand: $Y \cong I \oplus C$.
2. Let $p : Y \rightarrow C$ be the projection onto C .
3. Check $p \circ f = 0$: The image of f lies entirely within I . The projection p annihilates I . So $p \circ f = 0$.
4. Check Universality: Let $h : Y \rightarrow Z$ satisfy $h \circ f = 0$. This implies h restricted to I is zero ($h \circ \text{incl}_I = 0$). Since $p : Y \rightarrow C$ is an epimorphism (projection from a direct sum), it splits by the Key Consequence. Let $s : C \rightarrow Y$ be a section ($p \circ s = \text{id}_C$). Define $q = h \circ s : C \rightarrow Z$. We must show $q \circ p = h$.
Any map $h : Y \rightarrow Z$ corresponds to a pair (h_I, h_C) where $h_I = h \circ \text{incl}_I$ and $h_C = h \circ s$. The condition $h \circ f = 0$ implies $h_I = 0$. So h corresponds to $(0, h_C)$.
 $q = h \circ s = h_C$.
 $q \circ p$ takes $y = (y_I, y_C)$, maps via p to y_C , maps via q to $h_C(y_C)$. This corresponds to the map $(0, h_C)$, which is h . Yes. Uniqueness of q follows because p is epi.
5. Cokernels exist.

(D) Monos are Kernels, Epis are Cokernels:

- Let $m : A \rightarrow X$ be mono. It splits, so $X \cong A \oplus C$. Let $p : X \rightarrow C$ be the projection. By the kernel construction in (C), $K = \ker(p)$ is the sum of simple summands of X annihilated by p , which is exactly A . So
$$m = \ker(p) = \ker(\text{coker}(m)).$$
- Let $e : X \rightarrow C$ be epi. It splits, so $X \cong K \oplus C$. Let $k : K \rightarrow X$ be inclusion. By the cokernel construction in (C), $P = \text{coker}(k)$ is the projection onto the summand complementary to K , which is C . So
$$e = \text{coker}(k) = \text{coker}(\ker(e)).$$

Conclusion: Since axioms (A)–(D) hold, $\mathcal{C}$ is an abelian category.

5.5 The Hom bifunctor is left exact in both variables

For any object X in an abelian category $\mathcal{A}$, the functor $F = \text{Hom}(X, -) : \mathcal{A} \rightarrow \mathbf{Ab}$ and the functor $G = \text{Hom}(-, X) : \mathcal{A}^{op} \rightarrow \mathbf{Ab}$ are left exact.

To prove that $\text{Hom}(X, -)$ is left exact, we must show that for any short exact sequence $0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$ in $\mathcal{A}$, the following induced sequence of abelian groups is exact:

$$0 \rightarrow \text{Hom}(X, A) \xrightarrow{f_*} \text{Hom}(X, B) \xrightarrow{g_*} \text{Hom}(X, C)$$

Here, the maps are induced by post-composition: $f_*(\alpha) = f \circ \alpha$ and $g_*(\beta) = g \circ \beta$. We must prove exactness at $\text{Hom}(X, A)$ and at $\text{Hom}(X, B)$.

1. Exactness at $\text{Hom}(X, A)$: We must show that f_* is a monomorphism (injective). Let $\alpha \in \text{Hom}(X, A)$ be a morphism such that $f_*(\alpha) = 0$. By definition, $f_*(\alpha) = f \circ \alpha = 0$. Since f is a monomorphism (from the original exact sequence), it is left-cancellable. From the equation $f \circ \alpha = f \circ 0$, we can cancel f to obtain $\alpha = 0$. Thus, $\ker(f_*) = \{0\}$, and f_* is injective.

2. Exactness at $\text{Hom}(X, B)$: We must show that $\text{im}(f_*) = \ker(g_*)$.

Inclusion $\text{im}(f_*) \subseteq \ker(g_*)$: Let $\beta \in \text{im}(f_*)$. By definition, there exists an $\alpha \in \text{Hom}(X, A)$ such that $\beta = f_*(\alpha) = f \circ \alpha$. We apply g_* to β :

$$g_*(\beta) = g_*(f \circ \alpha) = g \circ (f \circ \alpha) = (g \circ f) \circ \alpha.$$

From the original short exact sequence, we know that $\text{im}(f) = \ker(g)$, which implies that the composition $g \circ f = 0$. Therefore, $g_*(\beta) = 0 \circ \alpha = 0$. This shows that $\beta \in \ker(g_*)$, so $\text{im}(f_*) \subseteq \ker(g_*)$.

Inclusion $\ker(g_*) \subseteq \text{im}(f_*)$: Let $\beta \in \ker(g_*)$. This means that $g_*(\beta) = g \circ \beta = 0$. In the original sequence, (A, f) is the kernel of g . The universal property of kernels states that any morphism into B that becomes zero when composed with g must factor uniquely through the kernel of g . Since $g \circ \beta = 0$, there exists a unique morphism $\alpha : X \rightarrow A$ such that $\beta = f \circ \alpha$. By the definition of f_* , this is equivalent to $\beta = f_*(\alpha)$, which proves that $\beta \in \text{im}(f_*)$. Thus, $\ker(g_*) \subseteq \text{im}(f_*)$.

Since we have shown both inclusions, we conclude that $\text{im}(f_*) = \ker(g_*)$. This completes the proof of left exactness for the covariant Hom functor.

Now we show that $G = \text{Hom}(-, Y) : \mathcal{A}^{op} \rightarrow \mathbf{Ab}$ is left exact.

To prove that $\text{Hom}(-, Y)$ is left exact, we must show that for any short exact sequence $0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$ in $\mathcal{A}$, the following induced sequence of abelian groups is exact:

$$0 \rightarrow \text{Hom}(C, Y) \xrightarrow{g^*} \text{Hom}(B, Y) \xrightarrow{f^*} \text{Hom}(A, Y)$$

Here, the maps are induced by pre-composition: $g^*(\gamma) = \gamma \circ g$ and $f^*(\delta) = \delta \circ f$. We must prove exactness at $\text{Hom}(C, Y)$ and at $\text{Hom}(B, Y)$.

1. Exactness at $\text{Hom}(C, Y)$: We must show that g^* is a monomorphism (injective). Let $\gamma \in \text{Hom}(C, Y)$ be a morphism such that $g^*(\gamma) = 0$. By definition, $g^*(\gamma) = \gamma \circ g = 0$. Since g is an epimorphism (from the original exact sequence), it is right-cancellable. From the equation $\gamma \circ g = 0 \circ g$, we can cancel g to obtain $\gamma = 0$. Thus, $\ker(g^*) = \{0\}$, and g^* is injective.

2. Exactness at $\text{Hom}(B, Y)$: We must show that $\text{im}(g^*) = \ker(f^*)$.

Inclusion $\text{im}(g^*) \subseteq \ker(f^*)$: Let $\delta \in \text{im}(g^*)$. By definition, there exists a $\gamma \in \text{Hom}(C, Y)$ such that $\delta = g^*(\gamma) = \gamma \circ g$. We apply f^* to δ :

$$f^*(\delta) = f^*(\gamma \circ g) = (\gamma \circ g) \circ f = \gamma \circ (g \circ f).$$

Since $g \circ f = 0$ in the original sequence, we have $f^*(\delta) = \gamma \circ 0 = 0$. This shows that $\delta \in \ker(f^*)$, so $\text{im}(g^*) \subseteq \ker(f^*)$.

Inclusion $\ker(f^*) \subseteq \operatorname{im}(g^*)$: Let $\delta \in \ker(f^*)$. This means that $f^*(\delta) = \delta \circ f = 0$. In the original sequence, (C, g) is the cokernel of f . The universal property of cokernels states that any morphism from B that is zero when pre-composed with f must factor uniquely through the cokernel of f . Since $\delta \circ f = 0$, there exists a unique morphism $\gamma : C \rightarrow Y$ such that $\delta = \gamma \circ g$. By the definition of g^* , this is equivalent to $\delta = g^*(\gamma)$, which proves that $\delta \in \operatorname{im}(g^*)$. Thus, $\ker(f^*) \subseteq \operatorname{im}(g^*)$.

Since we have shown both inclusions, we conclude that $\operatorname{im}(g^*) = \ker(f^*)$. This completes the proof of left exactness for the contravariant Hom functor.

5.6 The functor $- \otimes_R M$ is right exact

Suppose for a moment that we are in the category of R -modules. Then the functor $- \otimes M : R\text{-mod} \rightarrow R\text{-mod}$ is right exact. It is NOT left exact in general. When it is, M is called a flat module.

5.7 Flat module

That M for which $- \otimes M : R\text{-mod} \rightarrow R\text{-mod}$ is exact.

Examples of flat modules: free modules and projective modules.

5.8 Right exact functors that coincide on projective objects coincide everywhere

Theorem: Let $\mathcal{A}$ and $\mathcal{B}$ be abelian categories, and assume that $\mathcal{A}$ has enough projectives. Let $F, G : \mathcal{A} \rightarrow \mathcal{B}$ be two right-exact functors. If there exists a natural isomorphism $\phi : F|_{\operatorname{Proj}(\mathcal{A})} \rightarrow G|_{\operatorname{Proj}(\mathcal{A})}$ on the full subcategory of projective objects $\operatorname{Proj}(\mathcal{A})$, then F and G are naturally isomorphic.

That is, if they “coincide” on projective objects they “coincide” everywhere.

In other words, a right-exact functor from a category with enough projectives is determined by its restriction to projective objects.

Proof Our goal is to construct a natural isomorphism $\eta : F \rightarrow G$. This requires us to define, for each object $X \in \mathcal{A}$, an isomorphism $\eta_X : F(X) \rightarrow G(X)$ such that for any morphism $f : X \rightarrow Y$ in $\mathcal{A}$, the following diagram commutes:

$$\begin{array}{ccc} F(X) & \xrightarrow{\eta_X} & G(X) \\ \downarrow F(f) & & \downarrow G(f) \\ F(Y) & \xrightarrow{\eta_Y} & G(Y) \end{array}$$

Step 1: Construction of the isomorphism η_X .

Since $\mathcal{A}$ has enough projectives, for any object $X \in \mathcal{A}$, we can choose a projective presentation, which is an exact sequence:

$$P_1 \xrightarrow{d_1} P_0 \xrightarrow{d_0} X \rightarrow 0$$

where P_0 and P_1 are projective objects.

Applying the right-exact functors F and G to this sequence yields two exact sequences in $\mathcal{B}$:

$$\begin{aligned} F(P_1) &\xrightarrow{F(d_1)} F(P_0) \xrightarrow{F(d_0)} F(X) \rightarrow 0 \\ G(P_1) &\xrightarrow{G(d_1)} G(P_0) \xrightarrow{G(d_0)} G(X) \rightarrow 0 \end{aligned}$$

From the exactness of these sequences, we can identify $F(X)$ and $G(X)$ as the cokernels of $F(d_1)$ and $G(d_1)$ respectively:

$$\begin{aligned} F(X) &\cong \operatorname{coker}(F(d_1)) \\ G(X) &\cong \operatorname{coker}(G(d_1)) \end{aligned}$$

By hypothesis, we have a natural isomorphism $\phi : F|_{\operatorname{Proj}(\mathcal{A})} \rightarrow G|_{\operatorname{Proj}(\mathcal{A})}$. Since P_0 and P_1 are projective, we have isomorphisms $\phi_{P_0} : F(P_0) \rightarrow G(P_0)$ and $\phi_{P_1} : F(P_1) \rightarrow G(P_1)$. The naturality of ϕ with respect to the morphism $d_1 : P_1 \rightarrow P_0$ means that the following diagram commutes:

$$\begin{array}{ccc} F(P_1) & \xrightarrow{\phi_{P_1}} & G(P_1) \\ \downarrow F(d_1) & & \downarrow G(d_1) \\ F(P_0) & \xrightarrow{\phi_{P_0}} & G(P_0) \end{array}$$

Since ϕ_{P_0} and ϕ_{P_1} are isomorphisms, this commutative diagram induces a unique isomorphism between the cokernels. We define $\eta_X : F(X) \rightarrow G(X)$ to be this unique isomorphism.

Step 2: Independence of the choice of projective presentation.

We must show that η_X is independent of the chosen projective presentation. Suppose we have another projective presentation $P'_1 \rightarrow P'_0 \rightarrow X \rightarrow 0$. By the fundamental lemma of homological algebra (or properties of projective objects), there exists a chain map between these two presentations, which is unique up to chain homotopy. This chain map induces a commutative diagram:

$$\begin{array}{ccccccc} P_1 & \xrightarrow{d_1} & P_0 & \xrightarrow{d_0} & X & \longrightarrow & 0 \\ \downarrow h_1 & & \downarrow h_0 & & \downarrow \operatorname{id}_X & & \\ P'_1 & \xrightarrow{d'_1} & P'_0 & \xrightarrow{d'_0} & X & \longrightarrow & 0 \end{array}$$

Applying the functors F and G and the natural isomorphism ϕ results in a larger commutative prism. The universal property of the cokernel then guarantees that the induced isomorphism η_X is the same, making our definition well-defined.

Step 3: Naturality of η .

Now, let $f : X \rightarrow Y$ be an arbitrary morphism in $\mathcal{A}$. We need to show that $G(f) \circ \eta_X = \eta_Y \circ F(f)$.

Choose projective presentations for both X and Y :

$$\begin{aligned} P_1 &\xrightarrow{d_1} P_0 \xrightarrow{d_0} X \rightarrow 0 \\ Q_1 &\xrightarrow{e_1} Q_0 \xrightarrow{e_0} Y \rightarrow 0 \end{aligned}$$

Since P_0 is projective and $e_0 : Q_0 \rightarrow Y$ is an epimorphism, we can lift the morphism $f : X \rightarrow Y$ to a morphism $f_0 : P_0 \rightarrow Q_0$ such that $e_0 \circ f_0 = f \circ d_0$. Similarly, we can find a morphism $f_1 : P_1 \rightarrow Q_1$ that makes the following diagram commute:

$$\begin{array}{ccccccc} P_1 & \xrightarrow{d_1} & P_0 & \xrightarrow{d_0} & X & \longrightarrow & 0 \\ \downarrow f_1 & & \downarrow f_0 & & \downarrow f & & \\ Q_1 & \xrightarrow{e_1} & Q_0 & \xrightarrow{e_0} & Y & \longrightarrow & 0 \end{array}$$

Now, we apply the functors F and G . By functoriality, we get two commutative diagrams in $\mathcal{B}$. We then combine these with the natural isomorphisms given by ϕ to form the faces of a three-dimensional diagram (a cube). The commutativity of the diagram involving f_0 and f_1 for F and G

respectively, combined with the naturality of ϕ for f_0 and f_1 , leads to the following commutative diagram:

$$\begin{array}{ccccc}
F(P_1) & \xrightarrow{F(d_1)} & F(P_0) & \xrightarrow{\pi_F} & F(X) \\
\downarrow F(f_1) & & \downarrow F(f_0) & & \downarrow F(f) \\
F(Q_1) & \xrightarrow{F(e_1)} & F(Q_0) & \xrightarrow{\pi'_F} & F(Y) \\
\uparrow \phi_{P_1}^{-1} & & \uparrow \phi_{P_0}^{-1} & & \uparrow \eta_X^{-1} \\
G(P_1) & \xrightarrow{G(d_1)} & G(P_0) & \xrightarrow{\pi_G} & G(X) \\
\uparrow \phi_{Q_1}^{-1} & & \uparrow \phi_{Q_0}^{-1} & & \uparrow G(f) \\
G(Q_1) & \xrightarrow{G(e_1)} & G(Q_0) & \xrightarrow{\pi'_G} & G(Y)
\end{array}$$

$\eta_Y \circ F(f)$ (dashed arrow from $F(X)$ to $G(Y)$)
 η_Y (solid arrow from $F(Y)$ to $G(Y)$)
 $G(f)$ (solid arrow from $G(X)$ to $G(Y)$)

where $\pi_F, \pi_G, \pi'_F, \pi'_G$ are the cokernel maps.

From the universal property of the cokernel, $F(X) = \text{coker}(F(d_1))$, there is a unique map $F(f) : F(X) \rightarrow F(Y)$ satisfying $F(f) \circ \pi_F = \pi'_F \circ F(f_0)$. Similarly, a unique map $G(f) : G(X) \rightarrow G(Y)$ exists.

Our definition of η_X and η_Y is via the isomorphisms induced by ϕ on the cokernels. The commutativity of all the constituent squares (due to functoriality and the naturality of ϕ) forces the outer diagram involving $F(f), G(f), \eta_X$, and η_Y to commute.

Specifically, consider the composition $\eta_Y \circ F(f) \circ \pi_F$. By the diagram, this is equal to $\eta_Y \circ \pi'_F \circ F(f_0)$. By the definition of η_Y , this is $\pi'_G \circ \phi_{Q_0} \circ F(f_0)$. By the naturality of ϕ , this is $\pi'_G \circ G(f_0) \circ \phi_{P_0}$. By the definition of $G(f)$, this is $G(f) \circ \pi_G \circ \phi_{P_0}$. Finally, by the definition of η_X , this is $G(f) \circ \eta_X \circ \pi_F$. So we have $(\eta_Y \circ F(f)) \circ \pi_F = (G(f) \circ \eta_X) \circ \pi_F$. Since π_F is an epimorphism, we can cancel it from the right, which gives us:

$$\eta_Y \circ F(f) = G(f) \circ \eta_X$$

This establishes the naturality of η . Thus, $\eta : F \rightarrow G$ is a natural isomorphism.

5.9 Monoidal abelian category

A category which is monoidal and abelian with the following compatibility conditions: the tensor product $\otimes$ is a biadditive functor (i.e., additive in each variable).

5.10 Monoidal abelian $\mathbb{k}$ -linear category

A category which is monoidal, abelian and $\mathbb{k}$ -linear with the following compatibility conditions: the $\mathbb{k}$ -module enrichment extends the abelian group enrichment, the tensor product $\otimes$ is a $\mathbb{k}$ -bilinear functor, and the unit $\mathbf{1}$ is a simple object with $\text{End}(\mathbf{1}) = \mathbb{k}$.

5.11 Projective and injective objects

An object P in an abelian category $\mathcal{A}$ is *projective* if for every epimorphism $p : X \rightarrow Y$ and every morphism $f : P \rightarrow Y$, there is a morphism $f' : P \rightarrow X$ that lifts p , i.e., $f = p \circ f'$:

$$\begin{array}{ccc}
& & X \\
& \nearrow \exists f' & \downarrow p \\
P & \xrightarrow{f} & Y
\end{array}$$

An object I in an abelian category $\mathcal{A}$ is *injective* if for every monomorphism $i : Y \rightarrow X$ and every morphism $f : Y \rightarrow I$, there is a morphism $f' : X \rightarrow I$ that extends i , i.e., $f = f' \circ i$:

$$\begin{array}{ccc} Y & \xrightarrow{f} & I \\ i \downarrow & \nearrow \exists f' & \\ X & & \end{array}$$

One can show, as for the category of modules, that an object P is projective if and only if the functor

$$\mathrm{Hom}_{\mathcal{A}}(P, -) : \mathcal{A} \rightarrow \mathbf{Ab}$$

is exact. Similarly, an object I is injective if and only if the functor

$$\mathrm{Hom}_{\mathcal{A}}(-, I) : \mathcal{A}^{\mathrm{op}} \rightarrow \mathbf{Ab}$$

is exact.

Another characterisation: an object P is projective if and only if every short exact sequence

$$0 \rightarrow X \rightarrow Y \rightarrow P \rightarrow 0$$

is split. Similarly, an object I is injective if and only if every short exact sequence

$$0 \rightarrow I \rightarrow X \rightarrow Y \rightarrow 0$$

is split.

In the abelian category of R -modules, free $\Rightarrow$ projective.

In the abelian category of $\mathbb{Z}$ -modules (i.e., abelian groups), divisible $\Rightarrow$ injective (recall that an abelian group A is divisible if for every non-zero integer n , the canonical map $A \rightarrow A$, $a \mapsto na$ is surjective. If A is torsion-free, then this is the same as A being uniquely divisible, i.e., that map being an isomorphism, because being torsion-free means precisely that such a map is always injective.) So examples are $\mathbb{Q}$, $\mathbb{Q}/\mathbb{Z}$, etc.

5.12 Direct sums and summands of projective and injective objects

Proposition: Let $X := X_1 \oplus X_2$. Then we have that X is projective (resp. injective) if and only if so are X_1 and X_2 . I.e., direct sums and direct summands of projective and injective objects are of the same type.

Proof. We prove the statement for injective objects; the projective case is dual.

($\Rightarrow$) Suppose $X = X_1 \oplus X_2$ is injective. We show that X_1 and X_2 are injective. We treat X_1 ; the argument for X_2 is identical.

Let $j : A \hookrightarrow B$ be a monomorphism and $f : A \rightarrow X_1$ a morphism. Consider the composition

$$i_1 \circ f : A \rightarrow X,$$

where $i_1 : X_1 \rightarrow X_1 \oplus X_2$ is the canonical inclusion.

Since X is injective, there exists $g : B \rightarrow X$ such that

$$g \circ j = i_1 \circ f.$$

Let $p_1 : X \rightarrow X_1$ be the projection. Define

$$g_1 := p_1 \circ g : B \rightarrow X_1.$$

Then

$$g_1 \circ j = p_1 \circ g \circ j = p_1 \circ i_1 \circ f = \text{id}_{X_1} \circ f = f.$$

Thus f extends, and X_1 is injective.

($\Leftarrow$) Suppose X_1 and X_2 are injective. Let

$$0 \rightarrow X_1 \oplus X_2 \xrightarrow{i} M \rightarrow N \rightarrow 0$$

be a short exact sequence. Let $p_k : X_1 \oplus X_2 \rightarrow X_k$ be the projections, and set

$$i_k := p_k \circ i : X_1 \oplus X_2 \rightarrow X_k.$$

Then each i_k restricts to a monomorphism $X_k \hookrightarrow M$. Since X_k is injective, there exists a retraction

$$r_k : M \rightarrow X_k$$

such that $r_k \circ i = p_k$.

Define

$$r := (r_1, r_2) : M \rightarrow X_1 \oplus X_2.$$

Then

$$r \circ i = (r_1 \circ i, r_2 \circ i) = (p_1, p_2) = \text{id}_{X_1 \oplus X_2},$$

so the sequence splits. Hence $X_1 \oplus X_2$ is injective.

The proof for projective objects is dual: one replaces monomorphisms by epimorphisms and uses the lifting property for projectives. $\square$

5.13 Projective cover and injective hull

Let $\mathcal{C}$ be a finite category and $X \in \mathcal{C}$.

(Projective cover.) A *projective cover* of X is a projective object $P(X) \in \mathcal{C}$ together with an epimorphism

$$p : P(X) \twoheadrightarrow X$$

such that for every epimorphism $g : P \twoheadrightarrow X$ from a projective object P , there exists an epimorphism $h : P \twoheadrightarrow P(X)$ such that

$$ph = g.$$

If a projective cover $P(X)$ of X exists, then it is unique up to a (non-unique) isomorphism.

(Injective hull.) An *injective hull* of X is an injective object $I(X) \in \mathcal{C}$ together with a monomorphism

$$\iota : X \hookrightarrow I(X)$$

such that for every monomorphism $f : X \hookrightarrow I$ into an injective object I , there exists a monomorphism $k : I(X) \hookrightarrow I$ such that

$$k\iota = f.$$

If an injective hull $I(X)$ of X exists, then it is unique up to a (non-unique) isomorphism.

5.14 Socle and top

If $\mathcal{C}$ is a finite category and $X \in \mathcal{C}$, the *socle* of X is the maximal semisimple subobject of X , ie, the direct sum of all its simple subobjects,

$$\text{soc}(X) = \text{sum of all simple subobjects of } X.$$

Similarly, the *top* of X is the maximal semisimple quotient,

$$\text{top}(X) = X/\text{rad}(X).$$

Here $\text{rad}(X)$ is the intersection of all maximal subobjects of X ($M \subset X$ maximal if X/M is simple).

Proposition 5.1. *There are bijections*

$$\{ \text{simple objects of } \mathcal{C} \} \xlongequal{\quad} \left\{ \begin{array}{c} \text{indecomposable} \\ \text{projective objects of} \\ \mathcal{C} \end{array} \right\}$$

$$S \longrightarrow P(S)$$

$$\text{top}(P) \longleftarrow P$$

and

$$\{ \text{simple objects of } \mathcal{C} \} \xlongequal{\quad} \left\{ \begin{array}{c} \text{indecomposable} \\ \text{injective objects of} \\ \mathcal{C} \end{array} \right\}$$

$$S \longrightarrow I(S)$$

$$\text{soc}(P) \longleftarrow I$$

where $P(S)$ and $I(S)$ are the projective cover and injective hull of S .

For instance: in a finite tensor category, the socle of an indecomposable projective object is simple. For if P is indecomposable projective, in a finite tensor category it is also injective and indecomposable, ie there is a simple object (unique!) such that P is its injective hull. So an indecomposable projective has a unique simple subobject, ie the socle is that precise object ie simple.

5.15 Semisimple category

This is the analogue of pre-fusion categories for abelian categories (instead of for linear categories).

Let $\mathcal{A}$ be an abelian category and let Y be an object. A *subobject* of Y is a monomorphism $i : X \rightarrow Y$. A *quotient object* of Y is an epimorphism $\pi : Y \rightarrow Z$.

A non-zero object X is said to be

- (i) *simple* if the source objects of its subobjects are all isomorphic to X or 0,
- (ii) *semisimple* if it is a coproduct of simple objects,

(iii) *indecomposable* if it is not a coproduct of at least two non-zero subobjects.

An abelian category is *semisimple* if all its objects are semisimple.

Note: there are many other versions of what a semisimple category should be in other contexts different than abelian categories, see [here](#) for an overview and comparison.

5.16 Semisimplicity vs. Projectivity in Abelian Categories

Theorem: Let $\mathcal{C}$ be an abelian category. The following conditions are equivalent:

1. $\mathcal{C}$ is **semisimple** (every object is a direct sum of simple objects).
2. Every short exact sequence in $\mathcal{C}$ **splits**.
3. Every object in $\mathcal{C}$ is **projective**.
4. Every object in $\mathcal{C}$ is **injective**.
5. Every subobject of any object in $\mathcal{C}$ is a **direct summand**.

Proof Sketch for (1) $\iff$ (3)

(3) $\implies$ (1): Assume every object is projective. Show $\mathcal{C}$ is semisimple.

- It suffices to show that every short exact sequence (SES)

$$0 \rightarrow A \xrightarrow{i} B \xrightarrow{p} C \rightarrow 0$$

splits.

- By assumption, the object C is projective.
- The definition of projective means that for the epimorphism $p : B \rightarrow C$ and the identity morphism $\text{id}_C : C \rightarrow C$, there exists a lifting (a section) $s : C \rightarrow B$ such that $p \circ s = \text{id}_C$.
- The existence of such a section s means the SES splits.
- Since every SES splits, $\mathcal{C}$ is semisimple.

(1) $\implies$ (3): Assume $\mathcal{C}$ is semisimple. Show every object P is projective.

- We need to show that for any epimorphism $p : B \rightarrow C$ and any morphism $f : P \rightarrow C$, there exists $\tilde{f} : P \rightarrow B$ such that $p \circ \tilde{f} = f$.
- Since $\mathcal{C}$ is semisimple, every SES splits. Consider the SES

$$0 \rightarrow K \xrightarrow{i} B \xrightarrow{p} C \rightarrow 0,$$

where $K = \ker(p)$. This sequence splits.

- Splitting implies $B \cong K \oplus C$. Under this isomorphism, let p correspond to the projection

$$p_C : K \oplus C \rightarrow C.$$

- Given $f : P \rightarrow C$, define $\tilde{f}' : P \rightarrow K \oplus C$ by $\tilde{f}' = (0, f)$, where $0 : P \rightarrow K$ is the zero morphism.
- Let $\tilde{f} : P \rightarrow B$ correspond to $\tilde{f}'$ via the isomorphism $B \cong K \oplus C$.
- Then

$$p \circ \tilde{f} = p_C \circ \tilde{f}' = p_C \circ (0, f) = f.$$

- We have found the required lifting $\tilde{f}$, so P is projective.

Thus, an abelian category is semisimple if and only if all of its objects are projective.

5.17 Locally finite abelian $\mathbb{k}$ -linear category

An object A of an abelian category has *finite length* if there is a Jordan-Hölder series of finite length

$$0 = A_0 \rightarrow A_1 \rightarrow \cdots \rightarrow A_{n-1} \rightarrow A_n = A$$

of monomorphisms such that each object

$$A_{i+1}/A_i := \text{coker}(A_i \rightarrow A_{i+1})$$

is simple.

If S is a simple object, we say that a given Jordan-Hölder series contains S with multiplicity m if the number of values of i for which A_{i+1}/A_i is isomorphic to S is m . The Jordan-Hölder theorem states that all Jordan-Hölder series of a given object have the same length, and more particularly, that any two Jordan-Hölder series contain any simple object with the same multiplicity. We denote $[A : S] := m$.

Let $\mathbb{k}$ be a field. An abelian $\mathbb{k}$ -linear category $\mathcal{A}$ is *locally finite* if:

1. all $\text{Hom}_{\mathcal{A}}(X, Y)$ are finite-dimensional $\mathbb{k}$ -vector spaces,
2. every object has finite length.

5.18 Finite abelian $\mathbb{k}$ -linear category

Let $\mathbb{k}$ be a field. A locally finite abelian $\mathbb{k}$ -linear category $\mathcal{A}$ is *finite* if:

1. there are finitely many isomorphism classes of simple objects,
2. every simple object has a projective cover, that is, for every simple object A there exists a projective object P_A and an epimorphism $\pi : P_A \rightarrow A$ which is universal: any other epimorphism $p : P \rightarrow A$ from a projective object P factors through π , i.e., there exists an epimorphism $p' : P \rightarrow P_A$ such that $p = \pi \circ p'$.

2. can be replaced for the simpler condition that for every object X there exists an epimorphism $P \twoheadrightarrow X$ from a projective object. This is usually called "having enough projectives". This also means that every object has a projective presentation.

It is a theorem by Deligne that any finite abelian $\mathbb{k}$ -linear category is equivalent to the category Mod_A^{fd} of finite-dimensional A -modules for some finite-dimensional algebra A . More precisely, the given finite abelian $\mathbb{k}$ -linear category determines the Morita equivalence class of A (recall that two algebras are Morita equivalent if their categories of modules are equivalent).

5.19 A finite abelian category has a projective generator

Proof. Let $\mathcal{A}$ be a finite abelian category over a field k . By definition, $\mathcal{A}$ is

- k -linear and additive,
- of finite length (every object has a finite composition series),
- Hom-finite: $\dim_k \operatorname{Hom}(X, Y) < \infty$ for all $X, Y \in \mathcal{A}$,
- and every simple object has a projective cover.

Step 1: Finitely many simples and indecomposable projectives.

Let $\{S_1, \dots, S_n\}$ be a complete set of representatives of the simple objects of $\mathcal{A}$ (up to isomorphism). Each simple S_i has a projective cover $P_i \twoheadrightarrow S_i$, which is indecomposable.

Step 2: Every indecomposable projective is one of the P_i .

Let Q be an indecomposable projective in $\mathcal{A}$. Since $\mathcal{A}$ is finite length, Q has a simple quotient S . Let $P_S \twoheadrightarrow S$ be the projective cover of S . By projectivity of Q , the surjection $P_S \twoheadrightarrow S$ lifts to a morphism $P_S \rightarrow Q$. Since both P_S and Q are indecomposable and the map is nonzero, it must be an isomorphism. Hence $Q \cong P_S$.

Thus the set $\{P_1, \dots, P_n\}$ gives all indecomposable projectives up to isomorphism.

IN OTHER WORDS: THERE IS A BIJECTION BETWEEN SIMPLE OBJECTS AND INDECOMPOSABLE PROJECTIVE OBJECTS (and there are finitely many of them)

Step 3: Constructing a projective generator.

Consider the direct sum

$$P := \bigoplus_{i=1}^n P_i.$$

- P is projective because finite direct sums of projectives are projective.
- P is a generator: for any object $X \in \mathcal{A}$, take a composition series

$$0 = X_0 \subset X_1 \subset \dots \subset X_m = X, \quad X_j/X_{j-1} \cong S_{i_j}.$$

For each simple subquotient S_{i_j} , the projective P_{i_j} surjects onto it. By induction on the length of X , one can construct an epimorphism

$$\bigoplus_{j=1}^m P_{i_j} \twoheadrightarrow X.$$

Since each P_{i_j} is a summand of P , X is a quotient of a direct sum of copies of P .

Conclusion: The object P is a projective generator of $\mathcal{A}$. □

5.20 Left/right exactness, adjointness and (dual) representability

Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be linear functors between finite linear categories. The following statements are equivalent:

- (L1) F is left-exact.
- (L2) F admits a left adjoint.

Likewise, the following are equivalent:

(R1) G is right-exact.

(R2) G admits a right adjoint.

Moreover, for $\mathcal{D} = \mathbf{vect}$, in addition the following statements are equivalent to (L1) and (L2) and to (R1) and (R2), respectively:

(L3) F is representable, i.e. $F \cong \mathrm{Hom}_{\mathcal{C}}(c, -)$ for some $c \in \mathcal{C}$.

(R3) G is ‘dually representable’, i.e. $G \cong \mathrm{Hom}_{\mathcal{C}}(-, d)^*$ for some $d \in \mathcal{C}$.

5.21 Deligne product

Given $\mathbb{k}$ -linear abelian categories $\mathcal{C}, \mathcal{D}$, their Deligne product (if it exists) is another $\mathbb{k}$ -linear abelian category $\mathcal{C} \bullet \mathcal{D}$ together with a bilinear right exact functor

$$\mathcal{C} \times \mathcal{D} \rightarrow \mathcal{C} \bullet \mathcal{D}$$

that induces equivalences

$$\mathrm{Rex}[\mathcal{C} \bullet \mathcal{D}, \mathcal{E}] \simeq \mathrm{Rex}[\mathcal{C}, \mathcal{D}; \mathcal{E}]$$

for all abelian $\mathcal{E}$. The Deligne product might not always exist.

5.22 Kelly product

Given $\mathbb{k}$ -linear finitely cocomplete categories $\mathcal{C}, \mathcal{D}$, their Kelly product (if it exists) is another $\mathbb{k}$ -linear finitely cocomplete category $\mathcal{C} \boxtimes \mathcal{D}$ together with a bilinear right exact functor (i.e., finite colimit preserving)

$$\mathcal{C} \times \mathcal{D} \rightarrow \mathcal{C} \boxtimes \mathcal{D}$$

that induces equivalences

$$\mathrm{Rex}[\mathcal{C} \boxtimes \mathcal{D}, \mathcal{E}] \simeq \mathrm{Rex}[\mathcal{C}, \mathcal{D}; \mathcal{E}]$$

for all $\mathcal{E}$ finitely cocomplete. It always exists. One can realise

$$\mathcal{C} \boxtimes \mathcal{D} \subset \mathrm{Lex}[\mathcal{C}^{\mathrm{op}}, \mathcal{D}^{\mathrm{op}}; \mathbb{k}\text{-Mod}].$$

Example: If R, S are $\mathbb{k}$ -algebras, the functor

$$\otimes_{\mathbb{k}} : R\text{-Mod}_f \times S\text{-Mod}_f \rightarrow (R \otimes_{\mathbb{k}} S)\text{-Mod}_f$$

induces an equivalence

$$(R \otimes_{\mathbb{k}} S)\text{-Mod}_f \simeq R\text{-Mod}_f \boxtimes S\text{-Mod}_f.$$

Facts: (Franco, EGNO)

(1) If $\mathcal{C}, \mathcal{D}$ are abelian, then $\mathcal{C} \boxtimes \mathcal{D}$ is abelian if and only if $\mathcal{C} \bullet \mathcal{D}$ exists, and in that case:

$$\mathcal{C} \boxtimes \mathcal{D} \simeq \mathcal{C} \bullet \mathcal{D}.$$

(2) If $\mathcal{C}, \mathcal{D}$ are locally finite abelian $\mathbb{k}$ -linear categories, then so is $\mathcal{C} \boxtimes \mathcal{D}$.

Moral: Among locally finite abelian $\mathbb{k}$ -linear categories, the Kelly and Deligne products exist, and they coincide. This is called the *Deligne-Kelly tensor product*.

Further properties:

- (1) $\text{Hom}_{\mathcal{C} \boxtimes \mathcal{D}}(X \boxtimes X', Y \boxtimes Y') \cong \text{Hom}_{\mathcal{C}}(X, Y) \otimes_{\mathbb{k}} \text{Hom}_{\mathcal{D}}(X', Y')$;
- (2) A bilinear biexact functor $\mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ induces an exact functor $\mathcal{C} \boxtimes \mathcal{D} \rightarrow \mathcal{E}$;
- (3) Right exact functors $F_i : \mathcal{C}_i \rightarrow \mathcal{D}_i$ between locally finite abelian linear categories give rise to a right exact functor:

$$F_1 \boxtimes F_2 : \mathcal{C}_1 \boxtimes \mathcal{C}_2 \rightarrow \mathcal{D}_1 \boxtimes \mathcal{D}_2.$$

5.23 (Multi)tensor, fusion and ring categories

Let $\mathcal{C}$ be a locally finite abelian $\mathbb{k}$ -linear monoidal category.

- $\mathcal{C}$ is a **multiring** category if $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is bilinear and biexact.
- If $\text{End}(1) \cong \mathbb{k}$, it is a **ring** category.

If $\mathcal{C}$ is in addition rigid:

- **multitensor** if $\otimes$ is bilinear on morphisms,
- **multifusion** if it is finite semisimple multitensor,
- **tensor** if multitensor and $\text{End}(1) \cong \mathbb{k}$,
- **fusion** if multifusion and $\text{End}(1) \cong \mathbb{k}$.

Properties:

1. In a multiring category with left (resp. right) duals, the corresponding dualisation functor is exact.
2. A finite ring category with left duals is a tensor category.
3. If P is projective in a multiring category and X has a left (resp. right) dual, then $P \otimes X$ (resp. $X \otimes P$) is projective.
4. In a multiring category with left duals (e.g., multitensor),

$$1 \in \mathcal{C} \text{ is projective iff } \mathcal{C} \text{ is semisimple.}$$

5. In a ring category with left duals (e.g., tensor), $1 \in \mathcal{C}$ is simple; in general in multiring with left duals it is semisimple.
6. The Deligne-Kelly product of (multi)tensor, (multi)fusion, or (multi)ring categories is of such type.

5.24 Indecomposable projective objects in a finite tensor category

Let $\mathcal{C}$ be a finite multitensor category over a field k . Let $\{X_i\}_{i \in I}$ be the set of isomorphism classes of simple objects in $\mathcal{C}$, and let P_i be the projective cover of X_i .

Then for any object $Z \in \mathcal{C}$, we have the following isomorphisms in $\mathcal{C}$:

1. $P_i \otimes Z \cong \bigoplus_{j,k \in I} N_{k,j^*}^i [Z : X_j] P_k$
2. $Z \otimes P_i \cong \bigoplus_{j,k \in I} N_{*j,k}^i [Z : X_j] P_k$

where:

- $[Z : X_j]$ is the multiplicity of the simple object X_j in the Jordan-Hölder series of Z .
- j^* and $*j$ denote the indices for the right dual X_j^* and the left dual $*X_j$, respectively.
- $N_{ab}^c := [X_a \otimes X_b : X_c]$ is the fusion coefficient, i.e., the multiplicity of X_c in the Jordan-Hölder series of $X_a \otimes X_b$.

Detailed Proof of Formula 1 We want to prove that:

$$P_i \otimes Z \cong \bigoplus_{j,k \in I} N_{k,j^*}^i [Z : X_j] P_k.$$

Step 1: The object $P_i \otimes Z$ is projective.

In a tensor category, the tensor product of a projective object with any other object is projective. Since P_i is projective by definition, the object $P_i \otimes Z$ is also a projective object in $\mathcal{C}$.

Step 2: Decomposition of projective objects.

Every projective object in a finite abelian category (like $\mathcal{C}$) can be uniquely decomposed (up to isomorphism and permutation) into a direct sum of indecomposable projective objects. The indecomposable projectives in $\mathcal{C}$ are precisely $\{P_k\}_{k \in I}$.

More precisely, there is a bijection between simple objects of $\mathcal{C}$ and indecomposable projective objects of $\mathcal{C}$, the bijection being $X_i \mapsto P_i$ and $P \mapsto P/\text{rad}(P)$.

Therefore, we can write an isomorphism in $\mathcal{C}$:

$$P_i \otimes Z \cong \bigoplus_{k \in I} m_k P_k,$$

where m_k are non-negative integers representing the multiplicity of P_k in the decomposition. Our goal is to find an explicit formula for m_k .

Step 3: Calculating the multiplicities m_k .

The multiplicity m_k can be determined by computing the dimension of the space of morphisms from the object to the corresponding simple object X_k . This relies on the property of projective covers that $\text{Hom}_{\mathcal{C}}(P_l, X_k) \cong k$ if $l = k$ and is 0 otherwise. More generally:

$$\dim \text{Hom}_{\mathcal{C}} \left(\bigoplus_{l \in I} m_l P_l, X_k \right) = \sum_{l \in I} m_l \dim \text{Hom}_{\mathcal{C}}(P_l, X_k) = \sum_{l \in I} m_l \delta_{lk} = m_k$$

Therefore, we can calculate m_k as:

$$m_k = \dim \text{Hom}_{\mathcal{C}}(P_i \otimes Z, X_k)$$

Step 4: Using Tensor-Hom Adjunction.

The tensor functor $- \otimes Z$ has a right adjoint, which is $- \otimes Z^*$, where Z^* is the right dual of Z . This gives the following natural isomorphism of vector spaces (the "adjunction of rigidity"):

$$\mathrm{Hom}_{\mathcal{C}}(P_i \otimes Z, X_k) \cong \mathrm{Hom}_{\mathcal{C}}(P_i, X_k \otimes Z^*)$$

Combining this with the previous step, we get:

$$m_k = \dim \mathrm{Hom}_{\mathcal{C}}(P_i, X_k \otimes Z^*)$$

Step 5: Using the Defining Property of Projective Covers.

The projective cover P_i of the simple object X_i has the fundamental property that for any object $M \in \mathcal{C}$:

$$\dim \mathrm{Hom}_{\mathcal{C}}(P_i, M) = [M : X_i]$$

This means the dimension of the Hom space from P_i to M counts the number of times the simple object X_i appears in the composition series of M .

Let's apply this property with $M = X_k \otimes Z^*$:

$$m_k = [X_k \otimes Z^* : X_i]$$

as desired.

5.25 Nakayama functors: classical setting

Let A be a fin dim algebra and write $\mathrm{Hom}_A(-, -)$ for the hom of the category $A\text{-mod}$ of fin dim A -modules.

Both A and A^* can be viewed as a left/right/bi module with the regular action, ie $a \cdot b := ab$ for the left one; and the co-regular left/right/bimod action which is the dual of the regular one: for the left one this is multiplication by a in A^* is the map $(a \cdot -)^*$, ie $(a \cdot \omega)(b) = \omega(ab)$.

The left and right Nakayama functors of $A\text{-mod}$ are the functors $A\text{-mod} \rightarrow A\text{-mod}$ given by

$$\mathbf{N}_A^r := \mathrm{Hom}_A(-, A)^* \quad , \quad \mathbf{N}_A^l := \mathrm{Hom}_{A^{op}}(-^*, A).$$

This requires some explanation: first of all if V is a left module then $\mathrm{Hom}_A(V, A)$ is a right module with $(f \cdot a)(v) = f(a \cdot v)$. On the other hand if V is a right A -module then V^* is a left A -module with $(a \cdot \omega)(v) = \omega(v \cdot a)$. Also recall that $A^{op}\text{-mod} \cong \text{mod-}A$.

One can show, see [Ivanov prop 3.1] that

$$\mathbf{N}_A^r \cong A^* \otimes_A - \quad , \quad \mathbf{N}_A^l \cong \mathrm{Hom}_A(A^*, -).$$

A word on this: recall that given a right A -module M and a left A -module N , we have $M \otimes_A N := M \otimes_{\mathbb{K}} N / (ma \otimes n - m \otimes an)$. In particular if M is an (B, A) -bimodule then $M \otimes_A N$ is a left B -module.

Returning to Nakayama functors, if we view A^* as a A -bimodule with the left and right co-regular action, we get that $A^* \otimes_A N$ is a left module whenever so is N . In particular it is very easy to see that this functor is right exact ie preserves cokernels. A similar note goes for the left exact cousin.

Theorem 5.2 ([Iv, Sect. 3]). *We have*

1. *The right Nakayama functor takes projective objects to injective objects.*

2. The left Nakayama functor takes injective objects to projective objects.
3. Restriction induces an equivalence of categories

$$\mathbf{N}_A^r : \mathbf{Proj}(A\text{-mod}) \xrightarrow{\simeq} \mathbf{Inj}(A\text{-mod}) : \mathbf{N}_A^l.$$

Proof of (1). Let A be a finite-dimensional algebra over a field k , and let

$$\mathbf{N}_A^r = - \otimes_A A^*$$

be the right Nakayama functor, where $(-)^* = \text{Hom}_k(-, k)$.

Let P be a projective A -module. Since P is projective, it is a direct summand of a finite free module, i.e. there exists $n \in \mathbb{N}$ such that

$$P \oplus KA^{\oplus n}.$$

Since the functor $- \otimes_A A^*$ is additive and preserves direct summands, and direct sums and direct summands of injective objects are injective, it suffices to prove that

$$A \otimes_A A^* \cong A^*$$

is injective as an A -module.

We now show that A^* is injective. For any right A -module M , there is a natural isomorphism

$$\text{Hom}_A(M, A^*) \cong \text{Hom}_k(M, k) = M^*,$$

given by sending a morphism $f : M \rightarrow A^*$ to the linear functional

$$m \mapsto f(m)(1).$$

One checks that this construction is functorial and defines a natural isomorphism.

Since k is a field, the functor $(-)^* = \text{Hom}_k(-, k)$ is exact. Hence the functor

$$\text{Hom}_A(-, A^*)$$

is exact, and therefore A^* is an injective right A -module.

It follows that $A \otimes_A A^*$ is injective. Since injective modules are closed under finite direct sums and direct summands, we conclude that

$$P \otimes_A A^*$$

is injective for every projective module P .

This proves that the right Nakayama functor takes projective objects to injective objects. $\square$

5.26 The Nakayama permutation

A natural question is under which conditions the right and left Nakayama functors are equivalences.

Lemma 5.3. *The following properties of a finite-dimensional $\mathbb{k}$ -algebra A are equivalent:*

- (i) *The right and left Nakayama functors are quasi-inverse of each other.*
- (ii) *A is injective as a left A -module,*
- (iii) *A is injective as a right A -module,*

(iv) The class of (left or right) projective modules coincides with the class of (left or right) injective modules.

An algebra with any (and thus all) of these properties is called *self-injective* or *quasi-Frobenius*.

For self-injective A the right and left Nakayama functors are in particular exact and there are natural isomorphisms

$$\mathbf{N}_A^r \cong {}_A \langle {}_A \langle A^*, A \rangle, - \rangle \quad \text{and} \quad \mathbf{N}_A^l \cong {}_A \langle A^*, A \rangle \otimes_A -, \quad (39)$$

(here ${}_A \langle A^*, A \rangle, - \rangle$ is seen as a bimodule).

So if A is self-injective aka quasi-Frobenius, injective and projective objects are the same. So the bijections between simple objects and indecomposable inj/proj can be composed:

$$\begin{aligned} \{ \text{simple objects of } \mathcal{C} \} & \xlongequal{\quad} \left\{ \begin{array}{c} \text{indecomposable} \\ \text{projective objects} \\ = \text{indecomposable} \\ \text{injective objects of} \\ \mathcal{C} \end{array} \right\} \xlongequal{\quad} \{ \text{simple objects of } \mathcal{C} \} \\ S & \xrightarrow{\quad} P(S) \xrightarrow{\quad} \text{soc}(P(S)) \end{aligned}$$

This defines a permutation on the set of simples of $\mathcal{C}$ called the *Nakayama permutation* and denoted ν . Sometimes it is useful to write $\mathcal{S}(A)$ for the set of isoclasses of simple A -modules and

$$\nu : \mathcal{S}(A) \rightarrow \mathcal{S}(A) \quad , \quad S_i \mapsto S_{\nu(i)}$$

, ie viewing ν as permuting the indices of the simple objects. In this language if $P_i = P(S_i)$ and $I_j = I(S_j)$

$$\text{top}(P_i) = S_i \quad , \quad \text{soc}(P_i) = S_{\nu(i)}$$

and

$$\text{top}(I_j) = S_{\nu^{-1}(j)} \quad , \quad \text{soc}(I_j) = S_j.$$

It interacts nicely with the right Nakayama functor :

Proposition 5.4. *Let A be a self-injective fin dim algebra. Then*

1. $\mathbf{N}^r(P(S)) \cong I(S)$,
2. $\mathbf{N}^r(S) \cong \nu^{-1}(S)$, ie, $\mathbf{N}^r(S_i) \cong S_{\nu^{-1}(i)}$.

5.27 The Nakayama automorphism

Let A be a Frobenius algebra with counit ε . Let $\kappa = \langle -, - \rangle : A \otimes A \rightarrow k$ be the non-degenerate associative bilinear form defining the Frobenius structure, ie $\langle x, y \rangle = \varepsilon(xy)$. This κ is called the *Frobenius form*.

By non-degeneracy, there is a k -linear isomorphism

$$\Psi : A \longrightarrow A^*, \quad \Psi(z) = \langle -, z \rangle.$$

Now for $a \in A$, we have $\langle a, - \rangle \in A^*$. Therefore there is a unique element

$$a' := \Psi^{-1}(\langle a, - \rangle) \in A$$

such that

$$\langle a, - \rangle = \langle -, a' \rangle \quad \text{as elements of } A^*.$$

We define

$$\nu : A \longrightarrow A, \quad a \longmapsto a'.$$

By linearity of $\langle -, - \rangle$, this defines a k -linear map, and by the associativity of $\langle -, - \rangle$, it is in fact an algebra map.

To see that it is an automorphism, it suffices to check injectivity: if $\nu(a) = 0$, then

$$\langle a, - \rangle = \langle -, \nu(a) \rangle = 0,$$

and by non-degeneracy, $a = 0$. This is the *Nakayama automorphism* of A . Note that it is actually measuring how far is the Frobenius structure from being symmetric.

What is the relation between the Nakayama automorphism, the Nakayama functor and the Nakayama permutation? It turns out that the Nakayama functor, what it does is to twist the action of A on a given module by the Nakayama automorphism.

More precisely, if $\alpha : A \rightarrow A$ is an automorphism of A , and M is an A -module, let ${}_{\alpha}M$ be M as vector space but with A -module structure twisted by α ,

$$a * m := \alpha(a) \cdot m.$$

Let ν be the Nakayama automorphism as before. It turns out that the right Nakayama functor of A twists the A -action of the A -modules: for any A -module M we have

$$\mathbf{N}^r(M) \cong {}_{\nu}M. \tag{40}$$

so in view of Theorem 5.4 we have

$${}_{\nu}S_i \cong \mathbf{N}^r(S_i) \cong S_{\nu^{-1}(i)}.$$

In other words, for a simple S ,

$$\nu(S) \cong {}_{\nu^{-1}}S.$$

Same goes for the indecomposable projective / injective objects.

5.28 Comparison: quasi-Frobenius vs Frobenius algebras

Any Frob algebra is quasi-Frobenius, but the opposite is NOT true. What is true is that any quasi-Frob algebra is Morita equivalent to a Frob algebra, ie, for a finite category, being quasi-Frobenius = self injective, ie being equivalent to $A - \text{mod}$ for a quasi-Frobenius = self injective algebra, is the same as being equivalent to $A - \text{mod}$ for a Frobenius algebra.

When is a quasi-Frobenius algebra actually Frobenius?

Theorem 5.5. *A fin dim quasi-Frobenius algebra A is Frobenius if and only if*

$$\dim_{\mathbb{k}} \nu(S) = \dim_{\mathbb{k}} S$$

for all simple A -modules S .

5.29 Nakayama functors: Morita-independent version

Let $\mathcal{A}$ be a finite category and $X \in \mathcal{A}$. By Theorem 6.1, the representable functor $\mathrm{Hom}_{\mathcal{A}}(X, -) : \mathcal{A} \rightarrow \mathbf{vect}$ has a left adjoint that we denote $(-) \cdot X$. For $V \in \mathbf{vect}$, the object $V \cdot X$ is called the *copower* and exhibits $\mathcal{A}$ as a category tensored over $\mathbf{vect}$ or, in the language of Section 6.10, as a finite module category over $\mathbf{vect}$.

The aim of this subsection is describe the so-called Nakayama functors, that will play a major role in the sequel. To this end, we first recall the following

Theorem 5.6 ([?, Theorem 3.2]). *Let $\mathcal{A}, \mathcal{B}$ be finite categories. The functors*

$$\mathcal{A}^{\mathrm{op}} \boxtimes \mathcal{B} \longrightarrow \mathrm{Lex}(\mathcal{A}, \mathcal{B}) \quad , \quad X \boxtimes Y \longmapsto \mathrm{Hom}_{\mathcal{A}}(X, -) \cdot Y$$

and

$$\mathcal{A}^{\mathrm{op}} \boxtimes \mathcal{B} \longrightarrow \mathrm{Rex}(\mathcal{A}, \mathcal{B}) \quad , \quad X \boxtimes Y \longmapsto \mathrm{Hom}_{\mathcal{A}}(-, X)^* \cdot Y$$

define a zig-zag of adjoint equivalences,

$$\mathrm{Lex}(\mathcal{A}, \mathcal{B}) \xleftarrow{\simeq} \mathcal{A}^{\mathrm{op}} \boxtimes \mathcal{B} \xrightarrow{\simeq} \mathrm{Rex}(\mathcal{A}, \mathcal{B}). \quad (41)$$

The quasi-inverses of these functors can be described as follows: to every left exact functor F , we assign the coend $\int^{X \in \mathcal{A}} X \boxtimes F(X)$, and to every right exact functor G , the end $\int_{X \in \mathcal{A}} X \boxtimes G(X)$.

The previous theorem can be viewed as a Morita invariant version of the classical Eilenberg-Watts theorem,

$$\mathrm{Lex}(A\text{-mod}_{\mathbb{k}}, B\text{-mod}_{\mathbb{k}}) \simeq B\text{-bimod}_{\mathbb{k}}\text{-}A \simeq \mathrm{Rex}(A\text{-mod}_{\mathbb{k}}, B\text{-mod}_{\mathbb{k}})$$

as $B\text{-bimod}_{\mathbb{k}}\text{-}A \simeq A\text{-mod}_{\mathbb{k}}^{\mathrm{op}} \boxtimes B\text{-mod}_{\mathbb{k}}$.

Because the identity functor of $\mathcal{A}$ is both left and right exact, Theorem 5.6 gives rise to two canonical functors: the *left Nakayama functor* $\mathbf{N}^{\ell} \in \mathrm{Lex}(\mathcal{A}, \mathcal{A})$ and the *right Nakayama functor* $\mathbf{N}^r \in \mathrm{Rex}(\mathcal{A}, \mathcal{A})$. Explicitly,

$$\mathbf{N}^{\ell} = \int_{X \in \mathcal{A}} \mathrm{Hom}_{\mathcal{A}}(X, -) \cdot X \quad , \quad \mathbf{N}^r = \int^{X \in \mathcal{A}} \mathrm{Hom}_{\mathcal{A}}(-, X)^* \cdot X. \quad (42)$$

Here we have used that the Hom functor (resp. dual Hom functor) preserves ends (resp. coends) as it is left (resp. right) exact. It follows directly from the zig-zag (41) that we have the following isomorphisms in $\mathcal{A}^{\mathrm{op}} \boxtimes \mathcal{A}$:

$$\int^{X \in \mathcal{A}} X \boxtimes \mathbf{N}^{\ell}(X) \cong \int_{X \in \mathcal{A}} X \boxtimes X \quad , \quad \int_{X \in \mathcal{A}} X \boxtimes \mathbf{N}^r(X) \cong \int^{X \in \mathcal{A}} X \boxtimes X. \quad (43)$$

We remark that the left and right Nakayama functors of a finite category $\mathcal{A}$ are very closely related: the left (resp. right) Nakayama functor of $\mathcal{A}$ is precisely the right (resp. left) Nakayama functor of $\mathcal{A}^{\mathrm{op}}$, viewed as an endofunctor of $\mathcal{A}$. That is, $\mathbf{N}_{\mathcal{A}^{\mathrm{op}}}^{\ell} \cong (\mathbf{N}_{\mathcal{A}}^r)^{\mathrm{op}}$ and $\mathbf{N}_{\mathcal{A}^{\mathrm{op}}}^r \cong (\mathbf{N}_{\mathcal{A}}^{\ell})^{\mathrm{op}}$.

Another remarkable property is that the Nakayama functors of the Deligne product of two finite categories factor as the Deligne product of the Nakayama functors of the two categories,

$$\mathbf{N}_{\mathcal{A} \boxtimes \mathcal{B}}^{\ell} \cong \mathbf{N}_{\mathcal{A}}^{\ell} \boxtimes \mathbf{N}_{\mathcal{B}}^{\ell} \quad , \quad \mathbf{N}_{\mathcal{A} \boxtimes \mathcal{B}}^r \cong \mathbf{N}_{\mathcal{A}}^r \boxtimes \mathbf{N}_{\mathcal{B}}^r, \quad (44)$$

see [?, Proposition 3.20].

5.30 The canonical coend Δ

Let $\mathcal{A}$ be a finite tensor category. An object $X \in \mathcal{A}$ is said to be *invertible* if the evaluation $\text{ev}_X : X^\vee \otimes X \rightarrow \mathbb{1}$ and the coevaluation $\text{coev}_X : \mathbb{1} \rightarrow X \otimes X^\vee$ are isomorphisms. In that case, $X^\vee$ is often denoted as X^{-1} . The goal of this subsection is to introduce the so-called distinguished invertible element of a finite tensor category.

If $\mathcal{A}$ is a finite tensor category, its *envelope* is $\mathcal{A}^{\text{env}} := \mathcal{A} \boxtimes \mathcal{A}^{\text{rev}}$, which is a finite tensor category by Theorem 6.3. Note that because of (49), this category is in fact equivalent to $\mathcal{A} \boxtimes \mathcal{A}^{\text{op}}$. The category $\mathcal{A}$ can be viewed as a left $\mathcal{A}^{\text{env}}$ -module category with action functor determined by

$$(X \boxtimes Y) \otimes Z = X \otimes Z \otimes Y.$$

By the discussion of Section 6.11, we have that $\Delta := \underline{\text{End}}(\mathbb{1}) \in \mathcal{A}^{\text{env}}$ is an algebra object, that is usually called the *canonical algebra* of $\mathcal{A}$. By a result of Shimizu [?, Lemma 4.4], the canonical algebra can actually be expressed as the coend

$$\Delta \cong \int^{X \in \mathcal{A}} X^\vee \boxtimes X. \quad (45)$$

Moreover by a change of variables

$$\Delta \cong \int^{X \in \mathcal{C}} X^\vee \boxtimes X \cong \int^{X \in \mathcal{C}} X \boxtimes {}^\vee X$$

5.31 The distinguished invertible object (take 1)

Let $\mathcal{C}$ be a rigid monoidal category. An object $X \in \mathcal{C}$ is **invertible** if the evaluation and coevaluation maps give isomorphisms

$$X^* \otimes X \cong \mathbb{1} \quad \text{and} \quad \mathbb{1} \cong X \otimes X^*.$$

Example: In the category $\text{Rep}(G)$, the 1-dimensional representations are invertible.

Let $\mathcal{A}$ be a finite tensor category with (simple) unit $\mathbf{1}$. Define

$$\alpha := \text{soc}(P(\mathbf{1})), \quad (46)$$

ie the socle of the projective cover of the unit. In other words: if ν denotes the Nakayama permutation,

$$\alpha = \nu(\mathbf{1}).$$

In particular, this implies that

$$\mathbf{N}^r(\alpha) \cong \mathbf{1}.$$

5.32 Unimodular finite tensor category

A finite tensor category is **unimodular** if its distinguished invertible object α is the unit $\mathbf{1}$.

Example: In the category $\text{Rep}(G)$, the 1-dimensional representations are invertible.

Theorem 5.7 (ENO04). *For a finite tensor category,*

$$\text{semisimple} \Rightarrow \text{unimodular}.$$

5.33 The distinguished invertible object (take 2)

The monoidal unit $\mathbb{1} \in \mathcal{A}$ is well-known to be an $\mathcal{A}^{\text{env}}$ -progenerator: indeed by (54) we have $\underline{\text{Hom}}(\mathbb{1}, -) \cong (- \boxtimes \mathbb{1}) \otimes \underline{\text{End}}(\mathbb{1})$, which is right exact; and since $\underline{\text{End}}(\mathbb{1}) \neq 0$, the argument from [?, Lemma 6.11] tells us that $\underline{\text{Hom}}(\mathbb{1}, -)$ is also faithful. Therefore by Theorem 6.2 the functor

$$\underline{\text{Hom}}(\mathbb{1}, -) : \mathcal{A} \xrightarrow{\cong} \text{mod}_{\mathcal{A}^{\text{env}}-\Delta}$$

is an equivalence of $\mathcal{A}^{\text{env}}$ -module categories. Furthermore (54) implies that this functor is naturally isomorphic to the functor $(- \boxtimes \mathbb{1}) \otimes \Delta_{\Delta}$, where Δ_{Δ} denotes Δ viewed as a right Δ -module by the multiplication of Δ . By means of the equivalence, there exists an object α , unique up to isomorphism, such that

$$(\alpha \boxtimes \mathbb{1}) \otimes \Delta_{\Delta} \cong (\Delta \Delta)^{\vee}. \quad (47)$$

It can be shown that this object α is in fact invertible [?, §7.18] (see also [?, Lemma 4.3]), and will be called the *distinguished invertible object* of $\mathcal{A}$. When $\alpha \cong \mathbb{1}$, we will say that $\mathcal{A}$ is *unimodular*. This name comes from the theory of Hopf algebras: a finite-dimensional Hopf algebra H is unimodular (i.e., left- and right-integrals coincide) if and only if the category $H\text{-mod}_{\mathbb{k}}$ is unimodular. We refer the reader to [?, Theorem 4.10] for a list of different characterizations of unimodularity. Note that when $\mathcal{A}$ is unimodular, (62) implies the self-duality of Δ .

One of the main properties of the distinguished invertible object is that it implements the quadruple dual by conjugation [?, Theorem 3.3]: we have a natural isomorphism of monoidal functors

$$(-)^{\vee\vee\vee\vee} \cong \alpha \otimes - \otimes \alpha^{-1}. \quad (48)$$

In particular, if $\mathcal{A}$ is unimodular, then $(-)^{\vee\vee\vee\vee} \cong \text{id}_{\mathcal{A}}$.

We have two different versions of the distinguished invertible object in a finite tensor category. Both coincide:

Proposition 5.8 (Shimizu's unimodular Lemma 5.5). *Both definitions of α agree.*

Proof. Set $B(X, Y) = \text{Hom}_{\mathcal{C}}(X, \mathbb{1}) \cdot Y$. We denote by $\pi : D \rightrightarrows B$ the universal dinatural transformation of the end. For $X_1, X_2 \in \mathcal{C}$, we have

$$\begin{aligned} \pi_{X_1 \oplus X_2} &= B(\text{id}_{X_1 \oplus X_2}, i_1 p_1) \circ \pi_{X_1 \oplus X_2} + B(\text{id}_{X_1 \oplus X_2}, i_2 p_2) \circ \pi_{X_1 \oplus X_2} \\ &= B(p_1, i_1) \circ \pi_{X_1} + B(p_2, i_2) \circ \pi_{X_2}, \end{aligned}$$

where $i_r : X_r \rightarrow X_1 \oplus X_2$ and $p_r : X_1 \oplus X_2 \rightarrow X_r$ ($r = 1, 2$) are the inclusion and the projection, respectively. Hence, if $\pi_{X_1} = 0$ and $\pi_{X_2} = 0$, then $\pi_{X_1 \oplus X_2} = 0$.

Now let $V_0 = \mathbb{1}, V_1, \dots, V_n$ be a complete set of representatives of the isomorphism classes of simple objects of $\mathcal{C}$, and let P_i be the projective cover of V_i . Suppose that $\pi_{P_i} = 0$ for all i . Then, by the above argument, $\pi_P = 0$ for all projective object $P \in \mathcal{C}$. For each $X \in \mathcal{C}$, there are a projective object $P \in \mathcal{C}$ and an epimorphism $f : P \rightarrow X$. Since $B(f, X)$ is monic and $B(f, \text{id}_X) \circ \pi_X = B(\text{id}_P, f) \circ \pi_P = 0$, we have $\pi_X = 0$. Hence $D = 0$ by the universal property. This contradicts the fact that D is a simple object.

By the above argument, $\pi_{P_i} \neq 0$ for some i . Since $B(P_i, P_i) = 0$ for $i \neq 0$, the morphism π_{P_0} must be non-zero. Since D is a simple object, the morphism π_0 induces a monomorphism from D to the socle S of P_0 . On the other hand, it is known that S is simple [EO]. Thus $D \cong S$. $\square$

5.34 Self-duality of Δ in a unimodular ftc

TO BE FINISHED

Three different ways to see that Δ is self-dual:

- 1- the def of the distinguished invertible object
- 2- the lemma something in Shimizu's papepr

3- when the category is unimodular and pivotal, then the right Nakayama functor is nat isomorphic to the identity, and by FSS 3.52 we have

$$\int^{X \in \mathcal{C}} X \boxtimes X \cong \int_{X \in \mathcal{C}} X \boxtimes X.$$

Since the duality functor is an equivalence it preserves all limits and colimits and then applying $(-)^{\vee} \boxtimes id$ we get

$$\int^{X \in \mathcal{C}} X^{\vee} \boxtimes X \cong \int_{X \in \mathcal{C}} X^{\vee} \boxtimes X.$$

We conclude as

$$\Delta^{\vee} \cong \left(\int^{X \in \mathcal{C}} X^{\vee} \boxtimes X \right)^{\vee} \cong \int_{X \in \mathcal{C}} X^{\vee\vee} \boxtimes X^{\vee} \cong \int_{X \in \mathcal{C}} X^{\vee} \boxtimes X \cong \int^{X \in \mathcal{C}} X^{\vee} \boxtimes X \cong \Delta$$

5.35 Spherical finite tensor category à la Douglas-Schommer-Pries-Snyder

A pivotal finite tensor category is *spherical* (according to Douglas-Schommer-Pries-Snyder) if it is unimodular, ie $\alpha \cong \mathbf{1}$, and the trivialisation $(-)^{\vee\vee\vee\vee} \cong \text{Id}$ coming from the Radford isomorphism coincides with the trivialisation $(-)^{\vee\vee\vee\vee} \cong (-)^{\vee\vee} \cong \text{Id}$ induced by the pivotal structure.

This notion of sphericity is related to the sphericity in terms of quantum traces from Section 4.6 only in the semisimple case. Beyond semisimplicity, it is an entirely different notion. Most remarkably, sphericity, even in the non-semisimple situation, can be perfectly characterized through traces if one uses modified traces instead of quantum traces [SS21, Theorem 1.3].

Let us make this precise:

Definition 5.9. A pivotal tensor category $(\mathcal{C}, P : (-) \simeq (-)^{**})$ is *trace spherical* if for all objects $V \in \mathcal{C}$, the quantum trace of the pivotal structure on V agrees with the quantum trace of the pivotal structure on $*V$, that is $\text{Tr}(P_V) = \text{Tr}(P_{*V})$.

In the semisimple case, the two notions correspond precisely:

Proposition 5.10. *A semisimple pivotal finite tensor category with absolutely simple unit is spherical if and only if it is trace spherical.*

Proof. Both the property of sphericity and the property of trace sphericity are invariant under base change, so it suffices to compare them over an algebraically closed field. When the base field is algebraically closed, Etingof-Nikshych-Ostrik prove in [ENO04], Thm 7.3 and Cor 7.4] that the Radford equivalence $\mathcal{R}_{\mathcal{C}} : (-) \xrightarrow{\sim} (-)^{****}$ is characterized by a property that is equivalent to the following property: for any object $V \in \mathcal{C}$ and any morphism $f_V : V \rightarrow V^{**}$, there is an equality of traces $\text{Tr}(f_V) = \text{Tr}((f_V)^* \circ \mathcal{R}_{\mathcal{C}}(*V))$. Here we have abbreviated $\mathcal{R}_{\mathcal{C}}(*V)$ as $\mathcal{R}_{(*V)}$. When $(\mathcal{C}, P)$ is a pivotal tensor category, we have $(P_V)^* = (P_{*V})^{-1}$ [Sel11], Lemma 4.11; SR72, Prop 5.2.3]. If $(\mathcal{C}, P)$ is spherical, it follows that

$$\text{Tr}(P_V) = \text{Tr}((P_V)^* \circ \mathcal{R}_{(*V)}) = \text{Tr}(P_{V*}^{-1} \circ P_{V*} \circ P_{(*V)}) = \text{Tr}(P_{(*V)});$$

that is, $(\mathcal{C}, P)$ is trace spherical. The converse follows by cyclically permuting this equation. $\square$

WARNING: In the setting of non-semisimple finite tensor categories, it is neither the case that spherical implies trace spherical, nor the case that trace spherical implies spherical.

5.36 Eilenberg-Watts theorem

If $\mathcal{A}, \mathcal{B}$ are finite abelian categories, then there is a zig-zag of equivalences

$$\mathrm{Lex}(\mathcal{A}, \mathcal{B}) \simeq \mathcal{A}^{op} \boxtimes \mathcal{B} \simeq \mathrm{Rex}(\mathcal{A}, \mathcal{B})$$

TO BE FINISHED

5.37 The canonical Hopf algebra in a braided finite tensor category

Let $\mathcal{C}$ be a finite tensor category over a field k , braided or otherwise. We consider the bifunctor

$$T : \mathcal{C}^{op} \times \mathcal{C} \longrightarrow \mathcal{C}, \quad (Y, X) \longmapsto X \otimes Y^*.$$

We wish to form the coend

$$\mathbb{F} = \int^{X \in \mathcal{C}} X \otimes X^*.$$

This is the universal object equipped with a *dinatural transformation*

$$i_X : X \otimes X^* \longrightarrow \mathbb{F},$$

meaning that for every morphism $f : X \rightarrow Y$ in $\mathcal{C}$, the following diagram commutes:

$$\begin{array}{ccc} X \otimes X^* & \xrightarrow{f \otimes \mathrm{id}_{X^*}} & Y \otimes X^* \\ & \searrow i_X & \downarrow \\ & & \mathbb{F} \\ & \nearrow i_Y & \uparrow \\ X \otimes Y^* & \xrightarrow{\mathrm{id}_X \otimes f^*} & X \otimes X^* \end{array}$$

More precisely, $i_Y \circ (\mathrm{id}_Y \otimes f^*) = i_X \circ (f \otimes \mathrm{id}_{X^*})$ for all $f : X \rightarrow Y$.

Proposition 5.11. *The coend $\mathbb{F} = \int^{X \in \mathcal{C}} X \otimes X^*$ exists in $\mathcal{C}$.*

Proof. Since $\mathcal{C}$ is a finite tensor category, it has finitely many isomorphism classes of simple objects $L_1, \dots, L_n$ and all Hom-spaces are finite-dimensional over k . As a k -linear abelian category, $\mathcal{C}$ has all finite colimits (it is equivalent, as a k -linear category, to the category of finite-dimensional modules over a finite-dimensional k -algebra, which is cocomplete over finite diagrams).

Any coend can be computed as a coequalizer. Concretely, $\mathbb{F}$ is the coequalizer of the pair of maps

$$\bigoplus_{f: X \rightarrow Y} Y \otimes X^* \rightrightarrows \bigoplus_{Z \in \mathrm{Irr}(\mathcal{C})} Z \otimes Z^*,$$

where the two maps send the summand indexed by $f : X \rightarrow Y$ to the summands indexed by X and Y via $f \otimes \mathrm{id}_{X^*}$ and $\mathrm{id}_Y \otimes f^*$ respectively, and the coproduct on the right ranges over representatives of isomorphism classes of simple objects (by semisimplicity of the relevant Hom-spaces, one reduces the index set to the finitely many simples and the finitely many morphisms between them). Since both direct sums are *finite*, this coequalizer diagram is finite, and hence exists in $\mathcal{C}$. $\square$

The finiteness of $\mathcal{C}$ (finitely many simples, finite-dimensional hom-spaces) is essential: it reduces the a priori infinitely indexed coproducts in the coend formula to finite direct sums, which exist in any abelian category.

When $\mathcal{C}$ is **braided**, with braiding $c_{X,Y} : X \otimes Y \xrightarrow{\sim} Y \otimes X$, the object $\mathbb{F}$ acquires the structure of a **Hopf algebra in $\mathcal{C}$** . All structure maps are defined by specifying a dinatural family into (or out of) $\mathbb{F}$ and invoking the universal property of the coend:

- **Multiplication** $\mu : \mathbb{F} \otimes \mathbb{F} \rightarrow \mathbb{F}$:

Since $\mathbb{F} \otimes \mathbb{F} \cong \int^X \int^Y (X \otimes X^*) \otimes (Y \otimes Y^*)$ (the tensor product of coends is the iterated coend, by the Fubini theorem for coends), it suffices to give a dinatural family

$$\mu_{X,Y} : X \otimes X^* \otimes Y \otimes Y^* \longrightarrow \mathbb{F}.$$

Define it as the composite

$$X \otimes X^* \otimes Y \otimes Y^* \xrightarrow{\text{id}_X \otimes c_{X^*,Y} \otimes \text{id}_{Y^*}} X \otimes Y \otimes X^* \otimes Y^* \xrightarrow{\sim} (X \otimes Y) \otimes (X \otimes Y)^* \xrightarrow{i_{X \otimes Y}} \mathbb{F},$$

where the middle isomorphism uses $(X \otimes Y)^* \cong Y^* \otimes X^*$ and the braiding $c_{X^*,Y}$ is used to permute the middle factors. One verifies dinaturality in both X and Y using the naturality of the braiding and the dinaturality of $i_{(-)}$. The resulting morphism $\mu : \mathbb{F} \otimes \mathbb{F} \rightarrow \mathbb{F}$ is associative.

- **Unit** $\eta : \mathbf{1} \rightarrow \mathbb{F}$:

Set

$$\eta = i_{\mathbf{1}} \circ \lambda_{\mathbf{1}}^{-1} : \mathbf{1} \xrightarrow{\sim} \mathbf{1} \otimes \mathbf{1}^* \xrightarrow{i_{\mathbf{1}}} \mathbb{F},$$

using the canonical isomorphism $\mathbf{1}^* \cong \mathbf{1}$ and the unit constraint λ .

- **Comultiplication** $\Delta : \mathbb{F} \rightarrow \mathbb{F} \otimes \mathbb{F}$

Define the dinatural family $\delta_X : X \otimes X^* \rightarrow \mathbb{F} \otimes \mathbb{F}$ by

$$X \otimes X^* \xrightarrow{\text{id}_X \otimes \text{coev}_Y \otimes \text{id}_{X^*}} X \otimes Y \otimes Y^* \otimes X^* \xrightarrow{\text{id}_X \otimes \text{id}_Y \otimes c_{Y^*,X^*}} X \otimes Y \otimes X^* \otimes Y^* \xrightarrow{i_X \otimes i_Y} \mathbb{F} \otimes \mathbb{F},$$

summing over all Y (or more precisely, inserting a copy via coevaluation $\text{coev}_Y : \mathbf{1} \rightarrow Y \otimes Y^*$ for each Y and then using the braiding to sort the dual factors). The braiding c_{Y^*,X^*} rearranges $Y^* \otimes X^*$ into $X^* \otimes Y^*$ so that the two tensor factors land correctly in the two copies of $\mathbb{F}$. The result is coassociative.

- **Counit** $\varepsilon : \mathbb{F} \rightarrow \mathbf{1}$:

The evaluation morphism $\text{ev}_X : X \otimes X^* \rightarrow \mathbf{1}$ is natural (hence dinatural) in X . By the universal property of $\mathbb{F}$ it factors uniquely through a morphism

$$\varepsilon : \mathbb{F} \longrightarrow \mathbf{1}.$$

- **Antipode** $S : \mathbb{F} \rightarrow \mathbb{F}$:

In a pivotal braided category, define the dinatural family $s_X : X \otimes X^* \rightarrow \mathbb{F}$ by

$$X \otimes X^* \xrightarrow{c_{X,X^*}} X^* \otimes X \xrightarrow{\sim} X^* \otimes X^{**} \xrightarrow{i_{X^*}} \mathbb{F},$$

using the braiding to swap X and X^* , and the pivotal structure to identify $X \cong X^{**}$. This gives a well-defined map $S : \mathbb{F} \rightarrow \mathbb{F}$ satisfying the Hopf antipode axiom $\mu \circ (S \otimes \text{id}) \circ \Delta = \eta \circ \varepsilon = \mu \circ (\text{id} \otimes S) \circ \Delta$.

Furthermore, this Hopf algebra comes with a Hopf pairing

$$\omega_{\mathcal{C}} : \mathbb{F} \otimes \mathbb{F} \rightarrow \mathbb{F}.$$

As with the other structure maps, it is defined by specifying a dinatural family and invoking the universal property of $\mathbb{F} \otimes \mathbb{F}$ as an iterated coend. Define

$$\omega_{X,Y} : (X \otimes X^*) \otimes (Y \otimes Y^*) \longrightarrow \mathbf{1}$$

by the composite

$$X \otimes X^* \otimes Y \otimes Y^* \xrightarrow{\text{id}_X \otimes c_{X^*,Y} \otimes \text{id}_{Y^*}} X \otimes Y \otimes X^* \otimes Y^* \xrightarrow{\text{ev}_X \otimes \text{ev}_Y} \mathbf{1} \otimes \mathbf{1} \xrightarrow{\sim} \mathbf{1},$$

where we use the braiding $c_{X^*,Y}$ to bring X^* next to X and Y^* next to Y , and then evaluate both pairs simultaneously. More explicitly, this is the composite

$$X \otimes X^* \otimes Y \otimes Y^* \xrightarrow{\text{id} \otimes c_{X^*,Y} \otimes \text{id}} X \otimes Y \otimes X^* \otimes Y^* \xrightarrow{c_{X,Y} \otimes c_{X^*,Y^*}} Y \otimes X \otimes Y^* \otimes X^* \xrightarrow{\text{ev}_Y \otimes \text{ev}_X} \mathbf{1},$$

where one uses the braiding to arrange evaluation pairings (X, X^*) and (Y, Y^*) next to each other. One checks that this family is dinatural in both X and Y , so it descends to a well-defined morphism $\omega : \mathbb{F} \otimes \mathbb{F} \rightarrow \mathbf{1}$.

5.38 Non-degenerate braided finite tensor category

Let $\mathcal{C}$ be a braided finite tensor category. We say that $\mathcal{C}$ is **non-degenerate** if so is $\omega_{\mathcal{C}}$.

5.39 Weakly-factorisable braided finite tensor category

Let $\mathcal{C}$ be a braided finite tensor category. We say that $\mathcal{C}$ is **weakly-factorisable** if the map

$$\Omega_{\mathcal{C}} : \text{Hom}_{\mathcal{C}}(\mathbf{1}, \mathbb{F}) \rightarrow \text{Hom}_{\mathcal{C}}(\mathbb{F}, \mathbf{1}) \quad , \quad f \mapsto (f \otimes \text{id}_{\mathbb{F}}) \circ \omega_{\mathcal{C}}$$

is invertible.

5.40 Factorisable braided finite tensor category

Let $\mathcal{C}$ be a braided tensor category with braiding σ , and let $\bar{\mathcal{C}}$ denote the finite tensor category $\mathcal{C}^{\text{rev}}$ equipped with the braiding $\bar{\sigma}$ given by

$$(\bar{\sigma}_{V,W} : V \otimes^{\text{rev}} W \rightarrow W \otimes^{\text{rev}} V) := (\sigma_{V,W}^{-1} : W \otimes V \rightarrow V \otimes W)$$

for $V, W \in \mathcal{C}$. By using the universal property of the Deligne tensor product, we define the k -linear exact functor G by

$$G : \mathcal{C} \boxtimes \bar{\mathcal{C}} \rightarrow \mathcal{Z}(\mathcal{C}), \quad V \boxtimes W \mapsto (V \otimes W, c), \quad (3.10)$$

where the natural isomorphism c is given by

$$c_X : V \otimes W \otimes X \xrightarrow{\text{id}_V \otimes \sigma_{X,W}^{-1}} V \otimes X \otimes W \xrightarrow{\sigma_{V,X} \otimes \text{id}_W} X \otimes V \otimes W \quad (3.11)$$

for $X \in \mathcal{C}$. The functor G is in fact a braided tensor functor with the monoidal structure given by the braiding of $\mathcal{Z}(\mathcal{C})$.

Definition 5.12. The braided finite tensor category $\mathcal{C}$ is *factorizable* if the functor G from above is an equivalence.

IMPORTANT: Factorisable braided finite tensor category $\Rightarrow$ Unimodular by [EGNO, 8.10.10].

5.41 Modular category (non-ss)

The following is Theorem 1.1 in Shimizu's "Non-degeneracy conditions for braided..."

For a braided finite tensor category $\mathcal{C}$, the following assertions are equivalent:

1. $\mathcal{C}$ is non-degenerate.
2. $\mathcal{C}$ is factorizable.
3. $\mathcal{C}$ is weakly-factorizable.
4. The Müger center of $\mathcal{C}$ is trivial, ie $\mathcal{Z}_2(\mathcal{C}) \simeq \text{Vec}_{\mathbb{k}}$.

A ribbon finite tensor category satisfying any of the four equivalent conditions above is called a **modular tensor category**.

5.42 Hopf algebra terminology and relation with categorical ones

Let H be a finite-dimensional Hopf algebra.

1. H is **quasi-triangular** if....
2. H is **triangular** if it is quasi-triangular with $R^{-1} = R_{21}$.
3. H is **factorisable** if it is quasi-triangular and the map

$$H^* \rightarrow H \quad , \quad \omega \mapsto (\omega \otimes id)(R_{21}R)$$

is an isomorphism (for H not nec fin dim, simply surjective). This is the opposite of being triangular.

Equivalently, by a result of Schneider this can be characterised by $D(H) \cong (H \otimes H)^{F_R}$ (something is twisted by a cocycle), in particular $D(H) \cong H \otimes H$ as algebras.

4. H is **unimodular** if it has left and right integrals and they coincide.

Sweedler-Lardson theorem: H is semisimple if and only if there is a left integral Λ with $\varepsilon(\Lambda) \neq 0$.

Radford theorem: Factorisable $\Rightarrow$ Unimodular.

5. H is **ribbon** if

IMPORTANT NOTE: For a fd Hopf algebra,

- Semisimple implies unimodular
- Factorisable implies unimodular.

Proposition 5.13. *Let H be a fin dim Hopf algebra.*

1. $H\text{-mod}$ is a finite tensor category.
2. If H is quasi-triangular, then $H\text{-mod}$ is braided.
3. If H is triangular, then $H\text{-mod}$ is symmetric.

4. If H is factorisable, then $H\text{-mod}$ is factorisable. Roughly

$$H\text{-mod} \boxtimes H\text{-mod} \simeq (H \otimes H)\text{-mod} \simeq D(H)\text{-mod} \simeq \mathcal{Z}(H\text{-mod}).$$

5. If H is unimodular, then $H\text{-mod}$ is unimodular [EGNO, §6.5].

6. If H is ribbon, then $H\text{-mod}$ is also ribbon.

7. If H is a factorisable ribbon Hopf algebra, then $H\text{-mod}$ is a modular tensor category.

8. If H is a semisimple factorisable ribbon Hopf algebra, then $H\text{-mod}$ is a modular fusion category.

9. In particular if H is a semisimple fin dim Hopf algebra over a field of char 0, then $D(H)$ is a semisimple factorizable ribbon Hopf algebra (with balancing element 1) and then $D(H)\text{-mod}$ is a modular fusion category.

6 Module categories

6.1 Definition

Let $\mathcal{C} = (\mathcal{C}, \otimes, \mathbf{1}, a, l, r)$ be a monoidal category.

A **left module category** over $\mathcal{C}$ is a category $\mathcal{M}$ equipped with:

- a bifunctor $\otimes : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$,
- a natural isomorphism (module associativity constraint)

$$m_{X,Y,M} : (X \otimes Y) \otimes M \xrightarrow{\sim} X \otimes (Y \otimes M),$$

- a natural isomorphism (unit constraint)

$$l_M : \mathbf{1} \otimes M \xrightarrow{\sim} M,$$

satisfying two coherence conditions:

- **Pentagon axiom:** the diagram for all $X, Y, Z \in \mathcal{C}$, $M \in \mathcal{M}$,

$$\begin{aligned} ((X \otimes Y) \otimes Z) \otimes M &\xrightarrow{a_{X,Y,Z} \otimes \text{id}_M} (X \otimes (Y \otimes Z)) \otimes M \\ &\xrightarrow{m_{X,Y \otimes Z, M}} X \otimes ((Y \otimes Z) \otimes M) \\ &\xrightarrow{\text{id}_X \otimes m_{Y,Z,M}} X \otimes (Y \otimes (Z \otimes M)) \\ &= (X \otimes Y) \otimes (Z \otimes M) \\ &\xleftarrow{m_{X,Y,Z \otimes M}} ((X \otimes Y) \otimes Z) \otimes M \end{aligned}$$

- **Triangle axiom:** for all $X \in \mathcal{C}$, $M \in \mathcal{M}$,

$$(X \otimes \mathbf{1}) \otimes M \xrightarrow{m_{X,\mathbf{1},M}} X \otimes (\mathbf{1} \otimes M) \xrightarrow{\text{id}_X \otimes l_M} X \otimes M$$

equals the map

$$(X \otimes \mathbf{1}) \otimes M \xrightarrow{r_X \otimes \text{id}_M} X \otimes M$$

This categorifies the notion of a module over a monoid.

There is a bijective correspondence between:

- left $\mathcal{C}$ -module category structures on $\mathcal{M}$, and
- monoidal functors $F : \mathcal{C} \rightarrow \text{End}(\mathcal{M})$,

where $\text{End}(\mathcal{M})$ is the monoidal category of endofunctors of $\mathcal{M}$ with composition.

If $\mathcal{C}$ is rigid, then for all $X \in \mathcal{C}, M, N \in \mathcal{M}$ there is a canonical isomorphism:

$$\text{Hom}_{\mathcal{M}}(X^* \otimes M, N) \cong \text{Hom}_{\mathcal{M}}(M, X \otimes N)$$

natural in all variables.

6.2 $\mathcal{C}$ -module functor

Let $\mathcal{M}, \mathcal{N}$ be two left $\mathcal{C}$ -module categories. A **$\mathcal{C}$ -module functor** is:

- a functor $F : \mathcal{M} \rightarrow \mathcal{N}$,
- a natural isomorphism

$$s_{X,M} : F(X \otimes M) \xrightarrow{\sim} X \otimes F(M),$$

natural in $X \in \mathcal{C}, M \in \mathcal{M}$, such that for all $X, Y \in \mathcal{C}, M \in \mathcal{M}$ the diagram

$$\begin{aligned} F((X \otimes Y) \otimes M) &\xrightarrow{F(m_{X,Y,M})} F(X \otimes (Y \otimes M)) \\ &\xrightarrow{s_{X,Y \otimes M}} X \otimes F(Y \otimes M) \\ &\xrightarrow{\text{id}_X \otimes s_{Y,M}} X \otimes (Y \otimes F(M)) \\ &= (X \otimes Y) \otimes F(M) \\ &\xleftarrow{s_{X \otimes Y, M}} F((X \otimes Y) \otimes M) \end{aligned}$$

commutes.

6.3 Module category over a multitensor category

- A **module category** over a multitensor category $\mathcal{C}$ is a locally finite abelian category $\mathcal{M}$ over a field k , with a $\mathcal{C}$ -module category structure where the action functor

$$\otimes : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$$

is bilinear and exact in the first variable.

- Let $\text{Lex}(\mathcal{M}, \mathcal{M})$ denote the category of left exact endofunctors of $\mathcal{M}$. Then there is a bijection between $\mathcal{C}$ -module structures on $\mathcal{M}$ and tensor functors

$$F : \mathcal{C} \rightarrow \text{Lex}(\mathcal{M}, \mathcal{M})$$

- The direct sum $\mathcal{M}_1 \oplus \mathcal{M}_2$ of two $\mathcal{C}$ -module categories is again a $\mathcal{C}$ -module category.
- A $\mathcal{C}$ -module category $\mathcal{M}$ is *indecomposable* if it is not equivalent to a nontrivial direct sum of module categories.

Example: If $\mathcal{C}$ is a multitensor category, then $\mathcal{C}$ is a $\mathcal{C}^{env} := \mathcal{C} \boxtimes \mathcal{C}^{\text{op}}$ -module category with

$$(X \boxtimes Y) \otimes Z := X \otimes Z \otimes Y.$$

6.4 Finite module categories

TO BE FINISHED

6.5 Balanced Deligne product

TO BE FINISHED

6.6 Relative Serre functors

TO BE FINISHED

6.7 Pivotal module categories

TO BE FINISHED

6.8 Finite categories

Following [?], a *finite category* is an abelian category $\mathcal{A}$ such that

- (i) the hom-vector spaces are finite-dimensional,
- (ii) every object has finite length,
- (iii) there are finitely many isomorphism classes of simple objects,
- (iv) $\mathcal{A}$ has enough projectives.

It is well-known that an abelian category is finite if and only if it is equivalent to the category $A\text{-mod}$ of finite-dimensional modules over a finite-dimensional algebra A . Therefore, a given finite category determines the Morita equivalence class of a finite-dimensional algebra.

Recall that abelian categories are *finitely (co)complete*, that is, they have all finite (co)limits. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between abelian categories is *left* (resp. *right*) *exact* if it preserves finite limits (resp. colimits). Alternatively, F is left (resp. right) exact if and only if it preserves kernels (resp. cokernels). The functor F is *exact* if it is left and right exact.

The following version of the Eilenberg-Watts theorem will be heavily used in the sequel:

Lemma 6.1 ([?, Corollary 2.3]). *Let $F, G : \mathcal{A} \rightarrow \mathcal{B}$ be functors between finite categories. Then the following are equivalent:*

- (L1) F is left exact,
- (L2) F is right-adjoint,

and if $\mathcal{B} = \mathbf{vect}$ the category of finite-dimensional vector spaces, then they are likewise equivalent to

- (L3) F is representable, $F \cong \mathrm{Hom}_{\mathcal{A}}(X, -)$ for some object $X \in \mathcal{A}$.

Similarly, the following are equivalent:

- (R1) G is right exact,
- (R2) G is left-adjoint,

and if $\mathcal{B} = \mathbf{vect}$, then they are equivalent to

- (R3) G is “dually representable”, $G \cong \mathrm{Hom}_{\mathcal{A}}(-, X)^*$ for some object $X \in \mathcal{A}$.

6.9 Finite tensor categories

A finite category $\mathcal{A}$ equipped with a rigid bilinear monoidal product with simple unit is called a *finite tensor category* [?]. The left and right rigidity induce strong monoidal functors

$$(-)^\vee : \mathcal{A} \rightarrow \mathcal{A}^{\text{op,rev}} \quad , \quad {}^\vee(-) : \mathcal{A}^{\text{op,rev}} \rightarrow \mathcal{A} \quad (49)$$

respectively, which are mutually quasi-inverses. Here for a monoidal category $\mathcal{C} = (\mathcal{C}, \otimes, \mathbb{1})$, we have put

$$\mathcal{C}^{\text{op}} = (\mathcal{C}^{\text{op}}, \otimes, \mathbb{1}) \quad , \quad \mathcal{C}^{\text{rev}} = (\mathcal{C}, \otimes^{\text{rev}}, \mathbb{1}) \quad , \quad \mathcal{C}^{\text{op,rev}} = (\mathcal{C}^{\text{op}})^{\text{rev}}$$

with $X \otimes^{\text{rev}} Y := Y \otimes X$.

In a finite tensor category $\mathcal{A}$, we have adjunctions (e.g. [?, §2.7])

$$X^\vee \otimes (-) \dashv X \otimes (-) \dashv {}^\vee X \otimes (-) \quad , \quad (-) \otimes {}^\vee X \dashv (-) \otimes X \dashv (-) \otimes X^\vee \quad (50)$$

for every object $X \in \mathcal{A}$, so by Theorem 6.1 the monoidal product bifunctor $\otimes : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ is exact in each variable.

6.10 Finite module categories

Let $\mathcal{C}$ be a monoidal category. A *left $\mathcal{C}$ -module structure* on a category $\mathcal{M}$ is the data of an action bifunctor $\otimes : \mathcal{C} \times \mathcal{M} \rightarrow \mathcal{M}$ and a family of natural isomorphisms

$$\mathbb{1} \otimes M \cong M \quad , \quad (X \otimes Y) \otimes M \cong X \otimes (Y \otimes M)$$

satisfying a pentagon and triangle axioms similar to those for monoidal categories, see [?, §7.1] for details (a *right $\mathcal{A}$ -module category* is defined analogously with an action functor $\otimes : \mathcal{M} \times \mathcal{C} \rightarrow \mathcal{M}$). Similarly, an *$\mathcal{A}$ -module functor* $F : \mathcal{M} \rightarrow \mathcal{N}$ between $\mathcal{A}$ -module categories is a functor together with natural isomorphisms

$$F(X \otimes M) \cong F(X) \otimes M$$

satisfying similar axioms, cf. [?, §7.2].

If $\mathcal{A}$ is a finite tensor category, a finite category $\mathcal{M}$ equipped with a $\mathcal{A}$ -module structure is called a *finite $\mathcal{A}$ -module category* if the functors $- \otimes M : \mathcal{A} \rightarrow \mathcal{M}$ are right exact for any $M \in \mathcal{M}$. Similarly to the case of finite tensor categories, we have adjunctions

$$X^\vee \otimes (-) \dashv X \otimes (-) \dashv {}^\vee X \otimes (-) \quad (51)$$

so again Theorem 6.1 implies that the functor $X \otimes - : \mathcal{M} \rightarrow \mathcal{M}$ is always exact for any $X \in \mathcal{A}$.

6.11 Internal homs

Let $\mathcal{A}$ be a finite tensor category and $\mathcal{M}$ a finite $\mathcal{A}$ -module category. For every $M \in \mathcal{M}$, the functor $- \otimes M$ has a right-adjoint by Theorem 6.1, that we denote by $\underline{\text{Hom}}(M, -) : \mathcal{M} \rightarrow \mathcal{A}$. By definition, there are linear isomorphisms

$$\text{Hom}_{\mathcal{M}}(X \otimes M, N) \cong \text{Hom}_{\mathcal{A}}(X, \underline{\text{Hom}}(M, N)) \quad (52)$$

natural in $X \in \mathcal{A}$ and $N \in \mathcal{M}$. According to the parametrized adjunction theorem [?, Theorem IV.7.3], the right adjoints in this family of adjunctions assemble in a unique way into a bifunctor

$$\underline{\text{Hom}} : \mathcal{M}^{\text{op}} \times \mathcal{M} \rightarrow \mathcal{A} \quad (53)$$

such that the isomorphism (52) is natural in the three variables. We call (53) the *internal hom functor*. This bifunctor is, just like the hom functor, left exact in both variables: indeed by definition $\underline{\text{Hom}}(M, -)$ is right-adjoint to $- \otimes M$, and the isomorphisms

$$\text{Hom}_{\mathcal{M}^{\text{op}}}(X^{\vee} \otimes N, M) \stackrel{(51)}{\cong} \text{Hom}_{\mathcal{M}^{\text{op}}}(N, X \otimes M) \cong \text{Hom}_{\mathcal{M}}(X \otimes M, N) \cong \text{Hom}_{\mathcal{A}}(X, \underline{\text{Hom}}(M, N))$$

show that $\underline{\text{Hom}}(-, N)$ is right-adjoint to $(-)^{\vee} \otimes N$. Once again we conclude by Theorem 6.1.

We also record that for objects $X, Y \in \mathcal{A}$ and $M, N \in \mathcal{M}$ there is a natural isomorphism

$$\underline{\text{Hom}}(X \otimes M, Y \otimes N) \cong Y \otimes \underline{\text{Hom}}(M, N) \otimes X^{\vee}, \quad (54)$$

see [?, Lemma 7.9.4] or [?, (2.15)].

The aforementioned internal homs can be used to equip $\mathcal{M}$ with a structure of $\mathcal{A}$ -enriched category: for $L, M, N \in \mathcal{M}$ there is a composition map

$$\circ_{L,M,N} := \underline{\text{ev}}_{M,N} \circ (\text{id} \otimes \underline{\text{ev}}_{L,M}) : \underline{\text{Hom}}(M, N) \otimes \underline{\text{Hom}}(L, M) \longrightarrow \underline{\text{Hom}}(L, N) \quad (55)$$

where $\underline{\text{ev}}_{M,N} : \underline{\text{Hom}}(M, N) \otimes M \longrightarrow N$ is the counit of the adjunction (52). The identity arrow

$$\text{id}_M : \mathbb{1} \longrightarrow \underline{\text{End}}(M) := \underline{\text{Hom}}(M, M)$$

is the morphism corresponding to the isomorphism $\mathbb{1} \otimes M \cong M$ under the isomorphism (52). In particular, for any object $M \in \mathcal{M}$, we have that $\underline{\text{End}}(M)$ is an algebra object in $\mathcal{A}$, and for any $N \in \mathcal{M}$ the object $\underline{\text{Hom}}(M, N)$ is a right $\underline{\text{End}}(M)$ -module in $\mathcal{A}$ with structure morphism given by (55) with $L = M$.

We denote by $\text{mod}_{\mathcal{A}\text{-}\underline{\text{End}}(M)}$ the category of right $\underline{\text{End}}(M)$ -modules in $\mathcal{A}$. It is readily verified that the monoidal product $\otimes : \mathcal{A} \times \mathcal{A} \longrightarrow \mathcal{A}$ restricts to a functor $\mathcal{A} \times \text{mod}_{\mathcal{A}\text{-}\underline{\text{End}}(M)} \longrightarrow \text{mod}_{\mathcal{A}\text{-}\underline{\text{End}}(M)}$ turning it into a left $\mathcal{A}$ -module category. In particular, $\underline{\text{Hom}}(M, -)$ restricts to a functor

$$\underline{\text{Hom}}(M, -) : \mathcal{M} \longrightarrow \text{mod}_{\mathcal{A}\text{-}\underline{\text{End}}(M)}$$

for which the isomorphisms (54) make it an $\mathcal{A}$ -module functor.

Theorem 6.2 (Monadicity for module categories, e.g. [?, §7.10]). *Let $M \in \mathcal{M}$. Then the functor*

$$\underline{\text{Hom}}(M, -) : \mathcal{M} \longrightarrow \text{mod}_{\mathcal{A}\text{-}\underline{\text{End}}(M)}$$

is an equivalence of left $\mathcal{A}$ -module categories if and only if M is an $\mathcal{A}$ -progenerator. Moreover, such an object always exists.

In the statement above, an object $M \in \mathcal{M}$ is an $\mathcal{A}$ -progenerator if the functor $\underline{\text{Hom}}(M, -) : \mathcal{M} \longrightarrow \mathcal{A}$ is exact and faithful. When the functor $\underline{\text{Hom}}(M, -)$ is only exact (resp. faithful), then we say that $M \in \mathcal{M}$ is $\mathcal{A}$ -projective (resp. $\mathcal{A}$ -generator).

6.12 Deligne product

If $\mathcal{C}$ and $\mathcal{D}$ are abelian categories, let us denote $\text{Lex}(\mathcal{C}, \mathcal{D})$ (resp. $\text{Rex}(\mathcal{C}, \mathcal{D})$) the category of left (resp. right) exact functors $\mathcal{C} \longrightarrow \mathcal{D}$ and natural transformations between them. Because (co)limits in product categories are computed componentwise, note that if $\mathcal{C}'$ is another abelian category, then $\text{Lex}(\mathcal{C} \times \mathcal{C}', \mathcal{D})$ consists precisely of functors $\mathcal{C} \times \mathcal{C}' \longrightarrow \mathcal{D}$ that are left exact in each variable (and the same observation applies to right exact functors)

Now suppose that $\mathcal{A}, \mathcal{B}$ are finite categories. Its *Deligne product* [?, §5] is another such category $\mathcal{A} \boxtimes \mathcal{B}$ together with a functor

$$\boxtimes : \mathcal{A} \times \mathcal{B} \longrightarrow \mathcal{A} \boxtimes \mathcal{B} \quad (56)$$

which is right exact and is universal with respect to this property, in the sense that for any other finite category $\mathcal{E}$ the functor

$$\text{Rex}(\mathcal{A} \boxtimes \mathcal{B}, \mathcal{E}) \xrightarrow{\cong} \text{Rex}(\mathcal{A} \times \mathcal{B}, \mathcal{E}) \quad , \quad F \longmapsto F \circ \boxtimes$$

is an equivalence of categories. It is well-known that the Deligne product of finite categories exists and is of such kind, and moreover that it coincides with the Kelly product of finitely cocomplete categories in this case [?].

One can show that the functor (56) is in fact exact. Moreover, by the universal property of the tensor product of vector spaces, this (bilinear) functor induces a linear map

$$\text{Hom}_{\mathcal{A}}(X, Y) \otimes_{\mathbb{k}} \text{Hom}_{\mathcal{B}}(Y, Y') \xrightarrow{\cong} \text{Hom}_{\mathcal{A} \boxtimes \mathcal{B}}(X \boxtimes X', Y \boxtimes Y') \quad (57)$$

which is in fact a linear isomorphism for all $X, Y \in \mathcal{A}$ and $X', Y' \in \mathcal{B}$ [?, §1.11].

Proposition 6.3 ([?, Proposition 5.17]). *Let $\mathcal{A}, \mathcal{B}$ be finite tensor categories. Then their Deligne product $\mathcal{A} \boxtimes \mathcal{B}$ is again a finite tensor category with monoidal product determined by*

$$(X \boxtimes Y) \otimes (X' \boxtimes Y') = (X \otimes X') \boxtimes (Y \otimes Y'),$$

monoidal unit $\mathbb{1} \boxtimes \mathbb{1}$ and rigidity determined by $(X \boxtimes Y)^\vee = X^\vee \boxtimes Y^\vee$.

6.13 Balanced Deligne product

Let $\mathcal{A}$ be a finite tensor category and let $\mathcal{M}$ and $\mathcal{N}$ be finite right and left $\mathcal{A}$ -module categories, respectively. An *$\mathcal{A}$ -balanced functor* from $\mathcal{M} \times \mathcal{N}$ to a finite category $\mathcal{E}$ is a right exact functor $F : \mathcal{M} \times \mathcal{N} \longrightarrow \mathcal{E}$ together with a family of natural isomorphisms

$$F(M \otimes X, N) \cong F(M, X \otimes N)$$

satisfying the corresponding pentagon and triangle axioms. We denote by $\text{Bal}_{\mathcal{A}}(\mathcal{M} \times \mathcal{N}, \mathcal{E})$ the category of right exact $\mathcal{A}$ -balanced functors $\mathcal{M} \times \mathcal{N} \longrightarrow \mathcal{E}$ and natural transformations compatible with these natural isomorphisms.

If $\mathcal{A}, \mathcal{M}$ and $\mathcal{N}$ are as above, their *balanced Deligne product* is a finite category $\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}$ together with a functor

$$\boxtimes_{\mathcal{A}} : \mathcal{M} \times \mathcal{N} \longrightarrow \mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N} \quad (58)$$

which is right exact $\mathcal{A}$ -balanced and is universal with respect to this property, in the sense that for any finite category $\mathcal{E}$ the functor

$$\text{Rex}(\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}, \mathcal{E}) \xrightarrow{\cong} \text{Bal}_{\mathcal{A}}(\mathcal{M} \times \mathcal{N}, \mathcal{E}) \quad , \quad F \longmapsto F \circ \boxtimes_{\mathcal{A}}$$

is an equivalence of categories. The balanced product appeared first in [?] in the context of fusion categories. In the current setup, it is shown in [?, Theorem 3.3] that the balanced Deligne product always exists, that the functor (58) is in fact exact and moreover that we have the following linear isomorphism:

$$\text{Hom}_{\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}}(M \boxtimes_{\mathcal{A}} N, M' \boxtimes_{\mathcal{A}} N') \cong \text{Hom}_{\mathcal{A}}(\mathbb{1}, \underline{\text{Hom}}(M, M') \otimes \underline{\text{Hom}}(N, N')). \quad (59)$$

It will be useful to have a concrete realization of the balanced Deligne product. For $\mathcal{M}$ and $\mathcal{N}$ finite right and left $\mathcal{A}$ -module categories, Theorem 6.2 gives us (endomorphism) algebras objects $A, B \in \mathcal{A}$ together with equivalences $\mathcal{M} \simeq A\text{-mod}_{\mathcal{A}}$ and $\mathcal{N} \simeq \text{mod}_{\mathcal{A}-B}$. Then according to [?, Theorem 3.3], the balanced Deligned product $\mathcal{M} \boxtimes_{\mathcal{A}} \mathcal{N}$ can be realised as the category of A - B -bimodules in $\mathcal{A}$; more precisely, we have that the functor

$$\otimes : A\text{-mod}_{\mathcal{A}} \boxtimes_{\mathcal{A}} \text{mod}_{\mathcal{A}-B} \xrightarrow{\simeq} A\text{-mod}_{\mathcal{A}-B} \quad , \quad M \boxtimes N \mapsto M \otimes N \quad (60)$$

is an equivalence.

6.14 The distinguished invertible element

Let $\mathcal{A}$ be a finite tensor category. An object $X \in \mathcal{A}$ is said to be *invertible* if the evaluation $\text{ev}_X : X^\vee \otimes X \longrightarrow \mathbb{1}$ and the coevaluation $\text{coev}_X : \mathbb{1} \longrightarrow X \otimes X^\vee$ are isomorphisms. In that case, $X^\vee$ is often denoted as X^{-1} . The goal of this subsection is to introduce the so-called distinguished invertible element of a finite tensor category.

If $\mathcal{A}$ is a finite tensor category, its *envelope* is $\mathcal{A}^{\text{env}} := \mathcal{A} \boxtimes \mathcal{A}^{\text{rev}}$, which is a finite tensor category by Theorem 6.3. Note that because of (49), this category is in fact equivalent to $\mathcal{A} \boxtimes \mathcal{A}^{\text{op}}$. The category $\mathcal{A}$ can be viewed as a left $\mathcal{A}^{\text{env}}$ -module category with action functor determined by

$$(X \boxtimes Y) \otimes Z = X \otimes Z \otimes Y.$$

By the discussion of Section 6.11, we have that $\Delta := \underline{\text{End}}(\mathbb{1}) \in \mathcal{A}^{\text{env}}$ is an algebra object, that is usually called the *canonical algebra* of $\mathcal{A}$. By a result of Shimizu [?, Lemma 4.4], the canonical algebra can actually be expressed as the coend

$$\Delta \cong \int^{X \in \mathcal{A}} X^\vee \boxtimes X. \quad (61)$$

The monoidal unit $\mathbb{1} \in \mathcal{A}$ is well-known to be an $\mathcal{A}^{\text{env}}$ -progenerator: indeed by (54) we have $\underline{\text{Hom}}(\mathbb{1}, -) \cong (- \boxtimes \mathbb{1}) \otimes \underline{\text{End}}(\mathbb{1})$, which is right exact; and since $\underline{\text{End}}(\mathbb{1}) \neq 0$, the argument from [?, Lemma 6.11] tells us that $\underline{\text{Hom}}(\mathbb{1}, -)$ is also faithful. Therefore by Theorem 6.2 the functor

$$\underline{\text{Hom}}(\mathbb{1}, -) : \mathcal{A} \xrightarrow{\simeq} \text{mod}_{\mathcal{A}^{\text{env}}-\Delta}$$

is an equivalence of $\mathcal{A}^{\text{env}}$ -module categories. Furthermore (54) implies that this functor is naturally isomorphic to the functor $(- \boxtimes \mathbb{1}) \otimes \Delta_\Delta$, where Δ_Δ denotes Δ viewed as a right Δ -module by the multiplication of Δ . By means of the equivalence, there exists an object α , unique up to isomorphism, such that

$$(\alpha \boxtimes \mathbb{1}) \otimes \Delta_\Delta \cong (\Delta \Delta)^\vee. \quad (62)$$

It can be shown that this object α is in fact invertible [?, §7.18] (see also [?, Lemma 4.3]), and will be called the *distinguished invertible object* of $\mathcal{A}$. When $\alpha \cong \mathbb{1}$, we will say that $\mathcal{A}$ is *unimodular*. This name comes from the theory of Hopf algebras: a finite-dimensional Hopf algebra H is unimodular (i.e., left- and right-integrals coincide) if and only if the category $H\text{-mod}_{\mathbb{k}}$ is unimodular. We refer the reader to [?, Theorem 4.10] for a list of different characterizations of unimodularity. Note that when $\mathcal{A}$ is unimodular, (62) implies the self-duality of Δ .

One of the main properties of the distinguished invertible object is that it implements the quadruple dual by conjugation [?, Theorem 3.3]: we have a natural isomorphism of monoidal functors

$$(-)^{\vee\vee\vee\vee} \cong \alpha \otimes - \otimes \alpha^{-1}. \quad (63)$$

In particular, if $\mathcal{A}$ is unimodular, then $(-)^{\vee\vee\vee\vee} \cong \text{id}_{\mathcal{A}}$.

6.15 stuff

In the sequel it will be relevant to obtain a trivialization of the right Nakayama functor. To this end, recall from [?] that given a finite category $\mathcal{A}$ and an endofunctor $F : \mathcal{A} \rightarrow \mathcal{A}$, a *right F -twisted Calabi-Yau structure* on $\mathcal{A}$ is a family of natural isomorphisms

$$\mathrm{Hom}_{\mathcal{A}}(X, Y) \cong \mathrm{Hom}_{\mathcal{A}}(Y, F(X))^*.$$

If $F = \mathrm{id}_{\mathcal{A}}$ we will refer to it as a *Calabi-Yau structure* on $\mathcal{A}$. By a *(right F -twisted) projective Calabi-Yau structure* on $\mathcal{A}$ we will mean a (right F -twisted) Calabi-Yau structure on $\mathrm{Proj} \mathcal{A}$, the subcategory of projective objects of $\mathcal{A}$.

Lemma 6.4. *Let $\mathcal{A}$ be a finite category. Then $\mathcal{A}$ admits a projective Calabi-Yau structure if and only if the right Nakayama functor admits a trivialization $N^r \cong \mathrm{id}_{\mathcal{A}}$.*

Proof. By [?, Corollary 2.3], any finite category $\mathcal{A}$ is a right N^r -twisted projective Calabi-Yau category. Because $\mathcal{A}$ admits a projective Calabi-Yau structure, by the (enriched) Yoneda lemma and the fact that the hom-spaces are finite-dimensional, we obtain a natural isomorphism $N^r|_{\mathrm{Proj} \mathcal{A}} \cong \mathrm{id}_{\mathrm{Proj} \mathcal{A}}$. This trivialization lifts to a trivialization $N^r \cong \mathrm{id}_{\mathcal{A}}$ as both functors are right-exact and $\mathcal{A}$ has enough projectives. This demonstrates the necessity and the sufficiency follows directly from [?, Corollary 2.3]. $\square$

We say that a finite category $\mathcal{A}$ is *Frobenius* (resp. *symmetric Frobenius*) if $\mathcal{A}$ is equivalent to the category $A\text{-mod}_{\mathbb{k}}$ of finite-dimensional left A -modules for a Frobenius (resp. symmetric Frobenius) algebra A . Frobenius categories are also called quasi-Frobenius [?] or self-injective [?], and can be equivalently defined as those finite categories for which the classes of injective and projective objects coincide. It turns out that the Nakayama functors can be used to characterize Frobenius and symmetric Frobenius categories:

Proposition 6.5 ([?, Proposition 3.24]). *Let $\mathcal{A}$ be a finite category. Then*

1. *$\mathcal{A}$ is Frobenius if and only if the left and right Nakayama functors are equivalences. In this case, they are quasi-inverses to each other.*
2. *$\mathcal{A}$ is symmetric Frobenius if and only if the left and right Nakayama functors are isomorphic to the identity functor.*

Lastly, let us point out that if $\mathcal{M}$ is a finite left module category over a finite tensor category $\mathcal{A}$, then the left and right Nakayama functors of $\mathcal{M}$ have a natural structure of twisted $\mathcal{A}$ -module functors, in the sense that there are natural isomorphisms

$$N^\ell(X \otimes M) \cong X^{\vee\vee} \otimes N^\ell(M) \quad , \quad N^r(X \otimes M) \cong {}^{\vee\vee}X \otimes N^r(M), \quad (64)$$

see [?, Theorem 4.4] or [?, §2.5].

6.16 Relative Serre functors

Let $\mathcal{A}$ be a finite tensor category and let $\mathcal{M}$ be a finite left $\mathcal{A}$ -module category. A *left relative Serre functor* of $\mathcal{M}$ is an endofunctor $S^\ell : \mathcal{M} \rightarrow \mathcal{M}$ together with a family of natural isomorphisms in $\mathcal{A}$

$$\underline{\mathrm{Hom}}(S^\ell(M), N) \cong {}^\vee \underline{\mathrm{Hom}}(N, M). \quad (65)$$

Similarly, a *right relative Serre functor* of $\mathcal{M}$ is an endofunctor $S^\ell : \mathcal{M} \rightarrow \mathcal{M}$ together with a family of natural isomorphisms

$$\underline{\mathrm{Hom}}(M, S^r(N)) \cong \underline{\mathrm{Hom}}(N, M)^\vee. \quad (66)$$

If they exist, then they are essentially unique by the (enriched) Yoneda lemma. It is shown in [?, Proposition 4.24] that a finite left $\mathcal{A}$ -module category $\mathcal{M}$ has left and right relative Serre functors if and only if $\mathcal{M}$ is *exact*, that is, a finite left $\mathcal{A}$ -module category where all objects are $\mathcal{A}$ -projective. It is readily verified that in this case the left and right relative Serre functors are quasi-inverses to each other.

Example 6.6. Let $\mathcal{A}$ be a finite tensor category viewed as a finite left module category over itself. Then $S^\ell = {}^{\vee\vee}(-)$ and $S^r = (-)^{\vee\vee}$ are readily seen to be the left and right relative Serre functors (in particular $\mathcal{A}$ is exact). In particular, this says that these functors can be understood as the analogs of the double left and right dual functors in module categories.

Furthermore, according to [?, §3.3], a left (resp. right) relative Serre functor of $\mathcal{M}$ has a unique structure of twisted $\mathcal{A}$ -module functor such that the natural isomorphisms (65) (resp. (66)) are isomorphisms of $\mathcal{A}$ -bimodule functors. This means that there are natural isomorphisms (compare with (64))

$$S^\ell(X \otimes M) \cong {}^{\vee\vee}X \otimes S^\ell(M) \quad , \quad S^r(X \otimes M) \cong X^{\vee\vee} \otimes S^r(M), \quad (67)$$

see also [?, Lemma 4.23].

The relative Serre functors are in fact very closely related to the Nakayama functors:

Theorem 6.7 ([?, Theorem 4.26]). *Let $\mathcal{A}$ be a finite tensor category with distinguished invertible object α and let $\mathcal{M}$ be an exact left $\mathcal{A}$ -module category. Then we have natural isomorphisms*

$$N^\ell \cong \alpha \otimes S^\ell \quad , \quad N^r \cong \alpha^{-1} \otimes S^r$$

of twisted module functors.

Corollary 6.8. *Let $\mathcal{A}$ be a unimodular pivotal finite tensor category with a fixed trivialization $\alpha \cong \mathbb{1}$ of the distinguished invertible object, and let $\mathcal{M}$ be an exact left $\mathcal{A}$ -module category. Then the isomorphisms above induce canonical isomorphisms*

$$N^\ell \cong S^\ell \quad , \quad N^r \cong S^r$$

of untwisted module functors, with the untwisting induced by the pivotal structure.

Corollary 6.9. *For any finite tensor category $\mathcal{A}$, we have natural isomorphisms*

$$N^\ell \cong \alpha \otimes {}^{\vee\vee}(-) \quad , \quad N^r \cong \alpha^{-1} \otimes (-)^{\vee\vee}$$

From this last result we easily obtain two important properties about the distinguished invertible object: if $\mathcal{A}$ and $\mathcal{B}$ are finite tensor categories, the distinguished invertible object $\alpha_{\mathcal{A} \boxtimes \mathcal{B}}$ of $\mathcal{A} \boxtimes \mathcal{B}$ (which is a finite tensor category by Theorem 6.3) and that of $\mathcal{A}^{\mathrm{rev}}$ are given by

$$\alpha_{\mathcal{A} \boxtimes \mathcal{B}} \cong \alpha_{\mathcal{A}} \boxtimes \alpha_{\mathcal{B}} \quad , \quad \alpha_{\mathcal{A}^{\mathrm{rev}}} \cong \alpha_{\mathcal{A}}, \quad (68)$$

where $\alpha_{\mathcal{A}}$, $\alpha_{\mathcal{B}}$ and $\alpha_{\mathcal{A}^{\mathrm{rev}}}$ denote the distinguished invertible objects of $\mathcal{A}$, $\mathcal{B}$ and $\mathcal{A}^{\mathrm{rev}}$, respectively.

6.17 Pivotal module categories

Let $\mathcal{A}$ be a pivotal finite tensor category with pivotal structure $\omega : \text{id}_{\mathcal{A}} \xrightarrow{\cong} (-)^{\vee\vee}$, let $\mathcal{M}$ be an exact left $\mathcal{A}$ -module category and let S^r be a right relative Serre functor of $\mathcal{M}$. A *pivotal structure* for $\mathcal{M}$ is a natural isomorphism $\tilde{\omega} : \text{id}_{\mathcal{M}} \xrightarrow{\cong} S^r$ which is compatible with the pivotal structure of $\mathcal{A}$ in the sense that the diagram

$$\begin{array}{ccc} X \otimes M & \xrightarrow{\tilde{\omega}_{X \otimes M}} & S^r(X \otimes M) \\ & \searrow \omega_X \otimes \tilde{\omega}_M \quad \nearrow \xi_{X,M} & \\ & X^{\vee\vee} \otimes S^r(M) & \end{array} \quad (69)$$

commutes for all $X \in \mathcal{A}$ and $M \in \mathcal{M}$, where $\xi_{X,M}$ is the isomorphism from (67). An exact left $\mathcal{A}$ -module category $\mathcal{M}$ endowed with a pivotal structure will be called a *pivotal left $\mathcal{A}$ -module category*.

Corollary 6.10. *Let $\mathcal{A}$ be a unimodular pivotal finite tensor category with a fixed trivialization $\alpha \cong \mathbb{1}$ of the distinguished invertible object, and let $\mathcal{M}$ be an exact left $\mathcal{A}$ -module category. Then the isomorphisms above induce canonical isomorphisms*

$$N^\ell \cong S^\ell \quad , \quad N^r \cong S^r$$

of untwisted module functors, with the untwisting induced by the pivotal structure. A pivotal module structure on $\mathcal{M}$ is a trivialization of N^ℓ or N^r as module functor.

Example 6.11. Let $\mathcal{A}$ be a pivotal finite tensor category, viewed as a left $\mathcal{A}$ -module category (by left multiplication). By Theorem 6.6, $\mathcal{A}$ is exact and the left and right relative Serre functors can be taken to be the identity endofunctor, and the isomorphism $\xi_{X,Y}$ can be seen to be induced by the (inverse of the) pivotal structure. Then the identity natural transformation of the identity functor makes $\mathcal{A}$ a pivotal module category over itself (the commutativity of (69) is trivial in this case).

Next we will generalize the previous example. Before that let us introduce the following notation: if $\mathcal{A}$ is a finite tensor category and $p, q \geq 0$ are non-negative integers, let us denote by ${}_p\mathcal{A}_q$ the finite $\mathcal{A}^{\boxtimes p} \boxtimes (\mathcal{A}^{\text{rev}})^{\boxtimes q}$ -module category whose underlying finite category is $\mathcal{A}$ with module structure determined by

$$(X_1 \boxtimes \cdots \boxtimes X_p \boxtimes Y_1 \boxtimes \cdots \boxtimes Y_q) \otimes Z = X_1 \otimes \cdots \otimes X_p \otimes Z \otimes Y_q \otimes \cdots \otimes Y_1.$$

Lemma 6.12. *Let $\mathcal{A}$ be a unimodular pivotal finite tensor category with a fixed trivialization $\alpha \cong \mathbb{1}$ of the distinguished invertible object of $\mathcal{A}$. Then ${}_p\mathcal{A}_q$ inherits a structure of pivotal module category for every $p, q \geq 0$.*

Proof. By [?, Lemma 3.2], ${}_p\mathcal{A}_q$ is exact if for any projective object $P \in \mathcal{A}^{\boxtimes p} \boxtimes (\mathcal{A}^{\text{rev}})^{\boxtimes q}$ we have that $P \otimes X$ is projective for any $X \in {}_p\mathcal{A}_q$. By the properties of the Deligne product, P is a direct summand of a projective object W of the form $W = \bigoplus_i A_i \boxtimes B_i$ with A_i, B_i projective. Because in a finite tensor category, the monoidal product of a projective object and an arbitrary object is projective, $W \otimes X$ is projective, and since $- \otimes X$ is right exact, $P \otimes X$ is a direct summand of $W \otimes X$, hence again projective.

Next let us see how we can obtain a canonical trivialization of the right relative Serre functor of ${}_p\mathcal{A}_q$. On the one hand, the fixed trivialization of the distinguished invertible object and the

pivotal structure induce a trivialization of the right Nakayama functor of $\mathcal{A}$ (the underlying finite tensor category of ${}_p\mathcal{A}_q$) by Theorem 6.9. On the other hand, the distinguished invertible object of $\mathcal{A}^{\boxtimes p} \boxtimes (\mathcal{A}^{\text{rev}})^{\boxtimes q}$ is according to (68)

$$\alpha_{\mathcal{A}^{\boxtimes p} \boxtimes (\mathcal{A}^{\text{rev}})^{\boxtimes q}} \cong \alpha^{\boxtimes p+q},$$

which implies that $\mathcal{A}^{\boxtimes p} \boxtimes (\mathcal{A}^{\text{rev}})^{\boxtimes q}$ is unimodular as well with a trivialization induced by that of $\mathcal{A}$. Moreover the pivotal structure of $\mathcal{A}$ induces a canonical pivotal structure on $\mathcal{A}^{\boxtimes p} \boxtimes (\mathcal{A}^{\text{rev}})^{\boxtimes q}$. Therefore we obtain a trivialization $\text{id}_{\mathcal{A}} \cong \mathbf{N}^r \cong \mathbf{S}^r$ of the right relative Serre functor by Theorem 6.10. The commutativity of (69) holds by construction. $\square$

As it turns out, pivotal module categories can be characterized as categories of modules over symmetric Frobenius algebras:

Theorem 6.13 ([?, Corollary 3.16]). *If $\mathcal{M}$ is a pivotal left $\mathcal{A}$ -module category, then there is an equivalence of left $\mathcal{A}$ -module categories $\mathcal{M} \simeq \text{mod}_{\mathcal{A}}\text{-}A$ for a symmetric Frobenius algebra object A in $\mathcal{A}$.*

6.18 Bimodules and the hom

Setup. Let A and B be finite-dimensional algebras over a field k . Let

$$M, M' \in A\text{-Mod}, \quad N, N' \in \text{Mod-}B,$$

where M, M' are left A -modules and N, N' are right B -modules.

Then $M \otimes_k N$ and $M' \otimes_k N'$ carry natural A - B -bimodule structures via

$$a \cdot (m \otimes n) \cdot b := (am) \otimes (nb).$$

Claim (tensor–Hom adjunction for bimodules). There is a canonical isomorphism

$$\text{Hom}_{A\text{-}B}(M \otimes N, M' \otimes N') \cong \text{Hom}_A(M, \text{Hom}_B(N, M' \otimes N')).$$

Explanation. We first apply the usual tensor–Hom adjunction in Vect_k :

$$\text{Hom}_k(M \otimes N, M' \otimes N') \cong \text{Hom}_k(M, \text{Hom}_k(N, M' \otimes N')).$$

We then impose equivariance:

- A -linearity in the first variable replaces $\text{Hom}_k(M, -)$ by $\text{Hom}_A(M, -)$.
- B -linearity in the second variable replaces $\text{Hom}_k(N, -)$ by $\text{Hom}_B(N, -)$.

Also

$$\text{Hom}_B(N, M' \otimes N')$$

carries a natural structure of left A -module: For $f \in \text{Hom}_B(N, M' \otimes N')$ and $a \in A$, define

$$(a \cdot f)(n) := a \cdot f(n),$$

where the action on the right-hand side is the A -action on $M' \otimes N'$.

This yields the canonical identification

$$\boxed{\text{Hom}_{A\text{-}B}(M \otimes N, M' \otimes N') \cong \text{Hom}_A(M, \text{Hom}_B(N, M' \otimes N'))}.$$

Remark. There is a symmetric form obtained by currying in the other variable:

$$\mathrm{Hom}_{A-B}(M \otimes N, M' \otimes N') \cong \mathrm{Hom}_B(N, \mathrm{Hom}_A(M, M' \otimes N')).$$

If A, B are algebras in a finite tensor category $\mathcal{A}$, and M, M', N, N' are A/B modules in $\mathcal{A}$, then the formula becomes

$$\boxed{\mathrm{Hom}_{A-B}(M \otimes N, M' \otimes N') \cong \mathrm{Hom}_A(M, \underline{\mathrm{Hom}}_B(N, M' \otimes N'))},$$

where $\underline{\mathrm{Hom}}_B(N, M' \otimes N')$ is viewed as a left A -module with action map

$$A \otimes \underline{\mathrm{Hom}}_B(N, M' \otimes N') \cong \underline{\mathrm{Hom}}_B(N, A \otimes M' \otimes N') \rightarrow \underline{\mathrm{Hom}}_B(N, M' \otimes N').$$

7 Cyclic and modular operads

7.1 Classical operads and their algebras

A symmetric operad formalizes operations with multiple inputs and a single, distinguished output. The underlying combinatorics are those of rooted trees.

Let $\mathcal{V}$ be a symmetric monoidal category. A (uncolored) **symmetric** (or classical or ordinary) **operad** $\mathcal{P}$ in $\mathcal{V}$, or a $\mathcal{V}$ -valued operad, is a collection of $\{\mathcal{P}(n)\}_{n \geq 0}$ of objects in $\mathcal{V}$ endowed with the following structures:

1. **A Right Symmetric Group Action:** For each $n \geq 0$, there is a right action of the symmetric group $\mathfrak{S}_n$ on $\mathcal{P}(n)$.
2. **An Identity Element:** There exists a distinguished map $1 : \mathbf{1} \rightarrow \mathcal{P}(1)$.
3. **Composition Maps:** There are arrows in $\mathcal{V}$

$$\mathcal{P}(n) \otimes \bigotimes_{i=1}^n \mathcal{P}(k_i) \rightarrow \mathcal{P}(k_1 + \cdots + k_n),$$

(equivalently, for all integers $n, m \geq 0$ and each $i \in \{1, \dots, n\}$, there are linear maps, known as insertion maps:

$$\circ_i : \mathcal{P}(n) \otimes \mathcal{P}(m) \rightarrow \mathcal{P}(n + m - 1)$$

These structures must satisfy axioms of associativity, unitality, and equivariance with respect to the symmetric group action. For instance, associativity requires that for $p \in \mathcal{P}(n), q \in \mathcal{P}(m), r \in \mathcal{P}(k)$:

$$\begin{aligned} (p \circ_j r) \circ_i q &= (p \circ_i q) \circ_{j+m-1} r \quad \text{for } 1 \leq i < j \leq n \\ p \circ_i (q \circ_j r) &= (p \circ_i q) \circ_{i+j-1} r \quad \text{for } 1 \leq i \leq n, 1 \leq j \leq m \end{aligned}$$

Let $\mathcal{P} = \{\mathcal{P}(n)\}_{n \geq 0}$ and $\mathcal{Q} = \{\mathcal{Q}(n)\}_{n \geq 0}$ be two symmetric operads in the same symmetric monoidal category $\mathcal{V}$ (for instance, the category of vector spaces, $\mathbf{Vect}_{\mathbb{k}}$).

A **morphism of operads** $\Phi : \mathcal{P} \rightarrow \mathcal{Q}$ is a collection of morphisms, one for each arity $n \geq 0$:

$$\Phi_n : \mathcal{P}(n) \rightarrow \mathcal{Q}(n)$$

This collection of maps must satisfy the following three axioms:

1. **Compatibility with Composition:** The morphism must respect the operadic composition maps $\circ_i$. For any $p \in \mathcal{P}(n)$ and $q \in \mathcal{P}(m)$, the following diagram must commute for all $i \in \{1, \dots, n\}$:

$$\begin{array}{ccc} \mathcal{P}(n) \otimes \mathcal{P}(m) & \xrightarrow{\Phi_n \otimes \Phi_m} & \mathcal{Q}(n) \otimes \mathcal{Q}(m) \\ \circ_i \downarrow & & \downarrow \circ_i \\ \mathcal{P}(n+m-1) & \xrightarrow{\Phi_{n+m-1}} & \mathcal{Q}(n+m-1) \end{array}$$

This is equivalent to the equation:

$$\Phi_{n+m-1}(p \circ_i q) = \Phi_n(p) \circ_i \Phi_m(q)$$

2. **Compatibility with the Identity:** The morphism must map the identity element of $\mathcal{P}$ to the identity element of $\mathcal{Q}$.

$$\Phi_1(1_{\mathcal{P}}) = 1_{\mathcal{Q}}$$

3. **Equivariance:** The morphisms must respect the action of the symmetric group S_n . For any $p \in \mathcal{P}(n)$ and any permutation $\sigma \in S_n$:

$$\Phi_n(p \cdot \sigma) = \Phi_n(p) \cdot \sigma$$

This means the maps Φ_n are S_n -equivariant.

Let $\mathcal{P}$ be a symmetric operad. A **$\mathcal{P}$ -algebra** is an object V together with a collection of arrows (structure maps) for each $n \geq 0$:

$$\gamma_n : \mathcal{P}(n) \otimes V^{\otimes n} \rightarrow V$$

These maps must be compatible with the composition, identity, and group action of $\mathcal{P}$. Compatibility with composition, for instance, means that for any $p \in \mathcal{P}(n)$ and $q \in \mathcal{P}(m)$, the action of the composite operation $p \circ_i q$ is the composition of the individual actions. This is equivalent to requiring that the maps $\{\gamma_n\}$ constitute a morphism of operads $\gamma : \mathcal{P} \rightarrow \text{End}_V$, where End_V is the endomorphism operad of V with $\text{End}_V(n) = \text{Hom}(V^{\otimes n}, V)$. More below:

The Endomorphism Operad. Suppose that $\mathcal{V}$ is a **closed** symmetric monoidal category, aka enriched over itself. For any object V in $\mathcal{V}$, one can construct the **endomorphism operad**, denoted End_V .

- Its space of n -ary operations is $\text{End}_V(n) := \text{Hom}(V^{\otimes n}, V)$, the space of all arrows from n copies of V to V .
- Its composition is the standard composition of functions.

Definition of a $\mathcal{P}$ -Algebra. An **algebra over an operad $\mathcal{P}$** (or a **$\mathcal{P}$ -algebra**) on an object V is, by definition, a morphism of operads:

$$\Phi : \mathcal{P} \rightarrow \text{End}_V$$

The data of this morphism is a collection of maps $\Phi_n : \mathcal{P}(n) \rightarrow \text{Hom}(V^{\otimes n}, V)$. This is precisely the structure of a $\mathcal{P}$ -algebra: for each abstract n -ary operation $p \in \mathcal{P}(n)$, the morphism assigns a concrete multilinear map $\Phi_n(p)$ on V . The axioms for an operad morphism then guarantee that this assignment is compatible with composition, the identity, and permutation of inputs, as required for an algebra.

Operads in $\mathcal{V}$ and operad morphisms form a category $\text{Op}(\mathcal{V})$.

A more general definition of a $\mathcal{P}$ -Algebra. In the previous definition, what we have really used for it to make sense is that $\mathcal{V}$ is enriched over itself and then that the homs are objects in $\mathcal{V}$. But one could go a step further and consider an arbitrary symmetric monoidal category $\mathcal{C}$ enriched over $\mathcal{V}$. That is, if X is an object in $\mathcal{C}$, define the endomorphism operad (in $\mathcal{V}$) as $\text{End}_X(n) := \text{Hom}(X^{\otimes n}, X)$ (these are objects in $\mathcal{V}$ by the enrichment).

Then for an operad $\mathcal{P}$ in $\mathcal{V}$, a **$\mathcal{C}$ -valued $\mathcal{P}$ -algebra** is an operad morphism

$$\mathcal{P} \rightarrow \text{End}_X.$$

Main example. If $\mathcal{V} = \text{Cat}$ or Grp the category of categories or groupoids, and $\mathcal{C}$ is a $(2,1)$ -category (eg Rex^f or LFP_c), then for some object (category) $\mathcal{A}$ in $\mathcal{C}$ we can consider the operad $\text{End}_{\mathcal{A}}$ and a $\mathcal{C}$ -valued $\mathcal{P}$ -algebra $\mathcal{P} \rightarrow \text{End}_{\mathcal{A}}$.

7.2 Classical cyclic operads and their algebras

A cyclic operad removes the distinction between inputs and outputs, treating all connection points (punctures) symmetrically. The combinatorics are those of forests (graphs without cycles).

A **cyclic operad** $\mathcal{C}$ is a collection of vector spaces $\{\mathcal{C}(n)\}_{n \geq 0}$ where each $\mathcal{C}(n)$ is endowed with a right action of the symmetric group S_{n+1} . This extended group action allows the "output" (labeled $n+1$) to be permuted with the "inputs" (labeled $1, \dots, n$). The composition maps can be defined for any pair of punctures on two distinct operations:

$$\circ_{i,j} : \mathcal{C}(n) \otimes \mathcal{C}(m) \rightarrow \mathcal{C}(n+m)$$

Alternatively, a cyclic operad can be defined as a symmetric operad equipped with a compatible family of non-degenerate symmetric bilinear forms on its spaces, satisfying a cyclicity condition.

An algebra over a cyclic operad $\mathcal{C}$ is a vector space V that is an algebra over the underlying symmetric operad of $\mathcal{C}$ and is additionally equipped with a **non-degenerate symmetric bilinear form** $\langle -, - \rangle : V \otimes V \rightarrow \mathbb{k}$. This structure must satisfy a cyclicity condition relating the operad action to the bilinear form. For any $c \in \mathcal{C}(n)$ and $v_1, \dots, v_{n+1} \in V$:

$$\langle \gamma_n(c)(v_1, \dots, v_n), v_{n+1} \rangle = \langle \gamma_n(c \cdot \sigma)(v_2, \dots, v_{n+1}), v_1 \rangle$$

where σ is the cyclic permutation $(1, 2, \dots, n+1)$. A key example is a commutative Frobenius algebra.

7.3 Classical modular operads

A modular operad introduces the final layer of complexity by allowing self-composition, which creates loops. The underlying combinatorics are those of arbitrary graphs.

A **modular operad** $\mathcal{M}$, as defined by Getzler and Kapranov, is a collection of vector spaces indexed by a genus $g \geq 0$ and a number of punctures $n \geq 0$:

$$\{\mathcal{M}(g, n)\}_{(g,n) \in \mathbb{N}^2 \text{ such that } 2g-2+n > 0}$$

This collection is equipped with:

1. **A Right S_n Action** on each $\mathcal{M}(g, n)$.
2. **Composition Maps** of two types:

- **Gluing of distinct components:**

$$\circ : \mathcal{M}(g_1, n_1) \otimes \mathcal{M}(g_2, n_2) \rightarrow \mathcal{M}(g_1 + g_2, n_1 + n_2 - 2)$$

This map connects a puncture from an operation in $\mathcal{M}(g_1, n_1)$ with one from an operation in $\mathcal{M}(g_2, n_2)$.

- **Self-composition (contraction):**

$$\circ : \mathcal{M}(g, n) \rightarrow \mathcal{M}(g + 1, n - 2)$$

This map connects two distinct punctures on the *same* operation, thereby increasing the genus by one.

These maps must satisfy a set of coherence axioms ensuring that the result of a complex composition is independent of the order of elementary steps.

An algebra over a modular operad $\mathcal{M}$ is a $\mathbb{Z}_{\geq 0}$ -graded vector space $\mathbf{A} = \bigoplus_{g \geq 0} A_g$ equipped with structure maps that are compatible with the operad's composition rules. For an operation $m \in \mathcal{M}(g, n)$, the corresponding structure map is a multilinear map of the form:

$$\gamma(m) : A_{h_1} \otimes \cdots \otimes A_{h_n} \rightarrow A_{g+h_1+\cdots+h_n}$$

This structure must be compatible with both types of composition in $\mathcal{M}$. To realize the self-composition maps of the operad, the spaces A_g are typically required to be finite-dimensional, which allows for the definition of trace maps. The entire structure provides an algebraic formulation for systems like Topological Quantum Field Theories (TQFTs), where the grade g of the algebra corresponds to the genus of the underlying surfaces.


7.4 Ordinary, cyclic and modular operads à la Costello-Müller-Woike

(Following Müller-Woike)


In [Cos04] Costello gives a very efficient description of operads, cyclic operads and modular operads based on different categories of graphs. We will adopt this description and therefore briefly recall the most important definitions: A *graph* consists of a set H of *half edges* and a set V of *vertices* together with a map $H \rightarrow V$ and an involution $\iota : H \rightarrow H$ specifying how half edges are glued together. The orbits of the involution ι are the *edges* of the graph. Fixed points of ι are called *external legs* (legs, for short). We denote by $\text{Legs}(\Gamma)$ the set of external legs of a graph Γ . We may realize a graph Γ as a topological space $|\Gamma|$ with the vertices of Γ as the 0-cells and the edges of Γ as the 1-cells. A *corolla* is a graph with one vertex and only external legs. Often we will denote a graph as a pair $\Gamma = (V, H)$ of the set of vertices and the set of half edges suppressing all other parts of the structure in the notation.

Let $\Gamma = (V, H)$ and $\Gamma' = (V', H')$ be graphs. A *morphism of graphs* consists of maps $V \rightarrow V'$ and $H \rightarrow H'$ which are compatible with the graph structure in the obvious way.

Given a graph Γ we can form a new graph $\nu(\Gamma)$ by cutting open all internal edges. Formally, this replaces the involution on the half edges by the identity map. We can also form a graph $\pi_0(\Gamma)$ by contracting all internal edges. Below a pictorial presentation of these operations is given:



The category **Graphs** has as objects graphs which are finite disjoint unions of corollas. A morphism $\gamma_1 \rightarrow \gamma_2$ is given by an equivalence class of a graph Γ together with isomorphisms $\phi_1 : \gamma_1 \rightarrow \nu(\Gamma)$ and $\phi_2 : \gamma_2 \rightarrow \pi_0(\Gamma)$; note that here ϕ_1 and ϕ_2 are morphisms of graphs in the above sense, but not morphisms in the category **Graphs** (which is named after its morphisms). Two such triples (Γ, ϕ_1, ϕ_2) and $(\Gamma', \phi'_1, \phi'_2)$ are equivalent if there exists an isomorphism $\psi : \Gamma \rightarrow \Gamma'$ satisfying $\phi'_1 = \nu(\psi) \circ \phi_1$ and $\phi_2 = \pi_0(\psi) \circ \phi'_2$. The composition $\Gamma_2 \circ \Gamma_1$ is defined by replacing the vertices of Γ_2 by the graph Γ_1 , see below:



We denote by **Forests** the subcategory of **Graphs** whose objects are those objects of **Graphs** which do not contain a corolla with zero legs and whose morphisms are forests, i.e. disjoint unions of contractible graphs.

Finally, we define the category **RForests** of *rooted forests*: A *rooted graph* is a graph Γ equipped with a section $s : V(\pi_0(\Gamma)) \rightarrow \text{Legs}(\Gamma)$ of the obvious map $\text{Legs}(\Gamma) \rightarrow V(\pi_0(\Gamma))$, i.e. in each component of Γ we distinguish an external leg that we refer to as the *root*. Morphisms of rooted graphs are morphisms of the underlying graph which are compatible with the specified sections. Note that a rooted forest Γ induces the structure of a rooted graph on $\nu(\Gamma)$ by declaring for every vertex the edge in the direction of the root of Γ as the root of the vertex in $\nu(\Gamma)$. The category **RForests** is defined just like **Forests** with objects and morphisms replaced by their rooted version. There is a functor $\text{RForests} \rightarrow \text{Forests}$ which forgets the root.

We assume some familiarity with the theory of symmetric monoidal bicategories; we refer e.g. to [SP09, Chapter 2] for a detailed discussion and to [Lei98] for a short introduction (however without the treatment of monoidal structures). In particular, we rely on the following notions:

By a *bicategory* we mean a three-layered categorical structure with objects, 1-morphisms and 2-morphisms (sometimes also called 0-cells, 1-cells and 2-cells) in the weak sense. A *morphism between bicategories* (sometimes also referred to as (weak) 2-functor) will just be called *functor*. *Symmetric monoidal functors* between symmetric monoidal bicategories are to be understood in a strong (not in any kind of lax) sense unless otherwise stated. In particular, a symmetric monoidal functor between symmetric monoidal bicategories comprises various sorts of coherence data subject to coherence conditions. We will briefly illustrate this after the next definition.

Let $\mathcal{M}$ be a symmetric monoidal bicategory. An *operad* / *cyclic operad* / *modular operad* in $\mathcal{M}$ is a symmetric monoidal functor

$$\mathcal{O} : \mathbf{RForesets} / \mathbf{Forests} / \mathbf{Graph} \rightarrow \mathcal{M}$$

where we consider $\mathbf{RForesets} / \mathbf{Forests} / \mathbf{Graph}$ as a symmetric monoidal bicategory with only trivial 2-morphisms. See [MW, page 9] to see the equivalence with the classical definitions.

Let us emphasize that this defines operads *without* an operadic identity. However, all operads appearing in this paper have operadic unit, and it will be important to correctly keep track of those.

7.5 The correspondence between groupoids and 1-homotopy types

A 1-homotopy type is a topological space with $\pi_n(X, x_0) = 0$ for all basepoints $x_0 \in X$. A connected 1-homotopy type is an aspherical space, aka an Eilenberg-MacLane space.

There is a diagram of categories and functors that we explain below:

$$\begin{array}{ccccc}
 & & B & & \\
 & \nearrow & & \searrow & \\
 \text{Cat} & \xrightarrow{N} & \mathbf{sSet} & \xrightarrow{|\!-\!|} & \mathbf{Top}^{CW} \\
 \uparrow & & \uparrow & & \parallel \\
 \text{Gpd} & \xrightarrow{N} & \mathbf{Kan} & \xrightarrow{|\!-\!|} & \mathbf{Top}^{CW} \\
 & \nwarrow & \nwarrow & \swarrow & \\
 & \Pi_1 & & Sing & \\
 & \searrow & & \swarrow & \\
 & & \Pi & &
 \end{array}$$

Here

- N is the nerve. It is a fully faithful functor.
- $|\!-\!|$ is the geometric realisation functor.
- $Sing$ is the singular functor; $Sing(X)$ is the simplicial sets with $Sing(X)_n$ the set of singular n -chains.
- Π_1 is the fundamental group of a Kan complex.
- B (defined as the composition) is the classifying space functor.
- Π is the fundamental groupoid functor.

The main result is that up to homotopy we have that the the $Sing - |\!-\!|$ adjunction induces an equivalence

$$|\!-\!| : \mathbf{Ho}(\mathbf{Kan}) \simeq \mathbf{Ho}(\mathbf{Top}^{CW}) : Sing$$

The key point is that restricting (composing) to groupoids induces an equivalence

$$B : \text{Ho}(\mathbf{Gps}) \simeq \text{Ho}(\mathbf{Top}_{\leq 1}^{CW}) : \Pi$$

In particular this means that a groupoid is the same thing as a 1-homotopy type; and a **connected groupoid** the same thing as an aspherical space.

A way to see this equivalence is the following: if $\mathcal{C}$ is a groupoid, $B\mathcal{C} = |N\mathcal{C}|$ is a disjoint union the $K(\pi, 1)$'s. This is because every groupoid is a disjoint union of its connected components (a groupoid is connected is every object is connected through a morphism to every other object). Now $\mathcal{C} = \coprod_i \mathcal{C}_i$ with each $\mathcal{C}_i$ connected. Now for every i take an object $X_i \in \mathcal{C}_i$, and let $\mathcal{G}_i$ be the full subcategory on that only object. That is, $\mathcal{G}_i$ is a 1-object groupoid aka a group, and the inclusion $\mathcal{G}_i \hookrightarrow \mathcal{C}_i$ is an equivalence for all i as $\mathcal{C}_i$ is connected. So $\mathcal{C} \simeq \coprod_i \mathcal{G}_i$. It turns out that B commutes with Π so

$$B\mathcal{C} \simeq \coprod_i B\mathcal{G}_i \simeq \coprod_i K(G_i, 1).$$

7.6 The (framed) little disc operad E_n and $\mathfrak{f}E_n$

First I need to recall a few things:

- The (ordered) configuration space of a given space X is

$$\text{Conf}_n(X) := \{(x_1, x_2, \dots, x_n) \in X^n : x_i \neq x_j \text{ for all } i \neq j\}.$$

- The symmetric group $\mathfrak{S}_n$ naturally acts on $\text{Conf}_n(X)$. The unordered configuration space of X is

$$\text{UConf}_n(X) := \text{Conf}_n(X) / \mathfrak{S}_n,$$

ie this is the space of sets of n distinct points.

- We have (more or less by def) isomorphisms

$$PB_n \cong \pi_1(\text{Conf}_n(D^2)) \quad , \quad B_n \cong \pi_1(\text{UConf}_n(D^2)),$$

where B_n and PB_n are the braid and pure braid group on n strands, respectively.

- These two spaces $\text{Conf}_n(D^2)$ and $\text{UConf}_n(D^2)$ are aspherical by a classic theorem by Fox-Neuwirth. Therefore (as they are connected)

$$B(PB_n) \simeq \text{Conf}_n(D^2) \quad , \quad B(B_n) \simeq \text{UConf}_n(D^2).$$

There is a framed version of this story

- The (ordered) framed configuration space of a given manifold M is

$$\text{Conf}_n^{fr}(M) := \{((x_1, f_1), (x_2, f_2), \dots, (x_n, f_n)) \in (Fr(M))^n \mid x_i \neq x_j \text{ for all } i \neq j\}$$

where $Fr(M)$ is the frame bundle of M .

- The symmetric group $\mathfrak{S}_n$ naturally acts on $\text{Conf}_n^{fr}(M)$. The unordered framed configuration space of X is

$$\text{UConf}_n^{fr}(M) := \text{Conf}_n^{fr}(M) / \mathfrak{S}_n.$$

- We have (more or less by def) isomorphisms

$$PB_n^{fr} \cong \pi_1(\text{Conf}_n^{fr}(D^2)) \quad , \quad B_n^{fr} \cong \pi_1(\text{UConf}_n^{fr}(D^2)),$$

where B_n^{fr} and PB_n^{fr} are the framed braid and pure framed braid group on n strands, respectively. That is, we have ribbons instead of strings.

- These spaces are again aspherical and then

$$B(PB_n^{fr}) \simeq \text{Conf}_n^{fr}(D^2) \quad , \quad B(B_n^{fr}) \simeq \text{UConf}_n^{fr}(D^2).$$

The **little n -disc operad** E_n is a topological operad (ie an operad in the symmetric monoidal category of spaces) defined as

$$E_n(k) := \text{Emb}_{\text{rect}}([0, 1]^n)^{\amalg k}, [0, 1]^n$$

the space of rectilinear embeddings; see Example 1.8 in Heuts-Moerdijk for a precise definition. It is obvious that this space is isomorphic to $\text{Emb}_{\text{td}}((D^n)^{\amalg k}, D^n)$ the space of embeddings that are obtained by a composition of translations and dilatations, and D^n is the n -disc.

The **framed little n -disc operad** $\mathbf{f}E_n$ is the topological operad defined as

$$\mathbf{f}E_n(k) := \text{Emb}_{\text{tdr}}((D^n)^{\amalg k}, D^n)$$

the space of embeddings that are obtained by a composition of translations, dilatations and rotations. By ????? , the space $\text{Emb}_{\text{tdr}}((D^n)^{\amalg k}, D^n)$ is in fact homotopy equivalent to the space $\text{Emb}((D^n)^{\amalg k}, D^n)$ of all embeddings.

Key property: Evaluation at the centre induces homotopy equivalences

$$\text{Emb}_{\text{td}}((D^n)^{\amalg k}, D^n) \rightarrow \text{Conf}_k(D^n)$$

and

$$\text{Emb}_{\text{tdr}}((D^n)^{\amalg k}, D^n) \rightarrow \text{Conf}_k^{fr}(D^n).$$

Upshot: $E_2(k)$ and $\mathbf{f}E_2(k)$ are aspherical spaces, in particular we have

$$E_2(k) \simeq B(PB_k) \quad , \quad \mathbf{f}E_2(k) \simeq B(PB_k^{fr}).$$

In particular, by the equivalence between connected groupoids and aspherical spaces, we can regard $\mathbf{f}E_2$ and E_2 as operads in groupoids, not in spaces.

7.7 Disc algebras vs. framed E_n -algebras

I will describe here roughly the $(2,1)$ -categorical version; there is also an ∞ -categorical ones.

Let Disc_n the symmetric monoidal $(2,1)$ -category with objects disjoint union of oriented disks and hom groupoids orientation-preserving embeddings with isotopies, with disjoint union as monoidal product. If $\mathcal{C}$ is a symmetric monoidal $(2,1)$ -category, a **disk algebra** is a symmetric monoidal functor $\text{Disc}_n \rightarrow \mathcal{C}$.

There is a framed variant of this story: Disc_n^{fr} the symmetric monoidal $(2,1)$ -category with objects disjoint union of framed (=choice of trivialisation of the tangent bundle) disks and arrows embeddings equipped with a compatibility of framings (see Tanaka's notes for a precise statement), with disjoint union as monoidal product. If $\mathcal{C}$ is a symmetric monoidal $(2,1)$ -category, a **framed disk algebra** is a symmetric monoidal functor $\text{Disc}_n^{fr} \rightarrow \mathcal{C}$.

Theorem. The following three pieces of data are equivalent (inducing equivalences of the corresponding categories):

1. A 2-disk algebra in $\mathbf{Rex}^f$,
2. A $\mathbf{Rex}^f$ -valued framed E_2 -algebra,
3. A balanced monoidal category in $\mathbf{Rex}^f$.

($\mathbf{Rex}^f$ can be replaced by something more general like LFP_c).

The equivalence between 1 and 3 is explained with detail in Theorem 3.2 of Brochier's notes "Factorisation homology of braided tensor categories": for a functor $F : \mathbf{Disc}_n \rightarrow \mathbf{Rex}^f$ then $\mathcal{A} := F(D^2)$ is a balanced monoidal category.

I want to focus on the equivalence 1 - 2. The key word is **envelope construction** which is nothing but the adjunction between operads and PROPs (= symm mon cat gen by a single object)

$$F : \mathbf{Operads} \rightleftarrows \mathbf{PROPs} : U$$

and more concretely how to construct a symmetric monoidal category out of an operad. One builds the "free PROP" aka envelope of an operad putting together the trees; the right adjoint uses simply one output, so that's kinda a forgetful.

In the groupoid-valued setting, the adjunction reads

$$F : \mathbf{Op}(\mathbf{Gpd}) \rightleftarrows (2, 1) - \mathbf{PROPs} : U$$

Obviously the endomorphism operad $\mathbf{End}_{\mathcal{A}}$ can be viewed as the forgetful $\mathbf{End}_{\mathcal{A}} = U(\widetilde{\mathbf{End}_{\mathcal{A}}})$ (restriction) of a (2,1)-PROP $\widetilde{\mathbf{End}_{\mathcal{A}}}$ with $\widetilde{\mathbf{End}_{\mathcal{A}}}(n, m) := \mathbf{Rex}^f(\mathcal{A}^{\boxtimes n}, \mathcal{A}^{\boxtimes m})$. The adjunction says that there is an equivalence

$$\mathbf{Hom}_{\mathbf{PROP}}(F(\mathbf{f}E_2), \widetilde{\mathbf{End}_{\mathcal{A}}}) \simeq \mathbf{Hom}_{\mathbf{Op}}(\mathbf{f}E_2, \mathbf{End}_{\mathcal{A}}).$$

The RHS are precisely $\mathbf{Rex}^f$ -valued framed E_2 -algebras. Now the point is that the LHS are exactly the same as framed 2-disc algebras in $\mathbf{Rex}^f$, ie symm mon functors $\mathbf{Disc}_2 \rightarrow \mathbf{Rex}^f$. Modifying the target is easy: $\widetilde{\mathbf{End}_{\mathcal{A}}}$ is a full subcat of $\mathbf{Rex}^f$ so it changes nothing as $\mathbf{Disc}_2$ is generated by a single object (namely the 2-disc). So everything amounts to showing that $F(\mathbf{f}E_2) \simeq \mathbf{Disc}_2$ as (2,1)-categories.

Here one has to remember from the previous subsection that

$$\mathbf{f}E_2(n) = \mathbf{Emb}_{\mathbf{tdr}}((D^2)^{\amalg n}, D^2) \simeq \mathbf{Emb}((D^2)^{\amalg n}, D^2),$$

so viewed in groupoids, one applies by the correspondence explained above the fundamental groupoid functor,

$$\mathbf{f}E_2(n) = \amalg \mathbf{Emb}((D^2)^{\amalg n}, D^2)$$

as a groupoid-valued operad. If we then consider the free (2,1)-PROP, that is, the thing obtained by concatenating trees, then we get that

$$F(\mathbf{f}E_2)(n, m) = \amalg \mathbf{Emb}((D^2)^{\amalg n}, (D^2)^{\amalg m}),$$

that is $\mathbf{Disc}_2$.

There is also an "unframed" version of this theorem:

Theorem. The following three pieces of data are equivalent (inducing equivalences of the corresponding categories):

1. A framed 2-disk algebra in $\mathbf{Rex}^f$,
2. A $\mathbf{Rex}^f$ -valued E_2 -algebra,
3. A braided monoidal category in $\mathbf{Rex}^f$.

7.8 The cyclic and modular endomorphism operad

(Mostly taken from Woike’s reflection equivariance paper)

Classically, an algebra X over an operad $\mathcal{O}$ is an operad map

$$\mathcal{O} \rightarrow \text{End}_X$$

to the endomorphism operad of the underlying object of the algebra X ; recall that $\text{End}_X(n) = \text{Hom}(X^{\otimes n}, X)$. If X was a vector space for instance, in order to make this endomorphism operad cyclic (or more genrally modular), ie to exchange a copy of X to the target and the one from the target to the source, or to pair up two copies of X , we’d need an identification between X and its dual aka a non-deg pairing in X . We do this next in general (with $\mathcal{A}$ instead of X as we’re thinking of a symm mon bicat).

In order to generalize this to cyclic and modular operads, the endomorphism operad needs to be made modular (and therefore in particular cyclic). This is done in [MW23a, Section 2] for the bicategorical case, following the principles of [GK95, GK98, Cos04]: First we recall the notion of a **non-degenerate symmetric pairing** $\kappa : \mathcal{A} \boxtimes \mathcal{A} \rightarrow \mathcal{I}$ on an object $\mathcal{A}$ in a symmetric monoidal bicategory $\mathcal{S}$ with monoidal product $\boxtimes$ and monoidal unit $\mathcal{I}$, where

- *non-degeneracy* means that κ exhibits $\mathcal{A}$ as its own dual in the homotopy category of $\mathcal{S}$, that is, there is a coevaluation object $\Delta : \mathcal{I} \rightarrow \mathcal{A} \boxtimes \mathcal{A}$ that together with κ satisfies the zigzag identities up to isomorphism.
- and *symmetry* means that κ is a homotopy fixed point in $\mathcal{S}(\mathcal{A} \boxtimes \mathcal{A}, \mathcal{I})$ with respect to the $\mathbb{Z}_2$ -action coming from the symmetric braiding of $\mathcal{S}$.

Thanks to the non-degenerate symmetric pairing, we can define a symmetric monoidal functor

$$\text{End}_{\mathcal{A}, \kappa} : \text{Graphs} \rightarrow \text{Cat}$$

sending a corolla T to the morphism category

$$\text{End}_{\mathcal{A}, \kappa}(T) := \mathcal{S}(\mathcal{A}^{\boxtimes \text{Legs}(T)}, \mathcal{I}) \quad (70)$$

from the unordered monoidal product $\mathcal{A}^{\boxtimes \text{Legs}(T)}$ of $\mathcal{A}$ over the set $\text{Legs}(T)$ of legs of T to the monoidal unit $\mathcal{I} \in \mathcal{S}$. This Cat -valued modular operad is the *modular endomorphism operad* of $(\mathcal{A}, \kappa)$ and denoted by $\text{End}_{\mathcal{A}, \kappa}$.

If $\mathcal{O}$ is a Cat -valued modular operad $\mathcal{O} : \text{Graphs} \rightarrow \text{Cat}$, an $\mathcal{S}$ -valued modular $\mathcal{O}$ -algebra is now defined as an object $\mathcal{A}$ in $\mathcal{S}$ with a non-degenerate symmetric pairing κ plus a map $\mathcal{O} \rightarrow \text{End}_{\mathcal{A}, \kappa}$ of modular operads (aka a symm mon nat transf). A *cyclic algebra* is defined analogously. Both cyclic and modular algebras over a fixed cyclic or modular operad, respectively, with values in a symmetric monoidal bicategory form themselves bicategories in which all morphisms turn out to be invertible, i.e. they form 2-groupoids [MW23a, Proposition 2.18].

Unpacking the definition Let us partly unpack the definition of a Rex^f -valued modular $\mathcal{O}$ -algebra (all of this is worked out in detail in [MW23a, Section 2.5] where the dual case of Lex^f is treated): It has an underlying object $\mathcal{A} \in \text{Rex}^f$, i.e. a finite category, and a non-degenerate symmetric pairing $\kappa : \mathcal{A} \boxtimes \mathcal{A} \rightarrow \text{vect}$. By virtue of right exactness, we have

$$\text{Rex}^f(\mathcal{A} \boxtimes \mathcal{A}, \text{vect}) \simeq (\mathcal{A} \boxtimes \mathcal{A})^{op} \boxtimes \text{vect} \simeq \mathcal{A}^{op} \boxtimes \mathcal{A}^{op} \simeq \text{Rex}^f(\mathcal{A}, \mathcal{A}^{op})$$

(here we have used the Eilenberg-Watts theorem) ie κ determines an equivalence $D : \mathcal{A} \rightarrow \mathcal{A}^{\text{op}}$ via $\kappa(X, Y) = \mathcal{A}(X, DY)^*$, and the symmetry of κ induces an isomorphism $D^2 \simeq \text{id}_{\mathcal{A}}$. The coevaluation is a right exact functor $\Delta : \mathbf{vect} \rightarrow \mathcal{A} \boxtimes \mathcal{A}$ and is hence (again by the Eilenberg-Watts theorem) an object in $\mathcal{A} \boxtimes \mathcal{A}$ given by the end

$$\Delta = \int_{X \in \mathcal{A}} DX \boxtimes X;$$

we refer to [FSS20] for an introduction to (co)ends in finite linear categories. Moreover, the structure of a modular $\mathcal{O}$ -algebra on $\mathcal{A}$ provides us for each operation $o \in \mathcal{O}(T)$ for a corolla T with a right exact functor $\mathcal{A}_o : \mathcal{A}^{\boxtimes \text{Legs}(T)} \rightarrow \mathbf{vect}$. If $T \in \mathbf{Graphs}$ is not connected, i.e. $T = \coprod_{l \in L} T(l)$ with corollas $T(l)$, then to $o = (o_l)_{l \in L} \in \mathcal{O}(T) \simeq \prod_{l \in L} \mathcal{O}(T(l))$ the modular algebra associates the family of right exact functors $(\mathcal{A}_{o_l})_{l \in L}$; it gives us also a right exact functor $\bigoplus_{l \in L} \mathcal{A}_{o_l} \rightarrow \mathbf{vect}$ (by abuse of notation, we will not distinguish between these two objects). The data of a modular algebra includes also the compatibility with composition: Suppose that a morphism $\Gamma : T \rightarrow T'$ glues two legs (of T) together; that is, Γ is like the leftmost Γ' picture above if the legs belong to different corollas, or Γ is the result of taking a given corolla T and glue two legs (and then T' is just T minus the two legs that were glued).

Denote by $o' \in \mathcal{O}(T')$ the image of some $o \in \mathcal{O}(T)$ under the operadic composition $\mathcal{O}(\Gamma) : \mathcal{O}(T) \rightarrow \mathcal{O}(T')$. Then we obtain an isomorphism of right exact functors between $\mathcal{A}_{o'}$ and $\mathcal{A}_o$, but with Δ inserted into the two arguments of $\mathcal{A}_o$ corresponding to the legs that are glued together, i.e.

$$\mathcal{A}_{o'} \simeq \mathcal{A}_o(\dots, \Delta', \dots, \Delta'', \dots) \quad (2.2)$$

with Sweedler notation $\Delta = \Delta' \boxtimes \Delta''$ for the coevaluation object (as usual, the notion does not imply that $\mathcal{A}$ is a pure tensor). In some contexts, (2.2) is called *excision*.

7.9 Classifications of cyclic and modular algebras

The classical result:

$$\text{Alg}(\mathbf{f}E_2, \text{Rex}^{\mathbf{f}}) \simeq \text{balanced monoidal categories in } \text{Rex}^{\mathbf{f}}.$$

The intermediate result:

$$\text{CycAlg}(\text{As}, \text{Rex}^{\mathbf{f}}) \simeq \text{ModAlg}(\text{OSurf}, \text{Rex}^{\mathbf{f}}) \simeq \text{pivotal GV-categories in } \text{Rex}^{\mathbf{f}}.$$

(for these we have used

$$\text{OSurf}_{|g=0} \simeq \text{As} \quad , \quad \text{Hdby}_{|g=0} \simeq \mathbf{f}E_2$$

The classification of $\text{Rex}^{\mathbf{f}}$ -valued cyclic framed E_2 -algebras:

$$\text{CycAlg}(\mathbf{f}E_2, \text{Rex}^{\mathbf{f}}) \simeq \text{ModAlg}(\text{Hdby}, \text{Rex}^{\mathbf{f}}) \simeq \text{ribbon GV-categories in } \text{Rex}^{\mathbf{f}}$$

(a modular Hdby-algebra is called an ansular functor).

We also have

$$\text{Surf}_{|g=0} \simeq \mathbf{f}E_2,$$

ie $\mathbf{f}E_2$ is the “cyclic operad of genus zero surfaces”

8 Factorisation homology

8.1 Definition

$$\begin{array}{ccc}
 \text{Disc} & \xrightarrow{\mathcal{A}} & \text{Rex}^f \\
 \downarrow \iota & \nearrow f_{(-)} \mathcal{A} & \\
 \text{Surf} & &
 \end{array} \tag{71}$$

8.2 Properties

- We have $\int_{D^2} \mathcal{A} \simeq \mathcal{A}$, the equivalence being induced by any of the universal arrows $\mathcal{A} \rightarrow \int_{D^2} \mathcal{A}$ coming from the description of the homotopy colimit. This is true by construction: factorisation homology being a homotopy left Kan extension along an embedding means that the diagram (71) is actually commutative up to isomorphism. Since the image of the 2-disc under $\mathcal{A}$ is $\mathcal{A}$ as category, this yields the equivalence.

8.3 Quantum structure sheaf or distinguished object

Factorization homology is a *canonically pointed* theory in the following sense: for a surface Σ , the unique embedding $\emptyset \hookrightarrow \Sigma$ induces a functor

$$\mathcal{O}_\Sigma : \text{vect} \simeq \int_{\emptyset} \mathcal{A} \rightarrow \int_{\Sigma} \mathcal{A}.$$

This functor produces a distinguished object in $\int_{\Sigma} \mathcal{A}$, namely the image of the 1-dimensional vector space, $\mathcal{O}_{\mathcal{A},\Sigma} := \mathcal{O}_\Sigma(\mathbb{k})$, that is called the *quantum structure sheaf* in "Integrating Quantum groups".

Alternatively, $\mathcal{O}_{\mathcal{A},\Sigma}$ can be defined in a different way (up to isomorphism): as the image of the monoidal unit $\mathbf{1}$ of $\mathcal{A}$ under any of the universal functors $\mathcal{A} \rightarrow \int_{\Sigma} \mathcal{A}$ (a leg of the colimit cone) for any embedding $D^2 \hookrightarrow \Sigma$.

To see that the two descriptions agree, one argues as follows: the first observation is that if they agree for a given embedding then it is independent on the chosen embedding. Now the second observation is that it suffices to check this for the disc D^2 , this is because no matter for what embedding $D^2 \hookrightarrow \Sigma$ we have a commutative diagram of embeddings

$$\begin{array}{ccc}
 & \emptyset & \\
 \swarrow & & \searrow \\
 D^2 & \xrightarrow{\quad} & \Sigma
 \end{array}$$

since $\emptyset$ is initial; and then we get a commutative diagram (up to iso)

$$\begin{array}{ccc}
 & \text{vect} & \\
 \swarrow & & \searrow \\
 \int_{D^2} \mathcal{A} & \xrightarrow{\quad} & \int_{\Sigma} \mathcal{A}
 \end{array}$$

which means that any embedding $D^2 \hookrightarrow \Sigma$ induces a functor $\int_{D^2} \mathcal{A} \rightarrow \int_{\Sigma} \mathcal{A}$ that maps the distinguished object to the distinguished object. Therefore it suffices to check that under the equivalence $\int_{D^2} \mathcal{A} \simeq \mathcal{A}$, the distinguished object $\mathcal{O}_{\mathcal{A},D^2}$ corresponds precisely with the monoidal unit of $\mathcal{A}$.

Now the key is that a balanced monoidal category is a different name for a framed E_2 -algebra aka a 2-disc algebra, that is a functor $\text{Disk} \rightarrow \text{Rex}^f$. The monoidal unit is given by the canonical

embedding $\emptyset \hookrightarrow D^2$ inducing a map $\mathbf{vect} \rightarrow \mathcal{A}$ which maps the base field to the monoidal unit. By the commutativity of (71) we have that the functors $\mathbf{vect} \rightarrow \mathcal{A}$ and $\mathbf{vect} \rightarrow \int_{D^2} \mathcal{A}$ must be isomorphic (up to the equivalence in the target). Ie that $\mathcal{O}_{\mathcal{A}, D^2} \cong \mathbf{1}$ viewed in $\mathcal{A}$.

The same type of argument says that if $\Pi_k D^2 \hookrightarrow \Sigma$ is an embedding, with universal functor $\mathcal{A}^{\boxtimes k} \rightarrow \int_{\Sigma} \mathcal{A}$, and that embedding factors through a larger disc, ie there is a diagram

$$\begin{array}{ccc} & \Pi_k D^2 & \\ \swarrow & & \searrow \\ D^2 & \xrightarrow{\quad} & \Sigma \end{array}$$

then the universal functor $\mathcal{A}^{\boxtimes k} \rightarrow \int_{\Sigma} \mathcal{A}$ factors through the monoidal product of $\mathcal{A}$, ie there is a commutative diagram (up to iso)

$$\begin{array}{ccc} \mathcal{A}^{\boxtimes k} & \xrightarrow{\quad} & \int_{\Sigma} \mathcal{A} \\ \otimes^k \downarrow & \nearrow & \\ \mathcal{A} & & \end{array}$$

8.4 Moduli algebra

This is $A_{\Sigma} := \underline{\mathbf{End}}(\mathcal{O}_{\mathcal{A}, \Sigma})$. Not to be confused with $\mathcal{F}_{\mathcal{A}} := \otimes \underline{\mathbf{End}}(\mathbf{1})$ (the *reflection equation algebra*) or with $a_P := \mathcal{F}_{\mathcal{A}}^{\otimes n}$ where P is a pattern of rank n .

8.5 Skein algebra (categorical version)

This is the endomorphism algebra

$$\mathbf{SkAlg}_{\mathcal{A}}(\Sigma) := \mathbf{End}_{\int_{\Sigma} \mathcal{A}}(\mathcal{O}_{\mathcal{A}, \Sigma}).$$

9 Factorization homology

Factorization homology [?, ?] with coefficients in a given E_n -algebra is an invariant of n -dimensional manifolds that satisfies a generalization of the classical Eilenberg-Steenrod axioms for homology theories. In the present paper, we will be interested in oriented surfaces, so we will need framed E_2 -algebras, that for concreteness we will take in the symmetric monoidal $(2, 1)$ -category $\mathbf{Rex}^f$ of finite categories, right-exact functors and natural isomorphisms, with the Deligne product as monoidal product. We refer the reader to [?, Chapter 2] for the precise definition of a symmetric monoidal bicategory.

9.1 A brief review of factorization homology

Let $\mathbf{Surf}$ denote the symmetric monoidal $(2, 1)$ -category of compact, oriented surfaces (possibly with boundary) with hom-groupoids given by the fundamental groupoid $\Pi_1(\mathbf{Emb}(\Sigma, \Sigma'))$ of the space of smooth, oriented embeddings $\Sigma \hookrightarrow \Sigma'$. The monoidal product is given by disjoint union with unit the empty surface. We also write $\mathbf{Disk}$ for the full monoidal subcategory of $\mathbf{Surf}$ with objects being disjoint unions of disks.

It is well-known that a $\mathbf{Rex}^f$ -valued framed E_2 -algebra, viewed as a symmetric monoidal (pseudo)functor $\mathbf{Disk} \rightarrow \mathbf{Rex}^f$, is fully determined by the datum of a *balanced monoidal category* $\mathcal{A}$ in $\mathbf{Rex}^f$ (given by evaluation at the disk). This means that $\mathcal{A} \in \mathbf{Rex}^f$ is additionally equipped with a monoidal

structure $\otimes : \mathcal{A} \boxtimes \mathcal{A} \longrightarrow \mathcal{A}$, a *braiding* $c_{X,Y} : X \otimes Y \xrightarrow{\cong} Y \otimes X$ and a *twist* (or *balancing*) $\theta_X : X \xrightarrow{\cong} X$. The two latter are families of natural isomorphisms, where the braiding satisfies the so-called hexagon equations and the twist satisfies

$$\theta_{X \otimes Y} = c_{Y,X} c_{X,Y} (\theta_X \otimes \theta_Y)$$

If $\mathcal{A}$ is a finite balanced monoidal category (viewed as a symmetric monoidal functor $\mathcal{A} : \text{Disk} \longrightarrow \text{Rex}^f$), and $\iota : \text{Disk} \longrightarrow \text{Surf}$ is the canonical symmetric monoidal embedding, then *factorization homology* is defined as the homotopy left Kan extension of $\mathcal{A}$ along ι ,

$$\begin{array}{ccc} \text{Disk} & \xrightarrow{\mathcal{A}} & \text{Rex}^f \\ \downarrow \iota & \nearrow f_{(-)} \mathcal{A} & \\ \text{Surf} & & \end{array} \quad (72)$$

Given a compact, oriented surface Σ , the value of this symmetric monoidal functor on Σ will be denoted as $\int_{\Sigma} \mathcal{A}$. It follows from the theory of Kan extensions (e.g. [?, Theorem 6.2.1]) that this object can be described as the homotopy colimit (i.e., the $(2,1)$ -colimit)

$$\int_{\Sigma} \mathcal{A} \simeq \text{hocolim}_{\substack{(D^2)^{\sqcup n} \hookrightarrow \Sigma \\ n \geq 0}} \mathcal{A}^{\boxtimes n} \quad (73)$$

over all disk embeddings $(D^2)^{\sqcup n} \hookrightarrow \Sigma$ and all $n \geq 0$.

Here is a list of basic properties of factorization homology:

- (1) By definition (e.g. [?, Corollary 6.3.9]), there is an equivalence $\int_{D^2} \mathcal{A} \simeq \mathcal{A}$.
- (2) Factorization homology is a *canonically pointed* theory in the following sense: for a surface Σ , the unique embedding $\emptyset \hookrightarrow \Sigma$ induces a functor

$$\mathcal{O}_{\Sigma} : \text{vect} \simeq \int_{\emptyset} \mathcal{A} \longrightarrow \int_{\Sigma} \mathcal{A}.$$

This functor produces a distinguished object in $\int_{\Sigma} \mathcal{A}$, namely the image of the 1-dimensional vector space, $\mathcal{O}_{\mathcal{A},\Sigma} := \mathcal{O}_{\Sigma}(\mathbb{k})$, that is called the *quantum structure sheaf* in [?]. Alternatively, $\mathcal{O}_{\mathcal{A},\Sigma}$ can be defined (up to isomorphism) as the image of the monoidal unit $\mathbb{1}$ of $\mathcal{A}$ under any of the universal functors $\mathcal{A} \longrightarrow \int_{\Sigma} \mathcal{A}$ (a leg of the colimit cone) for any embedding $D^2 \hookrightarrow \Sigma$.

- (3) If $(D^2)^{\sqcup n} \hookrightarrow \Sigma$ is an embedding of a disjoint union of disks which factors through a bigger disk, then the corresponding universal morphism $\mathcal{A}^{\boxtimes n} \longrightarrow \int_{\Sigma} \mathcal{A}$ factors through the n -fold monoidal product functor (E_2 multiplication) $\mathcal{A}^{\boxtimes n} \longrightarrow \mathcal{A}$.
- (4) Given a compact, oriented 1-manifold P (that is, a disjoint union of intervals $I = [0,1]$ and circles S^1), the factorization homology $\int_{P \times I} \mathcal{A}$ carries a canonical monoidal structure. Indeed there is a natural stacking embedding

$$(P \times I) \amalg (P \times I) \longrightarrow P \times [0,2] \cong P \times I$$

which induces a functor

$$\otimes : \int_{P \times I} \mathcal{A} \boxtimes \int_{P \times I} \mathcal{A} \longrightarrow \int_{P \times I} \mathcal{A},$$

that defines a monoidal structure with unit $\mathcal{O}_{\mathcal{A},P \times I}$.

- (5) Given a surface Σ with boundary $\partial\Sigma$, we have that $\int_{\Sigma} \mathcal{A}$ is naturally a left $\int_{\partial\Sigma \times I} \mathcal{A}$ -module. Indeed for a collar neighborhood $\partial\Sigma \times I$ of $\partial\Sigma$, the inclusion map

$$(\partial\Sigma \times I) \amalg \Sigma \hookrightarrow \tilde{\Sigma} \cong \Sigma$$

induces the action functor

$$\int_{\partial\Sigma \times I} \mathcal{A} \boxtimes \int_{\Sigma} \mathcal{A} \longrightarrow \int_{\Sigma} \mathcal{A}.$$

- (6) In complete analogy to the Eilenberg-Steenrod axioms, factorization homology satisfies a local-to-global principle called *excision*, which is the main computational tool in the theory. Suppose Σ_1, Σ_2 are compact, oriented surfaces (with corners) let $\Sigma := \Sigma_1 \cup_{P \times I} \Sigma_2$ be the surface obtained by a collar gluing along a 1-manifold P . We choose the monoidal structure on $\int_{P \times I} \mathcal{A}$ so that $\int_{\Sigma_2} \mathcal{A}$ is a left $\int_{P \times I} \mathcal{A}$ -module category, and $\int_{\Sigma_1} \mathcal{A}$ is therefore a right $\int_{P \times I} \mathcal{A}$ -module category.

The excision axiom then affirms that the embedding $\Sigma_1 \amalg \Sigma_2 \hookrightarrow \Sigma$ induces an equivalence

$$\int_{\Sigma_1} \mathcal{A} \boxtimes_{\int_{P \times I} \mathcal{A}} \int_{\Sigma_2} \mathcal{A} \simeq \int_{\Sigma} \mathcal{A}. \quad (74)$$

In fact, it is shown in [?, §3.3] that the condition $\int_{D^2} \mathcal{A} \simeq \mathcal{A}$ and the excision axiom characterize a symmetric monoidal functor $\int_{(-)} \mathcal{A} : \mathbf{Surf} \longrightarrow \mathbf{Rex}^f$ as factorization homology.

9.2 The $\mathcal{A}^{\boxtimes n}$ -module structure

Let us now fix a non-negative integer $n \geq 0$ and a balanced monoidal category $\mathcal{A}$. Let us suppose that a connected, oriented, compact surface Σ comes with a choice of parametrized, oriented intervals $P = \cup_n I$ embedded in $\partial\Sigma$, that we call *marked boundary intervals*. That is, in the language of [?], Σ is an object of one of the groupoids of the open surface modular operad.

In this situation, $\int_{\Sigma} \mathcal{A}$ can be endowed with the structure of left $\mathcal{A}^{\boxtimes n}$ -module category. For choosing a collar neighborhood $P \times I \cong \amalg_n D^2$ of the embedded intervals, the embedding

$$(P \times I) \amalg \Sigma \hookrightarrow \Sigma$$

induces the action functor

$$\odot : \mathcal{A}^{\boxtimes n} \boxtimes \int_{\Sigma} \mathcal{A} \simeq \int_{P \times I} \mathcal{A} \boxtimes \int_{\Sigma} \mathcal{A} \longrightarrow \int_{\Sigma} \mathcal{A}.$$

In particular, if $\mathcal{A}$ is a finite tensor category, then $\int_{\Sigma} \mathcal{A}$ is a finite left $\mathcal{A}^{\boxtimes n}$ -module category. Indeed.....

Example 9.1. For a disk D^2 with n marked boundary intervals, the corresponding action functor

$$\odot : \mathcal{A}^{\boxtimes n} \boxtimes \mathcal{A} \simeq \mathcal{A}^{\boxtimes n+1} \longrightarrow \mathcal{A}$$

is naturally isomorphic, by property (3) of Section 9.1, to the $(n+1)$ -fold monoidal product of $\mathcal{A}$. Therefore, $\int_{D^2} \mathcal{A} \simeq {}_n\mathcal{A}$, that is, $\mathcal{A}$ with the left regular action of $\mathcal{A}^{\boxtimes n}$ on $\mathcal{A}$.

10 Modified traces

10.1 Twisted traces

Let $\mathcal{M}$ be a $\mathbb{k}$ -linear category, let $\mathcal{P}$ be a full subcategory of $\mathcal{M}$, and let $\Sigma : \mathcal{M} \rightarrow \mathcal{M}$ be a $\mathbb{k}$ -linear right exact endofunctor. A Σ -twisted trace on $\mathcal{P}$ is a family

$$\mathbf{t}_\bullet = \{\mathbf{t}_P : \mathrm{Hom}_{\mathcal{M}}(P, \Sigma(P)) \rightarrow \mathbb{k}\}_{P \in \mathcal{P}}$$

of $\mathbb{k}$ -linear maps parametrized by the objects of $\mathcal{P}$ such that the following Σ -cyclicity condition is satisfied:

Σ -cyclicity: For all objects $P, Q \in \mathcal{P}$ and morphisms $f : P \rightarrow Q$ and $g : Q \rightarrow \Sigma(P)$ in $\mathcal{M}$, the following equation holds:

$$\mathbf{t}_P\left(P \xrightarrow{f} Q \xrightarrow{g} \Sigma(P)\right) = \mathbf{t}_Q\left(Q \xrightarrow{g} \Sigma(P) \xrightarrow{\Sigma(f)} \Sigma(Q)\right). \quad (3.1)$$

We say that a Σ -twisted trace $\mathbf{t}_\bullet$ on $\mathcal{P}$ is *non-degenerate* if the pairing

$$\mathrm{Hom}_{\mathcal{M}}(M, \Sigma(P)) \times \mathrm{Hom}_{\mathcal{M}}(P, M) \rightarrow \mathbb{k}, \quad (f, g) \mapsto \mathbf{t}_P(fg)$$

is non-degenerate for all objects $M \in \mathcal{M}$ and $P \in \mathcal{P}$.

Important that Σ is not an endofunctor of $\mathcal{P}$ because a major case of interest is $\mathcal{P} = \mathrm{Proj}(\mathcal{M})$ and Σ the Nakayama functors (that takes proj obj to inj obj).

We write $\mathrm{Tr}(\mathcal{M})$ for the class of Σ -twisted traces on $\mathrm{Proj}(\mathcal{M})$.

Theorem 10.1 (SS). *There is a bijection*

$$\mathrm{Nat}(\Sigma, \mathbf{N}^r) \xrightarrow{\cong} \mathrm{Tr}(\mathcal{M})$$

via the “canonical Nakayama twisted trace”. Under this bijection, natural isomorphisms correspond to non-degenerate Σ -twisted traces.

10.2 Module-compatibility of twisted traces

Let $\mathcal{C}$ be a rigid monoidal category and let $\mathcal{M}$ be a finite category and a right $\mathcal{C}$ -module category. We write

$$\begin{aligned} \mathrm{cl}_{M, N|X} : \mathrm{Hom}_{\mathcal{M}}(M \otimes X, N \otimes X^{**}) &\rightarrow \mathrm{Hom}_{\mathcal{M}}(M, N) \quad (M, N \in \mathcal{M}, X \in \mathcal{C}), \\ \mathrm{cl}_{M, N|X}(f) &= (\mathrm{id}_N \otimes \mathrm{ev}_{X^*}) \circ (f \otimes \mathrm{id}_{X^*}) \circ (\mathrm{id}_M \otimes \mathrm{coev}_X). \end{aligned}$$

The linear map $\mathrm{cl}_{M, N|X}$ is called the partial trace in literature (at least when $\mathcal{M} = \mathcal{C}$ and $\mathcal{C}$ is pivotal), however, we call it the *closing operator* to prevent the proliferation of trace-like terms.

Now let $\Sigma : \mathcal{M} \rightarrow \mathcal{M}$ be a $(-)^{**}$ -twisted right exact $\mathcal{C}$ -module functor. In other words, Σ is a $\mathbb{k}$ -linear right exact endofunctor on $\mathcal{M}$ equipped with a natural isomorphism

$$\Psi_{M, X}^\Sigma : \Sigma(M) \otimes X^{**} \rightarrow \Sigma(M \otimes X) \quad (M \in \mathcal{M}, X \in \mathcal{C})$$

such that the equations $\Psi_{M, 1}^\Sigma = \mathrm{id}_{\Sigma(M)}$ and $\Psi_{M, X \otimes Y}^\Sigma = \Psi_{M \otimes X, Y}^\Sigma \circ (\Psi_{M, X}^\Sigma \otimes \mathrm{id}_{Y^{**}})$ hold for all objects $M \in \mathcal{M}$ and $X, Y \in \mathcal{C}$. Now we clarify what ‘compatible’ means:

Definition 10.2. A Σ -twisted trace $\mathbf{t}_\bullet$ on $\text{Proj}(\mathcal{M})$ is said to satisfy *the right partial trace property*, or to be *compatible with the right $\mathcal{C}$ -module structure* (or *module-compatible* for short) if the equation

$$\mathbf{t}_P(\text{cl}_{P, \Sigma(P)|X}(f)) = \mathbf{t}_{P \otimes X}(\Psi_{P, X}^\Sigma \circ f) \quad (5.3)$$

holds for all $P \in \text{Proj}(\mathcal{M})$, $X \in \mathcal{C}$ and $f \in \text{Hom}_{\mathcal{M}}(P \otimes X, \Sigma(P) \otimes X^{**})$.

Theorem 10.3 (SS). *Under the previous bijection*

$$\text{Nat}(\Sigma, \mathbf{N}^r) \xrightarrow{\cong} \text{Tr}(\mathcal{M}),$$

natural transformations which are module-compatible (ie natural transformations of module functors) correspond to Σ -twisted traces satisfying the right partial trace property. Natural isomorphisms of module functors correspond to non-degenerate traces.

10.3 Uniqueness

Let $\mathcal{M}$ be an **indecomposable** (ie not equivalent to a direct sum of categories) exact right $\mathcal{C}$ -module category, and let $\Sigma : \mathcal{M} \rightarrow \mathcal{M}$ be a $(-)^{**}$ -twisted right exact $\mathcal{C}$ -module functor.

Then there exists a non-degenerate module-compatible Σ -twisted trace on $\text{Proj}(\mathcal{M})$ if and only if $\Sigma \cong \mathbf{N}_{\mathcal{M}}$ as right $\mathcal{C}$ -module functors. If this is the case, a non-zero module-compatible Σ -twisted trace on $\text{Proj}(\mathcal{M})$ is unique up to scalar multiple and every such trace is non-degenerate.

10.4 Left/right modified traces

Let $\mathcal{C}$ be a pivotal finite tensor category. The identity of $\mathcal{C}$ becomes a $(-)^{**}$ -twisted left/right $\mathcal{C}$ -module functor thanks to the pivotal structure.

Definition 10.4. A *left* (resp. *right*) *modified trace* on $\text{Proj}(\mathcal{C})$ is an $\text{id}_{\mathcal{C}}$ -twisted trace on $\text{Proj}(\mathcal{C})$ compatible with the left (resp. right) $\mathcal{C}$ -module structure. We denote by $\mathcal{T}_{\mathcal{C}}^{(\ell)}$ and $\mathcal{T}_{\mathcal{C}}^{(r)}$ the spaces of left and right modified traces on $\text{Proj}(\mathcal{C})$, respectively.

In more detail: a left/right modified trace on $\text{Proj}(\mathcal{C})$ is a family

$$\mathbf{t}_\bullet = \{\mathbf{t}_P : \text{End}_{\mathcal{C}}(P) \rightarrow \mathbb{k}\}_{P \in \mathcal{P}}$$

of $\mathbb{k}$ -linear maps parametrized by the objects of $\mathcal{P}$ such that

1) Cyclicity: For all objects $P, Q \in \mathcal{P}$ and morphisms $f : P \rightarrow Q$ and $g : Q \rightarrow P$ in $\mathcal{C}$, the following equation holds:

$$\mathbf{t}_Q(f \circ g) = \mathbf{t}_P(g \circ f)$$

2) Left (resp. right) partial trace property: If $P \in \text{Proj}(\mathcal{C})$ and $X \in \mathcal{C}$,

$$\mathbf{t}_{V \otimes P}(f) = \mathbf{t}_P(\text{cl}_{P|X}(f)) \quad , \quad (\text{resp. } \mathbf{t}_{P \otimes V}(f) = \mathbf{t}_P(\text{cl}_{X|P}(f)))$$

Key observation: This closure operator, ie the left/right categorical aka pivotal trace, is a left/right modified trace. The issue is the following:

Proposition 10.5. *Let $\mathcal{C}$ be a pivotal finite tensor category over a field $\mathbb{k}$. The pivotal trace tr is non-zero on $\text{Proj}(\mathcal{C})$ if and only if $\mathcal{C}$ is semisimple.*

Proof. Recall that in a finite tensor category there is a distinguished projective object P_1 , the projective cover of the unit object $\mathbf{1}$. A fundamental theorem of finite tensor categories states that $\mathcal{C}$ is semisimple if and only if $\mathbf{1}$ is projective, that is, $P_1 \cong \mathbf{1}$.

We first observe that for any projective object $P \in \mathcal{C}$, the pivotal trace of any endomorphism $f \in \text{End}_{\mathcal{C}}(P)$ factors through P_1 . Indeed, by definition of the pivotal trace, $\text{tr}_P(f)$ is the composite

$$\mathbf{1} \xrightarrow{\text{coev}_P} P \otimes P^* \xrightarrow{f \otimes \text{id}_{P^*}} P \otimes P^* \xrightarrow{\widetilde{\text{ev}}_P} \mathbf{1},$$

where $\widetilde{\text{ev}}_P$ denotes the pivotal evaluation. This composite is a morphism $\mathbf{1} \rightarrow \mathbf{1}$, hence a scalar, and it factors as

$$\mathbf{1} \xrightarrow{f} P_1 \xrightarrow{g} \mathbf{1}$$

because P_1 is the projective cover of $\mathbf{1}$: the morphism $\mathbf{1} \xrightarrow{\text{coev}_P} P \otimes P^*$ lands in a projective object (since P projective implies $P \otimes P^*$ projective), and any morphism from a projective object into $\mathbf{1}$ factors through the projective cover $P_1 \rightarrow \mathbf{1}$. Thus $\text{tr}_P(f)$ factors through P_1 for every projective P and every f .

($\Leftarrow$) Suppose $\mathcal{C}$ is semisimple, so $P_1 \cong \mathbf{1}$. Then the factorisation

$$\mathbf{1} \xrightarrow{f} P_1 \xrightarrow{g} \mathbf{1}$$

through which $\text{tr}_P(\text{id}_P)$ factors has $P_1 = \mathbf{1}$, so $g \circ f$ is simply a scalar multiple of $\text{id}_{\mathbf{1}}$. In a semisimple pivotal category every simple object L satisfies $\dim(L) = \text{tr}(\text{id}_L) \neq 0$, since $\text{End}(L) \cong \mathbb{k}$ by Schur's lemma and the Hom-pairing is non-degenerate. As the trace is additive on direct sums, it follows that $\text{tr}_P \neq 0$ for every non-zero projective P .

($\Rightarrow$) We prove the contrapositive. Suppose $\mathcal{C}$ is not semisimple, so $P_1 \not\cong \mathbf{1}$, meaning the projective cover map $\pi : P_1 \rightarrow \mathbf{1}$ does not split. We claim that for every projective object P , the trace $\text{tr}_P(\text{id}_P) = 0$, that is, the composite

$$\mathbf{1} \xrightarrow{\text{coev}_P} P \otimes P^* \xrightarrow{\widetilde{\text{ev}}_P} \mathbf{1}$$

is zero. By the factorisation observation above, this composite factors as

$$\mathbf{1} \xrightarrow{f} P_1 \xrightarrow{g} \mathbf{1}$$

for some morphisms f and g . Since $\mathcal{C}$ is not semisimple, the sequence

$$0 \rightarrow \text{rad}(P_1) \rightarrow P_1 \xrightarrow{\pi} \mathbf{1} \rightarrow 0$$

is a non-split exact sequence with $\text{rad}(P_1) \neq 0$. Any morphism $g : P_1 \rightarrow \mathbf{1}$ is a scalar multiple of π , and any morphism $f : \mathbf{1} \rightarrow P_1$ would split π if $g \circ f \neq 0$. Since π does not split, there is no morphism $f : \mathbf{1} \rightarrow P_1$ with $g \circ f \neq 0$, and therefore $g \circ f = 0$ for all such f and g . Hence

$$\text{tr}_P(\text{id}_P) = g \circ f = 0$$

for every projective object P , and by linearity $\text{tr}_P \equiv 0$ on all of $\text{End}_{\mathcal{C}}(P)$. $\square$

Question: when does a pivotal finite tensor category admit modified traces on $\text{Proj}(\mathcal{C})$?

Theorem 10.6. *For a pivotal finite tensor category $\mathcal{C}$, the following assertions are equivalent:*

1. $\mathcal{C}$ is unimodular, that is, there is an isomorphism $\alpha_{\mathcal{C}} \cong \mathbf{1}$ in $\mathcal{C}$.
2. There is a non-zero right modified trace on $\text{Proj}(\mathcal{C})$.
3. There is a non-zero left modified trace on $\text{Proj}(\mathcal{C})$.

If these equivalent conditions are satisfied, then we have $\dim_{\mathbb{k}} \mathcal{T}_{\mathcal{C}}^{(r)} = \dim_{\mathbb{k}} \mathcal{T}_{\mathcal{C}}^{(\ell)} = 1$ and every non-zero left or right modified trace on $\text{Proj}(\mathcal{C})$ is non-degenerate.

10.5 Two-sided modified traces

Let $\mathcal{C}$ be a pivotal finite tensor category. A *two-sided modified trace* on $\text{Proj}(\mathcal{C})$ is an element of $\mathcal{T}_{\mathcal{C}}^{(\ell r)} := \mathcal{T}_{\mathcal{C}}^{(\ell)} \cap \mathcal{T}_{\mathcal{C}}^{(r)}$.

Recall that a pivotal finite tensor category is *spherical* (according to Douglas-Schommer-Pries-Snyder) if it is unimodular, ie $\alpha \cong \mathbf{1}$, and the trivialisation $(-)^{\vee\vee\vee\vee} \cong \text{Id}$ coming from the Radford isomorphism coincides with the trivialisation $(-)^{\vee\vee\vee\vee} \cong (-)^{\vee\vee} \cong \text{Id}$ induced by the pivotal structure.

Corollary 10.7. *Let $\mathcal{C}$ be a pivotal finite tensor category. Then*

$$\exists \text{ non-zero two-sided modified trace on } \text{Proj}(\mathcal{C}) \iff \mathcal{C} \text{ spherical}$$

If this is the case, a non-zero two-sided modified trace on $\text{Proj}(\mathcal{C})$ is unique up to scalar multiple and every such trace is non-degenerate.

If $\mathcal{C}$ is ribbon, the situation is even more interesting:

Corollary 10.8. *Let $\mathcal{C}$ be a ribbon finite tensor category. Then TFAE:*

1. $\mathcal{C}$ is spherical,
2. $\mathcal{C}$ is unimodular,
3. There exists a non-zero two-sided modified trace on $\text{Proj}(\mathcal{C})$ (in this case such a mod trace is always non-degenerate by the corollary above).

Corollary 10.9. *Any modular tensor categories admits a non-deg two-sided modified trace.*

Reason? Factorisable braided finite tensor category $\Rightarrow$ Unimodular by [EGNO, 8.10.10], and the corollary above.

11 Modular functors

11.1 Bakalov-Kirillov's modular functors

Definition 11.1 (Naïve version). Let Mnfd_d be the category with objects homeomorphism classes of d -manifolds and arrows only endomorphisms given by $\text{End}(M) = \text{MCG}(M)$. This is a symmetric monoidal category with respect to disjoint union.

A d -dimensional modular functor is a strong monoidal functor

$$\tau : \text{Mnfd}_d \rightarrow \text{vect}_{\mathbb{k}}$$

with the normalisation property that $\tau(S^d) = \mathbb{k}$.

Ie this is a system of MCG representations.

Now we put $d = 2$. Ideally one would like to glue surfaces and say that the MCG representations are coherent with respect to glueing. For this we need *extended surfaces*, ie surfaces with parametrised boundary (=circle) components. Homeomorphisms between extended surfaces are supposed to preserve the boundary parametrisations.

Definition 11.2. Let $\mathcal{C}$ be an abelian category over a field k , and let $R = \bigoplus V_i \boxtimes V_i^*$ be a symmetric object in $\text{ind-}\mathcal{C}^{\boxtimes 2}$ (see Section 2.4). Then a $\mathcal{C}$ -*extended modular functor* is the following collection of data:

- (i) To every extended surface Σ is assigned a polylinear functor

$$\tau(\Sigma): \mathcal{C}^{\boxtimes \pi_0(\partial\Sigma)} \longrightarrow \mathbf{vect}_{\mathbb{k}},$$

where $\pi_0(\partial\Sigma)$ is the set of boundary components (or punctures, depending on the point of view) of Σ . In other words, for every choice of objects $W_a \in \mathcal{C}$ attached to every boundary component of Σ (so, a runs through the set of connected components of $\partial\Sigma$) is assigned a finite-dimensional vector space $\tau(\Sigma; \{W_a\})$, and this assignment is functorial in W_a .

- (ii) To every homeomorphism $f: \Sigma \xrightarrow{\sim} \Sigma'$ is assigned a functorial isomorphism $f_*: \tau(\Sigma) \xrightarrow{\sim} \tau(\Sigma')$, which depends only on the isotopy class of f .
- (iii) Functorial isomorphisms $\tau(\emptyset) \xrightarrow{\sim} k$, $\tau(N_1 \sqcup N_2) \xrightarrow{\sim} \tau(N_1) \otimes \tau(N_2)$.
- (iv) **Gluing isomorphism:** Let $(\partial\Sigma)_\alpha, (\partial\Sigma)_\beta$ be two different boundary components of Σ , and let $\pi_\alpha: (\partial\Sigma)_\alpha \xrightarrow{\sim} S^1$, π_β be their parameterizations. Define a new surface $\sqcup_{\alpha,\beta}\Sigma$ by identifying all points $p \in (\partial\Sigma)_\alpha$ with $\pi_\beta^{-1}(-\pi_\alpha(p)) \in (\partial\Sigma)_\beta$. Let us write $R = \bigoplus_i A_i \boxtimes B_i$. Then we have a functorial isomorphism

$$G_{\alpha,\beta}: \bigoplus_i \tau(\Sigma; \{W_a\}, A_i, B_i) \xrightarrow{\sim} \tau(\sqcup_{\alpha,\beta}\Sigma; \{W_a\}), \quad (5.1.1)$$

where A_i, B_i correspond to the boundary components $(\partial\Sigma)_\alpha, (\partial\Sigma)_\beta$, respectively. In the notation of Section 2.4, this can be written as:

$$G_{\alpha,\beta}: \tau(\Sigma; \{W_a\}, R^{(1)}, R^{(2)}) \xrightarrow{\sim} \tau(\sqcup_{\alpha,\beta}\Sigma; \{W_a\}). \quad (5.1.2)$$

The above data have to satisfy the following axioms:

Multiplicativity: $(fg)_* = f_*g_*$, $\text{id}_* = \text{id}$.

Functoriality: All isomorphisms in parts (iii), (iv) above are functorial in Σ .

Compatibility: All isomorphisms in parts (iii), (iv) above are compatible with each other.

Symmetry of gluing: After the identification $R \simeq R^{\text{op}}$, we have $G_{\alpha,\beta} = G_{\beta,\alpha}$.

Normalization: $\tau(S^2) = k$.

As before, we leave it to the reader to write the explicit statements of all functoriality and compatibility axioms, taking as an example the definitions in Section 4.2. Let us give two of them.

Compatibility of gluing with disjoint union: Let Σ', Σ'' be two extended surfaces, and $(\partial\Sigma')_\alpha, (\partial\Sigma')_\beta$ be two different boundary components of Σ' . Then the following diagram is commutative:

$$\begin{array}{ccc} \tau(\Sigma' \sqcup \Sigma''; \{W'_a\}, R^{(1)}, R^{(2)}, \{W''_b\}) & \xrightarrow{G_{\alpha,\beta}} & \tau(\Sigma; \{W'_a\}, \{W''_b\}) \\ \downarrow & & \downarrow \\ \tau(\Sigma'; \{W'_a\}, R^{(1)}, R^{(2)}) \otimes \tau(\Sigma''; \{W''_b\}) & \xrightarrow{G_{\alpha,\beta} \otimes \text{id}} & \tau(\sqcup_{\alpha,\beta}\Sigma'; \{W'_a\}) \otimes \tau(\Sigma''; \{W''_b\}) \end{array}$$

where the vertical arrows are the isomorphisms from part (iii), and $\Sigma = \sqcup_{\alpha,\beta}(\Sigma' \sqcup \Sigma'') \simeq (\sqcup_{\alpha,\beta}\Sigma') \sqcup \Sigma''$ (with the homeomorphism isotopic to the identity).

Associativity of gluing: If $(\partial\Sigma)_\alpha, \dots, (\partial\Sigma)_\delta$ are four different boundary components of Σ , then:

$$G_{\alpha,\beta}G_{\gamma,\delta} = G_{\gamma,\delta}G_{\alpha,\beta}. \quad (5.1.3)$$

In more detail, the following diagram is commutative:

$$\begin{array}{ccc}
\tau(\Sigma; \{W_a\}, R'^{(1)}, R'^{(2)}, R''^{(1)}, R''^{(2)}) & \xrightarrow{G_{\alpha, \beta}} & \tau(\sqcup_{\alpha, \beta} \Sigma; \{W_a\}, R''^{(1)}, R''^{(2)}) \\
\downarrow G_{\gamma, \delta} & & \downarrow G_{\gamma, \delta} \\
\tau(\sqcup_{\gamma, \delta} \Sigma; \{W_a\}, R'^{(1)}, R'^{(2)}) & \xrightarrow{G_{\alpha, \beta}} & \tau(\Sigma'; \{W_a\})
\end{array}$$

where R', R'' are two copies of R , and $\Sigma' = \sqcup_{\alpha, \beta} \sqcup_{\gamma, \delta} \Sigma \simeq \sqcup_{\gamma, \delta} \sqcup_{\alpha, \beta} \Sigma$ (with the homeomorphism isotopic to the identity).

Remark Let us give a modification of Definition 5.1.13 which will be useful later. Instead of extended surfaces Σ , consider pairs (Σ, φ) where Σ is an extended surface and $\varphi: \pi_0(\partial\Sigma) \xrightarrow{\sim} S$ is a labeling of the boundary components of Σ with a finite set S . Then the functor $\tau(\Sigma): \mathcal{C}^{\boxtimes \pi_0(\partial\Sigma)} \rightarrow \mathcal{Vec}_f$ from Definition 5.1.13(i) gives rise to a functor $\tau(\Sigma, \varphi): \mathcal{C}^{\boxtimes S} \rightarrow \mathcal{Vec}_f$; explicitly, $\tau(\Sigma, \varphi) = \tau(\Sigma) \circ \varphi_*^{-1}$, where $\varphi_*: \mathcal{C}^{\boxtimes \pi_0(\partial\Sigma)} \rightarrow \mathcal{C}^{\boxtimes S}$ is the functor induced from φ . It is clear that for any bijection $\psi: S \xrightarrow{\sim} S'$, we have $\tau(\Sigma, \psi\varphi) = \tau(\Sigma, \varphi) \circ \psi_*^{-1}$. So a modular functor can be defined as a collection of functors $\tau(\Sigma, \varphi)$ satisfying obvious equivariance conditions.

11.2 Open modular functors

This is a simpler version of modular functor, where we do not glue along boundary components but along parametrised marked intervals.

The *open surface operad* $\mathcal{O}[\cdot, \cdot, \cdot]$ is the groupoid-valued operad whose groupoid $\mathcal{O}(n)$ of operations of arity n has as objects compact, oriented surfaces with at least one boundary component per connected component together with $n+1$ parametrized intervals embedded in their boundary; the morphisms are mapping classes (isotopy classes of diffeomorphisms preserving the orientation and the parametrized intervals), and operadic composition is given by gluing surfaces along the parametrized intervals. The open surface operad is in fact a *modular operad* in the sense of [?].


Definition 11.3. An *open modular functor*, also referred to as a *two-dimensional categorified open topological field theory*, is defined as a modular $\mathcal{O}$ -algebra that (at least in the context of this article) takes values in the symmetric monoidal bicategory $\mathbf{Lex}^f$ of finite categories, left exact functors and natural transformations.

Such a structure is exactly the mathematical description of the monodromy data of an *open conformal field theory*; this is informed by the notions [?, ?, ?] and defined in this form [?, Section 8] in the context of the modular microcosm principle.

Let us unpack the definition: a $\mathbf{Lex}^f$ -valued open modular functor $\mathfrak{B}$ consists of the data of a finite category $\mathcal{A}$ (ie the object underlying the algebra itself) and a rule that associates to every surface Σ with at least one boundary component per connected component with n marked boundary intervals a left exact functor (recall (70))

$$\mathfrak{B}(\Sigma; -) : \mathcal{A}^{\boxtimes n} \rightarrow \mathbf{vect},$$

ie given n objects $X_1, \dots, X_n \in \mathcal{A}$ attached to the boundary intervals, we get a vector space $\mathfrak{B}(\Sigma; X_1, \dots, X_n)$, the so-called *space of (open) conformal blocks*, which carries an action of the mapping class group of the surface and is compatible with gluing along parametrized intervals. By definition, the space of conformal blocks is left exact in the boundary labels.



Some boundary components may have no parametrized boundary intervals; they are called *free boundary circles*.

According to [?, Theorem 2.2], a $\mathbf{Lex}^f$ -valued open modular functor is, by evaluation on disks with marked intervals in their boundary, equivalent to the datum of a *pivotal Grothendieck-Verdier category* in $\mathbf{Lex}^f$ in the sense of [?]. In particular, the underlying pivotal finite tensor category of a finite ribbon category $\mathcal{A}$ extends to a unique open modular functor that we denote by $\mathcal{A}_!$.