

Lecture 1: Random variables, probability distributions and sampling EMAT30007 Applied Statistics

Nikolai Bode

Department of Engineering Mathematics



Outline of the lecture

In this lecture you will learn:

- How to use Random Variables to model empirical data.
- How to plot empirical probability density (PDF) and cumulative distribution (CDF) functions.
- ✓ How to compute probability distributions of functions of Random Variables.
- We How to generate synthetic data sampling from a Random Variable.



Lecture 1, Topic 1 – Random variables and probability distributions

@ntent covered

- Random variables
- Probability mass function (PMF)/Probability density function (PDF) and how to compute it for data
- Cumulative distribution function (CDF) and how to compute it for data
- Common probability distributions

What is statistics?

Statistics aims to describe and interpret empirical data using the mathematical objects of probability theory.

A statistical model is a mathematical representation of a real world process that has some degree randomness.

Example: Dice roll

We roll a 6-sided dice and write down each outcome in a list:

dice rolls
$$=\{ \textcircled{2}, \textcircled{2}, \textcircled{3}, \textcircled{4}, 5, 2, 6 \}$$

An empirical data set is a collection of outcomes of a specific real-world process or experiment (a dice roll, in this case).

We imagine that each outcome has a given *probability* to be observed.

In statistics, we interpret an empirical data set as a collection of outcomes generated by a Random Variable.



University of Sample Space = Collection of Applied Statistics - Lecture 1

BRISTOL

Spring Semester

Random Variables

A Random Variable (RV) is a function that maps any possible outcome of an experiment (sample space) to a number.

Example: if D is a RV for a dice roll, then $D(\odot) = 1, \dots, D(\boxdot) = 6$.

A random variable has an associated probability distribution that provides the probability of occurrence of all possible outcomes.

Example: if D is a fair dice, then $P_D(1) = \cdots = P_D(6) = 1/6$.

Example: Toss of a fair coin

Possible outcomes (sample space): $\Omega = \{\text{'head', 'tail'}\}.$

Random Variable:
$$X(\omega) = \begin{cases} 1 & \text{if } \omega = \text{`head'} \\ 0 & \text{if } \omega = \text{`tail'} \end{cases}$$


Probability distribution: $P_X(x) = p^x (1-p)^{1-x} = \begin{cases} p & \text{if } x = 1\\ 1-p & \text{if } x = 0 \end{cases}$

where p = 1/2 is the probability of 'head' ('head' and 'tail' are equally likely).



Discrete Random Variables

A discrete RV maps outcomes to discrete (natural) numbers.





Continuous Random Variables

A continuous RV maps outcomes to continuous (real) numbers.

Example: Diameter of a tennis ball

Sample space (possible outcomes): any real number larger than zero.

$$\Omega = [0, +\infty] = \mathbb{R}^+$$


RV: D is the measured diameter in centimetres.

Probability Density Function (PDF):

 $D \sim \mathsf{Gamma}(\alpha = 6540, \theta = 0.001)$

Gamma distribution:

Gamma distribution:
$$P_D(d;\alpha,\theta) = \frac{1}{\Gamma(\alpha)\theta^{\alpha}}d^{\alpha-1}e^{-d/\theta}$$






Cumulative Distribution Function (CDF)

The Cumulative Distribution Function (CDF) of a random variable X, $F_X(x)$, is the probability to get an outcome with value less or equal to x:


$$F_X(x) = Prob(X \le x) = \sum_{y \le x} P_X(y) \qquad \text{for discrete RVs} \qquad \text{(1)}$$

$$F_X(x) = \int_{-\infty}^x P_X(y) dy \qquad \text{for continuous RVs} \qquad \text{(2)}$$

Example: Sum of two dices



Example: Diameter of a tennis ball



$$F_D(d; \alpha, \theta) = \int_0^d \frac{1}{\Gamma(\alpha)\theta^{\alpha}} y^{\alpha - 1} e^{-y/\theta} dy$$



Properties of the CDF and PDF

- the CDF is a non decreasing function. (increases monitorically)
- the maximum value is 1 and the minimum is 0.
- $\bigvee Prob(X>x)=1-F_X(x)$ is the Complementary CDF (CCDF).
- From OF to PDF


 If the CDF of a continuous RV is differentiable, then the PDF is given by

$$P_X(x) = \frac{dF_X(x)}{dx} = \lim_{dx \to 0} \frac{F_X(x + dx) - F_X(x)}{dx}$$
 (3)

Example: Uniform distribution — equal possibility The continuous RV U defined over the interval

 $u \in [a, b]$ has CDF $F_U(u) = (u - a)/(b - a)$ and its PDF is the Uniform distribution:

$$P_U(u) = F'_U(u) = 1/(b-a).$$





How to compute the empirical CDF from data

Plot of the empirical CDF

- 1. Load data in a one-dim array, d.
- 2. Sort data in ascending order, x.
- 3. Create an array, y, of increasing integers from 1 to the length of x.
- 4. Divide each element of y by the length of x, so that the max of y is 1.
- 5. If there are (x_i, y_i) pairs with identical x-value, keep only the pair with the largest y-value. $y = F_X(x)$.

- 1. $d = \begin{bmatrix} 2, & 7, & 2, & 1, & 5 \end{bmatrix}$ 2. $x = \begin{bmatrix} 1, & 2, & 2, & 5, & 7 \end{bmatrix}$
- 3. $x = \begin{bmatrix} 1, & 2, & 2, & 5, & 7 \end{bmatrix}$ $y = \begin{bmatrix} 1, & 2, & 3, & 4, & 5 \end{bmatrix}$
- 4. $x = \begin{bmatrix} 1, & 2, & 2, & 5, & 7 \end{bmatrix}$ $y = \begin{bmatrix} 0.2, & 0.4, & 0.6, & 0.8, & 1 \end{bmatrix}$
- 5. $x = \begin{bmatrix} 1, & ., & 2, & 5, & 7 \end{bmatrix}$ $y = \begin{bmatrix} 0.2, & ., & 0.6, & 0.8, & 1 \end{bmatrix}$

In MATLAB, use stairs(x, y) or cdfplot(d) to plot the empirical CDF.

Spring Semester

How to compute the empirical PDF from data

Plot of the empirical PDF, using the definition in Eq. (3)

- 1. Divide the range of data d, $[\min(d), \max(d)]$, into bins, with bin edges b.
- 2. Compute the mid-point of each bin, $x_i = (b_{i+1} + b_i)/2.$

Count the number of data elements.

- that fall in each bin, $c_i = |\{d_i : b_i \le d_i < b_{i+1}\}|.$
- 4. Normalise. Rescale each element of c: $y_i = c_i/(|d|(b_{i+1} b_i))$, so that the PDF integral is 1. $y = P_X(x)$.

1. d = [2, 7, 2, 1, 5]b = [1, 3, 5, 7, 9]

 $2. \quad x = [2, 4, 6, 8]$

3. $x = \begin{bmatrix} 2, & 4, & 6, & 8 \end{bmatrix}$ $c = \begin{bmatrix} 3, & 0, & 1, & 1 \end{bmatrix}$

4. $x = \begin{bmatrix} 2, & 4, & 6, & 8 \end{bmatrix}$ $y = \begin{bmatrix} 0.3, & 0, & 0.1, & 0.1 \end{bmatrix}$

In MATLAB, use <u>histcounts</u> or <u>histogram(d, 'normalization', 'pdf')</u> to plot the empirical PDF of data in array d (use option 'BinEdges' to specify bin edges).

Common Random Variables

- Uniform (continuous outcomes, all equally likely)
- Bernoulli (coin toss)
- Binomial (# of 'heads' in n tosses)
- \blacktriangleright Multinomial (# of '1's, '2's, ..., '6's in n dice rolls)
- Exponential (time between consecutive independent arrivals/events)
- Normal (sum of many independent RVs)

- Beta (distribution of probabilities)
- Pareto (distribution of wealth, city populations)
- https://en.wikipedia.org/wiki/List_of_probability_distributions



Lecture 1, Topic 2 – Working with Random Variables

©ntent covered

- Expectation of RVs
- Functions of RVs
- Sampling from almost any RV

Expectation of RVs

The expectation or expected value, E(X), of Random Variable X with PDF $P_X(x)$ is defined as

$$E(X) \equiv \int P_X(x) x \, dx$$
 or $E(X) \equiv \sum_x P_X(x) x$ (4)

for continuous and discrete RVs. ${\cal E}(X)$ is also called first moment or mean of ${\cal X}.$

Properties of the expectation:

- Linearity of expectation: for any two RVs X and Y and any constant a E(X+Y)=E(X)+E(Y) and E(aX)=aE(X).
- Expectation of a product of **independent** RVs: if X and Y are <u>independent</u> RVs then $E(X \cdot Y) = E(X) \cdot E(Y)$. This is not true in general.
- Expectation of a function of a RV: $E(g(X)) = \int P_X(x) \, g(x) \, dx$. The sprad of volumes



Function of a Random Variable

Let X be a continuous RV with CDF $F_X(x)$ and PDF $P_X(x)$. If g is a strictly increasing or decreasing function with inverse $g^{-1}=h$, then the RV Y=g(X) has the following CDF and PDF (if F_X and h differentiable):

$$F_Y(y) = \begin{cases} F_X(h(y)) & \text{if } h \text{ and } g \text{ increasing} \\ 1 - F_X(h(y)) & \text{if } h \text{ and } g \text{ decreasing} \end{cases}$$
 (5)

$$P_Y(y) = P_X(h(y)) \cdot \left| \frac{dh(y)}{dy} \right|$$
 (6)

The domain of the new RV is transformed too: $Y \in [g(x_{min}), g(x_{max})]$ if g increasing or $Y \in [g(x_{max}), g(x_{min})]$ if g decreasing.




Example: Surface area of a tennis ball

Recall the RV D for the diameter of a tennis ball, which has a Gamma distribution: $P_D(d;\alpha,\theta)=\frac{1}{\Gamma(\alpha)\theta^\alpha}d^{\alpha-1}e^{-d/\theta}$. We can define the RV $S=\pi D^2$ for the surface of a tennis ball. What is the PDF of S?

We can apply the function $s=\pi d^2$ to our diameter data, obtaining the corresponding surface areas, and compute the empirical PDF (blue histogram).

We can also apply Eq. (6) and obtain $P_S(s) = P_D(\sqrt{s/\pi}) |\frac{1}{2\pi} \sqrt{\pi/s}|$ (red curve).



$$5 = g(d) = 11d^{2}$$

 $h'(s) = \sqrt{\frac{5}{11}}$
 $h'(s) = \frac{1}{2\pi} \sqrt{\frac{\pi}{5}}$

Sampling from a Random Variable

Sampling from a RV X with PDF P_X means to generate numbers such that the probability to generate number x is $P_X(x)$.

Being able to sample from a RV allows us to simulate real-world processes.

Computers have (pseudo) random numbers generators that produce numbers that are uniformly distributed in [0,1).

How can we use a computer's uniform random number generator to sample from a RV with a given distribution?

Example: Simulate a fair coin.

A fair coin can be modelled as a RV X with PDF $P_X(0)=1/2$ and $P_X(1)=1/2$, so we want to generate 0 ('tail') with probability 1/2 and 1 ('head') with probability 1/2.

We can sample a random number u from U(0,1), the computer's uniform random number generator: u will be smaller than 0.5 with probability 1/2 and larger than 0.5 with probability 1/2. To simulate the RV X we can transform u into x such that: x=0 if $0 \le u < 0.5$ and x=1 if $0.5 \le u < 1$.



Inverse Probability Integral Transform

The Inverse Probability Integral Transform (IPIT) is a method to transform uniformly distributed random numbers into numbers with a different distribution.

The IPIT is a sampling method based on Eq. (5) for the function of a RV: to sample numbers from RV X we only need to know the inverse of its CDF, F_X^{-1} :

Sampling using the Inverse Probability Integral Transform (IPIT)

- 1. Generate a number u from the uniform distribution U(0,1).
- 2. Compute $x = F_{\mathbf{v}}^{-1}(u)$


The CDF of X will be F_X because if $X = F_X^{-1}(U)$ then from Eq. (5) $F_X(x) = F_U((F_X^{-1})^{-1}(x)) = F_U(F_X(x)) = F_X(x)$.

The inverse of the CDF, F_X^{-1} , is called the quantile function.

Example: Exponential distribution


To sample from $P_X(x)=ae^{-ax}$ with CDF $F_X(x)=1-e^{-ax}$, compute the inverse of $u=F_X(x)=1-e^{-ax}$, obtaining $x=F_Y^{-1}(u)=-\ln(1-u)/a$.

Visual intuition for the IPIT




https://en.wikipedia.org/wiki/Inverse_transform_sampling





IPIT for the Exponential distribution, $X \sim Exp(1)$.



IPIT for the fair coin example, $X \sim Bernoulli(0.5)$.