

ON PROPER SPLINTERS IN POSITIVE CHARACTERISTIC

JOHANNES KRAH AND CHARLES VIAL

ABSTRACT. While the splinter property is a local property for Noetherian schemes in characteristic zero, Bhatt observed that it imposes strong conditions on the global geometry of proper schemes in positive characteristic. We show that if a proper scheme over a field of positive characteristic is a splinter, then its Nori fundamental group scheme is trivial and its Kodaira dimension is negative. In another direction, Bhatt also showed that any splinter in positive characteristic is a derived splinter. We ask whether the splinter property is a derived invariant for projective varieties in positive characteristic and give a positive answer for normal Gorenstein projective varieties with big anticanonical divisor. We also show that global F -regularity is a derived invariant for normal Gorenstein projective varieties in positive characteristic.

1. INTRODUCTION

A Noetherian scheme X is a *splinter* if for all finite surjective morphisms $f: Z \rightarrow X$ the map $\mathcal{O}_X \rightarrow f_*\mathcal{O}_Z$ splits in the category of coherent \mathcal{O}_X -modules. The direct summand conjecture, now a theorem due to André [And18], stipulates that any regular Noetherian affine scheme is a splinter. In characteristic zero, the splinter property is a local property: a Noetherian scheme over \mathbb{Q} is a splinter if and only if it is normal. In positive characteristic, the splinter property is no longer a local property in general. Bhatt’s beautiful [Bha12, Thm. 1.5], inspired from Hochster and Huneke’s [HH92, Thm. 1.2], shows that, for a proper scheme over an affine Noetherian scheme of positive characteristic, the positive-degree cohomology of the structure sheaf vanishes up to finite covers. Bhatt draws two consequences for splinters of positive characteristic: first that the positive-degree cohomology of semiample invertible sheaves on proper splinters vanishes, and second that splinters and derived splinters coincide. Our first aim is to provide further global constraints on proper splinters in positive characteristic. Our second aim is to study whether the splinter property is a derived invariant for projective varieties in positive characteristic.

Global constraints on proper splinters in positive characteristic. Recall that a smooth projective, separably rationally connected, variety over an algebraically closed field of positive characteristic has trivial Nori fundamental group scheme [Bis09], has negative Kodaira dimension [Kol96, Ch. IV, Cor. 1.11 & Prop. 3.3], and has no nonzero global differential forms [Kol96, Ch. IV, Cor. 3.8]. Motivated by the intriguing question whether proper splinters over an algebraically closed field of positive characteristic are separably rationally connected, we show:

Theorem. *Let X be a connected proper scheme over a field k of positive characteristic. Assume that X is a splinter.*

- (A) (Theorem 8.3) *If X has a k -rational point $x \in X(k)$, then the Nori fundamental group scheme $\pi_1^N(X, x)$ is trivial. In particular, if k is algebraically closed, then any finite torsor over X is trivial.*
- (B) (Theorem 9.1) *If X is positive-dimensional, then X has negative Kodaira dimension, i.e., $H^0(X, \mathcal{O}_X(nK_X)) = 0$ for all $n > 0$, where K_X denotes the canonical divisor of X .*
- (C) (Theorem 10.2) *If X is smooth, then $H^0(X, \Omega_X^1) = 0$.*

Date: March 18, 2025.

2020 Mathematics Subject Classification. 14G17, 14F35, 14L30, 14F08; (14J17, 14J26, 14E05).

Key words and phrases. Splinters, globally F -regular varieties, Nori fundamental group scheme, derived equivalence, K -equivalence, exceptional inverse image functor.

The research of both authors was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 491392403 – TRR 358.

Note from their respective proofs that **(C)**, and **(B)** in the Gorenstein case, follow from the known fact, recalled in [Proposition 9.4](#), that the Picard group of a proper splinter over a field of positive characteristic is torsion-free. The proofs of **(A)**, and of **(B)** in the general non-Gorenstein case, rely on a lifting property for splinters along finite torsors established in [Proposition 7.4](#), which itself relies on the more general key lifting property established in [Lemma 5.3\(b\)](#). The new idea, which makes it in particular possible to avoid any Gorenstein assumption, is the use of the exceptional inverse image functor for finite morphisms. By using the more general exceptional inverse image functor for proper morphisms, we further observe in [Proposition 5.11](#) that the derived splinter property for Noetherian schemes lifts along crepant morphisms. In [Lemma 5.3\(a\)](#) and building on [Lemma 5.3\(c\)](#), we establish notably the following lifting property for splinters:

Theorem (Theorem 7.7). *Let $\pi: Y \rightarrow X$ be a finite surjective morphism of normal Nagata Noetherian schemes. Assume that π satisfies either of the following conditions:*

(D) *π is quasi-étale.*

(E) *π is a quasi-torsor over a ring, and $\pi^\sharp: H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ has reduced closed fibers.*

If X is a splinter, then Y is a splinter.

Theorem **(D)** extends [\[DT23, Thm. A\]](#), where it is shown that any essentially étale cover of a Noetherian affine splinter is a splinter. Besides, Theorems **(D)** and **(E)** are stated more generally (by [Remark 3.3](#)) for globally $+$ -regular pairs as introduced in [\[Bha+23\]](#), and extend [\[Bha+23, Prop. 6.20\]](#) which deals with finite quasi-étale covers of normal excellent proper schemes over a Noetherian ring. Regarding Theorem **(A)**, we also show in [Theorem 8.3](#) that if X is a proper splinter over a separably closed field of positive characteristic, then its étale fundamental group is trivial. In that direction, we also refer to [\[Cai+23, Thm. 7.0.3\]](#), where it is in particular showed that the étale fundamental group of the regular locus of a normal projective globally $+$ -regular variety satisfying some additional technical assumptions is finite, but also to the references in the introduction of *loc. cit.* regarding the étale fundamental group of regular loci of near smooth Fano varieties.

Let k be an algebraically closed field of positive characteristic. It is well-known that a proper curve over k is a splinter if and only if it is isomorphic to \mathbb{P}_k^1 . For proper surfaces, we have the following results regarding splinters. In [Proposition 11.1](#), we use [Theorem 9.1](#) to show that if a proper surface over k is a splinter, then it is rational. We provide in [Proposition 11.4](#) new examples of proper rational surfaces that are splinters, by establishing that the blow-up of \mathbb{P}_k^2 in any number of points lying on a conic is a splinter. On the other hand, in [Proposition 11.10](#) and [Proposition 11.12](#), we give examples of proper rational surfaces that are not splinters. For instance, we show that over a finite field the blow-up of \mathbb{P}_k^2 in 9 points lying on a smooth cubic curve is not a splinter.

\mathcal{O} -invariance and D -invariance of the splinter property. The second aim of this paper is to study whether the splinter property, and the related notion of *global F -regularity*, is a derived invariant among projective varieties. We say that two projective varieties X and Y over a field k are D -equivalent if there is a k -linear equivalence $D^b(X) \cong D^b(Y)$ between their bounded derived categories of coherent sheaves. Given that a Gorenstein projective splinter, resp. a Gorenstein projective globally F -regular variety, in positive characteristic is expected to, resp. is known to, have big anticanonical divisor (see [Conjecture 3.12](#) due to [\[Bha+23\]](#), resp. [Proposition 3.11](#) due to [\[SS10, Cor. 4.5\]](#)), we obtain the following positive answer for varieties:

Theorem. *Let X and Y be normal Gorenstein projective varieties over a field k of positive characteristic. Assume that X and Y are D -equivalent. Then:*

(F) [\(Corollary 12.16\)](#) *X is a splinter if and only if Y is a splinter, provided $-K_X$ is big.*

(G) [\(Corollary 12.20\)](#) *X is globally F -regular if and only if Y is globally F -regular.*

For that purpose, we introduce in [Definition 12.6](#) the notion of \mathcal{O} -equivalence for separated schemes of finite type over a Noetherian base. By [Proposition 12.7](#), this notion coincides with the classical notion of K -equivalence in the case of normal Gorenstein varieties over a field. As before, but now for proper morphisms that are not necessarily finite, the new idea is to use the exceptional inverse image functor of Grothendieck, which allows for more flexibility. In [Proposition 12.12](#), we

observe that Kawamata's [Kaw02, Thm. 1.4(2)], stating that two D -equivalent smooth projective complex varieties X and Y with K_X or $-K_X$ big are K -equivalent, extends to the case of normal Gorenstein projective varieties over an arbitrary field. If in addition either X or Y has at worst canonical singularities, then X and Y are in fact, by Proposition 12.2(ii) and Proposition 12.7, strongly \mathcal{O} -equivalent in the sense of Definition 12.6. As is well-known, a splinter (resp. a globally F -regular variety) has pseudo-rational singularities, see Proposition 3.5, and a pseudo-rational normal Gorenstein variety has canonical singularities, see Proposition 12.5. As such, Theorem (F) and Theorem (G) follow from the following:

Theorem. *Let X and Y be proper varieties over a field k of positive characteristic. Assume that X and Y are strongly \mathcal{O} -equivalent. Then :*

(H) (Theorem 12.15) *X is a splinter if and only if Y is a splinter.*

Assume in addition that X and Y are normal projective over k . Then :

(I) (Theorem 12.18) *X is globally F -regular if and only if Y is globally F -regular.*

We observe that both Theorems (H) and (I) hold without any (\mathbb{Q} -)Gorenstein assumption and in fact without any restrictions on the singularities of X nor Y . Again, this is made possible by the systematic use of the exceptional inverse image functor. For the sake of illustration, we show in Corollary 12.3 that the splinter property is invariant under K -equivalence of *terminal* varieties, by utilizing the fact stated in Proposition 12.2(iii) that a K -equivalence between two such varieties induces a small birational map. Note however that, in positive characteristic, splinters and globally F -regular varieties may have worse singularities; in fact, both the splinter property and global F -regularity are expected to locally be the analogues of klt singularities in the complex setting.

Organization of the paper. In Sections 2 to 4, we mostly fix notation and collect basic and known facts about reflexive sheaves and Weil divisors, inverse image functors, traces, and splinters and globally F -regular varieties. Our first new contributions are contained in Sections 5 and 6 dealing respectively with splinters and globally F -regular varieties. Notably, the use of the exceptional inverse image functor makes its first appearance in Section 5.2, where we prove the key Lemma 5.3 stating in particular that the splinter property lifts along finite surjective morphisms $\pi: Y \rightarrow X$ of Noetherian schemes such that $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ as coherent sheaves on Y with $H^0(Y, \mathcal{O}_Y)$ a field or $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$.

In Section 7, we extend Lemma 5.3 to show in Proposition 7.4 that, under certain conditions, the splinter property lifts along finite torsors for Noetherian schemes, and establish Theorems (D) and (E).

In Sections 8 to 11, we explore global constraints on proper splinters in positive characteristic and establish Theorems (A) to (C). Their proof relies ultimately on Lemma 5.3(b). In Section 11, we show that proper splinter surfaces in positive characteristic are rational, and give examples of rational surfaces that are splinters as well as examples of rational surfaces that are not splinters.

In Section 12, which can be read mostly independently of the rest of the paper, we introduce the notion of \mathcal{O} -equivalence for separated schemes of finite type over a Noetherian base scheme and compare it, in the case of normal Gorenstein varieties, to the usual notions of K -equivalence and D -equivalence. We then establish Theorems (F) to (I).

Conventions. A *variety* is an integral separated scheme of finite type over a field. For a scheme X over the finite field \mathbb{F}_p with p elements, the *Frobenius* is denoted by $F: X \rightarrow X$; it is the identity on the underlying topological space and sends each local section of \mathcal{O}_X to its p -th power. A scheme X over \mathbb{F}_p is said to be *F -finite* if the Frobenius map $F: X \rightarrow X$ is finite. Note that a scheme of finite type over a field k is *F -finite* if and only if the field k is *F -finite*. For a Noetherian scheme X , we denote by X_{reg} its regular locus and by $X_{\text{sing}} := X \setminus X_{\text{reg}}$ its singular locus. If X is reduced, then X_{reg} contains the generic points of the irreducible components of X and hence is dense in X . If X is J-1, e.g. if X is excellent, then $X_{\text{reg}} \subseteq X$ is an open embedding [Sta24, Tag 07P6]. We denote by $\nu: X^\nu \rightarrow X$ the normalization of X ; if X is Nagata, e.g. if X is excellent, then ν is a finite morphism [Sta24, Tag 035S]. Both the Nagata and the excellence properties are stable under locally of finite type extensions; see [Sta24, Tag 0359] and [Sta24, Tag 07QS]. For a Noetherian scheme X , we denote by $D^b(X) := D^b(\text{Coh}(X))$ the bounded derived category of coherent sheaves on X and by

$D_{\text{Coh}}(\mathcal{O}_X)$ (resp. $D_{\text{Coh}}^b(\mathcal{O}_X)$, resp. $D_{\text{Coh}}^+(\mathcal{O}_X)$) the derived category of unbounded (resp. bounded, resp. bounded below) complexes of \mathcal{O}_X -modules with coherent cohomology sheaves. The functor $D^b(X) \rightarrow D_{\text{Coh}}(\mathcal{O}_X)$ is fully faithful with essential image $D_{\text{Coh}}^b(\mathcal{O}_X)$ [Sta24, Tag 08E0]. If X is not necessarily Noetherian, we use the notation $D_{\text{QCoh}}(\mathcal{O}_X)$ (resp. $D_{\text{QCoh}}^+(\mathcal{O}_X)$) for the derived category of unbounded (resp. bounded below) complexes of \mathcal{O}_X -modules with quasi-coherent cohomology sheaves. For a proper morphism $f: Z \rightarrow X$ of Noetherian schemes, we denote by $f^\#: \mathcal{O}_X \rightarrow f_*\mathcal{O}_Z$, resp. $f^\#: \mathcal{O}_X \rightarrow Rf_*\mathcal{O}_Z$, the canonical morphism in the category, resp. bounded derived category, of coherent sheaves on X . Given an effective Weil divisor D on a normal Noetherian scheme X , we write $\sigma_D: \mathcal{O}_X \rightarrow \mathcal{O}_X(D)$ for the morphism determined by D .

Acknowledgements. We thank Javier Carvajal-Rojas, Karl Schwede, and the anonymous referee, for useful comments. Lemma 5.3(c) started taking shape at the Fall School “Methods in Mixed Characteristic Geometry” in Mainz in October 2024 and we are grateful to the organizers and participants for the pleasant and stimulating atmosphere.

2. REFLEXIVE SHEAVES AND EXCEPTIONAL INVERSE IMAGE FUNCTORS

2.1. Reflexive sheaves and Weil divisors. Let X be an integral Noetherian scheme. Recall that a coherent sheaf \mathcal{F} on X is called *reflexive* if the canonical map $\mathcal{F} \rightarrow \mathcal{F}^{\vee\vee}$ is an isomorphism, where by definition $\mathcal{F}^\vee := \mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{O}_X)$.

- (i) For any coherent sheaf \mathcal{F} and any reflexive coherent sheaf \mathcal{G} the sheaf $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})$ is reflexive [Sta24, Tag 0AY4].
- (ii) If X is normal, then a coherent sheaf is reflexive if and only if it is S_2 [Har94, Thm. 1.9].
- (iii) If X is normal and $i: U \hookrightarrow X$ is an open immersion such that $\text{codim}_X(X \setminus U) \geq 2$, then $i_*i^*\mathcal{F} \cong \mathcal{F}$ for any reflexive coherent sheaf \mathcal{F} . Furthermore, the restriction i^* induces an equivalence of categories from reflexive coherent sheaves on X to reflexive coherent sheaves on U [Har94, Thm. 1.12].
- (iv) If X is normal and \mathcal{F} is a reflexive coherent sheaf, then there exists an open immersion $i: U \hookrightarrow X$ such that $\text{codim}_X(X \setminus U) \geq 2$ and such that $i^*\mathcal{F}$ is finite locally free [Sta24, Tag 0AY6].

To any Weil divisor D on an integral normal Noetherian scheme X with function field $K(X)$, one can associate a coherent sheaf $\mathcal{O}_X(D) \subseteq K(X)$ whose sections on open subsets $V \subseteq X$ are given by

$$\Gamma(V, \mathcal{O}_X(D)) := \{f \in K(X)^\times \mid \text{div}(f)|_V + D|_V \geq 0\} \cup \{0\}.$$

The sheaf $\mathcal{O}_X(D)$ is reflexive of rank 1 and it is invertible if and only if D is Cartier. Since any reflexive rank 1 sheaf is isomorphic to a subsheaf of the locally constant sheaf $K(X)$, we have [Sta24, Tag 0EBM] a 1-1 correspondence

$$\{\text{Weil divisors on } X \text{ up to linear equivalence}\} \leftrightarrow \{\text{reflexive sheaves of rank 1 on } X\} / \cong.$$

Moreover thanks to (iii) and (iv) this bijection turns out to be a group homomorphism, see [Har94, Prop. 2.8(d)], provided one takes the double dual of the usual tensor product, i.e.,

$$\mathcal{O}_X(D + D') \cong (\mathcal{O}_X(D) \otimes \mathcal{O}_X(D'))^{\vee\vee}.$$

A Weil divisor D on a normal Noetherian scheme is effective if and only if $\mathcal{O}_X \subseteq \mathcal{O}_X(D) \subseteq K(X)$. Thus a reflexive rank 1 sheaf \mathcal{F} corresponds to an effective Weil divisor D if and only if there is an injective morphism $\mathcal{O}_X \hookrightarrow \mathcal{F}$. For later use recall the following criterion.

Lemma 2.1. *Let X be an integral normal Noetherian scheme and assume that $H^0(X, \mathcal{O}_X)$ is a field. Then a Weil divisor D on X is linearly equivalent to zero if and only if both $\mathcal{O}_X(D)$ and $\mathcal{O}_X(-D)$ admit nonzero global sections.*

Proof. The only if part is obvious. Assume $s: \mathcal{O}_X \rightarrow \mathcal{O}_X(D)$ and $t: \mathcal{O}_X \rightarrow \mathcal{O}_X(-D)$ are nontrivial sections. We have isomorphisms of sheaves $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X(-D)) \cong \mathcal{H}om_{\mathcal{O}_X}(\mathcal{O}_X(D), \mathcal{O}_X)$ and $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{O}_X(D), \mathcal{O}_X(D)) \cong \mathcal{H}om_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X)$. Therefore we can interpret t as a global section of $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{O}_X(D), \mathcal{O}_X)$. The compositions $s \circ t$ and $t \circ s$ are both nonzero and give elements in $H^0(X, \mathcal{O}_X)$, thus they are isomorphisms. Hence $\mathcal{O}_X \cong \mathcal{O}_X(D)$, which shows that D is trivial. \square

Given two Weil divisors D and E on a normal Noetherian scheme X , we write $D \sim E$ if D and E are linearly equivalent. Likewise, for two \mathbb{Q} -Weil divisors D and E , we write $D \sim_{\mathbb{Q}} E$ if D and E are \mathbb{Q} -linearly equivalent.

Let $f: Y \rightarrow X$ be a finite surjective morphism of integral normal Noetherian schemes, and let D be a \mathbb{Q} -Weil divisor on X . Assume that X is universally catenary, e.g. Cohen–Macaulay, so that if $U \subseteq X$ is an open subset with $\text{codim}_X(X \setminus U) \geq 2$, then $\text{codim}_Y(Y \setminus f^{-1}(U)) \geq 2$. By (iv), choose such a U such that D restricts to a \mathbb{Q} -Cartier divisor. The pullback f^*D can be defined by restricting D to U , pulling back to $f^{-1}(U)$, and then extending the pullback uniquely to a \mathbb{Q} -Weil divisor on Y ; compare with [Har94, Prop. 2.18] and with the explicit construction of [ST14, §2.2].

2.2. Exceptional inverse image functor and dualizing complexes. Let $f: Y \rightarrow X$ be a morphism of schemes of finite type and separated over a Noetherian scheme S . We consider the *exceptional inverse image functor* $f^!: D_{\text{QCoh}}^+(\mathcal{O}_X) \rightarrow D_{\text{QCoh}}^+(\mathcal{O}_Y)$ as defined in [Sta24, Tag 0A9Y]. The functor $f^!$ sends $D_{\text{Coh}}^+(\mathcal{O}_X)$ to $D_{\text{Coh}}^+(\mathcal{O}_Y)$ [Sta24, Tag 0ATZ]. If $f: Y \rightarrow X$ is proper, then $f^!$ is right adjoint to $Rf_*: D_{\text{Coh}}^+(\mathcal{O}_Y) \rightarrow D_{\text{Coh}}^+(\mathcal{O}_X)$ [Sta24, Tag 0AU3].

Assume now that X admits a *dualizing complex* ω_X^\bullet as defined in [Sta24, Tag 0A85]. Then $\omega_Y^\bullet := f^! \omega_X^\bullet$ defines a dualizing complex on Y [Sta24, Tag 0AU3]. If X is equidimensional, $\omega_X := \mathcal{H}^{-\dim X}(\omega_X^\bullet)$ is an S_2 sheaf, called the *dualizing sheaf* [Sta24, Tag 0AWH]. Moreover, X is Cohen–Macaulay if and only if $\omega_X^\bullet = \omega_X[\dim X]$ [Sta24, Tag 0AWQ] and X is Gorenstein if and only if ω_X^\bullet is an invertible object [Sta24, Tag 0AWV]. The latter condition is further equivalent to $\omega_X^\bullet = \omega_X[\dim X]$ with ω_X an invertible sheaf [Sta24, Tag 0FPG]. If X is normal and equidimensional, then ω_X is a reflexive sheaf of rank 1 and there thus exists a unique (up to linear equivalence) Weil divisor K_X , called the *canonical divisor*, such that $\omega_X \cong \mathcal{O}_X(K_X)$.

Let now $h: X \rightarrow \text{Spec } k$ be a scheme of finite type and separated over a field k . Then $\omega_X^\bullet := h^! \mathcal{O}_{\text{Spec } k} \in D_{\text{Coh}}^+(\mathcal{O}_X)$ is a dualizing complex on X [Sta24, Tag 0AWJ]. If X is smooth over k , then the dualizing sheaf ω_X coincides with the *canonical sheaf* $\bigwedge^{\dim X} \Omega_{X/k}^1$ [Sta24, Tag 0E9Z]. If $h: X \rightarrow \text{Spec } k$ is proper and X is equidimensional, then Rh_* is left adjoint to $h^!$ and for every $K \in D^b(X)$, there is a functorial isomorphism $\text{Ext}_X^i(K, \omega_X^\bullet) = \text{Hom}_k(H^i(X, K), k)$ compatible with shifts and exact triangles, see, e.g., [Sta24, Tag 0FVU]. By Yoneda, the object ω_X^\bullet is unique up to unique isomorphism among all objects satisfying this universal property.

In the more general situation where $f: Y \rightarrow X$ is a morphism of quasi-compact quasi-separated schemes, $Rf_*: D_{\text{QCoh}}(\mathcal{O}_Y) \rightarrow D_{\text{QCoh}}(\mathcal{O}_X)$ still admits a right adjoint f^\times , see, e.g., [GW23, Thm. 25.17]. The functor f^\times sends $D_{\text{QCoh}}^+(\mathcal{O}_X)$ to $D_{\text{QCoh}}^+(\mathcal{O}_Y)$ and, if X is Noetherian and f is proper, agrees with $f^!$; see [GW23, Lem. 25.23 & Thm. 25.61].

The following general Lemma 2.2 is a consequence of the properties of the exceptional inverse image functor. In particular, choosing $q: Z \rightarrow Y$ in the statement of Lemma 2.2 to be an isomorphism, it shows that if $p: Z \rightarrow X$ is a separated morphism of finite type, then $p^! \mathcal{O}_X \cong \mathcal{O}_Z$ if and only if $Lp^* \omega_X^\bullet \cong \omega_Z^\bullet$.

Lemma 2.2. *Let X and Y be schemes of finite type and separated over a Noetherian scheme S such that S admits a dualizing complex ω_S^\bullet . Denote by $h_X: X \rightarrow S$ and $h_Y: Y \rightarrow S$ the structure morphisms. Let Z be a scheme of finite type and separated over S with S -morphisms $p: Z \rightarrow X$ and $q: Z \rightarrow Y$. Then in $D_{\text{Coh}}(\mathcal{O}_Z)$ we have*

$$Lp^* \omega_X^\bullet \cong Lq^* \omega_Y^\bullet \iff p^! \mathcal{O}_X \cong q^! \mathcal{O}_Y,$$

where $\omega_X^\bullet = h_X^! \omega_S^\bullet$ and $\omega_Y^\bullet = h_Y^! \omega_S^\bullet$. In particular, if X and Y are equidimensional and Gorenstein, then

$$p^* \omega_X[\dim X] \cong q^* \omega_Y[\dim Y] \iff p^! \mathcal{O}_X \cong q^! \mathcal{O}_Y,$$

where $\omega_X = \mathcal{H}^{-\dim X}(\omega_X^\bullet)$ and $\omega_Y = \mathcal{H}^{-\dim Y}(\omega_Y^\bullet)$.

Proof. Let $\omega_Z^\bullet := p^! \omega_X^\bullet = q^! \omega_Y^\bullet$. Recall, e.g., from [Sta24, Tag 0AU3], that the functor $R\mathcal{H}om_{\mathcal{O}_Z}(-, \omega_Z^\bullet)$ defines an involution of $D_{\text{Coh}}(\mathcal{O}_Z)$ and that we have the duality formula

$$R\mathcal{H}om_{\mathcal{O}_Z}(p^! M, \omega_Z^\bullet) = Lp^* R\mathcal{H}om_{\mathcal{O}_X}(M, \omega_X^\bullet)$$

which holds naturally in $M \in D_{\text{Coh}}^+(\mathcal{O}_X)$ and similarly for $q: Z \rightarrow Y$. Setting $M = \mathcal{O}_X$ or $M = \mathcal{O}_Y$ yields

$$R\mathcal{H}om_{\mathcal{O}_Z}(p^!\mathcal{O}_X, \omega_Z^\bullet) = Lp^*\omega_X^\bullet \quad \text{and} \quad R\mathcal{H}om_{\mathcal{O}_Z}(q^!\mathcal{O}_Y, \omega_Z^\bullet) = Lq^*\omega_Y^\bullet.$$

Therefore, $p^!\mathcal{O}_X \cong q^!\mathcal{O}_Y$ if and only if $Lp^*\omega_X^\bullet \cong Lq^*\omega_Y^\bullet$. \square

Remark 2.3. Let $\pi: Y \rightarrow X$ be a separated morphism of finite type of Noetherian schemes. If X admits a dualizing complex ω_X^\bullet and $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$, then [Lemma 2.2](#) implies that $L\pi^*\omega_X^\bullet \cong \omega_Y^\bullet$. Since a scheme admitting a dualizing complex is Gorenstein if and only if it admits an invertible dualizing complex, we observe that if X is Gorenstein, then Y is Gorenstein. Likewise, since a scheme admitting a dualizing complex is Cohen–Macaulay if and only if it admits a dualizing complex that is the shift of a sheaf, we observe that if X is Cohen–Macaulay and π is flat, then Y is Cohen–Macaulay.

Remark 2.4. A proper surjective morphism of integral excellent Noetherian schemes $\pi: Y \rightarrow X$ such that $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$ and such that X admits a dualizing complex on a dense open subset is generically finite. Indeed, the regular locus X_{reg} of X is open (since X is excellent) and dense (since X is integral) and, likewise, the regular locus U of $\pi^{-1}(X_{\text{reg}})$ is open and dense. Denote by $p: U \rightarrow X_{\text{reg}}$ the restriction of π . Since the restriction to open subsets commutes with exceptional inverse image functors [\[Sta24, Tag 0G4J\]](#), we have $p^!\mathcal{O}_{X_{\text{reg}}} \cong \mathcal{O}_U$. By [Lemma 2.2](#), we have an isomorphism $\omega_U[\dim Y] \cong p^*\omega_{X_{\text{reg}}}[\dim X]$, and thus $\dim Y = \dim X$.

2.3. Exceptional inverse image along finite morphisms. For the sake of the proof of the lifting [Lemma 5.3](#), we will need to consider the exceptional inverse functor for finite morphisms, and in particular for cases (a) and (c), for finite morphisms between possibly non-Noetherian schemes.

Let $\pi: Y \rightarrow X$ be a finite morphism of Noetherian schemes. The exact functor $\pi_*: \text{Coh}(Y) \rightarrow \text{Coh}(X)$ admits a right adjoint, which by abuse we denote by $\pi^!: \text{Coh}(X) \rightarrow \text{Coh}(Y)$, given by $\pi^!(\mathcal{F}) := \mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, \mathcal{F})$ considered as an \mathcal{O}_Y -module, see, e.g., [\[Har77, Ch. III, Ex. 6.10\]](#) or [\[Sta24, Tag 0AWZ\]](#). In the context of [Section 2.2](#), the right adjoint of $\pi_* = R\pi_*: D_{\text{Coh}}(\mathcal{O}_X) \rightarrow D_{\text{Coh}}(\mathcal{O}_Y)$ can be obtained from $\pi^! = \mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, -)$ by forming the right derived functor [\[Sta24, Tag 0AU3\]](#). If π is finite and flat, then $\pi_*\mathcal{O}_Y$ is a vector bundle and so $\mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, -) = R\mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, -)$, i.e., in that case, the notation $\pi^!$ is unambiguous. In general, in order to avoid any ambiguity, we will write $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$ in $\text{Coh}(Y)$ if there is an isomorphism of \mathcal{O}_Y -modules $\mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, \mathcal{O}_X) \cong \mathcal{O}_Y$, and we will write $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$ in $D_{\text{Coh}}(\mathcal{O}_Y)$ if there is an isomorphism $R\mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, \mathcal{O}_X) \cong \mathcal{O}_Y$ in $D_{\text{Coh}}(\mathcal{O}_Y)$.

If $\pi: Y \rightarrow X$ is a finite morphism of quasi-compact quasi-separated schemes, $R\pi_*: D_{\text{QCoh}}(\mathcal{O}_Y) \rightarrow D_{\text{QCoh}}(\mathcal{O}_X)$ admits π^\times as a right adjoint. If π is finite and *pseudo-coherent*, then π^\times coincides with $R\mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, -)$ on $D_{\text{QCoh}}^+(\mathcal{O}_X)$ [\[Sta24, Tag 0AX2\]](#). A morphism of finite type of Noetherian schemes is pseudo-coherent [\[Sta24, Tag 0684\]](#) and a flat base change of a pseudo-coherent morphism is pseudo coherent [\[Sta24, Tag 0680\]](#). We will use this more general adjoint in the proof of [Lemma 5.3](#) for a flat base change of a finite morphism of Noetherian schemes.

Remark 2.5. Let $\pi: Y \rightarrow X$ be a finite morphism of Noetherian schemes. By taking the cohomology sheaves in degree zero, one sees that if $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$ in $D_{\text{Coh}}(\mathcal{O}_Y)$, then $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$ in $\text{Coh}(Y)$.

Remark 2.6. A finite morphism $\pi: Y \rightarrow X$ of Noetherian schemes with X integral and such that $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$ in $\text{Coh}(Y)$ is surjective. Indeed, the question is local on X and we may assume both X and Y are affine, say $X = \text{Spec } A$ and $Y = \text{Spec } B$. Let $I := \ker(\pi^\sharp: A \rightarrow B)$. Then any $\varphi \in \text{Hom}_A(B, A)$ has image contained in $\text{Ann}_A(I)$. Since A is a domain, $\text{Hom}_A(B, A) \cong B \neq 0$ forces I to be zero and hence π to be surjective.

2.4. Traces for finite morphisms. The *classical trace map* $\text{tr}_{Y/X}: \pi_*\mathcal{O}_Y \rightarrow \mathcal{O}_X$ for a finite surjective morphism $\pi: Y \rightarrow X$ of integral schemes with X normal is defined as follows. Denote $K(X)$ and $K(Y)$ the function fields of X and Y , respectively. On affine open subsets $U = \text{Spec } A \subseteq X$ with $\pi^{-1}(U) = \text{Spec } B \subseteq Y$, $\text{tr}_{B/A}: B \rightarrow A$ sends $b \in B$ to the trace of the multiplication by b map on $K(Y)$ viewed as a $K(X)$ -vector space and, by normality of A , defines an element of A .

If $U \subseteq X$ is any open subset in an integral scheme X , then $H^0(U, \mathcal{O}_X) = \bigcap_{x \in U} \mathcal{O}_{X,x} \subseteq K(X)$, and if X is further assumed to be normal, then $H^0(U, \mathcal{O}_X)$ is integrally closed in $K(X)$. As such,

the classical trace map $\mathrm{tr}_{Y/X}: \pi_* \mathcal{O}_Y \rightarrow \mathcal{O}_X$ is given on open subsets $U \subseteq X$ as the map sending a section $s \in H^0(\pi^{-1}(U), \mathcal{O}_Y)$ to the trace of the multiplication by s map on $K(Y)$ viewed as a $K(X)$ -vector space. By Lemma 2.7 below, such a section s is integral over $H^0(U, \mathcal{O}_X)$ and hence, the trace is an element of $H^0(U, \mathcal{O}_X)$.

Lemma 2.7. *Let $\pi: Y \rightarrow X$ be an integral surjective morphism of integral schemes with X normal. Then $\pi^\sharp: H^0(U, \mathcal{O}_X) \rightarrow H^0(\pi^{-1}(U), \mathcal{O}_Y)$ is an integral ring extension for every open subset $U \subseteq X$. Moreover, if Y is normal, then π is the normalization of X in $K(Y)$ and for every open subset $U \subseteq X$, $H^0(\pi^{-1}(U), \mathcal{O}_Y)$ is the integral closure of $H^0(U, \mathcal{O}_X)$ in the function field $K(Y)$.*

Proof. Without loss of generality we can assume $U = X$. Let $\mathrm{Spec} A \subseteq X$ be an affine open subset and let $\mathrm{Spec} B := \pi^{-1}(\mathrm{Spec} A) \subseteq Y$ be the pre-image under π . Let $s \in S := H^0(Y, \mathcal{O}_Y)$ be nonzero. Since $A \rightarrow B$ is integral, there exists a monic irreducible polynomial $P_A \in A[T]$ such that $P_A(s) = 0$ in B . By normality of X , A is integrally closed in $K(X)$, and it follows that P_A is irreducible as a polynomial in $K(X)[T]$. Indeed, if $P_A = QR$ in $K(X)[T]$ with Q and R monic, then denoting C the integral closure of A in a splitting field of P_A over $K(X)$, we have $Q, R \in C[T] \cap K(X)[T] = A[T]$. Let now $\mathrm{Spec} A' \subseteq X$ be another affine open subset and let $P \in K(X)[T]$ be the gcd of P_A and $P_{A'}$. Then $P(s) = 0$, so P is not constant as $s \neq 0$ in $K(Y)$ by integrality of Y . As P_A and $P_{A'}$ are irreducible, we obtain $P_A = P = P_{A'}$. It follows that $P = P_A$ has coefficients in $R = H^0(X, \mathcal{O}_X) = \bigcap A$, where the intersection runs over all affine open subsets of X . Thus s is integral over R .

If Y is normal, then $H^0(V, \mathcal{O}_Y) = \bigcap_{y \in V} \mathcal{O}_{Y,y} \subseteq K(Y)$ is integrally closed for every open subset $V \subseteq Y$. Hence, for every open subset $U \subseteq X$, $H^0(\pi^{-1}(U), \mathcal{O}_Y)$ is the integral closure of $H^0(U, \mathcal{O}_X)$ in $K(Y)$. This coincides with the relative normalization in [Sta24, Tag 0BAK] if U is affine. \square

The following lemma is a slight generalization of [Spe20, Lem. 9] (we do not assume that the fields are the fraction fields of the integral rings involved). It will be used in the proofs of Lemma 5.3(a) with $R = H^0(X, \mathcal{O}_X)$, $S = H^0(Y, \mathcal{O}_Y)$, $K = K(X)$, and $L = K(Y)$.

Lemma 2.8. *Let $R \hookrightarrow S$ be an integral extension of integral domains fitting into a commutative diagram*

$$\begin{array}{ccc} R & \hookrightarrow & S \\ \downarrow & & \downarrow \\ K & \hookrightarrow & L \end{array}$$

with $K \rightarrow L$ a finite field extension such that R and S are integrally closed in K and L , respectively. Then the classical trace $\mathrm{tr}_{L/K}: L \rightarrow K$ sends S to R , and sends $\sqrt{\mathfrak{p}S}$ to \mathfrak{p} for every prime ideal $\mathfrak{p} \subseteq R$.

Proof. If L/K is not separable, then $\mathrm{tr}_{L/K}$ is zero. Thus we can and do assume that L/K is separable. Let M be the Galois closure of L over K , denote by $G = \mathrm{Gal}(M/K)$ the Galois group, and let $H \subseteq G$ be the stabilizer of L . Let $T \subseteq M$ be the integral closure of R in M . Since $R \hookrightarrow S$ is integral, $S \subseteq T$. Note that for any $s \in L$, we have $\mathrm{tr}_{L/K}(s) = \sum_{g \in G/H} g(s)$, where g runs over a set of coset representatives of G/H . By definition of the integral closure, any automorphism $g \in G$ satisfies $g(T) = T \subseteq M$, so $\mathrm{tr}_{L/K}(s)$ lies in $T \cap K = R$ for any $s \in S$. Similarly, any $g \in G$ preserves $\sqrt{T\mathfrak{p}} \subseteq T$. Hence, for any $s \in \sqrt{S\mathfrak{p}} \subseteq \sqrt{T\mathfrak{p}}$, $\mathrm{tr}_{L/K}(s)$ lies in $\sqrt{T\mathfrak{p}} \cap R = \mathfrak{p}$. \square

Remark 2.9. Note that in the situation of Lemma 2.8, the classical trace map $\mathrm{tr}_{L/K}: L \rightarrow K$ need not restrict to the classical trace map $\mathrm{tr}_{S/R}: S \rightarrow R$. Consider for instance the Frobenius $F: \mathbb{P}_{\mathbb{F}_p}^1 \rightarrow \mathbb{P}_{\mathbb{F}_p}^1$. It induces the identity on global sections but on function fields induces the purely inseparable field extension $\mathbb{F}_p(t) \rightarrow \mathbb{F}_p(t^{1/p})$.

The classical trace map $\mathrm{tr}_{Y/X}: \pi_* \mathcal{O}_Y \rightarrow \mathcal{O}_X$ for a finite flat morphism $\pi: Y \rightarrow X$ locally of finite presentation is given by the \mathcal{O}_X -linear map sending a local section s of the finite locally free \mathcal{O}_X -module $\pi_* \mathcal{O}_Y$ to the trace of the multiplication by s on $\pi_* \mathcal{O}_Y$. If X is further assumed to be normal, and X and Y to be integral, the classical trace map $\mathrm{tr}_{Y/X}$ coincides with the previous

construction (hence the same nomenclature for both trace maps). Indeed, a local basis of $\pi_*\mathcal{O}_Y$ as an \mathcal{O}_X -module gives a local basis of $K(Y)$ as a $K(X)$ -vector space.

Let $\pi: Y \rightarrow X$ be a finite surjective morphism of Noetherian schemes. To give an isomorphism $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$ in $\text{Coh}(Y)$ is equivalent to give an isomorphism of \mathcal{O}_Y -modules

$$\mathcal{O}_Y \xrightarrow{\cong} \mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, \mathcal{O}_X),$$

or equivalently, if π is further assumed to be flat, to produce an \mathcal{O}_X -linear map $\text{Tr}_{Y/X}: \pi_*\mathcal{O}_Y \rightarrow \mathcal{O}_X$ such that the symmetric bilinear form $\text{Tr}_{Y/X}(\alpha \cdot \beta)$ on the locally free sheaf $\pi_*\mathcal{O}_Y$ with values in \mathcal{O}_X is nonsingular. Such a nonsingular trace map $\text{Tr}_{Y/X}$ then spans $\pi^!\mathcal{O}_X$ freely as an \mathcal{O}_Y -module.

If π is finite flat, or if Y is integral and X integral normal, then the classical trace map provides a canonical morphism of \mathcal{O}_Y -modules

$$\mathcal{O}_Y \longrightarrow \mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, \mathcal{O}_X), \quad 1 \mapsto \text{tr}_{Y/X}.$$

We will write $\pi^!\mathcal{O}_X = \mathcal{O}_Y$ in $\text{Coh}(Y)$ if this canonical morphism is an isomorphism.

The classical trace map $\text{tr}_{Y/X}: \pi_*\mathcal{O}_Y \rightarrow \mathcal{O}_X$ for a finite flat morphism $\pi: Y \rightarrow X$ of schemes is nonsingular if and only if π is finite étale [Sta24, Tag 0BVH]. In fact, the nonsingularity of $\text{tr}_{Y/X}$ characterizes finite surjective quasi-étale morphisms, i.e., finite surjective morphisms $\pi: Y \rightarrow X$ such that there exists an open subset $U \subseteq X$ with $\text{codim}_X(X \setminus U) \geq 2$ and such that $\pi|_U: \pi^{-1}(U) \rightarrow U$ is étale:

Lemma 2.10. *Let $\pi: Y \rightarrow X$ be a finite surjective morphism of integral normal Noetherian schemes. Consider the following statements:*

- (i) π is quasi-étale.
- (ii) $\pi^!\mathcal{O}_X = \mathcal{O}_Y$ in $\text{Coh}(Y)$, i.e., the classical trace is nonsingular.

Then (i) \Rightarrow (ii), and (ii) \Rightarrow (i) holds if X is universally catenary, e.g., if X is Cohen–Macaulay.

Proof. Assume π is quasi-étale. Let $U \subseteq X$ be an open subset with $\text{codim}_X(X \setminus U) \geq 2$ such that $\pi|_U: \pi^{-1}(U) \rightarrow U$ is finite étale. The map $\mathcal{O}_Y \rightarrow \pi^!\mathcal{O}_X = \mathcal{H}om_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, \mathcal{O}_X), 1 \mapsto \text{tr}_{Y/X}$ is a map of quasi-coherent sheaves on Y that restricts to an isomorphism on $\pi^{-1}(U)$, and hence is an isomorphism on Y since both sheaves are reflexive after applying π_* . Conversely, we claim that if $\pi: Y \rightarrow X$ is a finite surjective morphism with Y integral, and X normal and universally catenary, then there exists an open subset $U \subseteq X$ with $\text{codim}_X(X \setminus U) \geq 2$ such that $\pi|_U: V := \pi^{-1}(U) \rightarrow U$ is finite flat. This suffices to conclude since by [Sta24, Tag 0BVH] the classical trace for $\pi|_U$ is nonsingular if and only if $\pi|_U$ is étale. Let $V \subseteq Y$ be the flat locus of π , which by [Sta24, Tag 0398] is open, and let $Z := Y \setminus V$. Since X is universally catenary, any codimension-1 point of Y is mapped to a codimension-1 point of X by [Sta24, Tag 02JT], and by [Har77, Ch. III, Prop. 9.7], V contains all codimension-1 points of Y . Hence Z has codimension at least 2 in Y , and so $\pi(Z)$ defines a closed subscheme of codimension at least 2 in X . We may then choose $U := X \setminus \pi(Z)$. \square

3. PRELIMINARIES ON SPLINTERS AND GLOBALLY F -REGULAR VARIETIES

3.1. Splinters. We review the notion of *splinter* for Noetherian schemes, the local constraints it imposes, as well as the global constraints it imposes on proper schemes over a field of positive characteristic.

Definition 3.1. A Noetherian scheme X is a *splinter* if for any finite surjective morphism $f: Y \rightarrow X$ the canonical map $\mathcal{O}_X \rightarrow f_*\mathcal{O}_Y$ splits in the category $\text{Coh}(X)$ of coherent sheaves on X . A Noetherian scheme X is a *derived splinter* if for any proper surjective morphism $f: Y \rightarrow X$ the map $\mathcal{O}_X \rightarrow \text{R}f_*\mathcal{O}_Y$ splits in the bounded derived category $\text{D}^b(X)$ of coherent sheaves on X .

Note that a derived splinter is a splinter, so that being a derived splinter is *a priori* more restrictive than being a splinter. In the recent work [Bha+23], the notion of splinter has been extended to pairs. Precisely:

Definition 3.2 ([Bha+23, Def. 6.1]). Let (X, Δ) be a pair consisting of a normal excellent Noetherian scheme X and of an effective \mathbb{Q} -Weil divisor Δ . The pair (X, Δ) is called *globally $+$ -regular* if for any finite surjective morphism $f: Y \rightarrow X$ with Y normal, the natural map $\mathcal{O}_X \rightarrow f_*\mathcal{O}_Y(\lfloor f^*\Delta \rfloor)$ splits in $\text{Coh}(X)$.

Remark 3.3. A normal excellent Noetherian scheme X is a splinter if and only if $(X, 0)$ is globally $+$ -regular. The “only if” statement is obvious. For the converse, recall from [Sta24, Tag 035S] that the normalization of an excellent scheme is a finite morphism. Thus, if $\pi: Y \rightarrow X$ is a finite surjective morphism, then Y is excellent so that the normalization $\nu: Y^\nu \rightarrow Y$ is finite. Any splitting of $\mathcal{O}_X \rightarrow (\pi \circ \nu)_*\mathcal{O}_{Y^\nu}$ provides a splitting of $\mathcal{O}_X \rightarrow \pi_*\mathcal{O}_Y$.

Remark 3.4. Note that [Bha+23, Def. 6.1] only consider schemes whose closed points have residue fields of positive characteristic. A reason for this is outlined in [Bha+23, Rmk. 6.3]. Since for our applications this condition is not necessary, we deviate from that convention. We also note that in this work the excellence condition on X can be weakened to requiring X to be universally catenary (in order to define pullbacks of \mathbb{Q} -Weil divisors along finite surjective morphisms) and Nagata (in order to dominate any finite cover of X by a normal finite cover).

In general, if a Noetherian scheme X is a splinter, then it is a basic fact that any open $U \subseteq X$ is a splinter; see, e.g., Lemma 4.1(i) below. Hence, if X is a splinter, then all of its local rings are splinters. Moreover, if X is in addition assumed to be affine, X is a splinter if and only if all its local rings are splinters [DT23, Lem. 2.1.3].

In characteristic zero, a Noetherian scheme X is a splinter if and only if it is normal [Bha12, Ex. 2.1], and it is a derived splinter if and only if it has rational singularities [Bha12, Thm. 2.12]. In particular, in characteristic zero, the splinter and derived splinter properties are distinct, and they both define local properties.

In positive characteristic, Bhatt showed that the splinter and the derived splinter properties agree [Bha12, Thm. 1.4] and observed that, in contrast to the affine setting, the splinter property is not a local property for proper schemes. The following proposition summarizes the known local constraints on splinters in positive characteristic.

Proposition 3.5 ([Bha12, Ex. 2.1, Rmk. 2.5, Cor. 6.4], [Bha21, Rmk. 5.14], [Sin99], [Smi00, §2.2]). *Let X be a scheme of finite type over a field of positive characteristic. If X is a splinter, then*

- (i) *X is normal;*
- (ii) *X is Cohen–Macaulay;*
- (iii) *X is pseudo-rational;*
- (iv) *X is F -rational.*

Moreover, if X is \mathbb{Q} -Gorenstein and F -finite, then its local rings are strongly F -regular.

Recall from [HH89] that a ring R of positive characteristic is *strongly F -regular* if it is F -finite and if for any $c \in R$ not belonging to any minimal prime ideal of R there exists $e > 0$ such that the inclusion of R -modules $R \hookrightarrow F_*^e R$ which sends 1 to $F_*^e c$ splits as a map of R -modules. The ring R is strongly F -regular if and only if its local rings are strongly F -regular. If R is strongly F -regular, then the affine scheme $X = \text{Spec } R$ is a splinter; see, e.g., [DT23, Rmk. 2.13(2)].

The splinter property also imposes strong constraints on the global geometry of proper varieties in positive characteristic. For example, from Bhatt’s “vanishing up to finite cover in positive characteristic” [Bha12, Thm. 1.5], we have:

Proposition 3.6 ([Bha12]). *Let X be a proper variety over a field of positive characteristic and let \mathcal{L} be a semiample invertible sheaf on X . If X is a splinter, then $H^i(X, \mathcal{L}) = 0$ for all $i > 0$. In particular, $H^i(X, \mathcal{O}_X) = 0$ for all $i > 0$.*

Proof. For a proper variety X over a field of positive characteristic, there exists, by [Bha12, Prop. 7.2], for any $i > 0$ a finite surjective morphism $\pi: Y \rightarrow X$ such that the induced map $H^i(X, \mathcal{L}) \rightarrow H^i(Y, \pi^*\mathcal{L})$ is zero. If now X is a splinter, the pullback map $\mathcal{O}_X \rightarrow \pi_*\mathcal{O}_Y$ admits a splitting s , i.e., we have

$$\text{id}: \mathcal{O}_X \rightarrow \pi_*\mathcal{O}_Y \xrightarrow{s} \mathcal{O}_X.$$

Tensoring with \mathcal{L} and using the projection formula, we obtain

$$\mathrm{id}: H^i(X, \mathcal{L}) \xrightarrow{0} H^i(Y, \pi^* \mathcal{L}) = H^i(X, \pi_* \pi^* \mathcal{L}) \rightarrow H^i(X, \mathcal{L}),$$

where the equality in the middle uses that π is finite, in particular affine. We conclude that $H^i(X, \mathcal{L}) = 0$. \square

As a direct consequence, we have the following useful constraint, which will be refined in [Theorem 9.1](#), on proper splinters in positive characteristic:

Lemma 3.7. *Let X be a proper scheme over a field of positive characteristic, with positive-dimensional irreducible components. If X is a splinter, then its canonical divisor K_X is not effective, in particular its dualizing sheaf ω_X is nontrivial.*

Proof. Since a splinter is normal, by working on each connected component of X separately, we can assume that X is of pure positive dimension, say n . By [Proposition 3.5](#), X is Cohen–Macaulay, so $\omega_X^\bullet = \omega_X[n]$. By [Proposition 3.6](#) and Serre duality for Cohen–Macaulay schemes, we obtain $H^0(X, \omega_X)^\vee \cong H^n(X, \mathcal{O}_X) = 0$. This shows that the Weil divisor K_X is not effective. \square

3.2. Globally F -regular varieties. Let p be a prime number. We recall the notion of *global F -regularity* for normal varieties over an F -finite field of characteristic p , and review the local constraints it imposes, as well as the global constraints it imposes on proper varieties.

Definition 3.8 ([\[Smi00; SS10\]](#)). A normal scheme X over \mathbb{F}_p is called *globally F -regular* if it is F -finite and if for any effective Weil divisor D on X there exists a positive integer $e \in \mathbb{Z}_{>0}$ such that the map

$$\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X \xrightarrow{F_*^e(\sigma_D)} F_*^e \mathcal{O}_X(D)$$

of \mathcal{O}_X -modules splits. Here $\sigma_D: \mathcal{O}_X \rightarrow \mathcal{O}_X(D)$ is the morphism determined by the Weil divisor D . A normal scheme X over \mathbb{F}_p is called *F -split* if it is F -finite and if $\mathcal{O}_X \rightarrow F_* \mathcal{O}_X$ splits.

A pair (X, Δ) consisting of a normal scheme X over \mathbb{F}_p and an effective \mathbb{Q} -Weil divisor Δ is called *globally F -regular* if X is F -finite and if for any effective Weil divisor D on X there exists an integer $e > 0$ such that the natural map

$$\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X((p^e - 1)\Delta + D)$$

of \mathcal{O}_X -modules splits. In particular, X is globally F -regular if and only if $(X, 0)$ is globally F -regular.

The following well-known proposition gives local constraints on a normal variety to be globally F -regular and echoes [Proposition 3.5](#). For the convenience of the reader, we outline the proof.

Proposition 3.9. *Let X be a normal scheme over \mathbb{F}_p . If X is globally F -regular, then its local rings are strongly F -regular.*

Proof. Fix a point $x \in X$. If $c \in \mathcal{O}_{X,x}$ is a nonzero element, c defines an effective divisor in the neighborhood of x . By taking the Zariski closure, this divisor extends to a Weil divisor D on X such that the map $\mathcal{O}_X \rightarrow \mathcal{O}_X(D)$ localizes to the map $\mathcal{O}_{X,x} \rightarrow \mathcal{O}_{X,x}[c^{-1}]$ sending $1 \mapsto 1$. Since X is globally F -regular, there exists $e > 0$ such that $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ splits. Thus, localizing yields a splitting of the map $\mathcal{O}_{X,x} \rightarrow F_*^e(\mathcal{O}_{X,x}[c^{-1}])$ sending $1 \mapsto 1$. Multiplying by c yields a splitting of the map $\mathcal{O}_{X,x} \rightarrow F_*^e \mathcal{O}_{X,x}$ sending $1 \mapsto F_*^e c$. \square

Global F -regularity is a local property for normal affine varieties. Indeed, a normal affine variety over an F -finite field k is globally F -regular if and only if it is strongly F -regular if and only if all its local rings are strongly F -regular; see [\[HH89, Thm. 3.1\]](#). By [Proposition 3.5](#) and the discussion that follows, we see that a \mathbb{Q} -Gorenstein normal affine variety over an F -finite field is globally F -regular if and only if it is a splinter. In fact, any normal globally F -regular variety is a splinter:

Proposition 3.10 ([\[Bha12, Prop. 8.9\]](#), [\[Bha+23, Lem. 6.14\]](#)). *Let X be a normal excellent Noetherian scheme over \mathbb{F}_p and let Δ be an effective \mathbb{Q} -Weil divisor. If (X, Δ) is globally F -regular, then (X, Δ) is globally $+$ -regular.*

In particular, by [Remark 3.3](#), assuming X is a normal scheme of finite type over a field, if X is globally F -regular, then X is a splinter.

As [Proposition 3.6](#) in the splinter case, global F -regularity imposes strong constraints on the global geometry of normal *projective* varieties:

Proposition 3.11 ([[Smi00](#), Cor. 4.3], [[SS10](#), Thm. 1.1]). *Let X be a normal projective variety over a field of positive characteristic. Assume that X is globally F -regular. Then:*

- (i) *For all nef invertible sheaves \mathcal{L} on X , $H^i(X, \mathcal{L}) = 0$ for all $i > 0$.*
- (ii) *X is log Fano; in particular, if X is in addition \mathbb{Q} -Gorenstein, then $-K_X$ is big.*

A proper normal curve over an algebraically closed field k of positive characteristic is globally F -regular if and only if it is splinter if and only if it is isomorphic to the projective line. It follows from the proof of [[KT25](#), Thm. 6.2] that a smooth projective Fano variety over a perfect field of positive characteristic is globally F -regular if and only if it is a splinter if and only if it is F -split. The following conjecture stems from [Proposition 3.11\(ii\)](#) and the folklore expectation (see, e.g., [[Bha+23](#), Rmk. 6.16]) that splinters should be globally F -regular.

Conjecture 3.12 ([[Bha+23](#), Conj. 6.17]). *Let X be a \mathbb{Q} -Gorenstein projective scheme over a field of positive characteristic. If X is a splinter, then $-K_X$ is big.*

4. INVARIANCE UNDER SMALL BIRATIONAL MAPS

We start by describing the behavior of the splinter property under open embeddings. Recall that the normalization of a Nagata scheme is finite [[Sta24](#), [Tag 035S](#)], and that, if $Y \rightarrow X$ is locally of finite type and X is Nagata, then Y is Nagata [[Sta24](#), [Tag 0359](#)]. Moreover, any excellent scheme is Nagata [[Sta24](#), [Tag 07QS](#)].

Lemma 4.1. *Let X be a Noetherian scheme and let $U \subseteq X$ be an open dense subset.*

- (i) *If X is a splinter, then U is a splinter.*
- (ii) *Assume that X is normal and Nagata, and that $\text{codim}_X(X \setminus U) \geq 2$. If U is a splinter, then X is a splinter.*

Proof. To prove (i) consider a finite cover $f: Y \rightarrow U$. By Zariski's Main Theorem [[Sta24](#), [Tag 05K0](#)], this cover extends to a finite morphism $\bar{f}: \bar{Y} \rightarrow X$. Since X is a splinter, we obtain a section s such that the composition

$$\mathcal{O}_X \rightarrow \bar{f}_* \mathcal{O}_{\bar{Y}} \xrightarrow{s} \mathcal{O}_X$$

is the identity. By restricting to U , we obtain the desired section of $\mathcal{O}_U \rightarrow f_* \mathcal{O}_Y$.

For (ii) consider a finite cover $f: Y \rightarrow X$. Since X is Nagata, so is Y . By possibly replacing Y by its normalization, we can assume that Y is normal. The sheaf $f_* \mathcal{O}_Y$ satisfies the property S_2 by [[Gro65](#), Prop. 5.7.9] and is therefore reflexive. Since U is a splinter, we obtain a splitting of $\mathcal{O}_U \rightarrow f_* \mathcal{O}_Y|_U$ and this extends to a splitting of $\mathcal{O}_X \rightarrow f_* \mathcal{O}_Y$ as X is normal and all the involved sheaves are reflexive. \square

Remark 4.2. Let Δ be an effective \mathbb{Q} -Weil divisor on a normal excellent Noetherian scheme X and let $U \subseteq X$ be an open dense subset. As the proof is analogous to the one of [Lemma 4.1](#), we leave it to the reader to verify that if (X, Δ) is globally $+$ -regular, then $(U, \Delta|_U)$ is globally $+$ -regular. Conversely, if $\text{codim}_X(X \setminus U) \geq 2$ and if $(U, \Delta|_U)$ is globally $+$ -regular, then (X, Δ) is globally $+$ -regular.

Lemma 4.3. *Let $\pi: X \rightarrow Y$ be a surjective morphism of Noetherian schemes. Assume that X is a splinter and that Y is integral with generic point η , then the generic fiber X_η is a splinter.*

Proof. Let $f: Z \rightarrow X_\eta$ be a finite surjective morphism. Since f is of finite presentation, it spreads to a morphism of finite presentation $\mathcal{Z} \rightarrow X|_V$ [[Gro66](#), Thm. 8.8.2] for some dense affine open $V \subseteq Y$, and the latter restricts on a dense affine open $U \subseteq V$ to a finite surjective morphism $\mathcal{Z}|_U \rightarrow X|_U$ [[Gro66](#), Thm. 8.10.5]. By [Lemma 4.1](#) the scheme X_U is a splinter. Thus $\mathcal{O}_{X_U} \rightarrow f_{U*} \mathcal{O}_{\mathcal{Z}|_U}$ admits a section. By flat base change, restricting to the generic fiber yields the desired splitting. \square

We now turn to the analogues of [Lemma 4.1](#) and [Lemma 4.3](#) in the global F -regular setting. The following elementary lemma appears, e.g., in [[Gon+15](#), Lem. 1.5].

Lemma 4.4. *Let (X, Δ) be a pair consisting of a normal variety over a field of positive characteristic and of an effective \mathbb{Q} -Weil divisor Δ , and let $U \subseteq X$ be an open subset.*

- (i) *If (X, Δ) is globally F -regular, then $(U, \Delta|_U)$ is globally F -regular.*
- (ii) *Assume that $U \subseteq X$ is dense and $\text{codim}_X(X \setminus U) \geq 2$. If $(U, \Delta|_U)$ is globally F -regular, then (X, Δ) is globally F -regular.*

Proof. For the proof of (i), consider an effective Weil divisor D_0 on U . Let D be the Zariski closure of D_0 in X . By assumption there exists $e > 0$ such that $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(\lceil (p^e - 1)\Delta \rceil + D)$ splits. Restricting to U yields the desired splitting of $\mathcal{O}_U \rightarrow F_*^e \mathcal{O}_U(\lceil (p^e - 1)\Delta|_U \rceil + D_0)$.

To prove (ii), note that, since F is finite, F_* preserves S_2 sheaves by [Gro65, Prop. 5.7.9], so F_* sends reflexive sheaves to reflexive sheaves. Furthermore, the restriction along $U \hookrightarrow X$ induces a bijection between Weil divisors of X and Weil divisors of U . Statement (ii) follows since, for any Weil divisor D , the map $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(\lceil (p^e - 1)\Delta \rceil + D)$ of reflexive sheaves splits if and only if its restriction to U splits. \square

Lemma 4.5. *Let $\pi: X \rightarrow Y$ be a surjective morphism of normal varieties over a field of positive characteristic. Assume that (X, Δ) is globally F -regular and that Y is integral with generic point η . Then the generic fiber X_η is normal and (X_η, Δ_η) is globally F -regular.*

Proof. By [SS10, Lem. 3.5], X is globally F -regular. It follows from Proposition 3.10 that X is a splinter and then from Lemma 4.3 that X_η is a splinter, so X_η is normal. Given an effective Weil divisor D_0 on X_η , we denote by D its Zariski closure in X . Since (X, Δ) is globally F -regular, there exists $e > 0$ such that $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(\lceil (p^e - 1)\Delta \rceil + D)$ splits. The lemma follows by restricting the splitting along $X_\eta \hookrightarrow X$. \square

Finally, we draw as a consequence of the above that both the splinter property and the global F -regular property are invariant under *small birational maps* of normal varieties.

Definition 4.6 (Small birational map). A rational map $f: X \dashrightarrow Y$ between Noetherian schemes is said to be a *small birational map* if there exist nonempty open subsets $U \subseteq X$ and $V \subseteq Y$ with $\text{codim}_X(X \setminus U) \geq 2$ and $\text{codim}_Y(Y \setminus V) \geq 2$ such that f induces an isomorphism $U \xrightarrow{\sim} V$.

Proposition 4.7. *Let $f: X \dashrightarrow Y$ be a small birational map between normal schemes of finite type over a Noetherian scheme S . The following statements hold:*

- (i) *Assuming S is Nagata, X is a splinter if and only if Y is a splinter.*
- (ii) *Assuming $S = \text{Spec } k$ with k of positive characteristic, X is globally F -regular if and only if Y is globally F -regular.*

Proof. Statement (i) follows directly from Lemma 4.1 and statement (ii) from Lemma 4.4. \square

5. LIFTING AND DESCENDING THE SPLINTER PROPERTY

5.1. Descending the splinter property. If $\pi: Y \rightarrow X$ is a morphism of varieties such that $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$ is an isomorphism, e.g., if π is flat proper with geometrically connected and geometrically reduced fibers [Sta24, Tag 0E0L], or if $\pi: Y \rightarrow X$ is proper birational and X is a normal variety, it is a formal consequence that the splinter property descends along π . Indeed, we have the following lemma.

Lemma 5.1. *Let $\pi: Y \rightarrow X$ be a morphism of Noetherian schemes.*

- (i) *If Y is a splinter and the map $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$ is split, then X is a splinter.*
- (ii) *If Y is a derived splinter and $\mathcal{O}_X \rightarrow R\pi_* \mathcal{O}_Y$ is split, then X is a derived splinter.*

Assume further that X and Y are normal and excellent, and let Δ be an effective \mathbb{Q} -Cartier divisor on X , or let Δ be an effective \mathbb{Q} -Weil divisor and assume that π is finite.

- (iii) *If $(Y, \pi^* \Delta)$ is globally $+$ -regular and $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$ is split, then (X, Δ) is globally $+$ -regular.*

Proof. We only prove (iii) since (i) and (ii) admit similar simpler proofs. First note that the condition that $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$ splits forces π to be dominant. We can assume that X and Y are integral. Thus, the pullback $\pi^* \Delta$ is well-defined if Δ is \mathbb{Q} -Cartier [Sta24, Tag 0200], or if π is finite. Let $f: Z \rightarrow X$ be a finite surjective morphism with Z normal. Let $Z' \rightarrow Y \times_X Z$ be the

normalization of the fiber product, which is finite over Y since Y is excellent. (This step is not needed for the proofs of (i) and (ii) where one simply takes $Z' = Y \times_X Z$.) We obtain a commutative square

$$\begin{array}{ccc} Z' & \xrightarrow{\pi'} & Z \\ \downarrow f' & & \downarrow f \\ Y & \xrightarrow{\pi} & X. \end{array}$$

Since $(Y, \pi^* \Delta)$ is globally $+$ -regular, we have a splitting

$$\mathcal{O}_Y \rightarrow f'_* \mathcal{O}_{Z'}(\lfloor f'^* \pi^* \Delta \rfloor) \xrightarrow{t} \mathcal{O}_Y.$$

Fixing a splitting $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y \xrightarrow{s} \mathcal{O}_X$, we obtain a factorization

$$\mathrm{id}_{\mathcal{O}_X} : \mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y \rightarrow \pi_* f'_* \mathcal{O}_{Z'}(\lfloor f'^* \pi^* \Delta \rfloor) = f_* \pi'_* \mathcal{O}_{Z'}(\lfloor \pi'^* f^* \Delta \rfloor) \xrightarrow{\pi_* t} \pi_* \mathcal{O}_Y \xrightarrow{s} \mathcal{O}_X.$$

Note that $\pi'^* \lfloor f^* \Delta \rfloor \leq \lfloor \pi'^* f^* \Delta \rfloor$. Hence, $\mathcal{O}_X \rightarrow f_* \pi'_* \mathcal{O}_{Z'}(\lfloor \pi'^* f^* \Delta \rfloor)$ factors through $\mathcal{O}_X \rightarrow f_* \mathcal{O}_Z(\lfloor f^* \Delta \rfloor) \rightarrow f_* \pi'_* \pi'^* \mathcal{O}_Z(\lfloor f^* \Delta \rfloor)$ and the lemma follows. \square

Remark 5.2. Assume X and Y are schemes over a field k . If $X \times_k Y$ is a splinter, then so are X and Y . Indeed, if $\pi : X \times_k Y \rightarrow X$ denotes the first projection, then $\pi_* \mathcal{O}_{X \times_k Y} = \mathcal{O}_X \otimes_k H^0(Y, \mathcal{O}_Y)$ and any splitting of the k -linear map $k \rightarrow H^0(Y, \mathcal{O}_Y), 1 \mapsto 1_Y$ provides a splitting of the natural map $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_{X \times_k Y}$.

5.2. Lifting the splinter property. We refer to [Section 2.3](#) for an overview regarding the exceptional inverse image functor for finite morphisms. We will say that a ring homomorphism $A \rightarrow B$ has *reduced closed fibers* if the fibers over closed points of the corresponding map on spectra are reduced; this happens for instance when either B is a field or $A = B$. The following lemma in case (a) extends [DT23, Thm. A] and [Bha+23, Prop. 6.20], where it is shown respectively that the splinter property lifts along étale morphisms of affine Noetherian schemes and along finite étale covers of normal excellent proper schemes over a Noetherian affine base. [Lemma 5.3](#) in cases (a) and (c) is key to the proof of Theorems (D) and (E), while [Lemma 5.3](#) in case (b) (whose proof is much more elementary than in cases (a) and (c)) is key to the proofs of Theorems (A) to (C). Recall from [Section 2.4](#) that, for a finite surjective morphism $\pi : Y \rightarrow X$ of integral normal Noetherian schemes, we denote $\mathrm{tr}_{Y/X}$ the classical trace. Recall from [Lemma 2.10](#) that if π is quasi-étale, then $\pi^! \mathcal{O}_X = \mathcal{O}_Y$ in $\mathrm{Coh}(Y)$ (i.e., $\mathrm{tr}_{Y/X}$ freely generates $\pi^! \mathcal{O}_X := \mathcal{H}om_{\mathcal{O}_X}(\pi_* \mathcal{O}_Y, \mathcal{O}_X)$ as an \mathcal{O}_Y -module), and that the converse implication holds if X is a splinter (since splinters are Cohen–Macaulay).

Lemma 5.3 (Lifting the splinter property). *Let $\pi : Y \rightarrow X$ be a finite surjective morphism of Noetherian schemes. Assume either of the following conditions :*

- (a) *X and Y are integral normal and Nagata, and $\pi^! \mathcal{O}_X = \mathcal{O}_Y$ in $\mathrm{Coh}(Y)$, e.g., π is quasi-étale.*
- (b) *$\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ in $\mathrm{Coh}(Y)$, and either $H^0(Y, \mathcal{O}_Y)$ is a field or $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$.*
- (c) *$\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ in $\mathrm{Coh}(Y)$, X is Nagata, Y is normal, and $\pi^\sharp : H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ has reduced closed fibers.*

If X is a splinter, then Y is a splinter.

Proof. Let $f : Z \rightarrow Y$ be a finite surjective morphism. The splitting of $f^\sharp : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_Z$ is equivalent to the surjectivity of

$$\mathrm{Hom}_{\mathcal{O}_Y}(f_* \mathcal{O}_Z, \mathcal{O}_Y) \xrightarrow{- \circ f^\sharp} \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{O}_Y).$$

The adjunction $\pi_* \dashv \pi^!$ provides the following commutative diagram

$$(5.4) \quad \begin{array}{ccc} \mathrm{Hom}_{\mathcal{O}_Y}(f_*\mathcal{O}_Z, \mathcal{O}_Y) & \xrightarrow{-\circ f^\sharp} & \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{O}_Y) \\ \downarrow \cong & & \downarrow \cong \\ \mathrm{Hom}_{\mathcal{O}_Y}(f_*\mathcal{O}_Z, \pi^!\mathcal{O}_X) & \xrightarrow{-\circ f^\sharp} & \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \pi^!\mathcal{O}_X) \\ \downarrow = & & \downarrow = \\ \mathrm{Hom}_{\mathcal{O}_X}(\pi_*f_*\mathcal{O}_Z, \mathcal{O}_X) & \xrightarrow{-\circ \pi_*f^\sharp} & \mathrm{Hom}_{\mathcal{O}_X}(\pi_*\mathcal{O}_Y, \mathcal{O}_X) \\ \downarrow -\circ(\pi \circ f)^\sharp & & \downarrow -\circ \pi^\sharp \\ \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X) & \xlongequal{\quad} & \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X). \end{array} \quad \begin{array}{c} \text{Tr}_{Y/X} \end{array}$$

Here, $\mathrm{Tr}_{Y/X}$ is simply defined as the composite of the right vertical arrows. In case (a), we choose the isomorphism $\pi^!\mathcal{O}_X \cong \mathcal{O}_Y$ to be the one induced by the classical trace map $\mathrm{tr}_{Y/X}$; under the identifications $\mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{O}_Y) = H^0(Y, \mathcal{O}_Y)$ and $\mathrm{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X) = H^0(X, \mathcal{O}_X)$, the map $\mathrm{Tr}_{Y/X}$ agrees with $\mathrm{tr}_{Y/X}$. Since X is a splinter, the bottom-left vertical arrow $-\circ(\pi \circ f)^\sharp$ is surjective. This implies on the one hand that $-\circ f^\sharp$ is nonzero, and on the other hand that $-\circ \pi^\sharp$ is surjective. This is already sufficient to establish Lemma 5.3 under condition (b). Assuming that $H^0(Y, \mathcal{O}_Y)$ is a field, that $-\circ f^\sharp$ is nonzero yields that $-\circ f^\sharp$ is surjective since it is a map of $H^0(Y, \mathcal{O}_Y)$ -modules. Assuming that $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$, that $-\circ \pi^\sharp$ is surjective yields that the composition of the right vertical arrows, which is $H^0(X, \mathcal{O}_X)$ -linear, is bijective and hence that $-\circ f^\sharp$ is surjective.

We now prove Lemma 5.3 under conditions (a) or (c). The proof works uniformly for both cases, but we refer to Remark 5.8 for a direct proof in case π is étale. We write $R = H^0(X, \mathcal{O}_X)$ and $S = H^0(Y, \mathcal{O}_Y)$. Recall that the map $\coprod \mathrm{Spec} R_z^h \rightarrow \mathrm{Spec} R$, where the coproduct runs through all closed points z of $\mathrm{Spec} R$ and where R_z^h is the henselization of R_z , is faithfully flat [Sta24, Tag 07QM]. We thus have to prove that for all closed points of $\mathrm{Spec} R$ the map $-\circ f^\sharp$ is surjective after base change to R_z^h . Write $\pi^h: Y^h \rightarrow X^h$ and $f^h: Z^h \rightarrow Y^h$ for the pullbacks of π and f along $R \rightarrow R_z^h$, respectively. It is a priori not clear that R is Noetherian; however, since $\mathrm{Spec} R_z^h \rightarrow \mathrm{Spec} R$ is affine, X^h and Y^h are quasi-compact and quasi-separated. By base change for the exceptional inverse image [GW23, Thm. 25.31], we have $h^*\pi^!\mathcal{O}_X = (\pi^h)^\times \mathcal{O}_{X^h}$ in $\mathrm{D}_{\mathrm{QCoh}}(\mathcal{O}_{Y^h})$, where $h: Y^h \rightarrow Y$ is the natural map and $(\pi^h)^\times$ is the right adjoint to $R\pi_*^h: \mathrm{D}_{\mathrm{QCoh}}^+(\mathcal{O}_Y) \rightarrow \mathrm{D}_{\mathrm{QCoh}}^+(\mathcal{O}_X)$. Since h is flat, we have $\mathcal{H}^0(h^*\pi^!\mathcal{O}_X) = h^*\mathcal{H}^0(\pi^!\mathcal{O}_X)$, and so $\mathcal{H}^0((\pi^h)^\times \mathcal{O}_{X^h}) \cong \mathcal{O}_{Y^h}$. As outlined in Section 2.3, $(\pi^h)^\times$ is the right derived functor of $\mathcal{H}om_{\mathcal{O}_{X^h}}(\pi_*^h \mathcal{O}_{Y^h}, -)$ and we thus have $\mathcal{H}om_{\mathcal{O}_{X^h}}(\pi_*^h \mathcal{O}_{Y^h}, \mathcal{O}_{X^h}) \cong \mathcal{O}_{Y^h}$ as \mathcal{O}_{Y^h} -modules. On the other hand, for sheaves $\mathcal{F}, \mathcal{G} \in \mathrm{Coh}(X)$ we have $\mathrm{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) = H^0(X, R\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}))$ [Sta24, Tag 08DK]. Therefore, by flat base change of quasi-coherent cohomology and the compatibility of $R\mathcal{H}om(-, -)$ with flat pullback [GW23, Prop. 22.70], we obtain the same diagram as (5.4) with X replaced by X^h , Y by Y^h , Z by Z^h , $\pi^!$ by $(\pi^h)^\times$, and where $-\circ(\pi^h \circ f^h)^\sharp$ is surjective. We are thus reduced to showing that $-\circ(f^h)^\sharp$ is surjective for every closed point $z \in \mathrm{Spec} R$.

Since a splinter is normal, we have in both cases (a) and (c) that X and Y are normal. Working componentwise, we may and do assume that X and Y are both integral. In particular, as discussed in Section 2.4, both R and S are integrally closed domains in $K(X)$ and $K(Y)$, respectively. Normality of rings is preserved under smooth base change [Sta24, Tag 033C] and under filtered colimits [Sta24, Tag 037D], and hence X^h and Y^h are normal. Since Y^h is quasi-compact, Y^h has only finitely many connected components. Let $Y^h = Y_1 \amalg \cdots \amalg Y_r$ be the decomposition into connected components and let $S \otimes_R R_z^h = H^0(Y^h, \mathcal{O}_{Y^h}) = S_1 \times \cdots \times S_r$, where $S_i = H^0(Y_i, \mathcal{O}_{Y_i})$. By Lemma 2.7, $R \hookrightarrow S$ is integral, hence each S_i is integral over R_z^h .

Recall that R_z^h is the filtered colimit of the pointed étale ring maps $(R_z, z) \rightarrow (T, \mathfrak{q})$ such that $\kappa(z) = \kappa(\mathfrak{q})$. Each projector $S \otimes_R R_z^h \rightarrow S_i$ is defined by an idempotent in $S \otimes_R R_z^h$. Thus, there exists a pointed étale ring map $(R_z, z) \rightarrow (T, \mathfrak{q})$ such that each projector $S \otimes_R R_z^h \rightarrow S_i$ is already defined in $S \otimes_R T$, i.e., $S \otimes_R T = S'_1 \times \cdots \times S'_r$ with $S_i = S'_i \otimes_T R_z^h$. We then have $Y_T := Y \times_{\mathrm{Spec} R} \mathrm{Spec} T = Y'_1 \amalg \cdots \amalg Y'_r$ with $Y_i = Y'_i \times_{\mathrm{Spec} T} S_i$. Note that Y_T is normal Noetherian so that each Y'_i is an integral normal Noetherian scheme. Using that for any normal R -algebra A

and any étale ring morphism $T \rightarrow T'$ the base change map $A \otimes_R T \rightarrow A \otimes_R T'$ is injective (by openness of flat morphisms locally of finite presentation [Sta24, Tag 01UA]), and that a filtered colimit of integral domains is an integral domain, we see that any affine open cover of Y'_i yields an affine open cover of Y_i by integral domains and hence that Y_i is integral.

In particular, the rings S_i are normal domains. We claim that each S_i is a local domain. Indeed, $R_z^h \rightarrow S_i$ is an integral ring map and S_i contains no nontrivial idempotent elements. Since R_z^h is henselian, any finite R_z^h -subalgebra of S_i is a finite product of local rings [Sta24, Tag 04GG (10)]. Since any such subalgebra contains no nontrivial idempotents it is a local ring. Hence, S_i is a filtered colimit of local rings and because S_i is integral over R_z^h , each morphism in the directed system is a morphism of local rings and so S_i is a local ring.

Moving forward, since we are assuming X to be Nagata, after possibly normalizing Z and working on each component separately, we may and do assume that Z is integral and normal. By further taking the normalization of Z in a normal closure of $K(Z)$ over $K(X)$ we may and do assume that $K(Z)/K(X)$ is normal. We summarize the situation in the commutative diagram

$$\begin{array}{ccc} Z_1 \amalg \cdots \amalg Z_r & \xrightarrow{f_1 \amalg \cdots \amalg f_r} & Y_1 \amalg \cdots \amalg Y_r \longrightarrow \mathrm{Spec} S_1 \amalg \cdots \amalg \mathrm{Spec} S_r \\ & \searrow g_1 \amalg \cdots \amalg g_r & \downarrow \pi_1 \amalg \cdots \amalg \pi_r \quad \downarrow \\ & & X^h \longrightarrow \mathrm{Spec} R_z^h \end{array}$$

where $f_i: Z_i \rightarrow Y_i$ is the base change of Z^h along $S \otimes_R R_z^h \rightarrow S_i$ and $g_i = \pi_i \circ f_i: Z_i \rightarrow X^h$.

Claim. The map $\mathcal{O}_{X^h} \rightarrow (g_i)_* \mathcal{O}_{Z_i}$ splits for every $1 \leq i \leq r$.

Proof of the Claim. Let G be the Galois group of $K(Z)$ over $K(X)$ and let $T := H^0(Z, \mathcal{O}_Z)$ be the integral closure of R in $K(Z)$. The field extension $K(Z)^G/K(X)$ is purely inseparable and the ring homomorphism $R \rightarrow T^G$ to the invariant subring T^G , which is the integral closure of R in $K(Z)^G$, is a homeomorphism on spectra. By a straightforward variant of [Sta24, Tag 0BRK] for domains that are integrally closed in a field, G acts transitively on the maximal ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_s$ of T lying over z . As discussed previously in the case of the integral extension $R \rightarrow S$, $T \otimes_R R_z^h$ splits into a product of local rings $T_1 \times \cdots \times T_s$. Let Z'_j be the connected component of Z^h with $H^0(Z'_j, \mathcal{O}_{Z'_j}) = T_j$. The induced action of G on Z^h then permutes the morphisms $g'_j: Z'_j \rightarrow X^h$ and so induces isomorphisms $(g'_j)_* \mathcal{O}_{Z'_j} \cong (g'_k)_* \mathcal{O}_{Z'_k}$ of \mathcal{O}_{X^h} -algebras. Since X is a splinter, the pullback $\mathcal{O}_X \rightarrow \pi_* f_* \mathcal{O}_Z$ splits. By base change, such a splitting induces a splitting

$$\mathcal{O}_{X^h} \rightarrow \bigoplus_{j=1}^r (g'_j)_* \mathcal{O}_{Z'_j} \xrightarrow{\sum_j s_j} \mathcal{O}_{X^h}.$$

Since $\sum_j s_j$ sends the constant section $1_{X^h} \mapsto 1_{X^h} \in R_z^h$, there exists j_0 such that $s_{j_0}(1_{X^h}) \notin \mathfrak{m}_z^h$, where \mathfrak{m}_z^h denotes the maximal ideal of R_z^h . Thus, $\mathcal{O}_{X^h} \rightarrow (g'_{j_0})_* \mathcal{O}_{Z'_{j_0}}$ splits. Using the isomorphisms $(g'_j)_* \mathcal{O}_{Z'_j} \cong (g'_k)_* \mathcal{O}_{Z'_k}$, we see that every $\mathcal{O}_{X^h} \rightarrow (g'_j)_* \mathcal{O}_{Z'_j}$ splits. The claim then follows, as each Z_i is a finite disjoint union of some connected components Z'_j . \square

Now the map $- \circ (f^h)^\#$ decomposes as the direct sum :

$$- \circ \oplus_i f_i^\#: \bigoplus_{i=1}^r \mathrm{Hom}_{\mathcal{O}_{Y_i}}((f_i)_* \mathcal{O}_{Z_i}, \mathcal{O}_{Y_i}) = \mathrm{Hom}_{\mathcal{O}_{Y^h}}(f_* \mathcal{O}_{Z^h}, \mathcal{O}_{Y^h}) \rightarrow \mathrm{Hom}_{\mathcal{O}_{Y^h}}(\mathcal{O}_{Y^h}, \mathcal{O}_{Y^h}) = \bigoplus_{i=1}^r S_i.$$

By the claim above, $\mathcal{O}_{X^h} \rightarrow (g_i)_* \mathcal{O}_{Z_i}$ splits for every i . Thus, the left-vertical arrow in (5.4) base changed to R_z^h remains surjective after restricting to a component Y_i of Y^h . Hence, it is enough to show that any morphism of rings $\varphi_i: T_i \rightarrow S_i$ such that the composite $\mathrm{Tr}_{Y_i/X^h} \circ \varphi_i: T_i \rightarrow R_z^h$ is surjective is also surjective. Denote $\mathfrak{m} \subseteq R$ and $\mathfrak{n}_i \subseteq S_i$ the maximal ideals.

Under condition (c), we still have that $R_z^h \rightarrow S_i$ has reduced closed fibers, and hence $\mathfrak{m}S_i = \mathfrak{n}_i$. By R_z^h -linearity of the map Tr_{Y_i/X^h} , we get $\mathrm{Tr}_{Y_i/X^h}(\mathfrak{n}_i) \subseteq \mathfrak{m}$. Thus, any element $s \in S_i$ such that $\mathrm{Tr}_{Y_i/X^h}(s) = 1$ is a unit in S_i . Since φ_i is S_i -linear, we conclude that φ_i is surjective.

Assume now condition (a). Note that the restriction of the classical trace for Y/X is the classical trace for Y_i/X^h ; indeed, they both agree on the base change along $R \rightarrow R_z^h$ of the flat locus

$U \subseteq X$ of π and then use that X^h is integral. Note that for any integral extension of local rings $\varphi: (A, \mathfrak{a}) \hookrightarrow (B, \mathfrak{b})$ we have $\mathfrak{b} = \sqrt{\mathfrak{a}B}$; this is indeed clear if φ is finite and the general case follows by a limit argument. By [Lemma 2.8](#) and the discussion in [Section 2.4](#), the trace $\mathrm{tr}_{Y_i/X^h}: S_i \rightarrow R_z^h$ sends $\mathfrak{n}_i = \sqrt{\mathfrak{m}S}$ into \mathfrak{m} . We can then conclude as before. \square

Remark 5.5. There are finite étale maps $\pi: Y \rightarrow X$ such that $\pi^\sharp: H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ admits non-reduced closed fibers. Consider for instance a smooth projective surface S over a field k admitting a free action by a finite cyclic group $\mathbb{Z}/n\mathbb{Z}$ with $n > 1$ invertible in k , and consider the action of $\mathbb{Z}/n\mathbb{Z}$ on \mathbb{A}_k^1 given by multiplication by the powers of a primitive n -th root of unity. Then the diagonal action of $\mathbb{Z}/n\mathbb{Z}$ on $Y := S \times_k \mathbb{A}_k^1$ is free and the quotient map π is étale, but the induced map π^\sharp on global sections is $k[T] \rightarrow k[T]$, $T \mapsto T^n$, which has a non-reduced fiber over 0.

Remark 5.6. The proof of [Lemma 5.3\(a\)](#) and [\(c\)](#) simplifies if X is proper over a Noetherian ring as in this case (1) R^h is Noetherian, so that X^h is Noetherian, and (2) $R \hookrightarrow S$ is finite, so that $S^h = S \otimes_R R^h$ splits as a finite product of local domains by [\[Sta24, Tag 04GG \(10\)\]](#).

Remark 5.7. Let $R \hookrightarrow S$ be a finite extension of normal Gorenstein Nagata Noetherian domains with reduced closed fibers. If R is a splinter, then S is a splinter. Indeed, we have to show that for any finite ring extension $S \hookrightarrow T$ the evaluation at 1 map $\mathrm{Hom}_S(T, S) \rightarrow S$ is surjective, or equivalently, that its base change to R_m^h is surjective for any maximal ideal \mathfrak{m} of R . Since R is Nagata, we can assume that T is integrally closed. Writing $S \otimes_R R_m^h = S_1 \times \cdots \times S_r$, with each S_i a henselian domain, we are reduced to showing that each evaluation at 1 map $\mathrm{Hom}_{S_i}(T \otimes_S S_i, S_i) \rightarrow S_i$ is surjective. Since a Noetherian local ring is Gorenstein if and only if its henselization is, and since the Picard group of a local ring is trivial, we have $\pi_i^! R_m^h \cong S_i$, where $\pi_i: R_m^h \hookrightarrow S_i$ is the natural finite ring extension. By the Claim in the proof of [Lemma 5.3](#), the composition $\mathrm{Hom}_{S_i}(T \otimes_S S_i, S_i) \rightarrow S_i \xrightarrow{\mathrm{Tr}_i} R_m^h$ is surjective for all i , where Tr_i is the R_m^h -linear map induced by the isomorphism $\pi_i^! R_m^h \cong S_i$. Since henselization of local rings preserves the closed fiber, we have $\mathfrak{m}S_i = \mathfrak{n}_i$, where \mathfrak{n}_i is the maximal ideal of S_i , and we can conclude as in the proof of [Lemma 5.3\(c\)](#).

Remark 5.8 (Lifting the splinter property along finite étale covers). As mentioned to us by Karl Schwede, [Lemma 5.3](#) admits the following simpler proof in case $\pi: Y \rightarrow X$ is a finite étale morphism of Nagata Noetherian schemes. Assume that X is a splinter and let $f: Z \rightarrow Y$ be a finite surjective morphism. To show that $f^\sharp: \mathcal{O}_Y \rightarrow f_* \mathcal{O}_Z$ splits, it suffices to show that $(f \circ f')^\sharp$ splits for a further finite cover $f': Z' \rightarrow Z$. By dominating Z with the disjoint union of the reductions of its irreducible components, we may assume that Z is integral. By further replacing Z with its normalization in the normal closure of $K(Z)$ over $K(X)$, which remains finite over X since X is assumed Nagata, we may and do assume that Z is normal and that $K(Z)/K(X)$ is a finite normal extension of fields. In that situation, the base change $g_Y: Z \times_X Y \rightarrow Y$ of $g = \pi \circ f: Z \rightarrow X$ identifies with the morphism $\coprod_\sigma g_\sigma: \coprod_\sigma Z_\sigma \rightarrow Y$, where the coproduct runs through all embeddings $\sigma: K(Y) \hookrightarrow K(Z)$ that are the identity on $K(X)$ and where g_σ is the integral closure of Y with respect to the embedding σ . A splitting of $\mathcal{O}_X \rightarrow g_* \mathcal{O}_Z$ provides by base change a splitting of $\mathcal{O}_Y \rightarrow (g_Y)_* \mathcal{O}_W$. If $H^0(Y, \mathcal{O}_Y)$ is a local ring, then arguing as in the proof of the claim there exists σ such that $\mathcal{O}_Y \rightarrow (g_\sigma)_* \mathcal{O}_{Z_\sigma}$ splits, and using the action of $\mathrm{Gal}(K(Z)/K(X))$, which permutes the g_σ 's transitively, we find that $\mathcal{O}_Y \rightarrow (g_\sigma)_* \mathcal{O}_{Z_\sigma}$ splits for all σ . In general, we can reduce to this case by using that $\mathcal{O}_Y \rightarrow (g_\sigma)_* \mathcal{O}_{Z_\sigma}$ splits if and only if its base change to the localizations of all closed points of $\mathrm{Spec} H^0(Y, \mathcal{O}_Y)$ splits; see the proof of [Lemma 7.3](#) below. In particular, since $g_{\mathrm{id}} = f: Z_{\mathrm{id}} = Z \rightarrow Y$, we find that f^\sharp splits, as desired.

Remark 5.9 (Lifting the F -split property). Let $\pi: Y \rightarrow X$ be a finite surjective morphism of Noetherian schemes over \mathbb{F}_p . Assume either that π is étale with X normal, or that condition [\(b\)](#) is fulfilled with the additional assumption that $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$ splits. If X is F -split, then Y is F -split.

In the étale case, this is [\[PZ20, Lem. 11.1\]](#). In the other case, we argue as in the proof of [Lemma 5.3\(b\)](#) with $f: Z \rightarrow Y$ replaced by the Frobenius $F: Y \rightarrow Y$. If $s: \pi_* \mathcal{O}_Y \rightarrow \mathcal{O}_X$ is a splitting of the map $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$, then, with a Frobenius splitting of X , we obtain a diagram

$$\mathcal{O}_X \rightarrow \pi_* F_* \mathcal{O}_Y = F_* \pi_* \mathcal{O}_Y \xrightarrow{F_*(s)} F_* \mathcal{O}_X \rightarrow \mathcal{O}_X,$$

where the composition is the identity. This proves that the bottom-left vertical arrow $- \circ (\pi \circ F)^\sharp$ of (5.4) is surjective. As in the splinter case, we deduce that $- \circ F^\sharp$ is surjective, i.e., that Y is F -split.

Lemma 5.10 (Lifting the derived splinter property). *Let $\pi: Y \rightarrow X$ be a proper surjective morphism of Noetherian schemes. Assume $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ in $\mathbf{D}_{\text{Coh}}(\mathcal{O}_Y)$, and either $H^0(Y, \mathcal{O}_Y)$ is a field or $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$. If X is a derived splinter, then Y is a derived splinter.*

Proof. This is proved as in Lemma 5.3(b) by using the adjunction $R\pi_* \dashv \pi^!$ for a proper morphism $\pi: Y \rightarrow X$ instead of the adjunction $\pi_* \dashv \pi^!$ for a finite morphism $\pi: Y \rightarrow X$. \square

A morphism of schemes $\pi: Y \rightarrow X$ is said to be a *crepant morphism* if it is proper, birational, and is such that $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$. Note that, by Lemma 2.2, if X is assumed to be Gorenstein, the latter condition is equivalent to $\pi^* \omega_X = \omega_Y$.

Proposition 5.11 (The splinter property in positive characteristic lifts along crepant morphisms). *Let $\pi: Y \rightarrow X$ be a crepant morphism of Noetherian schemes.*

- (i) *If $\pi: Y \rightarrow X$ is defined over \mathbb{F}_p and if X is a splinter, then Y is a splinter.*
- (ii) *If X is a derived splinter, then Y is a derived splinter.*
- (iii) *If X is F -split and Y is integral, then Y is F -split.*

Proof. If X is a derived splinter, then it is normal so that $\pi_* \mathcal{O}_Y = \mathcal{O}_X$ and hence $\pi^\sharp: H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ is an isomorphism. We can then apply Lemma 5.10. The statement (i) follows from the fact [Bha12, Thm. 1.4] that splinters in positive characteristic agree with derived splinters.

For (iii), choose a splitting $\mathcal{O}_X \rightarrow F_* \mathcal{O}_X \xrightarrow{s} \mathcal{O}_X$. Let \tilde{s} be the image of s under

$$\begin{aligned} \text{Hom}_{\mathcal{O}_X}(F_* \mathcal{O}_X, \mathcal{O}_X) &= \text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, F^! \mathcal{O}_X) \rightarrow \text{Hom}_{\mathcal{O}_Y}(\pi^! \mathcal{O}_X, \pi^! F^! \mathcal{O}_X) \\ &= \text{Hom}_{\mathcal{O}_Y}(\pi^! \mathcal{O}_X, F^! \pi^! \mathcal{O}_X) \cong \text{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, F^! \mathcal{O}_Y) = \text{Hom}_{\mathcal{O}_Y}(F_* \mathcal{O}_Y, \mathcal{O}_Y). \end{aligned}$$

Then $\mathcal{O}_Y \rightarrow F_* \mathcal{O}_Y \xrightarrow{\tilde{s}} \mathcal{O}_Y$ is the identity. Indeed, since Y is integral, this can be verified after restricting to the dense open subset where π is an isomorphism so that the claim follows from the fact that s is a splitting of $\mathcal{O}_X \rightarrow F_* \mathcal{O}_X$. \square

Remark 5.12. First, Proposition 5.11(ii) should be compared to Theorem 12.15 below. Second, Proposition 5.11(iii) notably removes the Gorenstein assumption in [BK05, Lem. 1.3.13] where it is shown that if $\pi: Y \rightarrow X$ is a crepant morphism of normal F -finite schemes over \mathbb{F}_p with X Gorenstein and F -split, then Y is F -split. Third, regarding Proposition 5.11(i), we have the following related result. Let $\pi: Y \rightarrow X$ be a birational morphism of integral normal schemes proper over a complete Noetherian local domain with positive characteristic residue field. Assume that X is \mathbb{Q} -Gorenstein and that $\pi^* K_X = K_Y$ in $\text{Pic}(Y) \otimes \mathbb{Q}$. It is shown in [Bha+23, Lem. 4.19] that if X is a splinter, then Y is a splinter.

For the sake of completeness, we mention that Lemma 5.3 also holds for globally $+$ -regular pairs:

Lemma 5.13 (Lifting global $+$ -regularity). *Let $\pi: Y \rightarrow X$ be a finite surjective morphism of normal excellent Noetherian schemes and let Δ be an effective \mathbb{Q} -Weil divisor on X . Assume either of the following conditions:*

- (a) *π is quasi-étale.*
- (b) *$\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ in $\text{Coh}(Y)$, and $\pi^\sharp: H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ has reduced closed fibers.*

If (X, Δ) is globally $+$ -regular, then $(Y, \pi^ \Delta)$ is globally $+$ -regular.*

Proof. Let $f: Z \rightarrow Y$ be a finite surjective morphism with Z normal. We have to show that the map $f_{\pi^* \Delta}^\sharp: \mathcal{O}_Y \rightarrow f_* \mathcal{O}_Z([f^* \pi^* \Delta])$ splits, or equivalently, that

$$\text{Hom}_{\mathcal{O}_Y}(f_* \mathcal{O}_Z([f^* \pi^* \Delta]), \mathcal{O}_Y) \xrightarrow{- \circ f_{\pi^* \Delta}^\sharp} \text{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{O}_Y)$$

is surjective. Note that the map $(\pi \circ f)_\Delta^\sharp: \mathcal{O}_X \rightarrow \pi_* f_* \mathcal{O}_Z(\lfloor f^* \pi^* \Delta \rfloor)$ factors through $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$. Hence, the following diagram commutes:

$$\begin{array}{ccc}
\mathrm{Hom}_{\mathcal{O}_Y}(f_* \mathcal{O}_Z(\lfloor f^* \pi^* \Delta \rfloor), \mathcal{O}_Y) & \xrightarrow{- \circ f_{\pi^* \Delta}^\sharp} & \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{O}_Y) \\
\downarrow \cong & & \downarrow \cong \\
\mathrm{Hom}_{\mathcal{O}_Y}(f_* \mathcal{O}_Z(\lfloor f^* \pi^* \Delta \rfloor), \pi^! \mathcal{O}_X) & \xrightarrow{- \circ f_{\pi^* \Delta}^\sharp} & \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \pi^! \mathcal{O}_X) \\
\downarrow = & & \downarrow = \\
\mathrm{Hom}_{\mathcal{O}_X}(\pi_* f_* \mathcal{O}_Z(\lfloor f^* \pi^* \Delta \rfloor), \mathcal{O}_X) & \xrightarrow{- \circ \pi_* f_{\pi^* \Delta}^\sharp} & \mathrm{Hom}_{\mathcal{O}_X}(\pi_* \mathcal{O}_Y, \mathcal{O}_X) \\
\downarrow - \circ (\pi \circ f)_\Delta^\sharp & & \downarrow - \circ \pi^\sharp \\
\mathrm{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X) & \xlongequal{\quad} & \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X).
\end{array}$$

We then proceed as in the proof of [Lemma 5.3](#). \square

5.3. The splinter property and base change of field. The following proposition establishes the invariance of the splinter property for proper schemes over a field under algebraic base change. We refer to [Remark 6.10](#) for base change along any field extension.

Proposition 5.14. *Let X be a scheme such that $H^0(X, \mathcal{O}_X)$ is a field, e.g., X is proper, connected, and reduced over a field k . Then, for any algebraic field extension L of $H^0(X, \mathcal{O}_X)$, the scheme $X_L := X \times_{H^0(X, \mathcal{O}_X)} L$ is a splinter if and only if X is a splinter.*

Proof. The “only if” part of the proposition follows from [Lemma 5.1](#). Let $K := H^0(X, \mathcal{O}_X)$. First assume L is a finite extension of K and let $h: \mathrm{Spec} L \rightarrow \mathrm{Spec} K$. Then

$$h^! \mathcal{O}_{\mathrm{Spec} K} = \mathrm{Hom}_K(L, K) \cong L = \mathcal{O}_{\mathrm{Spec} L}$$

as L -vector spaces. By base change for the exceptional inverse image [\[Sta24, Tag 0E9U\]](#), there exists an isomorphism $\pi^! \mathcal{O}_X \cong \mathcal{O}_{X_L}$, where $\pi: X_L \rightarrow X$ is the base change of h along $X \rightarrow \mathrm{Spec} K$. We conclude by [Lemma 5.3](#) that X_L is a splinter, since

$$H^0(X_L, \mathcal{O}_{X_L}) = H^0(X, \mathcal{O}_X) \times_K L = L$$

by flat base change. Now assume $K \rightarrow L$ is any algebraic field extension and take a finite cover $f: Y \rightarrow X_L$. Since f is defined by finitely many equations and L is algebraic over K , we can find a finite cover $f': Y' \rightarrow X_{L'}$ such that $K \subseteq L' \subseteq L$ is an intermediate extension, finite over K , and $f = f' \times_{L'} L$ is the base change of f' . By the previous argument $X_{L'}$ is a splinter, thus $\mathcal{O}_{X_{L'}} \rightarrow f'_* \mathcal{O}_{Y'}$ admits a section $s: f'_* \mathcal{O}_{Y'} \rightarrow \mathcal{O}_{X_{L'}}$. Now pulling back to X_L and using flat base change, we obtain the desired section of $\mathcal{O}_{X_L} \rightarrow f_* \mathcal{O}_Y$. \square

Corollary 5.15. *Let X be a connected proper scheme over a field k such that $H^0(X, \mathcal{O}_X)$ is a separable extension of k . Then, for any algebraic field extension K of k , the scheme $X_K := X \times_k K$ is a splinter if and only if X is a splinter.*

Proof. The “only if” part of the corollary follows from [Lemma 5.1](#). Let \bar{k} be an algebraic closure of k . By the assumption on $H^0(X, \mathcal{O}_X)$, $X_{\bar{k}} := X \times_k \bar{k}$ is isomorphic to the disjoint union of $\dim_k H^0(X, \mathcal{O}_X)$ copies of $X \times_{H^0(X, \mathcal{O}_X)} \bar{k}$. By the “if” part of [Proposition 5.14](#), if X is a splinter, then $X_{\bar{k}}$ is a splinter. By [Lemma 5.1](#), we get that X_K is a splinter. \square

Corollary 5.16. *Let X be a connected proper scheme over a field k . If X is a splinter, then $H^0(X, \mathcal{O}_X)$ is a field, and X , considered as a scheme over $\mathrm{Spec} H^0(X, \mathcal{O}_X)$, is geometrically normal.*

Proof. By [Proposition 3.5](#), a splinter is normal. In particular, $H^0(X, \mathcal{O}_X)$ is a field. The corollary then follows from [Proposition 5.14](#). \square

Remark 5.17 (Algebraic base change for globally $+$ -regular pairs). Let X be a connected normal proper scheme over a field k such that $H^0(X, \mathcal{O}_X)$ is a field, and let Δ be an effective \mathbb{Q} -Weil divisor on X . Let L be an algebraic field extension of $H^0(X, \mathcal{O}_X)$. The arguments of the proof of

Proposition 5.14 show that if (X, Δ) is globally $+$ -regular, then $X_L := X \times_{H^0(X, \mathcal{O}_X)} L$ is normal and the pair (X_L, Δ_L) is globally $+$ -regular.

Assume in addition that $H^0(X, \mathcal{O}_X)$ is a separable extension of k , and let K be any algebraic extension of k . As in [Corollary 5.15](#), we have that if (X, Δ) is globally $+$ -regular, then $X_K := X \times_k K$ is normal and the pair (X_K, Δ_K) is globally $+$ -regular.

6. LIFTING AND DESCENDING GLOBAL F -REGULARITY

The first aim of this section is to show that the results of [Section 5](#) regarding splinters, notably [Lemma 5.3](#), extend to the globally F -regular setting; see [Proposition 6.4](#). The second aim is to show how the criterion of Schwede–Smith (recalled in [Theorem 6.1](#)) can be used to establish further results in the global F -regular setting; for instance, we show in [Proposition 6.7](#) that global F -regularity is stable under product and, combined with [Proposition 6.4](#), use it to recover in [Proposition 6.9](#) a result of [\[Gon+15\]](#) stating that global F -regularity for normal proper schemes over a field is stable under base change of F -finite fields. Except for [Theorem 6.1](#), which is due to Schwede–Smith, the results of this section will not be used in the rest of the paper.

6.1. A criterion for global F -regularity. The following criterion of Schwede–Smith makes it possible in practice to reduce checking that a variety is globally F -regular to simply check that $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ splits for one specific Weil divisor D .

Theorem 6.1 ([\[SS10, Thm. 3.9\]](#)). *Let X be a normal variety over an F -finite field of positive characteristic. Then X is globally F -regular if and only if there exists an effective Weil divisor D on X such that*

- (i) *there exists an $e > 0$ such that the natural map $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ splits, and*
- (ii) *the variety $X \setminus D$ is globally F -regular.*

Remark 6.2. Suppose X is a normal projective variety over an F -finite field of positive characteristic. If D is an ample divisor on X , the variety $X \setminus D$ is affine and therefore globally F -regular if and only if its local rings are strongly F -regular. Since regular local rings are strongly F -regular, a smooth projective variety X over an F -finite field of positive characteristic is globally F -regular if and only if $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ splits for some ample divisor D .

6.2. Descending global F -regularity. The following [Lemma 6.3](#), which is due to Schwede–Smith, holds in particular when the map $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$ is an isomorphism, e.g., when $\pi: Y \rightarrow X$ is flat proper with geometrically connected and geometrically reduced fibers [\[Sta24, Tag 0E0L\]](#), or when $\pi: Y \rightarrow X$ is proper birational and X is a normal variety.

Lemma 6.3 ([\[SS10, Cor. 6.4\]](#)). *Let $\pi: Y \rightarrow X$ be a morphism of varieties over a field of positive characteristic. If Y is normal globally F -regular and if the map $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_Y$ is split, then X is normal globally F -regular.*

Proof. By [Proposition 3.10](#), Y is a splinter and, by [Lemma 5.1](#), X is a splinter. Hence, by [Proposition 3.5](#), X is normal. We can now apply [\[SS10, Cor. 6.4\]](#). \square

6.3. Lifting global F -regularity. The following [Proposition 6.4](#) is the analogue of [Lemma 5.3](#) and [Proposition 5.11](#). [Proposition 6.4\(a\)](#) was previously established in [\[PZ20, Lem. 11.1\]](#) and, under the more restrictive assumptions that X is projective and Gorenstein and that π is birational, [Proposition 6.4\(d\)](#) was established in [\[GT16, Lem. 3.3\]](#).

Proposition 6.4. *Let $\pi: Y \rightarrow X$ be a proper surjective morphism of varieties over a field k of characteristic $p > 0$. Assume either of the following conditions:*

- (a) *π is finite quasi-étale, and Y is normal.*
- (b) *π is finite, $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ in $\text{Coh}(Y)$, and either $H^0(Y, \mathcal{O}_Y)$ is a field or $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$.*
- (c) *π is finite, $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ in $\text{Coh}(Y)$, Y is normal, and $\pi^\sharp: H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ has reduced closed fibers.*
- (d) *X is quasi-projective, $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ in $\text{D}_{\text{Coh}}(\mathcal{O}_Y)$, and either $H^0(Y, \mathcal{O}_Y)$ is a field or $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$.*

If X is a normal globally F -regular variety, then Y is a normal globally F -regular variety.

Proof. We first prove (a), (b) and (c). By Proposition 3.10, X is a splinter, and it follows from Lemma 5.3 that Y is a splinter and hence is normal. By normality of X , the complement of X_{reg} has codimension at least 2, and by finiteness of π , the complement of $\pi^{-1}(X_{\text{reg}})$ in Y has codimension at least 2. Therefore, by Lemma 4.4, Y is globally F -regular if and only if $\pi^{-1}(X_{\text{reg}}) \subseteq Y$ is globally F -regular. Replacing X by X_{reg} we can and do assume that X is regular. Since Y is normal, Y_{sing} is a proper closed subset of Y and since π is finite, $\pi(Y_{\text{sing}}) \subseteq X$ is also a proper closed subset. Let $U \subseteq X$ be an affine open subset in the complement of $\pi(Y_{\text{sing}})$. By [Sta24, Tag 0BCU], $D := X \setminus U$ has codimension 1, so defines a Weil divisor on X . Since X is regular, D is further Cartier and $\pi^*\mathcal{O}_X(D)$ is invertible. Note that the pullback $\pi^*D = \pi^{-1}(D)$ of D is a Cartier divisor [Sta24, Tag 0200]. Let $\sigma_D: \mathcal{O}_X \rightarrow \mathcal{O}_X(D)$ be the global section defined by the divisor D . Then the pullback $\pi^*\sigma_D$ defines a global section of $\pi^*\mathcal{O}_X(D)$ whose zero locus is precisely $\pi^{-1}(D)$. Thus $\pi^*\sigma_D$ is the global section of $\pi^*\mathcal{O}_X(D) = \mathcal{O}_Y(\pi^*D)$ defined by the divisor π^*D [Sta24, Tag 0C4S].

Now $\pi^{-1}(U) = Y \setminus \pi^*D$ is an affine open subset contained in the regular locus of Y , so is strongly F -regular. By Theorem 6.1, it is enough to show that there exists an $e > 0$ such that the map $\mathcal{O}_Y \rightarrow F_*^e \mathcal{O}_Y(\pi^*D)$ splits. Since X is globally F -regular, there exists an integer $e > 0$ and a splitting s such that

$$\text{id}_{\mathcal{O}_X}: \mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X \xrightarrow{F_*^e(\sigma_D)} F_*^e \mathcal{O}_X(D) \xrightarrow{s} \mathcal{O}_X.$$

Since X is a splinter, we have a splitting

$$\text{id}_{\mathcal{O}_X}: \mathcal{O}_X \xrightarrow{\pi^\sharp} \pi_* \mathcal{O}_Y \xrightarrow{t} \mathcal{O}_X.$$

Note that by the projection formula, t induces a splitting of $\mathcal{O}_X(D) \rightarrow \pi_* \pi^* \mathcal{O}_X(D) = \pi_* \mathcal{O}_Y(\pi^*D)$ which by abuse we still denote by t . The composite map

$$(6.5) \quad \mathcal{O}_X \xrightarrow{\pi^\sharp} \pi_* \mathcal{O}_Y \xrightarrow{\pi_* F_*^e} \pi_* F_*^e \mathcal{O}_Y \xrightarrow{\pi_* F_*^e(\pi^* \sigma_D)} \pi_* F_*^e \mathcal{O}_Y(\pi^* D)$$

then admits

$$\pi_* F_*^e \mathcal{O}_Y(\pi^* D) = F_*^e \pi_* \mathcal{O}_Y(\pi^* D) \xrightarrow{F_*^e(t)} F_*^e \mathcal{O}_X(D) \xrightarrow{s} \mathcal{O}_X$$

as a splitting. As in Lemma 5.3, we consider the diagram

$$\begin{array}{ccc} \text{Hom}_{\mathcal{O}_Y}(F_*^e \mathcal{O}_Y(\pi^* D), \mathcal{O}_Y) & \xrightarrow{-\circ \pi^* \sigma_D} & \text{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{O}_Y) \\ \downarrow \cong & & \downarrow \cong \\ \text{Hom}_{\mathcal{O}_Y}(F_*^e \mathcal{O}_Y(\pi^* D), \pi^! \mathcal{O}_X) & \xrightarrow{-\circ \pi^* \sigma_D} & \text{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \pi^! \mathcal{O}_X) \\ \downarrow = & & \downarrow = \\ \text{Hom}_{\mathcal{O}_X}(\pi_* F_*^e \mathcal{O}_Y(\pi^* D), \mathcal{O}_X) & \xrightarrow{-\circ \pi_* F_*^e(\pi^* \sigma_D) \circ \pi_* F_*^e} & \text{Hom}_{\mathcal{O}_X}(\pi_* \mathcal{O}_Y, \mathcal{O}_X) \\ \downarrow -\circ \pi_* F_*^e(\pi^* \sigma_D) \circ \pi_* F_*^e \circ \pi^\sharp & & \downarrow -\circ \pi^\sharp \\ \text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X) & \xlongequal{\quad} & \text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X). \end{array}$$

Since the splitting of (6.5) is equivalent to the left-vertical map $-\circ \pi_* F_*^e(\pi^* \sigma_D) \circ \pi_* F_*^e \circ \pi^\sharp$ being surjective, in case (b) we conclude, as in the proof of Lemma 5.3, that the map $-\circ \pi^* \sigma_D$ is surjective, or equivalently, that $\mathcal{O}_Y \rightarrow F_*^e \mathcal{O}_Y(\pi^* D)$ splits. In cases (a) and (c), we argue again as in the proof of Lemma 5.3, whose notation we take up. Let $R = H^0(X, \mathcal{O}_X)$ and $z \in \text{Spec } R$ be a closed point. Since henselization is a filtered colimit of étale ring maps, the base change of the absolute Frobenius $F: Y \rightarrow Y$ along $Y^h \rightarrow Y$ identifies with the absolute Frobenius of Y^h . Let $\pi': Y' \rightarrow Y$ be the normalization of Y in a normal closure of $K(Y)$ over $K(X)$. Arguing as above we obtain a splitting of the map $\mathcal{O}_X \rightarrow \pi_* \pi'^* F_*^e \mathcal{O}_{Y'}(\pi'^* \pi^* D)$. This provides a splitting of the base change $\mathcal{O}_{X^h} \rightarrow \pi_*^h \pi'^h F_*^e \mathcal{O}_{Y'^h}(\pi'^h \pi^h D|_{X^h})$. We have a decomposition into connected components $Y^h = Y_1 \amalg \dots \amalg Y_r$ and $Y'^h = Y'_1 \amalg \dots \amalg Y'_s$, where the induced action of $\text{Gal}(K(Y')/K(X))$ permutes the morphisms $Y'_j \rightarrow X^h$. Hence, we obtain a splitting of $\mathcal{O}_{X^h} \rightarrow \pi_*^h \pi'^h F_*^e \mathcal{O}_{Y'^h}(\pi'^h \pi^h D|_{X^h})|_{Y'_j}$ for every j . As every Y_i is dominated by one Y'_j , we obtain that $\mathcal{O}_{X^h} \rightarrow \pi_*^h F_*^e \mathcal{O}_{Y^h}(\pi^h D|_{X^h})|_{Y_i}$ splits for every i . We then conclude as in the proof of Lemma 5.3.

We now prove (d). By Proposition 3.10, X is a splitter, so a derived splinters since k is of positive characteristic [Bha12, Thm. 1.4]. By Remark 2.4, π is generically finite. Let $U \subseteq X$ be an affine regular dense open subset such that the restriction of π to U is finite. After possibly further shrinking U we can assume that $\pi^{-1}(U)$ is also regular. Since restriction to open subschemes commutes with exceptional inverse image functors [Sta24, Tag 0G4J], we have $\pi|_U^! \mathcal{O}_U \cong \mathcal{O}_{\pi^{-1}(U)}$. Since X is quasi-projective, any Weil divisor on X is dominated by a Cartier divisor. Thus, by possibly further shrinking U , we may and do assume that $D := X \setminus U$ is a Cartier divisor. Since $\pi^{-1}(U)$ is affine and regular, it is enough to show, by Theorem 6.1, that there exists an $e > 0$ such that the map $\mathcal{O}_Y \rightarrow F_*^e \mathcal{O}_Y(\pi^* D)$ splits. By arguing as in case (b) with π_* replaced by $R\pi_*$, we conclude that a splitting of $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ induces a splitting of $\mathcal{O}_Y \rightarrow F_*^e \mathcal{O}_Y(\pi^* D)$. \square

Remark 6.6. By using the version of Theorem 6.1 for pairs, i.e., the original [SS10, Thm. 3.9], we leave it to the reader to show the following version of Proposition 6.4 for pairs. Let $\pi: Y \rightarrow X$ be a finite surjective morphism of varieties over a field of positive characteristic that satisfies (a), (b), or (c), and let Δ be an effective \mathbb{Q} -Weil divisor on X . If X is normal and if (X, Δ) is globally F -regular, then Y is normal and $(Y, \pi^* \Delta)$ is globally F -regular.

6.4. Products of globally F -regular varieties. As far as we know, it is unknown whether the splitter property is stable under product. On the other hand, global F -regularity for products is more tractable since the Frobenius of a product is the product of the Frobenius and since one may use the criterion of Theorem 6.1 to check global F -regularity for one specific divisor. The following Proposition 6.7 generalizes [Has03, Thm. 5.2], where the case of products of projective globally F -regular varieties was dealt with by taking affine cones.

Proposition 6.7. *Let X and Y be normal varieties over a perfect field k of positive characteristic. Then, X and Y are globally F -regular if and only if their product $X \times_k Y$ is globally F -regular.*

Proof. Denote by π_X and π_Y the natural projections from $X \times_k Y$ to X and Y , respectively. Since X and Y are normal and k is perfect, their product $X \times_k Y$ is normal, see [Sta24, Tag 038L]. Moreover, $X \setminus X_{\text{reg}}$ and $Y \setminus Y_{\text{reg}}$ both have codimension ≥ 2 and thus $X \times_k Y \setminus X_{\text{reg}} \times_k Y_{\text{reg}}$ has codimension ≥ 2 . Therefore, by Lemma 4.4, we can and do assume without loss of generality that X and Y are smooth over k .

Assume first that $X \times_k Y$ is globally F -regular. As in Remark 5.2, since $\pi_* \mathcal{O}_{X \times_k Y} = \mathcal{O}_X \otimes_k H^0(Y, \mathcal{O}_Y)$, any splitting of the k -linear map $k \rightarrow H^0(Y, \mathcal{O}_Y), 1 \mapsto 1_Y$ provides a splitting to the natural map $\mathcal{O}_X \rightarrow \pi_* \mathcal{O}_{X \times_k Y}$. By Lemma 6.3, it follows that X is globally F -regular.

For the converse, we first note that there exist effective Cartier divisors D on X and E on Y such that $X \setminus D$ and $Y \setminus E$ are affine. Indeed, since X and Y are normal and k is perfect, they admit dense affine open subsets, and then use the fact that the complement of a dense affine open subset is a divisor by [Sta24, Tag 0BCU]. The divisors obtained this way are a priori Weil divisors, but since X and Y are smooth, they are actually Cartier divisors. Since k is assumed to be perfect,

$$X \setminus D \times_k Y \setminus E = X \times_k Y \setminus \pi_X^* D \cup \pi_Y^* E$$

is smooth and affine, so in particular strongly F -regular. By Theorem 6.1 it is enough to show that the map

$$\mathcal{O}_{X \times Y} \rightarrow F_*^e \mathcal{O}_{X \times Y} \rightarrow F_*^e \mathcal{O}_{X \times Y}(\pi_X^* D + \pi_Y^* E)$$

splits for some $e > 0$. Since X is globally F -regular, we can find an $e > 0$ such that $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ splits and similarly for Y . As remarked in [Smi00, p. 558], if $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ splits for some $e > 0$, then $\mathcal{O}_X \rightarrow F_*^{e'} \mathcal{O}_X(D)$ splits for all $e' \geq e$. Thus, there exists an integer $e > 0$ such that both $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ and $\mathcal{O}_Y \rightarrow F_*^e \mathcal{O}_Y(E)$ split. The morphism

$$\sigma_{\pi_X^* D + \pi_Y^* E}: \mathcal{O}_{X \times Y} \rightarrow \mathcal{O}_{X \times Y}(\pi_X^* D + \pi_Y^* E)$$

can be identified with the tensor product $\pi_X^* \sigma_D \otimes \pi_Y^* \sigma_E$, where

$$\sigma_D: \mathcal{O}_X \rightarrow \mathcal{O}_X(D) \quad \text{and} \quad \sigma_E: \mathcal{O}_Y \rightarrow \mathcal{O}_Y(E)$$

denote the corresponding morphisms on X and Y . Pushing forward along Frobenius, we obtain

$$F_*^e \sigma_{\pi_X^* D + \pi_Y^* E} = \pi_X^* F_*^e \sigma_D \otimes \pi_Y^* F_*^e \sigma_E.$$

We conclude, by taking the tensor product of the sections of $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ and $\mathcal{O}_Y \rightarrow F_*^e \mathcal{O}_Y(E)$, that

$$\mathcal{O}_{X \times_k Y} \rightarrow F_*^e \mathcal{O}_{X \times_k Y}(\pi_X^* D + \pi_Y^* E)$$

splits. Hence $X \times_k Y$ is globally F -regular. \square

Remark 6.8. By using the version of [Theorem 6.1](#) for pairs, i.e., the original [\[SS10, Thm. 3.9\]](#), we leave it to the reader to show the following version of [Proposition 6.7](#) for pairs: Let X and Y be normal varieties over a perfect field k of positive characteristic, and denote by $\pi_X: X \times_k Y \rightarrow X$ and $\pi_Y: X \times_k Y \rightarrow Y$ the natural projections. Let Δ_X and Δ_Y be effective \mathbb{Q} -Weil divisors on X and Y , respectively. Then, (X, Δ_X) and (Y, Δ_Y) are globally F -regular if and only if $(X \times_k Y, \pi_X^* \Delta_X + \pi_Y^* \Delta_Y)$ is globally F -regular.

6.5. Global F -regularity and base change of field. By using the lifting [Proposition 6.4](#), it is possible to show that the base change results for splinters along algebraic field extensions of [Section 5.3](#) also hold for normal global F -regular varieties. However, by using the criterion of Schwede–Smith [\[SS10, Thm. 3.9\]](#), Gongyo–Li–Patakfalvi–Schwede–Tanaka–Zong [\[Gon+15\]](#) have established a more general base change results that deals with not necessarily algebraic extensions.

Proposition 6.9 (Gongyo–Li–Patakfalvi–Schwede–Tanaka–Zong [\[Gon+15\]](#)). *Let X be a normal proper scheme over a field k of positive characteristic. Assume that (X, Δ) is globally F -regular. Then, for any F -finite field extension L of k with a morphism $\text{Spec } L \rightarrow \text{Spec } H^0(X, \mathcal{O}_X)$, the scheme $X_L := X \times_{H^0(X, \mathcal{O}_X)} L$ is normal and the pair (X_L, Δ_L) is globally F -regular.*

Proof. Let us provide an alternate proof. We may and do assume X is connected. Since any divisor on X_L is defined over a finitely generated field extension of the field $H^0(X, \mathcal{O}_X)$, we may assume that L is a simple extension of $H^0(X, \mathcal{O}_X)$. If L is algebraic, we can apply [Proposition 6.4](#), while if L is purely transcendental, we can apply [Proposition 6.7](#) (or [Remark 6.8](#) in case $\Delta \neq 0$) and [Lemma 4.5](#) to $X \times_{H^0(X, \mathcal{O}_X)} \mathbb{A}^1 \rightarrow \mathbb{A}^1$. Note that in this situation it is not necessary to assume that $H^0(X, \mathcal{O}_X)$ is perfect as $X_{\text{reg}} \times_{H^0(X, \mathcal{O}_X)} \mathbb{A}^1$ is regular. \square

Remark 6.10. Our proof of [Proposition 6.9](#) shows that one could extend [Proposition 5.14](#) concerned with base change of splinters along algebraic field extensions to arbitrary field extensions if one could establish that the splinter property is stable under taking product with the affine line \mathbb{A}^1 .

Corollary 6.11 ([\[Gon+15, Cor. 2.8\]](#)). *Let X be a connected normal proper scheme over an F -finite field k . Assume that $H^0(X, \mathcal{O}_X)$ is a separable extension of k and that (X, Δ) is globally F -regular. Then, for any F -finite field extension K of k , the scheme $X_K := X \times_k K$ is normal and the pair (X_K, Δ_K) is globally F -regular.*

7. FINITE TORSORS OVER SPLINTERS

We say that a morphism $\pi: Y \rightarrow X$ of schemes over a scheme S is a *finite torsor* over S if it is a fppf torsor under a finite locally free group scheme G over S . (The reason for not restricting ourselves to finite flat group schemes over a Noetherian scheme is that $H^0(X, \mathcal{O}_X)$ might fail to be Noetherian although X is Noetherian.) The aim of this section is to prove [Theorems \(D\)](#) and [\(E\)](#). Recall from [Section 2.4](#) that a finite étale morphism $\pi: Y \rightarrow X$ satisfies $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$. First, in order to apply our lifting [Lemma 5.3](#) to finite torsors over splinters, we have:

Lemma 7.1. *Let $\pi: Y \rightarrow X$ be a morphism of Noetherian schemes over a ring R such that $\text{Pic}(\text{Spec } R) = 0$; e.g., R is a local ring or a UFD. If π is a finite torsor over R , then $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$.*

Proof. As discussed in [Section 2.4](#), in order to show that $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$, it is equivalent to produce an \mathcal{O}_X -linear map $\text{Tr}_{Y/X}: \pi_* \mathcal{O}_Y \rightarrow \mathcal{O}_X$ such that the symmetric bilinear form $\text{Tr}_{Y/X}(\alpha \cdot \beta)$ on the locally free sheaf $\pi_* \mathcal{O}_Y$ with values in \mathcal{O}_X is nonsingular. In case π is étale, this is achieved by considering the classical trace. In general, the classical trace need not be nonsingular, but the existence of a nonsingular trace is claimed in [\[BM76, p. 222\]](#) in the special case where R is a field and a proof is provided in [\[CR22, Thm. 3.9\]](#). (Note from e.g. [\[Bri17, Prop. 2.6.4 & 2.6.5\(i\)\]](#) that any finite G -torsor is a finite G -quotient in the sense of [\[CR22, Rmk. 2.3\]](#).) We now consider the more general case where G is a finite locally free group scheme over a ring R . In that case,

$H := H^0(G, \mathcal{O}_G)$ is a finitely generated projective Hopf algebra with antipode. The dual Hopf algebra $H^\vee = \text{Hom}_R(H, R)$ is also a finitely generated projective Hopf algebra with antipode. If $\text{Pic}(\text{Spec } R) = 0$, then H^\vee admits the additional structure of a Frobenius algebra [Par71, Thm. 1]. By [Par71, Thm. 3 & Discussion on p. 596] the R -submodule $\int_{H^\vee}^l \subseteq H^\vee$ of left integrals is freely generated by a nonsingular left integral $\text{Tr}_G \in H^\vee$. If $\pi: Y \rightarrow X$ is a G -torsor over R , by pulling back Tr_G along $Y \rightarrow \text{Spec } R$ and composing with $\pi_* \mathcal{O}_Y \rightarrow \pi_* a_* \mathcal{O}_{Y \times_R G}$, where $a: Y \times_R G \rightarrow Y$ is the action of G on Y , one obtains as in [CR22, §3.1, p. 12] an \mathcal{O}_X -linear map $\pi_* \mathcal{O}_Y \rightarrow \pi_* \mathcal{O}_Y$ that factors through $\mathcal{O}_X = (\pi_* \mathcal{O}_Y)^G$ to an \mathcal{O}_X -linear map $\text{Tr}_{Y/X}: \pi_* \mathcal{O}_Y \rightarrow \mathcal{O}_X$. Since Tr_G is nonsingular, arguing as in [CR22, Thm. 3.9] shows that the bilinear form $(\alpha, \beta) \mapsto \text{Tr}_{Y/X}(\alpha \cdot \beta)$ is nonsingular. \square

Remark 7.2. Let $\pi: Y \rightarrow X$ be a finite torsor over a ring R . Assume that $\text{Pic}(\text{Spec } R) = 0$ and that X is Noetherian and admits a dualizing complex. As explained in Remark 2.3, Lemma 7.1 implies that $\pi^* \omega_X^\bullet \cong \omega_Y^\bullet$. In particular, if X is Gorenstein, then Y is Gorenstein. Likewise, if X is Cohen–Macaulay, then Y is Cohen–Macaulay.

We have the following lemma from [Bha+23]:

Lemma 7.3 ([Bha+23, Lem. 6.6]). *Let $X \rightarrow \text{Spec } R$ be a Noetherian scheme over a ring R . Then the following are equivalent:*

- (i) *The scheme X is a splinter.*
- (ii) *For each closed point $z \in \text{Spec } R$ the base change to the localization X_{R_z} is a splinter.*

Further, assume X is in addition normal and excellent and let Δ be an effective \mathbb{Q} -Weil divisor on X . Then the following are equivalent:

- (i) *The pair (X, Δ) is globally $+$ -regular.*
- (ii) *For each closed point $z \in \text{Spec } R$ the base change to the localization (X_{R_z}, Δ_{R_z}) is globally $+$ -regular.*

Proof. Our assumptions are less restrictive than the setup of [Bha+23, §6], but the proof of [Bha+23, Lem. 6.6] works as we outline below. Note that in any case X is normal. Working on each connected component of X separately, we can and do assume that X is integral. Let $f: Y \rightarrow X$ be a finite cover (with Y normal if we assume (X, Δ) globally $+$ -regular). By flat base change, and as argued in the proof of [Bha+23, Lem. 6.6], the evaluation at 1 map

$$\text{Hom}_{\mathcal{O}_X}(f_* \mathcal{O}_Y(\lfloor f^* \Delta \rfloor), \mathcal{O}_X) \rightarrow H^0(X, \mathcal{O}_X)$$

is surjective if and only if the evaluation-at-1 map

$$\text{Hom}_{\mathcal{O}_{X_{R_z}}}(f_* \mathcal{O}_{R_z}(\lfloor f^* \Delta|_{X_{R_z}} \rfloor), \mathcal{O}_{X_{R_z}}) \rightarrow H^0(X_{R_z}, \mathcal{O}_{X_{R_z}})$$

is surjective for every closed point $z \in \text{Spec } R$. This shows the implications (ii) \Rightarrow (i). For the converse, it is enough to observe that any finite surjective morphism $Y' \rightarrow X_{R_z}$ is the localization of a finite surjective morphism $Y \rightarrow X$, and in case X is excellent that any finite surjective morphism $Y' \rightarrow X_{R_z}$ with Y' normal and integral is the localization of a finite surjective morphism $Y \rightarrow X$ with Y normal. To see this, use that $Y' \rightarrow X_{R_z}$ is of finite presentation and hence spreads to a morphism of finite presentation $Y'' \rightarrow X|_V$ for some dense affine open $V \subseteq \text{Spec } R$ [Gro66, Thm. 8.8.2]. The latter restricts on a dense affine open $U \subseteq V$ to a finite surjective morphism $Y''|_U \rightarrow X|_U$ [Gro66, Thm. 8.10.5], which then by Zariski’s Main Theorem [Sta24, Tag 05K0] extends to a finite surjective morphism $Y \rightarrow X$. In case X is excellent, X is in particular Nagata, so is X_{R_z} [Sta24, Tag 032U]. Therefore, a finite surjective morphism $Y \rightarrow X$ with Y normal is provided by taking the normalization of \mathcal{O}_X in the fraction field $K(Y')$, see [Sta24, Tag 0AVK]. \square

The following proposition extends Lemma 5.3(a) from lifting the splinter property along finite étale covers to finite torsors.

Proposition 7.4 (Lifting the splinter property along finite torsors). *Let $\pi: Y \rightarrow X$ be a morphism of Noetherian schemes. Assume that π is a finite torsor over a ring R , and that it satisfies either of the following conditions:*

- (a) *$H^0(Y, \mathcal{O}_Y)$ is a field.*

- (b) $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$.
- (c) X is Nagata, Y is normal, and $\pi^\#: H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ has reduced closed fibers.

If X is a splinter, then Y is a splinter.

Proof. Let $z \in \operatorname{Spec} R$ be a closed point. Note that all three conditions (a) to (c) are stable under base change along $R \rightarrow R_z$, so that by Lemma 7.3 we can reduce to the case where R is local. From Lemma 7.1 we know that $\pi^! \mathcal{O}_X \cong \mathcal{O}_Y$ for any finite torsor morphism $\pi: Y \rightarrow X$ over a local ring. Lemma 5.3 shows that Y is a splinter. \square

Remark 7.5. Proposition 7.4 does not hold without any assumptions on global sections. Consider indeed the trivial torsor given by a finite non-reduced group scheme G over a field k : $\operatorname{Spec} k$ is a splinter but G fails to be a splinter as it is non-reduced.

Proposition 7.6 (Lifting the derived splinter property along finite torsors). *Let $\pi: Y \rightarrow X$ be a morphism of Noetherian schemes. Assume that π is a finite torsor over a ring R , and that either $H^0(Y, \mathcal{O}_Y)$ is a field or $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$. If X is a derived splinter, then Y is a derived splinter.*

Proof. One argues as in the proof of Proposition 7.4 by reducing to R local (using Nagata compactification [Sta24, Tag 0F41] instead of Zariski's Main Theorem in the proof of Lemma 7.3) and by replacing the use of Lemma 5.3 with Lemma 5.10. \square

We say that a morphism of schemes $\pi: Y \rightarrow X$ is a *quasi-torsor* over a ring R if it is a morphism over R and there exists $U \subseteq X$ open with $\operatorname{codim}_X(X \setminus U) \geq 2$ such that $f|_{f^{-1}(U)}$ is a torsor under a finite locally free group scheme G over R . The following theorem, which generalizes [Bha+23, Prop. 6.20], establishes Theorems (D) and (E).

Theorem 7.7 (Theorems (D) and (E)). *Let $\pi: Y \rightarrow X$ be a finite surjective morphism of normal Nagata Noetherian schemes. Assume that π satisfies either of the following conditions:*

- (a) π is quasi-étale.
- (b) π is a quasi-torsor over a ring R , and $\pi^\#: H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ has reduced closed fibers.

If X is a splinter, then Y is a splinter.

Assume in addition that X is excellent, and let Δ be an effective \mathbb{Q} -Weil divisor on X . If (X, Δ) is globally $+$ -regular, then $(Y, \pi^* \Delta)$ is globally $+$ -regular.

Proof. The theorem in case (a) is Lemma 5.3(a) and Lemma 5.13(a). We now prove case (b). Using Lemma 7.3 we can and do reduce to the case where R is local so that $\operatorname{Pic}(\operatorname{Spec} R) = 0$. By assumption, there exists an open subset $U \subseteq X$ such that $\operatorname{codim}_X(X \setminus U) \geq 2$ and such that $\pi|_U: V := \pi^{-1}(U) \rightarrow U$ is a finite torsor over R . By Lemma 7.1, we have $\pi_U^! \mathcal{O}_U \cong \mathcal{O}_V$. Since Y is assumed to be normal, we can work on each connected component of Y separately and assume without loss of generality that Y is connected. Since X is a splinter, it is Cohen–Macaulay and hence universally catenary [Sta24, Tag 02II]. From the fact that π is finite surjective, it follows from [Sta24, Tag 02II] that $Y \setminus V$ has codimension at least 2 in Y . Since X and Y are normal, we have $H^0(U, \mathcal{O}_U) = H^0(X, \mathcal{O}_X)$ and $H^0(V, \mathcal{O}_V) = H^0(Y, \mathcal{O}_Y)$. If X is a splinter, Lemma 5.3(c) shows that V is a splinter and we conclude with Lemma 4.1 that Y is a splinter. On the other hand, if (X, Δ) is globally $+$ -regular, Lemma 5.13 shows that $(V, \pi^* \Delta|_V)$ is globally $+$ -regular and we conclude with Remark 4.2 that $(Y, \pi^* \Delta)$ is globally $+$ -regular. \square

Remark 7.8. By replacing the use of Lemma 5.13 with Proposition 6.4 (or rather Remark 6.6) and the use of Remark 4.2 with Lemma 4.4 in the proof of Theorem 7.7 in the global $+$ -regular case, one obtains the following statement. Let $\pi: Y \rightarrow X$ be a finite morphism of normal varieties over a field of positive characteristic. Assume that π is quasi-étale, or that π is a quasi-torsor over a ring such that $\pi^\#: H^0(X, \mathcal{O}_X) \rightarrow H^0(Y, \mathcal{O}_Y)$ has reduced closed fibers. Let Δ be an effective \mathbb{Q} -Weil divisor on X . If (X, Δ) is globally F -regular, then $(Y, \pi^* \Delta)$ is globally F -regular. This generalizes [PZ20, Lem. 11.1] where the quasi-étale case was treated.

8. FINITE TORSORS OVER PROPER SPLINTERS OVER A FIELD

We now focus on finite torsors over proper splinters over a field and prove Theorem (A). The following lemma can be found in [Mum70, Thm. 2, p. 121] (we thank Michel Brion for bringing this reference to our attention). We provide an alternate proof based on Hirzebruch–Riemann–Roch for (not necessarily smooth) proper schemes over a field.

Lemma 8.1. *Let X be a proper scheme over a field k and let $\pi: Y \rightarrow X$ be a morphism of schemes over k . Assume that π satisfies either of the following conditions:*

- (a) π is finite étale.
- (b) π is a finite torsor.

Then $\chi(\mathcal{O}_Y) = \deg(\pi)\chi(\mathcal{O}_X)$.

Proof. We first establish (b). Recall that there is a Hirzebruch–Riemann–Roch formula

$$\chi(\mathcal{E}) = \int_X \text{ch}(\mathcal{E}) \cap \text{td}(X)$$

for any vector bundle \mathcal{E} on a proper scheme X over a field; see [Ful98, Cor. 18.3.1]. In particular, the Euler characteristic only depends on the class of the Chern character $\text{ch}(\mathcal{E}) \in A^*(X)_{\mathbb{Q}}$, where $A^*(X)_{\mathbb{Q}}$ denotes the Chow cohomology [Sta24, Tag 0FDV] with \mathbb{Q} -coefficients. Assume $\pi: Y \rightarrow X$ is a torsor under a finite group scheme G over k . By definition of a G -torsor, the product $Y \times_X Y \rightarrow Y$ is isomorphic to $G \times_k Y \rightarrow Y$ as schemes over Y . This gives an isomorphism

$$\pi_* \mathcal{O}_Y \otimes \pi_* \mathcal{O}_Y \cong \pi_* \mathcal{O}_Y^{\oplus n},$$

where $n = \deg(\pi)$ is the order of G . Since π is finite flat, $\pi_* \mathcal{O}_Y$ is a vector bundle (of rank n) on X , and hence by [Sta24, Tag 02UM] we have the identity

$$\text{ch}(\pi_* \mathcal{O}_Y) \cdot \text{ch}(\pi_* \mathcal{O}_Y) = n \text{ch}(\pi_* \mathcal{O}_Y) \quad \text{in } A^*(X)_{\mathbb{Q}}.$$

Since $\text{ch}(\pi_* \mathcal{O}_Y)$ is a unit in $A^*(X)_{\mathbb{Q}}$, we obtain $\text{ch}(\pi_* \mathcal{O}_Y) = n \text{ch}(\mathcal{O}_X)$. By Hirzebruch–Riemann–Roch, the equality $\chi(\pi_* \mathcal{O}_Y) = \chi(\mathcal{O}_Y) = n\chi(\mathcal{O}_X)$ follows. If π is finite étale, one can use Grothendieck–Riemann–Roch for proper schemes over a field [Ful98, Thm. 18.3], while noting that the relative Todd class $\text{td}(T_{\pi})$ is equal to 1. Alternatively one can reduce to case (b) by using the fact [Sta24, Tag 03SF] that there exists a Galois cover $Y' \rightarrow X$ dominating π . \square

Theorem 8.2. *Let X be a proper scheme over an integral Noetherian scheme S of positive characteristic and let $\pi: Y \rightarrow X$ be a morphism of schemes over S . Assume that $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$. In addition, assume either of the following conditions:*

- (a) π is finite étale.
- (b) π is a finite torsor.

If X is a splinter, then π is an isomorphism.

Proof. Let η be the generic point of S . It is enough to show that the restriction $\pi_{\eta}: Y_{\eta} \rightarrow X_{\eta}$ of π to η is an isomorphism. By Lemma 4.3, if X is a splinter, then X_{η} is a splinter. Therefore, we may and do assume that S is the spectrum of a field. Moreover, since a splinter is normal, we may and do assume that X is connected, in which case $H^0(X, \mathcal{O}_X)$ is a field. On the other hand, by Lemma 5.3(a), or more simply Lemma 5.3(b), in the étale case and by Proposition 7.4 in the torsor case, Y is a splinter. Since the structure sheaf of a proper splinter in positive characteristic has trivial positive cohomology by Proposition 3.6, we have $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_Y) = 1$, where the dimension is taken with respect to the field $H^0(X, \mathcal{O}_X)$ and we conclude with Lemma 8.1 that π is an isomorphism. \square

Theorem 8.3. *Let X be a connected proper scheme over a field k of positive characteristic with a k -rational point $x \in X(k)$. Assume that X is a splinter.*

- (i) *If k is separably closed, then the étale fundamental group $\pi_1^{\text{ét}}(X, x)$ of X is trivial.*
- (ii) *(Theorem (A)) The Nori fundamental group scheme $\pi_1^N(X, x)$ of X is trivial.*

Proof. Statement (i) follows from [Theorem 8.2](#) since for k separably closed any connected finite étale cover $\pi: Y \rightarrow X$ satisfies $H^0(Y, \mathcal{O}_Y) = H^0(X, \mathcal{O}_X) = k$. For statement (ii), first note that the Nori fundamental group scheme $\pi_1^N(X, x)$ is well-defined as a splinter is reduced. Assume for contradiction that $\pi_1^N(X, x)$ is nontrivial. Since $\pi_1^N(X, x)$ is pro-finite, there is a surjective group scheme homomorphism $\pi_1^N(X, x) \twoheadrightarrow G$ to a nontrivial finite group scheme G . By [\[Nor82, Prop. 3, p. 87\]](#), there exists a G -torsor $Y \rightarrow X$ with $H^0(Y, \mathcal{O}_Y) = H^0(X, \mathcal{O}_X)$. This contradicts [Theorem 8.2](#). \square

Remark 8.4 (On the triviality of the Nori fundamental group scheme). As mentioned in [Corollary 5.16](#), a connected proper splinter X is geometrically normal, hence geometrically reduced, over the field $K := H^0(X, \mathcal{O}_X)$; in particular it acquires a rational point after some finite separable field extension of K . Moreover, recall the general facts that the Nori fundamental group scheme is invariant under separable base change, and that the triviality of the Nori fundamental group scheme is independent of the choice of base point.

We say that a finite étale cover $\pi: Y \rightarrow X$ is trivial if it is isomorphic over X to a disjoint union of copies of X . We say that a finite torsor $\pi: Y \rightarrow X$ under a finite group scheme G over k is trivial if it is isomorphic to $X \times_k G$ over X . An immediate consequence of [Theorem 8.3](#) is the following:

Corollary 8.5. *Let X be a connected proper scheme over a field k of positive characteristic. Assume that X is a splinter.*

- (i) *If k is separably closed, then any finite étale cover of X is trivial.*
- (ii) *If k is algebraically closed, then any finite torsor over X is trivial.*

Proof. Statement (i) is clear from (the proof of) [Theorem 8.3](#), while statement (ii) follows from the fact [\[Nor82\]](#) that for a k -point of $x \in X(k)$ there is an equivalence of categories between the category of finite torsors $Y \rightarrow X$ equipped with a k -point $y \in Y(k)$ mapping to x and the category of finite group schemes G over k equipped with a k -group scheme homomorphism $\pi_1^N(X, x) \rightarrow G$. \square

9. PROPER SPLINTERS HAVE NEGATIVE KODAIRA DIMENSION

Let X be a Gorenstein projective scheme over a field k of positive characteristic. Since $-K_X$ big implies that X has negative Kodaira dimension, it is expected in view of [Conjecture 3.12](#) that, if X is a splinter, then its Kodaira dimension is negative. In this section, we confirm this expectation (without assuming X to be Gorenstein) and prove [Theorem \(B\)](#).

Let X be a normal proper variety over a field k and let D be a Weil divisor on X . We define the *Iitaka dimension* of D to be

$$\kappa(X, D) := \min \{ r \in \mathbb{Z}_{\geq 0} \mid (h^0(X, \mathcal{O}_X(dD)) / d^r)_{d \geq 0} \text{ is bounded} \}.$$

By convention, if $h^0(X, \mathcal{O}_X(dD)) := \dim_k H^0(X, \mathcal{O}_X(dD)) = 0$ for all $d > 0$, we set $\kappa(X, D) = -\infty$. Beware that we deviate from usual conventions as the Iitaka dimension is usually defined for invertible sheaves on projective varieties. If X is a smooth projective variety over k , $\kappa(X, K_X)$ agrees with the Kodaira dimension of X . The following proposition refines the observation from [Lemma 3.7](#) showing that if X is a proper splinter in positive characteristic, then K_X is not effective.

Theorem 9.1 ([Theorem \(B\)](#)). *Let X be a positive-dimensional connected proper scheme over a field of positive characteristic. If X is a splinter, then $\kappa(X, K_X) = -\infty$.*

First we have the following variant of a well-known lemma; see, e.g., [\[PST17, Ex. 2.12\]](#).

Lemma 9.2. *Let X be a proper scheme over a field k of positive characteristic $p > 0$. Assume either of the following conditions:*

- (a) *X is a splinter.*
- (b) *X is normal and F -split.*

Then the Weil divisor $(1 - p)K_X$ is effective. In particular, either $\kappa(X, K_X) = -\infty$, or K_X is torsion (in which case $\kappa(X, K_X) = 0$).

Proof. First assume that X is a splinter. Since X is normal, we can and do assume that X is connected. After replacing the base field k by $H^0(X, \mathcal{O}_X)$, we consider the base change $\pi: X_{\bar{k}} \rightarrow X$ along an algebraic closure $k \rightarrow \bar{k}$. By [Proposition 5.14](#), $X_{\bar{k}}$ is a splinter. The base change formula for the exceptional inverse image [\[Sta24, Tag 0E9U\]](#) shows that $\pi^*\omega_X = \omega_{X_{\bar{k}}}$. Thus, it is enough to show the statement for $X_{\bar{k}}$, since π flat implies $H^0(X_{\bar{k}}, \pi^*\mathcal{O}_X((1-p)K_X)) = H^0(X, \mathcal{O}_X((1-p)K_X)) \otimes_k \bar{k}$. Since \bar{k} is F -finite, $X_{\bar{k}}$ is in particular F -split. Thus, it is enough to show the statement under the assumptions (b).

Assume that X is normal and F -split. Parts of the arguments below can for example be found in [\[SS10, §4.2\]](#). We provide nonetheless a proof for the sake of completeness. Choosing a (non-canonical) isomorphism $F^!k = \text{Hom}_k(F_*k, k) \cong k$ as k -vector spaces, we obtain an isomorphism $F^!\omega_X^\bullet \cong \omega_X^\bullet$ in $\text{D}_{\text{Coh}}(\mathcal{O}_X)$ and, by restricting to X_{reg} and taking cohomology sheaves, an isomorphism $F^!\omega_{X_{\text{reg}}} \cong \omega_{X_{\text{reg}}}$ in $\text{Coh}(X_{\text{reg}})$. Since the involved sheaves are reflexive, $F^!\omega_X \cong \omega_X$ holds in $\text{Coh}(X)$ and we obtain $\mathcal{H}om_{\mathcal{O}_X}(F_*\mathcal{O}_X, \omega_X) \cong F_*\omega_X$.

By assumption, there exists a map s such that the composition

$$\mathcal{O}_X \rightarrow F_*\mathcal{O}_X \xrightarrow{s} \mathcal{O}_X$$

is the identity. We apply $\mathcal{H}om_{\mathcal{O}_X}(-, \omega_X)$ to obtain

$$\omega_X \leftarrow F_*\omega_X \xleftarrow{s^\vee} \omega_X,$$

where the composition is the identity. After restricting to the regular locus, we can twist with $\omega_{X_{\text{reg}}}^{-1}$ and apply the projection formula to obtain a diagram

$$\mathcal{O}_{X_{\text{reg}}} \leftarrow F_*\mathcal{O}_{X_{\text{reg}}}((1-p)K_{X_{\text{reg}}}) \xleftarrow{s^\vee} \mathcal{O}_{X_{\text{reg}}},$$

where the composition is the identity. Using that the involved sheaves are reflexive, we obtain a nonzero global section of $F_*\mathcal{O}_X((1-p)K_X)$. This gives a nonzero element of $H^0(X, \mathcal{O}_X((1-p)K_X))$, hence $(1-p)K_X$ is effective. If no positive multiple of K_X is effective, then $\kappa(X, K_X) = -\infty$. So assume that nK_X is effective for some $n > 0$. Since $n(p-1)K_X$ and $n(1-p)K_X$ are both effective, $n(1-p)K_X$ is trivial by [Lemma 2.1](#). \square

Remark 9.3. Note that F -split varieties may have trivial canonical divisor. For instance, ordinary elliptic curves and ordinary $K3$ or abelian surfaces are F -split; see [\[BK05, Rmk. 7.5.3\(i\)\]](#).

In order to prove [Theorem 9.1](#), it remains to show that the canonical divisor of a proper splinter is not torsion. First we note that the Picard group of a proper splinter is torsion-free; this is a small generalization of a result of Carvajal-Rojas [\[CR22, Cor. 5.4\]](#) who showed that a globally F -regular projective variety has torsion-free Picard group. In particular, this provides a proof of [Theorem 9.1](#) if K_X is Cartier, e.g., if X is in addition assumed to be Gorenstein.

Proposition 9.4. *Let X be a proper scheme over a field of positive characteristic. If X is a splinter, then $\text{Pic}(X)$ is torsion-free.*

Proof. We argue as in [\[CR22, Rmk. 5.6\]](#) which is concerned with the globally F -regular case. If \mathcal{L} is a torsion invertible sheaf, then it is in particular semiample and therefore if X is a splinter, then we have $\chi(X, \mathcal{L}) = h^0(X, \mathcal{L})$ by [\[Bha12, Prop. 7.2\]](#) (recalled in [Proposition 3.6](#)). But then $0 = \chi(X, \mathcal{L}) = \chi(X, \mathcal{O}_X) = 1$ if \mathcal{L} is nontrivial. This is impossible.

Alternately, one can argue using [Theorem 8.2](#) as follows. An n -torsion invertible sheaf \mathcal{L} on X gives rise to a nontrivial μ_n -torsor $\pi: Y \rightarrow X$, where Y is defined to be the relative spectrum of the finite \mathcal{O}_X -algebra $\mathcal{O}_X \oplus \mathcal{L} \oplus \cdots \oplus \mathcal{L}^{n-1}$. If $n > 1$, we must have $H^0(X, \mathcal{L}) = 0$, since any nontrivial section $s: \mathcal{O}_X \rightarrow \mathcal{L}$ would also give a nontrivial section s^n of \mathcal{O}_X , so s would be nowhere vanishing and therefore \mathcal{L} would be trivial. This yields the equality

$$H^0(Y, \mathcal{O}_Y) = H^0(X, \pi_*\mathcal{O}_Y) = H^0(X, \mathcal{O}_X \oplus \mathcal{L} \oplus \cdots \oplus \mathcal{L}^{n-1}) = H^0(X, \mathcal{O}_X).$$

We conclude from [Theorem 8.2](#) that if X is a splinter, then $\deg(\pi) = 1$, i.e., $n = 1$ and \mathcal{L} is trivial. \square

To deal with the non-Gorenstein case, we have:

Proposition 9.5. *Let X be a connected positive-dimensional proper scheme over a field of positive characteristic. If X is a splinter, then the canonical divisor K_X is not torsion.*

Proof. Assume for contradiction that K_X is torsion of order r , i.e., that $\omega_X|_{X_{\text{reg}}}$ is a torsion invertible sheaf of order r . By considering the relative spectrum, we obtain a μ_r -quasi-torsor

$$\pi: Y = \text{Spec}_X \left(\bigoplus_{i=0}^{r-1} \mathcal{O}_X(iK_X) \right) \rightarrow X,$$

which, over X_{reg} , restricts to a μ_r -torsor $\pi|_U: V := \pi^{-1}(X_{\text{reg}}) \rightarrow X_{\text{reg}} =: U$.

In addition, $\pi_* \mathcal{O}_Y = \bigoplus_{i=0}^{r-1} \mathcal{O}_X(iK_X)$, and by Lemma 2.1 the sheaves $\mathcal{O}_X(iK_X)$ have no nonzero global sections for $1 \leq i \leq r-1$. Thus $H^0(V, \mathcal{O}_V) = H^0(X_{\text{reg}}, \mathcal{O}_{X_{\text{reg}}}) = H^0(X, \mathcal{O}_X)$, where the second equality holds by normality of X , is a field and we conclude by Proposition 7.4(a) or (b) that V is a splinter. In particular, V is normal and therefore, by, e.g., [Sta24, Tag 035E], the normalization $Y^\nu \rightarrow Y$ is an isomorphism over V . Since normalization is finite, $Y^\nu \setminus V$ has codimension ≥ 2 and thus Y^ν is a splinter by Lemma 4.1.

Now $\pi_U^! \mathcal{O}_{X_{\text{reg}}} = \mathcal{O}_V$ holds by Lemma 7.1 and this implies $\pi_U^* \omega_{X_{\text{reg}}} = \omega_V$. On the other hand, we have isomorphisms

$$(\pi_U)_* \pi_U^* \omega_{X_{\text{reg}}} \cong \omega_{X_{\text{reg}}} \otimes_{\mathcal{O}_{X_{\text{reg}}}} \bigoplus_{i=0}^{r-1} \omega_{X_{\text{reg}}}^i = \bigoplus_{i=1}^r \omega_{X_{\text{reg}}}^i \cong \bigoplus_{i=0}^{r-1} \omega_{X_{\text{reg}}}^i \cong (\pi_U)_* \mathcal{O}_V$$

as $(\pi_U)_* \mathcal{O}_V$ -modules. Hence $V = \text{Spec}_{X_{\text{reg}}} \bigoplus_{i=0}^{r-1} \omega_{X_{\text{reg}}}^i$ has trivial dualizing sheaf. Consequently, since ω_{Y^ν} is a reflexive sheaf, Y^ν has trivial dualizing sheaf. But, by Lemma 3.7, a proper splinter cannot have trivial canonical sheaf. \square

Proof of Theorem 9.1. The canonical divisor K_X is not torsion by Proposition 9.5 (or more simply by Proposition 9.4 if K_X is Cartier, e.g. if X is Gorenstein), and it follows from Lemma 9.2 that $\kappa(X, K_X) = -\infty$. \square

Remark 9.6 (Torsion Weil divisors on splinters). Proper splinters may have nontrivial torsion Weil divisor classes. Indeed, projective toric varieties are globally F -regular and in particular splinters, and Carvajal-Rojas [CR22, Ex. 5.7] gives an example of a projective toric surface that admits a nontrivial 2-torsion Weil divisor class.

10. VANISHING OF GLOBAL DIFFERENTIAL FORMS

Fix a perfect field k of positive characteristic p and let X be a smooth proper variety over k . Let $\Omega_{X/k}^\bullet$ be the de Rham complex and recall, e.g., from [Kat70, Thm. 7.2], that there exists an isomorphism of graded \mathcal{O}_X -modules

$$C_X^{-1}: \bigoplus_{j \geq 0} \Omega_X^j \rightarrow \bigoplus_{j \geq 0} \mathcal{H}^j(F_* \Omega_X^\bullet),$$

whose inverse C_X is the so-called *Cartier operator*. It gives rise for all $j \geq 0$ to short exact sequences of \mathcal{O}_X -modules

$$(10.1) \quad 0 \rightarrow B_X^j \rightarrow Z_X^j \xrightarrow{C_X^j} \Omega_X^j \rightarrow 0,$$

where B_X^j denotes the j -th coboundaries and Z_X^j the j -th cocycles of $F_* \Omega_X^\bullet$. Note that these coincide with the image under F_* of the coboundaries and cocycles of Ω_X^\bullet . Moreover, there is a short exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow F_* \mathcal{O}_X \rightarrow B_X^1 \rightarrow 0.$$

The proof of the following theorem is inspired by the proof of [AWZ21, Lem. 6.3.1].

Theorem 10.2 (Theorem (C)). *Let X be a smooth proper variety over a field k of positive characteristic p . If X is a splinter, then $H^0(X, \Omega_X^1) = 0$.*

Proof. Clearly, we may and do assume that X is connected. Let \bar{k} be an algebraic closure of k . It is enough to show that $H^0(X_{\bar{k}}, \Omega_{X_{\bar{k}}}^1) = 0$. Since X is in particular geometrically reduced over k , $H^0(X, \mathcal{O}_X)$ is a finite separable extension of k ; see, e.g., [Sta24, Tag 0BUG]. It follows that $X_{\bar{k}}$ is the disjoint union of $\dim_k H^0(X, \mathcal{O}_X)$ copies of $X \times_{H^0(X, \mathcal{O}_X)} \bar{k}$. From Corollary 5.15, we find that $X_{\bar{k}}$ is a splinter. Therefore it is enough to establish the theorem in case k is algebraically closed. So assume k is algebraically closed, in which case the p -th power map $k = H^0(X, \mathcal{O}_X) \rightarrow H^0(X, F_* \mathcal{O}_X) = k$ is an isomorphism. Since by Proposition 3.6, for a splinter X , \mathcal{O}_X has zero cohomology in positive degrees, the long exact sequence associated to

$$0 \rightarrow \mathcal{O}_X \rightarrow F_* \mathcal{O}_X \rightarrow B_X^1 \rightarrow 0$$

gives that $H^j(X, B_X^1) = 0$ for all $j \geq 0$. From the long exact sequence associated to (10.1), we find that the Cartier operator induces an isomorphism $H^0(X, Z_X^1) \rightarrow H^0(X, \Omega_X^1)$. In particular, $\dim H^0(X, Z_X^1) = \dim H^0(X, \Omega_X^1)$. The inclusion of closed 1-forms $\ker(d: \Omega_X^1 \rightarrow \Omega_X^2) \subseteq \Omega_X^1$ yields an injection $H^0(X, \ker(d: \Omega_X^1 \rightarrow \Omega_X^2)) \subseteq H^0(X, \Omega_X^1)$. Since $H^0(X, \ker(d: \Omega_X^1 \rightarrow \Omega_X^2)) = H^0(X, Z_X^1)$, the above inclusion is in fact an equality. In other words, any global 1-form on X is closed. By [GK03, Prop. 4.3], there exists an isomorphism

$$H^0(X, \Omega_X^1) \cong \text{Pic}(X)[p] \otimes_{\mathbb{Z}} k,$$

where $\text{Pic}(X)[p]$ denotes the p -torsion invertible sheaves on X . By Proposition 9.4, $\text{Pic}(X)$ is torsion-free. \square

11. ON THE SPLINTER PROPERTY FOR PROPER SURFACES

A proper curve over an algebraically closed field of positive characteristic is a splinter if and only if it is isomorphic to the projective line. In this section, we investigate which proper surfaces over an algebraically closed field of positive characteristic are splinters. First we show in Proposition 11.1 that proper surface splinters are rational. We then show in Proposition 11.4 that the blow-up of the projective plane in any number of closed points lying on a given conic is a splinter. On the other hand, we give examples of rational surfaces that are not splinters in Section 11.3.

11.1. Proper splinter surfaces are rational. The fact proved in Theorem 9.1 that proper splinters in positive characteristic have negative Kodaira dimension can be used to show that proper surface splinters over an algebraically closed field of positive characteristic are rational:

Proposition 11.1. *Let X be an irreducible proper surface over an algebraically closed field of positive characteristic. If X is a splinter, then X is rational.*

Proof. If X is not smooth, choose a resolution of singularities $\pi: \tilde{X} \rightarrow X$ such that π is an isomorphism over the regular locus X_{reg} , which exists by [Lip69, §2]. It suffices to show that \tilde{X} is rational. Note that Grauert–Riemenschneider vanishing holds for surfaces, see, e.g., [Sta24, Tag 0AXD]. Thus, the proof of [Bha12, Thm. 2.12] shows that the above resolution of singularities satisfies $R\pi_* \mathcal{O}_{\tilde{X}} = \mathcal{O}_X$, that is, that X has rational singularities. Therefore

$$\chi(\mathcal{O}_{\tilde{X}}) = \chi(R\pi_* \mathcal{O}_{\tilde{X}}) = \chi(\mathcal{O}_X) = 1.$$

By Castelnuovo’s rationality criterion it remains to show that $\omega_{\tilde{X}}^2$ has no nonzero global sections. So assume that there is a nonzero section $s \in H^0(\tilde{X}, \omega_{\tilde{X}}^2)$. Then s is nonzero after restriction to the open subset $\pi^{-1}(X_{\text{reg}})$. Since π is an isomorphism over X_{reg} , the section s would provide a nonzero section of $\mathcal{O}_X(2K_X)$, contradicting Theorem 9.1. \square

11.2. Examples of projective rational surfaces that are splinters. We give examples of projective rational surfaces that are splinters; in all cases, this is achieved by showing that they are globally F -regular. We start with already known examples.

Example 11.2 (Del Pezzo surfaces, [Har98, Ex. 5.5]). Let X be a smooth projective del Pezzo surface over an algebraically closed field of characteristic $p > 0$. Then X is globally F -regular if one of the following conditions holds:

- (i) $K_X^2 > 3$.
- (ii) $K_X^2 = 3$ and $p > 2$.

- (iii) $K_X^2 = 2$ and $p > 3$.
- (iv) $K_X^2 = 1$ and $p > 5$.

Moreover, if none of the above conditions are satisfied, there are globally F -regular and non globally F -regular cases; for instance, the Fermat cubic surface in characteristic 2 is not globally F -regular.

Example 11.3 (Hirzebruch surfaces, [GT16, Prop. 3.1]). If X is a Hirzebruch surface $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-n))$ over a perfect field of positive characteristic, then X is globally F -regular.

To the above list of examples, we can add blow-ups of \mathbb{P}^2 in any number of points lying on a conic:

Proposition 11.4. *Let k be an algebraically closed field of characteristic $p > 0$ and let p_1, \dots, p_n be distinct closed points in \mathbb{P}_k^2 . Assume either of the following conditions:*

- (a) *The points p_1, \dots, p_n lie on a line $L \subseteq \mathbb{P}_k^2$.*
- (b) *The points p_1, \dots, p_n lie on a (possibly singular) conic $C \subseteq \mathbb{P}_k^2$.*

then the blow-up $X := \text{Bl}_{\{p_1, \dots, p_n\}} \mathbb{P}_k^2$ is globally F -regular. In particular, the blow-up of \mathbb{P}_k^2 in at most 5 closed points is globally F -regular.

In some sense Proposition 11.4 is optimal since by Example 11.2 the Fermat cubic surface, which is the blow-up of \mathbb{P}_k^2 in 6 points, is not globally F -regular if $\text{char } k = 2$. The proof adapts a strategy which is similar to the arguments of [BK05, §§1.3-1.4]. First we determine, under the isomorphism $\text{Hom}_X(F_*\mathcal{O}_X, \mathcal{O}_X) \cong H^0(X, \mathcal{O}_X((1-p)K_X))$, which global sections correspond to splittings of $\mathcal{O}_X \rightarrow F_*\mathcal{O}_X$ in the case where $X = \mathbb{P}_k^n$. For the sake of completeness, we state and prove the following lemma, which appears as an exercise in [BK05].

Lemma 11.5 (cf. [BK05, Ex. 1.3.E(1)]). *Let $X = \mathbb{P}_k^n = \text{Proj } k[X_0, \dots, X_n]$ for a field k of characteristic $p > 0$. Let $F^{e\sharp}: \mathcal{O}_X \rightarrow F_*^e\mathcal{O}_X$ be the canonical map and consider the following chain of isomorphisms*

$$\Phi: \text{Hom}_X(F_*^e\mathcal{O}_X, \mathcal{O}_X) \xrightarrow{\cong} \text{Ext}_X^n(\mathcal{O}_X, (F_*^e\mathcal{O}_X) \otimes \omega_X)^\vee \xrightarrow{\cong} H^n(X, \omega_X^{p^e})^\vee \xrightarrow{\cong} H^0(X, \omega_X^{1-p^e})$$

where the first and last isomorphism are given by Serre duality and the second isomorphism follows from the projection formula. Then the following diagram commutes

$$\begin{array}{ccc} \text{Hom}_X(F_*^e\mathcal{O}_X, \mathcal{O}_X) & \xrightarrow{-\circ F^{e\sharp}} & \text{Hom}_X(\mathcal{O}_X, \mathcal{O}_X) \\ \downarrow \Phi & & \downarrow \text{ev}_1 \\ H^0(X, \mathcal{O}_X(\omega_X^{1-p^e})) & \xrightarrow{\tau} & k, \end{array}$$

where ev_1 is the evaluation at the constant global section 1 and τ is the map sending a homogeneous polynomial $P \in H^0(X, \mathcal{O}_X((n+1)(p^e-1)))$ of degree $(n+1)(p^e-1)$ to the coefficient of the monomial $(X_0 \cdots X_n)^{(p^e-1)}$ in P . In particular, $\Phi^{-1}(P)$ provides a splitting of $F^{e\sharp}$ if and only if $\tau(P) = 1$.

Proof. Recall, e.g., from [Har77, Ch. III, Thm. 5.1 & Thm. 7.1], that Serre duality for invertible sheaves on \mathbb{P}^n is given by the bilinear form

$$\begin{aligned} H^0(X, \mathcal{O}_X(a)) \otimes H^n(X, \mathcal{O}_X(-(n+1)-a)) &\rightarrow H^n(X, \mathcal{O}_X(-(n+1))) = k \frac{1}{X_0 \cdots X_n} \\ (P, Q) &\mapsto \text{coefficient of } (X_0 \cdots X_n)^{-1} \text{ in } PQ, \end{aligned}$$

where, for $b < 0$, we identify $H^n(X, \mathcal{O}_X(b))$ with the degree b part of the negatively graded k -algebra $(X_0 \cdots X_n)^{-1}k[X_0^{-1}, \dots, X_n^{-1}]$. Consider the following commutative diagram of isomorphisms

$$\begin{array}{ccccc} \text{Hom}_X(F_*^e\mathcal{O}_X, \mathcal{O}_X) & \xrightarrow{-\circ F^{e\sharp}} & \text{Hom}_X(\mathcal{O}_X, \mathcal{O}_X) & \xrightarrow{\text{ev}_1} & k \\ \downarrow \text{SD} & & \downarrow \text{SD} & & \\ H^0(X, \mathcal{O}_X((1-p^e)K_X)) & \xrightarrow{\text{SD}} & H^n(X, \mathcal{O}_X(p^e K_X))^\vee & \xrightarrow{(F^{e*})^\vee} & H^n(X, \mathcal{O}_X(K_X))^\vee, \end{array}$$

where SD stands for Serre duality. Since $F^{e*}: H^n(X, \mathcal{O}_X(-(n+1))) \rightarrow H^n(X, \mathcal{O}_X(-p^e(n+1)))$ raises a polynomial to its p^e -th power, we find that a monomial in $H^0(X, \mathcal{O}_X((1-p^e)K_X))$ is sent to 1 in k along the above diagram if it is $(X_0 \cdots X_n)^{p^e-1}$ and to zero otherwise. \square

For X the blow-up of \mathbb{P}_k^2 in n distinct closed points $p_1, \dots, p_n \in \mathbb{P}_k^2$, we denote by E_i the exceptional curve over the point p_i and we let $H \in \text{Pic}(X)$ be the pullback of the class of a hyperplane in \mathbb{P}_k^2 . Then

$$\text{Pic}(X) = \mathbb{Z}H \oplus \mathbb{Z}E_1 \oplus \cdots \oplus \mathbb{Z}E_n,$$

is an orthogonal decomposition with respect to the intersection pairing, and we have $H^2 = 1$ and $E_i^2 = -1$ for all $1 \leq i \leq n$. The canonical class of X is $K_X = -3H + \sum_i E_i$. If $C \subseteq \mathbb{P}_k^2$ is an irreducible curve, its strict transform $\tilde{C} \subseteq X$ has class

$$\tilde{C} = dH - \sum_{i=1}^n m_i E_i \in \text{Pic}(X),$$

where d is the degree of C and m_i is the multiplicity of C at p_i . Any irreducible curve in X is either one of the exceptional curves E_i , or the strict transform of an irreducible curve in \mathbb{P}_k^2 . Note that for $d > n$, the divisor $dH - \sum_i E_i$ is ample, since it has positive square and since the intersection with any integral curve in X is positive.

Proof of Proposition 11.4. Since X is smooth, it suffices by Theorem 6.1 (see Remark 6.2) to prove that there exists an ample divisor D on X such that $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ splits for some $e > 0$.

Let $d > 0$ be an integer such that the divisor $D := dH - \sum_i E_i$ on X is ample and fix a global section $\sigma: \mathcal{O}_X \rightarrow \mathcal{O}_X(D)$. We can interpret σ as a homogeneous polynomial of degree d vanishing at the points p_1, \dots, p_n . Now, as in Lemma 11.5, we have an isomorphism

$$\Psi: \text{Hom}_X(F_*^e \mathcal{O}_X(D), \mathcal{O}_X) \xrightarrow{\cong} H^0(X, \mathcal{O}_X((1-p^e)K_X - D))$$

and global sections of $\mathcal{O}_X((1-p^e)K_X - D)$ correspond again to polynomials of a certain degree, vanishing to some certain order at the points p_i . The following claim is similar to [BK05, p. 39].

Claim. A section $\varphi \in H^0(X, \mathcal{O}_X((1-p^e)K_X - D))$ defines a section of $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ if and only if $\Psi^{-1}(\varphi)\sigma \in H^0(X, \mathcal{O}_X((1-p^e)K_X))$ defines a splitting of $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X$, where $\Psi^{-1}(\varphi)\sigma$ is the usual product of polynomials.

Proof of the claim. A map $\varphi \in \text{Hom}_X(F_*^e \mathcal{O}_X(D), \mathcal{O}_X)$ is a section of $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(D)$ if and only if $\varphi \circ F_*^e(\sigma)$ is a section of $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X$. Thus, we have to check that the composition $\varphi \circ F_*^e(\sigma)$ corresponds to the product $\Psi(\varphi)\sigma$. This is done by verifying, that the following diagram commutes

$$\begin{array}{ccc} \text{Hom}_X(F_*^e \mathcal{O}_X(D), \mathcal{O}_X) & \xrightarrow{- \circ F_*^e(\sigma)} & \text{Hom}_X(F_*^e \mathcal{O}_X, \mathcal{O}_X) \\ \downarrow \text{SD} & & \downarrow \text{SD} \\ H^n(X, \omega_X \otimes F_*^e \mathcal{O}_X(D))^\vee & \xrightarrow{(\omega_X \otimes F_*^e(\sigma))^\vee} & H^n(X, \omega_X \otimes F_*^e \mathcal{O}_X)^\vee \\ \downarrow \text{projection formula} & & \downarrow \text{projection formula} \\ H^n(X, F_*^e \mathcal{O}_X(p^e K_X + D))^\vee & \xrightarrow{(F_*^e(\mathcal{O}_X(p^e K_X) \otimes \sigma))^\vee} & H^n(X, F_*^e \mathcal{O}_X(p^e K_X))^\vee \\ \downarrow & & \downarrow \\ H^n(X, \mathcal{O}_X(p^e K_X + D))^\vee & \xrightarrow{(\mathcal{O}_X(p^e K_X) \otimes \sigma)^\vee} & H^n(X, \mathcal{O}_X(p^e K_X))^\vee \\ \downarrow \text{SD} & & \downarrow \text{SD} \\ H^0(X, \mathcal{O}_X((1-p^e)K_X - D)) & \xrightarrow{- \cdot \sigma} & H^0(X, \mathcal{O}_X((1-p^e)K_X)). \quad \square \end{array}$$

Denote by $\mu: X \rightarrow \mathbb{P}^2$ the blow-up map. Since $\mu_* \mathcal{O}_X = \mathcal{O}_{\mathbb{P}^2}$, μ_* induces an isomorphism $\text{End}(\mathcal{O}_X) \rightarrow \text{End}(\mathcal{O}_{\mathbb{P}^2})$. Since any morphism of schemes commutes with the Frobenius, a map $\varphi: F_*^e \mathcal{O}_X \rightarrow \mathcal{O}_X$ is a splitting of $\mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X$ if and only if $\mu_*(\varphi)$ is a splitting of $\mathcal{O}_{\mathbb{P}^2} \rightarrow F_*^e \mathcal{O}_{\mathbb{P}^2}$. The proposition will follow if we can find suitable polynomials $\varphi \in H^0(X, \mathcal{O}_X((1-p^e)K_X - D))$ and $\sigma \in H^0(X, \mathcal{O}_X(D))$ such that $\mu_*(\varphi\sigma)$ defines a splitting of $\mathcal{O}_{\mathbb{P}^2} \rightarrow F_*^e \mathcal{O}_{\mathbb{P}^2}$. In terms of Lemma 11.5,

the monomial $(XYZ)^{p^e-1}$ has to occur with coefficient 1 in $\varphi\sigma$ (here we are using coordinates $\mathbb{P}_k^2 = \text{Proj } k[X, Y, Z]$).

We first compute

$$(1-p^e)K_X - D = 3(p^e-1)H - (p^e-1)\sum_i E_i - dH + \sum_i E_i = (3(p^e-1)-d)H - (p^e-2)\sum_i E_i.$$

If all the points p_i lie on a line L , we can assume without loss of generality that $L = V(Z)$. Consider the polynomials $\tilde{\varphi} := X^{(p^e-1)-(d-1)}Y^{p^e-1}$ and $\tilde{\sigma} := X^{d-1}$. Moreover, if we set $\varphi := \tilde{\varphi}Z^{p^e-2}$ and $\sigma := \tilde{\sigma}Z$, then $\varphi \in H^0(X, \mathcal{O}_X((1-p^e)K_X - D))$ and $\sigma \in H^0(X, \mathcal{O}_X(D))$ and the coefficient of $(XYZ)^{p^e-1}$ in $\varphi\sigma$ is 1.

If the points lie on a conic C , we can assume after possible change of coordinates that the conic is given by an equation of the form $XY - Z^2$, XY , or X^2 ; see the elementary [Lemma 11.7](#) below. In the last case, the points lie on the line $X = 0$, thus we may assume that C is given by one of the equations $\tilde{\sigma} = XY - Z^2$ or $\tilde{\sigma} = XY$. Now set $\sigma := Z^{d-2}\tilde{\sigma}$ and $\varphi := Z^{(p^e-1)-(d-2)}\tilde{\varphi}Z^{p^e-2}$ and observe that $(XYZ)^{p^e-1}$ occurs with coefficient 1 in $\varphi\sigma$. \square

Remark 11.6. Recall, e.g. from [[Har77](#), Ch. IV, Prop. 4.21], that an elliptic curve $C = V(P) \subseteq \mathbb{P}_k^2$ is *ordinary* if $(XYZ)^{p-1}$ occurs with nonzero coefficient in P^{p-1} . The above [Lemma 11.5](#) can also be used to show, similarly to [[Har15](#), Rmk. 6.3], that the blow-up of \mathbb{P}_k^2 in any number of points, which lie on an ordinary elliptic curve, is F -split.

Lemma 11.7. *Let k be an algebraically closed field. After suitable coordinate transform a conic in \mathbb{P}_k^2 is given one of the following equations :*

$$XY - Z^2, \text{ } XY, \text{ or } X^2.$$

Proof. First recall that $\text{PGL}_3(k)$ acts 2-transitively on $\mathbb{P}_k^2 = \text{Proj } k[X, Y, Z]$. In particular, $\text{PGL}_3(k)$ acts also 2-transitively on the set of lines in \mathbb{P}_k^2 , which is $\mathbb{P}(H^0(\mathbb{P}_k^2, \mathcal{O}_{\mathbb{P}_k^2}(1))) = \mathbb{P}(k[X, Y, Z]_{\deg 1})$. If a conic $C \subseteq \mathbb{P}_k^2$ is reducible, then it is either a double line or the union of two lines and we can assume C to be defined by the equations $X^2 = 0$ or $XY = 0$, resp. It remains to show that an integral conic C is defined by the equation $XY - Z^2 = 0$ after suitable coordinate transformation. We follow the arguments of [[Kir92](#), Cor. 3.12]. Since C has only finitely many singular points, we can assume without loss of generality that $[0 : 1 : 0]$ is a smooth point of C and the line $Z = 0$ is tangent to C at $[0 : 1 : 0]$. This means that C is given by an equation of the form

$$(11.8) \quad aYZ + bX^2 + cXZ + dZ^2,$$

since the line tangent to C at the point $[0 : 1 : 0]$ is precisely the line given by the linear factors in the dehomogenized equation defining C on the affine open $\{Y \neq 0\} \cong \text{Spec } k[X, Z]$. By assumption (11.8) is an irreducible polynomial. This implies that $b \neq 0$ and $a \neq 0$. We conclude by noting that $XY - Z^2$ is mapped to (11.8) under the coordinate transformation

$$[x, y, z] \mapsto [\sqrt{b}x, ay + cx + dz, -z]. \quad \square$$

11.3. Examples of projective rational surfaces that are not splinters. In this paragraph, we use Bhatt's [Proposition 3.6](#) to show that certain projective rational surfaces are not splinters. First, we have the following example of surfaces that are not globally F -regular.

Example 11.9 ([[SS10](#), Ex. 6.6]). Let k be an algebraically closed field of positive characteristic. If X is the blow-up of \mathbb{P}_k^2 in 9 closed points in general position, then $-K_X$ is not big. Therefore, by [Proposition 3.11](#), X is not globally F -regular. Furthermore, this shows that the blow-up of \mathbb{P}^2 in at least 9 points in general position is not globally F -regular, as global F -regularity descends along birational morphisms ([Lemma 6.3](#)).

On the other hand, we note that if X is the blow-up of \mathbb{P}^2 in 9 points lying on an ordinary elliptic curve, then X is F -split; see [Remark 11.6](#). Since being ordinary is an open property, it follows that the blow-up of \mathbb{P}^2 in 9 points in general position is F -split.

We can extend [Example 11.9](#) and show that in some cases the blow-up of \mathbb{P}^2 in 9 points is not a splinter :

Proposition 11.10. *Let X be the blow-up of \mathbb{P}_k^2 in 9 distinct k -rational points. Assume either of the following conditions:*

- (i) *The base field k is the algebraic closure of a finite field and the 9 points lie on a smooth cubic curve (e.g., the 9 points are in general position).*
- (ii) *The base field k has positive characteristic and the 9 points lie at the transverse intersection of two cubic curves in \mathbb{P}_k^2 .*

Then X is not a splinter.

Proof. In both cases, we show that the anticanonical divisor $-K_X$ is semiample and satisfies $H^1(X, \mathcal{O}_X(-nK_X)) \neq 0$ for some $n > 0$. It follows from Proposition 3.6 that X is not a splinter.

In case (i), the anticanonical divisor $-K_X$ is the strict transform of the smooth cubic curve and is therefore smooth of genus 1. Since $(-K_X)^2 = 0$, we get from [Tot09, Thm. 2.1] that $-K_X$ is semiample, that is, there exists a positive integer n such that $-nK_X$ is basepoint free. In particular, since K_X is not torsion, $h^0(-nK_X) \geq 2$. By Riemann–Roch $\chi(-nK_X) = 1$, and hence $h^1(-nK_X) \neq 0$.

In case (ii), $-K_X$ is basepoint free (in particular semiample) and satisfies $h^0(-K_X) \geq 2$. Indeed, $-K_X$ admits two sections, corresponding to the strict transforms of the cubics, which do not meet in any point in the blow-up as they pass through the 9 points in \mathbb{P}_k^2 from different tangent directions. On the other hand, we have $\chi(-K_X) = 1$ and it follows that $h^1(-K_X) \neq 0$. \square

Remark 11.11. An alternative proof of Proposition 11.10(ii) can be obtained by using Lemma 4.3. Indeed, if the 9 blown up points lie in the intersection of two distinct smooth cubic curves, the set of cubic curves passing through the 9 points forms a pencil and we obtain an elliptic fibration $X \rightarrow \mathbb{P}^1$. Since the generic fiber of this fibration is not a splinter, X is not a splinter.

Proposition 11.10 leaves open the question of whether the blow-up of \mathbb{P}_k^2 in 9 closed points in very general position over an uncountable algebraically closed field k of positive characteristic is a splinter. Finally, further examples of rational surfaces over $\overline{\mathbb{F}}_p$ that are not splinters are provided by the following:

Proposition 11.12. *Let k be the algebraic closure of a finite field. Let $d \geq 4$ be an integer and let C be an irreducible curve of degree d in \mathbb{P}_k^2 . Let $n := \binom{2+d}{2} = \frac{(d+2)(d+1)}{2}$; e.g., $n = 15$ if $d = 4$. The blow-up X of \mathbb{P}_k^2 in n distinct smooth points of C is not a splinter.*

Proof. By Proposition 3.6, it suffices to construct a semiample invertible sheaf \mathcal{L} on X such that $H^1(X, \mathcal{L}) \neq 0$. For that purpose, we consider the class D of the strict transform of the curve C . We have $D = dH - \sum_{i=1}^n E_i$, where H is the pullback of the hyperplane class in \mathbb{P}_k^2 and E_i are the exceptional classes lying above the n blown up points. We claim that the invertible sheaf $\mathcal{L} := \mathcal{O}_X(D)$ is semiample and satisfies $H^1(X, \mathcal{L}) \neq 0$. On the one hand, D is nef since it is effective and satisfies $D^2 = d^2 - n > 0$ for $d \geq 4$. Being nef and having positive self intersection, it is also big, see, e.g., [Kol96, Cor. 2.16]. By Keel’s [Kee99, Cor. 0.3] any nef and big invertible sheaf on a surface over the algebraic closure of a finite field is semiample. On the other hand, by Riemann–Roch

$$\chi(D) = 1 + \frac{1}{2}(D^2 - K_X \cdot D) = 1 + \frac{1}{2}(d^2 + 3d - 2n) = 0,$$

for $K_X = -3H + \sum_i E_i$ the canonical divisor on X . Since D is effective, $H^0(X, \mathcal{L}) \neq 0$, and we conclude that $H^1(X, \mathcal{L}) \neq 0$. \square

12. K -EQUIVALENCE, \mathcal{O} -EQUIVALENCE, AND D -EQUIVALENCE

The aim of this section is to study the derived invariance of the (derived) splinter property and of global F -regularity for projective varieties over a field of positive characteristic. For that purpose, we introduce the notion of \mathcal{O} -equivalence, which is closely related to K -equivalence but offers more flexibility, and explore whether the (derived) splinter property and global F -regularity are preserved under (strong) \mathcal{O} -equivalence.

12.1. K -equivalence. In this paragraph, we fix an excellent Noetherian scheme S admitting a dualizing complex ω_S^\bullet . Any scheme X over S with structure morphism $h: X \rightarrow S$ of finite type and separated will be endowed with the dualizing complex $\omega_X^\bullet := h^! \omega_S^\bullet$. If X is normal, there is a unique, up to linear equivalence, Weil divisor K_X on X such that $\omega_X \cong \mathcal{O}_X(K_X)$. The following notions are classical, at least in the case of smooth varieties over a field (where they agree):

Definition 12.1 (K -equivalence and strong K -equivalence). Let X and Y be integral normal \mathbb{Q} -Gorenstein schemes of finite type and separated over S . We say X and Y are *K -equivalent* if there exists a normal scheme Z over S with proper birational S -morphisms $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ such that p^*K_X and q^*K_Y are \mathbb{Q} -linearly equivalent.

If in addition X and Y are Gorenstein, we say X and Y are *strongly K -equivalent* if there exists a normal scheme Z over S with proper birational S -morphisms $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ such that p^*K_X and q^*K_Y are linearly equivalent.

Obviously, if X and Y are Gorenstein and strongly K -equivalent, then they are K -equivalent. The converse holds provided X and Y are not too singular; see [Proposition 12.2\(i\)](#) below. The following [Proposition 12.2\(iii\)](#) says, in particular, that K -equivalent normal terminal varieties are isomorphic in codimension 1. This is certainly well-known, at least in characteristic zero, see, e.g., [\[Kaw02, Lem. 4.2\]](#), but as we were not able to find a suitable reference for our more general setting, we provide a proof. An integral normal excellent scheme X of finite type and separated over S is said to be *terminal* (resp. *canonical*) if X is \mathbb{Q} -Gorenstein and for any proper birational S -morphism $f: X' \rightarrow X$ with X' normal, the discrepancies of the exceptional divisors are all positive (resp. nonnegative).

Proposition 12.2. *Let X and Y be integral normal schemes of finite type and separated over S . Assume that X and Y are K -equivalent.*

- (i) *If X and Y are canonical and Gorenstein, then X and Y are strongly K -equivalent.*
- (ii) *If X and Y are Gorenstein proper varieties over $S = \operatorname{Spec} k$ a field, and either X or Y is canonical, then X and Y are strongly K -equivalent.*
- (iii) *If X and Y are terminal, then the induced birational map $X \dashrightarrow Y$ is small in the sense of [Definition 4.6](#).*

Proof. We first prove (i) and (iii). Let Z be a normal scheme over S with proper birational morphisms $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ such that $p^*K_X \sim_{\mathbb{Q}} q^*K_Y$. Let $E_i \subseteq \operatorname{Exc}(p)$ and $F_j \subseteq \operatorname{Exc}(q)$ be the irreducible components, endowed with their reduced structure, of codimension 1 of the exceptional loci of p and q , respectively. Let $K_{Z/X} = \sum_i a_i E_i$ and $K_{Z/Y} = \sum_j b_j F_j$ be the relative canonical \mathbb{Q} -divisors, where $a_i, b_j \in \mathbb{Q}_{\geq 0}$ (resp. $a_i, b_j \in \mathbb{Q}_{> 0}$) if X and Y are canonical (resp. terminal). Let

$$D := K_{Z/Y} - K_{Z/X} = \sum b_j F_j - \sum a_i E_i.$$

The divisor D is \mathbb{Q} -Cartier and we have $D \sim_{\mathbb{Q}} p^*K_X - q^*K_Y \sim_{\mathbb{Q}} 0$. Note that $-D$ is p -nef and $p_*D = \sum_j b_j p_*F_j$ is effective. By the Negativity Lemma [\[Bha+23, Lem. 2.16\]](#) (which can be applied since S is assumed to be excellent and since we can assume that p and q are projective after applying Chow's Lemma [\[Sta24, Tag 02O2\]](#) and normalizing [\[Sta24, Tag 035E\]](#)), D is effective. (For the classical version of the Negativity Lemma in characteristic zero, see [\[KM98, Lem. 3.39\]](#).) Arguing similarly for $-D$ shows that $-D$ is effective, and therefore that $D = 0$. This establishes (i), as under the additional assumption that X and Y are Gorenstein we have that $a_i, b_i \in \mathbb{Z}_{\geq 0}$ and $p^*K_X - q^*K_Y \sim D = 0$. Regarding (iii), if X and Y are terminal, we have $\operatorname{Supp}(K_{Z/X}) = \operatorname{Exc}(p)$ and $\operatorname{Supp}(K_{Z/Y}) = \operatorname{Exc}(q)$ up to some locally closed subsets of codimension ≥ 2 . Therefore, $D = 0$ implies that $\operatorname{Exc}(p) = \operatorname{Exc}(q)$ up to some locally closed subsets of codimension ≥ 2 . This proves the statement, since $p(\operatorname{Exc}(p))$ and $q(\operatorname{Exc}(q))$ both have codimension ≥ 2 (a proper birational morphism to a normal Noetherian scheme has geometrically connected fibers by [\[Sta24, Tag 03H0\]](#)).

We now prove (ii). Assume that Y is canonical and keep the notation as above. Then $b_j \in \mathbb{Z}_{\geq 0}$ for all j so that p_*D is effective and we can conclude as above that D is effective. Thus, D is an effective divisor with $nD \sim 0$ for some $n > 0$. But we can identify the linear system $|nD| = \mathbb{P}(H^0(Z, \mathcal{O}_Z)) = \{0\}$ as sets, showing that $nD = 0$ and consequently that $D = 0$. \square

Together with [Lemma 4.1](#), we obtain that both the splinter property and global F -regularity for normal terminal varieties over a field of positive characteristic are invariant under K -equivalence:

Corollary 12.3. *Let X and Y be K -equivalent integral normal terminal schemes of finite type and separated over S . The following statements hold.*

- (i) *X is a splinter if and only if Y is a splinter.*
- (ii) *X is globally F -regular if and only if Y is globally F -regular, provided $S = \operatorname{Spec} k$ with k an F -finite field.*

Proof. This is the combination of [Lemma 4.4](#), [Proposition 4.7](#), and [Proposition 12.2\(iii\)](#). \square

Remark 12.4. [Proposition 12.2\(iii\)](#) does not hold without restrictions on the singularities of X and Y . Consider indeed any crepant morphism $p: Y \rightarrow X$ to a Gorenstein variety X over a field k , with exceptional locus containing a divisor. Then $Y = Z \rightarrow X$ provides a strong K -equivalence between the Gorenstein proper varieties X and Y that does not induce a small birational map. Nonetheless, we know from [Proposition 5.11](#) that if $p: Y \rightarrow X$ is a crepant morphism of normal varieties over a field k of positive characteristic, then X is a splinter if and only if Y is a splinter. In the next paragraph, we will introduce the notion of \mathcal{O} -equivalence and will use it to circumvent going through small birational maps to improve upon [Corollary 12.3](#) and show that the derived splinter property for proper varieties is invariant under strong \mathcal{O} -equivalence; see [Theorem 12.15](#).

The following proposition in characteristic zero is [\[KM98, Cor. 5.24\]](#). We provide an analogous proof in the general situation of normal excellent Noetherian schemes admitting a dualizing complex.

Proposition 12.5. *Let X be a normal scheme of finite type and separated over S . If X is Gorenstein, then X is pseudo-rational if and only if X is canonical.*

Proof. Recall from [\[LT81, Cor. on p.107\]](#) that a normal excellent Cohen–Macaulay Noetherian scheme X admitting a dualizing complex is pseudo-rational if and only if for every proper birational morphism $p: Y \rightarrow X$ with Y normal, the canonical map $p_*\omega_Y \rightarrow \omega_X$ is an isomorphism. Let $p: Y \rightarrow X$ be a proper birational map with Y normal and denote by $\operatorname{Exc}(p) \subseteq Y$ the exceptional locus so that p is an isomorphism over $U := X \setminus p(\operatorname{Exc}(p))$. Since X is normal, $\operatorname{codim}_X p(\operatorname{Exc}(p)) \geq 2$, and the map $p_*\omega_Y \rightarrow \omega_X$ can be identified with

$$H^0(V, p_*\omega_Y) \rightarrow H^0(V \cap U, p_*\omega_Y) = H^0(V \cap U, \omega_X) = H^0(U, \omega_X)$$

for any open subset $V \subseteq X$, where the first map is the restriction map and the last equality holds since ω_X is reflexive; cf. [\[LT81, p. 109\]](#). Assume now that X is Gorenstein, and let $K_Y \sim p^*K_X + E$ with E an effective divisor supported on $\operatorname{Exc}(p)$. Via the projection formula, the canonical map $p_*\omega_Y \rightarrow \omega_X$ identifies with the map $\omega_X \otimes p_*\mathcal{O}_Y(E) \rightarrow \omega_X \otimes \mathcal{O}_X = \omega_X$ obtained by tensoring the identity on ω_X with the map $p_*\mathcal{O}_Y(E) \rightarrow \mathcal{O}_X$ given by restricting a local section of $p_*\mathcal{O}_Y(E)$ along $U \hookrightarrow X$. Since E is supported on $\operatorname{Exc}(p)$, this map $p_*\mathcal{O}_Y(E) \rightarrow \mathcal{O}_X$ is an isomorphism if and only if E is effective. Since ω_X is invertible, this proves the statement of the proposition. \square

12.2. \mathcal{O} -equivalence. In this paragraph, we use the formalism of the exceptional inverse image functor, as described in [\[Sta24, Tag 0A9Y\]](#) (under the name of *upper shriek functor*), for separated schemes of finite type over a fixed Noetherian base; see [Section 2.2](#).

Definition 12.6 (\mathcal{O} -equivalence and strong \mathcal{O} -equivalence). Let S be a Noetherian scheme and let X and Y be schemes of finite type and separated over S .

We say X and Y are *strongly \mathcal{O} -equivalent*, if there exists a scheme Z over S with proper birational S -morphisms $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ such that $p^!\mathcal{O}_X \cong q^!\mathcal{O}_Y$ holds in $\operatorname{D}_{\operatorname{Coh}}(\mathcal{O}_Z)$.

We say X and Y are *\mathcal{O} -equivalent*, if there exists a scheme Z over S with proper birational S -morphisms $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ and a proper surjective morphism $\mu: \tilde{Z} \rightarrow Z$ such that $\mu^!p^!\mathcal{O}_X \cong \mu^!q^!\mathcal{O}_Y$ holds in $\operatorname{D}_{\operatorname{Coh}}(\mathcal{O}_{\tilde{Z}})$.

Obviously, strong \mathcal{O} -equivalence implies \mathcal{O} -equivalence. As will become clear, \mathcal{O} -equivalence offers more generality and more flexibility than K -equivalence: in particular, it does not involve any (\mathbb{Q}) -Gorenstein assumption. For integral normal Gorenstein schemes, (strong) \mathcal{O} -equivalence and (strong) K -equivalence relate as follows:

Proposition 12.7. *Let X and Y be integral normal Gorenstein schemes of finite type and separated over an excellent Noetherian scheme S admitting a dualizing complex ω_S^\bullet . Consider the following statements :*

- (i) X and Y are strongly K -equivalent.
- (ii) X and Y are strongly \mathcal{O} -equivalent.
- (iii) X and Y are K -equivalent.
- (iv) X and Y are \mathcal{O} -equivalent.

Then (i) \Leftrightarrow (ii) \Rightarrow (iii) \Leftrightarrow (iv). If in addition X and Y are canonical, then (i) \Leftrightarrow (ii) \Leftrightarrow (iii) \Leftrightarrow (iv).

Proof. Let $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ be birational morphisms of schemes of finite type and separated over a Noetherian scheme S admitting a dualizing complex. Recall from Lemma 2.2 that if X and Y are Gorenstein, then $p^!\mathcal{O}_X \cong q^!\mathcal{O}_Y$ if and only if $p^*\omega_X \cong q^*\omega_Y$. If X is reduced, then Z is generically reduced and hence [Sta24, Tag 0BXC] the normalization $Z^\nu \rightarrow Z$ is birational. We thus see that strong \mathcal{O} -equivalence coincides with strong K -equivalence, i.e., that (i) \Leftrightarrow (ii).

The implication (ii) \Rightarrow (iv) is clear and holds without assuming X and Y to be normal Gorenstein.

For (iii) \Rightarrow (iv), consider the cyclic covering associated to the torsion invertible sheaf $\mathcal{L} := p^*\omega_X \otimes q^*\omega_Y^{-1}$, i.e.,

$$\mu: \tilde{Z} := \operatorname{Spec}_Z \left(\bigoplus_{i=0}^{r-1} \mathcal{L}^i \right) \rightarrow Z,$$

where r is the torsion index of \mathcal{L} . As in the proof of Proposition 9.5, we have that $\mu^*\mathcal{L} = \mathcal{O}_{\tilde{Z}}$, i.e., $\mu^*p^*\omega_X \cong \mu^*q^*\omega_Y$. One concludes with Lemma 2.2 that $\mu^!p^!\mathcal{O}_X \cong \mu^!q^!\mathcal{O}_Y$.

For (iv) \Rightarrow (iii), recall, e.g., from [Sta24, Tag 02U9], that for a proper morphism $\mu: \tilde{Z} \rightarrow Z$ and an invertible sheaf \mathcal{L} on Z we have

$$(12.8) \quad \mu_*(c_1(\mu^*\mathcal{L}) \cap [\tilde{Z}]) = c_1(\mathcal{L}) \cap \mu_*[\tilde{Z}] \quad \text{in } \operatorname{CH}_{\dim Z - 1}(Z).$$

Assume now that X and Y are \mathcal{O} -equivalent and let $\mu: \tilde{Z} \rightarrow Z$, $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ be as in Definition 12.6. Up to normalizing \tilde{Z} and Z , we may and do assume that Z is normal. By Chow's Lemma [Sta24, Tag 02O2] and by taking hyperplane sections, we may assume that μ is generically finite. The condition $\mu^!p^!\mathcal{O}_X \cong \mu^!q^!\mathcal{O}_Y$ together with (12.8) then imply that the divisor class associated to $p^*\omega_X \otimes q^*\omega_Y^{-1}$ is torsion, and hence by normality of Z that $p^*\omega_X \otimes q^*\omega_Y^{-1}$ is a torsion invertible sheaf on Z , i.e. $p^*K_X \sim_{\mathbb{Q}} q^*K_Y$.

Finally if X and Y are canonical, then (i) \Leftrightarrow (iii) by Proposition 12.2. \square

Moreover, under some regularity assumption, the cohomology of the structure sheaf is invariant under strong \mathcal{O} -equivalence:

Proposition 12.9. *Let X and Y be excellent regular schemes of finite type and separated over a Noetherian scheme S . If there exists an excellent regular scheme Z of finite type and separated over S with projective birational morphisms $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ over S such that $p^!\mathcal{O}_X \cong q^!\mathcal{O}_Y$, then $H^i(X, \mathcal{O}_X) \cong H^i(Y, \mathcal{O}_Y)$ as $H^0(S, \mathcal{O}_S)$ -modules for all $i \geq 0$.*

Proof. By [CR15, Thm. 1.1], both canonical maps $\mathcal{O}_X \rightarrow \operatorname{Rp}_*\mathcal{O}_Z$ and $\mathcal{O}_Y \rightarrow \operatorname{Rq}_*\mathcal{O}_Z$ are isomorphisms. Using the fact that the exceptional inverse image functor commutes with shifts, we obtain the chain of isomorphisms

$$\begin{aligned} H^i(Y, \mathcal{O}_Y) &= \operatorname{Hom}_{\operatorname{D}_{\operatorname{Coh}}(\mathcal{O}_Y)}(\mathcal{O}_Y, \mathcal{O}_Y[i]) = \operatorname{Hom}_{\operatorname{D}_{\operatorname{Coh}}(\mathcal{O}_Y)}(\operatorname{Rq}_*\mathcal{O}_Z, \mathcal{O}_Y[i]) \\ &= \operatorname{Hom}_{\operatorname{D}_{\operatorname{Coh}}(\mathcal{O}_Z)}(\mathcal{O}_Z, q^!\mathcal{O}_Y[i]) \cong \operatorname{Hom}_{\operatorname{D}_{\operatorname{Coh}}(\mathcal{O}_Z)}(\mathcal{O}_Z, p^!\mathcal{O}_X[i]) \\ &= \operatorname{Hom}_{\operatorname{D}_{\operatorname{Coh}}(\mathcal{O}_X)}(\operatorname{Rp}_*\mathcal{O}_Z, \mathcal{O}_X[i]) = \operatorname{Hom}_{\operatorname{D}_{\operatorname{Coh}}(\mathcal{O}_X)}(\mathcal{O}_X, \mathcal{O}_X[i]) = H^i(X, \mathcal{O}_X). \quad \square \end{aligned}$$

Remark 12.10. Let X and Y be regular schemes of finite type and separated over a perfect field k . If X and Y are strongly \mathcal{O} -equivalent and if resolution of singularities holds for reduced schemes of dimension $\dim X$ over k , then by applying Chow's Lemma to Z as in Definition 12.6 and then resolving singularities, one may choose Z to be regular. Hence, provided resolution of singularities holds for reduced schemes of dimension $\dim X$ over k (which is the case if $\dim X \leq 3$ by [CP09]), Proposition 12.9 shows that if X and Y are strongly \mathcal{O} -equivalent, then $H^i(X, \mathcal{O}_X) \cong H^i(Y, \mathcal{O}_Y)$ for all $i \geq 0$.

12.3. D -equivalence. The following is classical.

Definition 12.11 (D -equivalence). Two proper varieties X and Y over a field k are said to be D -equivalent (or *derived equivalent*) if there is a k -linear equivalence of categories $D^b(X) \cong D^b(Y)$ between their bounded derived categories of coherent sheaves.

Kawamata [Kaw02, Thm. 2.3(2)] showed that if X and Y are D -equivalent smooth projective varieties over an algebraically closed field of characteristic zero and if K_X or $-K_X$ is big, then X and Y are K -equivalent. We have the same result in the broader context of normal Gorenstein projective varieties over any field:

Proposition 12.12. *Let X and Y be normal Gorenstein projective varieties over a field k . Assume that K_X or $-K_X$ is big. If X and Y are D -equivalent, then X and Y are K -equivalent (equivalently, \mathcal{O} -equivalent).*

Proof. By [LO10, Cor. 9.17], a k -linear equivalence $D^b(X) \cong D^b(Y)$ is induced by a Fourier–Mukai transform with kernel $K \in D^b(X \times_k Y)$. Denote by $\pi_X: X \times_k Y \rightarrow X$ and $\pi_Y: X \times_k Y \rightarrow Y$ the projections to X and Y . By [Her+09, Prop. 4.2], we have a natural isomorphism

$$(12.13) \quad R\mathcal{H}om_{\mathcal{O}_{X \times_k Y}}(K, \pi_X^! \mathcal{O}_X) \cong R\mathcal{H}om_{\mathcal{O}_{X \times_k Y}}(K, \pi_Y^! \mathcal{O}_Y).$$

(Note that the proof of [Her+09, Prop. 4.2] does not require that the base field be algebraically closed.) On the other hand, by base change [Sta24, Tag 0E9U] and since X and Y are Gorenstein, $\pi_X^! \mathcal{O}_X \cong L\pi_Y^* \omega_Y^\bullet = \pi_Y^* \omega_Y[\dim Y]$ and $\pi_Y^! \mathcal{O}_Y \cong L\pi_X^* \omega_X^\bullet = \pi_X^* \omega_X[\dim X]$. Thus, (12.13) is equivalent to

$$K^\vee \cong K^\vee \otimes \pi_X^* \omega_X \otimes \pi_Y^* \omega_Y^{-1}[\dim X - \dim Y] \quad \text{where } K^\vee := R\mathcal{H}om_{\mathcal{O}_{X \times_k Y}}(K, \mathcal{O}_{X \times_k Y}).$$

As shown in [Her+09, Proof of Prop. 2.10], K^\vee lies in $D^b(X \times_k Y)$, and hence $\dim X = \dim Y$.

Let $\nu: Z^\nu \rightarrow Z$ be the normalization of an irreducible component Z of $\text{Supp}(K^\vee)$ and set $p = \pi_X \circ \nu$ and $q = \pi_Y \circ \nu$. Then, there exists $i \in \mathbb{Z}$ such that $\nu^* \mathcal{H}^i(K^\vee)|_Z$ generically has positive rank $r > 0$. Arguing as in [Huy06, Lem. 6.9], we obtain

$$\mathcal{O}_{Z^\nu}(rp^* K_X) \cong \mathcal{O}_{Z^\nu}(rq^* K_Y).$$

Since K_X or $-K_X$ is big, arguing as in the proof of [Kaw02, Thm. 2.3(2)] (see also [Huy06, Prop. 6.19]) shows that there exists a component Z that dominates X and Y and is such that p and q are birational morphisms. This proves that X and Y are K -equivalent. By Proposition 12.7, X and Y are also \mathcal{O} -equivalent. \square

Remark 12.14. In the same way that D -equivalent smooth proper complex varieties are not necessarily K -equivalent [Ueh04], D -equivalent smooth proper varieties in positive characteristic are not necessarily \mathcal{O} -equivalent. Indeed, in [ABP23], Addington and Bragg have produced D -equivalent smooth projective threefolds over \mathbb{F}_3 with different Hodge numbers $h^{0,i}$ for $i = 1$ and 2 . Such varieties are not \mathcal{O} -equivalent by Proposition 12.9 and Remark 12.10.

12.4. Invariance of the splinter property under strong \mathcal{O} -equivalence. Since a crepant morphism $p: Y \rightarrow X$ provides a strong \mathcal{O} -equivalence between Y and X , the following theorem extends Proposition 5.11(ii) in the case of proper varieties over a field.

Theorem 12.15 (Theorem (H), Derived splinters are stable under strong \mathcal{O} -equivalence). *Let X and Y be proper varieties over a field. If X and Y are strongly \mathcal{O} -equivalent, then X is a derived splinter if and only if Y is a derived splinter.*

Proof. Suppose X is a derived splinter. Let $f: B \rightarrow Y$ be a proper surjective morphism. We have to show that $f^\sharp: \mathcal{O}_Y \rightarrow Rf_* \mathcal{O}_B$ splits in $D_{\text{Coh}}(\mathcal{O}_Y)$. Consider the diagram

$$\begin{array}{ccccc} & & Z & \xleftarrow{f'} & B' \\ & p \swarrow & \searrow q & & \downarrow q' \\ X & & Y & \xleftarrow{f} & B \end{array}$$

where p and q are proper birational over S with $p^! \mathcal{O}_X \cong q^! \mathcal{O}_Y$, and where $B' := B \times_Y Z$. We then have the following commutative diagram:

$$\begin{array}{ccccc}
\mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_X)}(\mathrm{Rp}_* \mathrm{R}f'_* \mathcal{O}_{B'}, \mathcal{O}_X) & \xrightarrow{-\circ \mathrm{Rp}_* f'^{\sharp}} & \mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_X)}(\mathrm{Rp}_* \mathcal{O}_Z, \mathcal{O}_X) & \xrightarrow{-\circ p^{\sharp}} & \mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_X)}(\mathcal{O}_X, \mathcal{O}_X) \\
\downarrow = & & \downarrow = & & \\
\mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_Z)}(\mathrm{R}f'_* \mathcal{O}_{B'}, p^! \mathcal{O}_X) & \xrightarrow{-\circ f'^{\sharp}} & \mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_Z)}(\mathcal{O}_Z, p^! \mathcal{O}_X) & & \\
\downarrow \cong & & \downarrow \cong & & \\
\mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_Z)}(\mathrm{R}f'_* \mathcal{O}_{B'}, q^! \mathcal{O}_Y) & \xrightarrow{-\circ f'^{\sharp}} & \mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_Z)}(\mathcal{O}_Z, q^! \mathcal{O}_Y) & & \\
\downarrow = & & \downarrow = & & \\
\mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_Y)}(\mathrm{R}q_* \mathrm{R}f'_* \mathcal{O}_{B'}, \mathcal{O}_Y) & \xrightarrow{-\circ \mathrm{R}q_* f'^{\sharp}} & \mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_Y)}(\mathrm{R}q_* \mathcal{O}_Z, \mathcal{O}_Y) & \xrightarrow{-\circ q^{\sharp}} & \mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_Y)}(\mathcal{O}_Y, \mathcal{O}_Y).
\end{array}$$

Let $s: \mathrm{Rp}_* \mathrm{R}f'_* \mathcal{O}_{B'} \rightarrow \mathcal{O}_X$ be a section of $(p \circ f')^{\sharp}: \mathcal{O}_X \rightarrow \mathrm{Rp}_* \mathrm{R}f'_* \mathcal{O}_{B'}$ and let $t: \mathrm{R}q_* \mathrm{R}f'_* \mathcal{O}_{B'} \rightarrow \mathcal{O}_Y$ be the image of s under composition of the left vertical arrows above. Let $s' := s \circ \mathrm{Rp}_* f'^{\sharp}$ and $t' := t \circ \mathrm{R}q_* f'^{\sharp}$. Since $\mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{O}_Y, \mathcal{O}_Y)$ is a field, it is enough to show that $t' \circ q^{\sharp}$ is nonzero. Choose dense open subsets $U \subseteq X$ and $V \subseteq Y$ such that $p^{-1}(U) = q^{-1}(V)$ and such that p and q restrict to isomorphisms over U and V , respectively. Since Y is integral, it is enough to show that $t' \circ q^{\sharp}|_V$ is nonzero. As $-\circ q^{\sharp}: \mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_V)}(\mathrm{R}q_* \mathcal{O}_Z|_V, \mathcal{O}_Y|_V) \rightarrow \mathrm{Hom}_{\mathrm{D}_{\mathrm{Coh}}(\mathcal{O}_V)}(\mathcal{O}_Y|_V, \mathcal{O}_Y|_V)$ is an isomorphism, the latter is equivalent to $t'|_V$ being nonzero. Since s' is a section of $\mathcal{O}_X \rightarrow \mathrm{Rp}_* \mathcal{O}_Z$, $s'|_U$ is nonzero. This shows that $t'|_V$ is nonzero, since $t'|_V$ is the image of $s'|_U$ under the middle vertical isomorphism when restricted to $U \leftarrow p^{-1}(U) = q^{-1}(V) \rightarrow V$. Hence, $\mathcal{O}_Y \rightarrow \mathrm{R}q_* \mathrm{R}f'_* \mathcal{O}_{B'} = \mathrm{R}f_* \mathrm{R}q'_* \mathcal{O}_{B'}$ splits and thus, $\mathcal{O}_Y \rightarrow \mathrm{R}f_* \mathcal{O}_B$ splits. \square

Combined with [Proposition 12.12](#), we obtain a partial answer to the question, whether the splinter property for smooth projective schemes over a field is stable under derived equivalence:

Corollary 12.16 (Theorem [\(F\)](#)). *Let X and Y be normal Gorenstein projective varieties over a field of positive characteristic. Assume that $-K_X$ is big. If X and Y are D -equivalent, then X is a splinter if and only if Y is a splinter.*

Proof. Under the assumption that $-K_X$ is big, we have from [Proposition 12.12](#) that X and Y are \mathcal{O} -equivalent. Assume now that X and Y are \mathcal{O} -equivalent and that X is a splinter. Recall from [Proposition 3.5](#) that a splinter is normal pseudo-rational, and from [Proposition 12.5](#) that a normal Gorenstein pseudo-rational variety has canonical singularities. Therefore X and Y are strongly \mathcal{O} -equivalent by [Proposition 12.2\(ii\)](#) and [Proposition 12.7](#). We may now conclude with [Theorem 12.15](#) that Y is a splinter by using the fact due to Bhatt [[Bha12](#), Thm. 1.4] that a Noetherian scheme of positive characteristic is a splinter if and only if it is a derived splinter. \square

Remark 12.17. According to [Conjecture 3.12](#), the assumption that $-K_X$ is big in [Corollary 12.16](#) is conjecturally superfluous.

12.5. Invariance of global F -regularity under strong \mathcal{O} -equivalence and D -equivalence.

Theorem 12.18 (Theorem [\(I\)](#), Global F -regularity is stable under strong \mathcal{O} -equivalence). *Let X and Y be projective varieties over a field k of positive characteristic. If X and Y are strongly \mathcal{O} -equivalent, then X is normal globally F -regular if and only if Y is normal globally F -regular.*

Proof. Let $p: Z \rightarrow X$ and $q: Z \rightarrow Y$ be as in [Definition 12.6](#). After possibly normalizing Z , we may and do assume that Z is normal. Assume that X is normal globally F -regular. By [Proposition 3.10](#), X is a splinter. By [[Bha12](#), Thm. 1.4], X is a derived splinter; in particular the map $\mathcal{O}_X \rightarrow \mathrm{Rp}_* \mathcal{O}_Z$ splits. Moreover, by [Theorem 12.15](#), Y is also a derived splinter and hence is normal. Fix nonempty regular affine open subsets $U \subseteq X$ and $V \subseteq Y$ such that $p^{-1}(U) = q^{-1}(V)$ and such that $p|_U$ and $q|_V$ are isomorphisms. By [[Sta24](#), Tag 0BCU], $D := X \setminus U$ and $E := Y \setminus V$ are divisors. Since Y is projective, any Weil divisor on Y is dominated by a Cartier divisor and we may thus further assume that E is Cartier. By [Theorem 6.1](#) it suffices to show that there exists $e > 0$ such that $\mathcal{O}_Y \rightarrow F_*^e \mathcal{O}_Y(E)$ splits. Let $D' \geq D$ be a Cartier divisor on X . Since $q^* E = \sum_i a_i E_i$ for some $a_i \in \mathbb{Z}_{>0}$ with E_i the irreducible components of $\mathrm{Supp}(q^* E) \subseteq \mathrm{Supp}(p^* D')$, there exists $n \in \mathbb{Z}_{>0}$

such that $q^*E \leq np^*D'$. Let $U' := X \setminus D'$ and set $V' := q(p^{-1}(U'))$. Then U' and V' are regular affine open subsets such that $p|_{U'}$ and $q|_{V'}$ are isomorphisms.

Since X is globally F -regular, there exists an integer $e > 0$ such that $\sigma_{nD'}: \mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(nD')$ splits. By the projection formula, the splitting of $\mathcal{O}_X \rightarrow R p_* \mathcal{O}_Z$ gives a splitting of $\text{id}_{\text{Coh}(X)} \rightarrow R p_* L p^*$. Thus, we obtain a splitting s of

$$\alpha: \mathcal{O}_X \rightarrow F_*^e \mathcal{O}_X(nD') \rightarrow F_*^e R p_* L p^* \mathcal{O}_X(nD') = R p_* F_*^e \mathcal{O}_Z(np^*D').$$

Note that the map α can also be obtained as the composition

$$\mathcal{O}_X \rightarrow R p_* \mathcal{O}_Z \rightarrow R p_* F_*^e \mathcal{O}_Z \rightarrow R p_* F_*^e \mathcal{O}_Z(np^*D').$$

We obtain a commutative diagram (where $D(-)$ stands for $D_{\text{Coh}}(-)$):

$$\begin{array}{ccccc} \text{Hom}_{D(\mathcal{O}_X)}(R p_* F_*^e \mathcal{O}_Z(np^*D'), \mathcal{O}_X) & \xrightarrow{-\circ R p_* \sigma_{np^*D'}^\#} & \text{Hom}_{D(\mathcal{O}_X)}(R p_* \mathcal{O}_Z, \mathcal{O}_X) & \xrightarrow{-\circ p^\#} & \text{Hom}_{D(\mathcal{O}_X)}(\mathcal{O}_X, \mathcal{O}_X) \\ \downarrow = & & \downarrow = & & \\ \text{Hom}_{D(\mathcal{O}_Z)}(F_*^e \mathcal{O}_Z(np^*D'), p^! \mathcal{O}_X) & \xrightarrow{-\circ \sigma_{np^*D'}^\#} & \text{Hom}_{D(\mathcal{O}_Z)}(\mathcal{O}_Z, p^! \mathcal{O}_X) & & \\ \downarrow \cong & & \downarrow \cong & & \\ \text{Hom}_{D(\mathcal{O}_Z)}(F_*^e \mathcal{O}_Z(np^*D'), q^! \mathcal{O}_Y) & \xrightarrow{-\circ \sigma_{np^*D'}^\#} & \text{Hom}_{D(\mathcal{O}_Z)}(\mathcal{O}_Z, q^! \mathcal{O}_Y) & & \\ \downarrow \gamma & & \downarrow = & & \\ \text{Hom}_{D(\mathcal{O}_Z)}(F_*^e \mathcal{O}_Z(q^*E), q^! \mathcal{O}_Y) & \xrightarrow{-\circ \sigma_{q^*E}^\#} & \text{Hom}_{D(\mathcal{O}_Z)}(\mathcal{O}_Z, q^! \mathcal{O}_Y) & & \\ \downarrow = & & \downarrow = & & \\ \text{Hom}_{D(\mathcal{O}_Y)}(R q_* F_*^e \mathcal{O}_Z(q^*E), \mathcal{O}_Y) & \xrightarrow{-\circ R q_* \sigma_{q^*E}^\#} & \text{Hom}_{D(\mathcal{O}_Y)}(R q_* \mathcal{O}_Z, \mathcal{O}_Y) & \xrightarrow{-\circ q^\#} & \text{Hom}_{D(\mathcal{O}_Y)}(\mathcal{O}_Y, \mathcal{O}_Y), \end{array}$$

where s maps to the identity under the composition of the top horizontal arrows. Here the vertical map γ is the precomposition with $F_*^e \mathcal{O}_Z(q^*E) \rightarrow F_*^e \mathcal{O}_Z(np^*D')$.

Let $t: R q_* F_*^e \mathcal{O}_Z(q^*E) \rightarrow \mathcal{O}_Y$ be the image of s under the composition of the left vertical arrows. As in the proof of [Theorem 12.15](#), we conclude that t maps to a nonzero element of $\text{Hom}_{D(\mathcal{O}_Y)}(\mathcal{O}_Y, \mathcal{O}_Y)$, thereby providing a splitting of $\mathcal{O}_Y \rightarrow F_*^e \mathcal{O}_Y(E) \rightarrow F_*^e R q_* L q^* \mathcal{O}_Y(E) = R q_* F_*^e \mathcal{O}_Z(q^*E)$. \square

Remark 12.19 (The F -split property is stable under strong \mathcal{O} -equivalence). Assume that X and Y are normal proper varieties over a field k of positive characteristic. If X and Y are strongly \mathcal{O} -equivalent, then X is F -split if and only if Y is F -split. This follows indeed by considering in the proof of [Theorem 12.18](#) dense opens $U' = U \subseteq X$ and $V' = V \subseteq Y$ such that $p^{-1}(U) = q^{-1}(V)$ and such that $p|_{U'}$ and $q|_{V'}$ are isomorphisms, and by setting $D = 0$ and $E = 0$.

Corollary 12.20 (Theorem [\(G\)](#), Global F -regularity is stable under D -equivalence). *Let X and Y be normal Gorenstein projective varieties over a field k of positive characteristic. If X and Y are D -equivalent, then X is globally F -regular if and only if Y is globally F -regular.*

Proof. Since normal globally F -regular projective varieties have big anticanonical class, this follows as in [Corollary 12.16](#) from [Proposition 12.2\(ii\)](#), [Propositions 12.5](#), [12.7](#) and [12.12](#), combined with [Theorem 12.18](#). \square

Remark 12.21 (The F -split property and D -equivalence). As asked by Zsolt Patakfalvi, we are unaware whether the F -split property is stable under D -equivalence. As a partial result, we mention that under the assumptions of [Corollary 12.20](#), if X and Y are D -equivalent and if $-K_X$ is big, then X is F -split if and only if Y is F -split. For this, one argues as in the proof of [Corollary 12.16](#) by using [Remark 12.19](#).

Moreover, if X and Y are D -equivalent abelian varieties or D -equivalent strict Calabi–Yau varieties over a perfect field, then X is F -split if and only if Y is F -split. In the case of abelian varieties, X and Y are then isogenous by [\[Orl02, Thm. 2.19\]](#) and therefore have the same height. In the case of strict Calabi–Yau varieties, X and Y have the same height by [\[AB22, Thm. 5.1\]](#). One concludes with the fact that such varieties have height 1 if and only if they are F -split; this is

classical in the case of abelian varieties and is [GK03, Thm. 2.1] in the case of strict Calabi–Yau varieties.

Remark 12.22. It would be interesting to know whether the derived splinter property is stable under (strong) \mathcal{O} -equivalence for schemes of finite type and separated over a Noetherian base, and whether Theorems **(F)** and **(G)** hold without assuming X and Y to be Gorenstein.

REFERENCES

- [AWZ21] Piotr Achinger, Jakub Witaszek, and Maciej Zdanowicz. “Global Frobenius liftability I”. *J. Eur. Math. Soc. (JEMS)* 23.8 (2021), pp. 2601–2648.
- [ABP23] Nicolas Addington, Daniel Bragg, and Alexander Petrov. “Hodge numbers are not derived invariants in positive characteristic”. *Math. Ann.* 387.1-2 (2023), pp. 847–878.
- [And18] Yves André. “La conjecture du facteur direct”. *Publ. Math. Inst. Hautes Études Sci.* 127 (2018), pp. 71–93.
- [AB22] Benjamin Antieau and Daniel Bragg. “Derived invariants from topological Hochschild homology”. *Algebr. Geom.* 9.3 (2022), pp. 364–399.
- [Bha12] Bhargav Bhatt. “Derived splinters in positive characteristic”. *Compos. Math.* 148.6 (2012), pp. 1757–1786.
- [Bha21] Bhargav Bhatt. *Cohen-Macaulayness of absolute integral closures*. 2021. arXiv: [2008.08070v2](#).
- [Bha+23] Bhargav Bhatt, Linqun Ma, Zsolt Patakfalvi, Karl Schwede, Kevin Tucker, Joe Waldron, and Jakub Witaszek. “Globally $+$ -regular varieties and the minimal model program for threefolds in mixed characteristic”. *Publ. Math. Inst. Hautes Études Sci.* 138 (2023), pp. 69–227.
- [Bis09] Indranil Biswas. “On the fundamental group-scheme”. *Bull. Sci. Math.* 133.5 (2009), pp. 477–483.
- [BM76] Enrico Bombieri and David B. Mumford. “Enriques’ classification of surfaces in char. p . III”. *Invent. Math.* 35 (1976), pp. 197–232.
- [Bri17] Michel Brion. “Some structure theorems for algebraic groups”. *Algebraic groups: structure and actions*. Vol. 94. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2017, pp. 53–126.
- [BK05] Michel Brion and Shrawan Kumar. *Frobenius splitting methods in geometry and representation theory*. Vol. 231. Progress in Mathematics. Birkhäuser Boston Inc., Boston, MA, 2005. x+250.
- [Cai+23] Hanlin Cai, Seungsu Lee, Linqun Ma, Karl Schwede, and Kevin Tucker. *Perfectoid signature, perfectoid Hilbert-Kunz multiplicity, and an application to local fundamental groups*. 2023. arXiv: [2209.04046v2](#).
- [CR22] Javier A. Carvajal-Rojas. “Finite torsors over strongly F -regular singularities”. *Épjournal Géom. Algébrique* 6 (2022), Art. 1, 30.
- [CR15] Andre Chatzistamatiou and Kay Rülling. “Vanishing of the higher direct images of the structure sheaf”. *Compos. Math.* 151.11 (2015), pp. 2131–2144.
- [CP09] Vincent Cossart and Olivier Piltant. “Resolution of singularities of threefolds in positive characteristic. II”. *J. Algebra* 321.7 (2009), pp. 1836–1976.
- [DT23] Rankeya Datta and Kevin Tucker. “On Some Permanence Properties of (Derived) Splinters”. *Michigan Math. J.* 73.2 (2023), pp. 371–400.
- [Ful98] William Fulton. *Intersection theory*. Vol. 2. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics. Springer-Verlag, Berlin, 1998. xiv+470.
- [GK03] Gerard van der Geer and Toshiyuki Katsura. “On the height of Calabi-Yau varieties in positive characteristic”. *Doc. Math.* 8 (2003), pp. 97–113.
- [Gon+15] Yoshinori Gongyo, Zhiyuan Li, Zsolt Patakfalvi, Karl Schwede, Hiromu Tanaka, and Runhong Zong. “On rational connectedness of globally F -regular threefolds”. *Adv. Math.* 280 (2015), pp. 47–78.
- [GT16] Yoshinori Gongyo and Shunsuke Takagi. “Surfaces of globally F -regular and F -split type”. *Math. Ann.* 364.3-4 (2016), pp. 841–855.
- [GW23] Ulrich Görtz and Torsten Wedhorn. *Algebraic geometry II: Cohomology of schemes—with examples and exercises*. Springer Studium Mathematik—Master. 2023. vii+869.
- [Gro65] Alexander Grothendieck. “Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II”. *Inst. Hautes Études Sci. Publ. Math.* 24 (1965).
- [Gro66] Alexander Grothendieck. “Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III”. *Inst. Hautes Études Sci. Publ. Math.* 28 (1966).
- [Har98] Nobuo Hara. “A characterization of rational singularities in terms of injectivity of Frobenius maps”. *Amer. J. Math.* 120.5 (1998), pp. 981–996.
- [Har15] Nobuo Hara. “Looking out for Frobenius summands on a blown-up surface of \mathbb{P}^2 ”. *Illinois J. Math.* 59.1 (2015), pp. 115–142.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977, pp. xvi+496.
- [Har94] Robin Hartshorne. “Generalized divisors on Gorenstein schemes”. *Proceedings of Conference on Algebraic Geometry and Ring Theory in honor of Michael Artin, Part III (Antwerp, 1992)*. Vol. 8. 3. 1994, pp. 287–339.
- [Has03] Mitsuyasu Hashimoto. “Surjectivity of multiplication and F -regularity of multigraded rings”. *Commutative algebra (Grenoble/Lyon, 2001)*. Vol. 331. Contemp. Math. Amer. Math. Soc., Providence, RI, 2003, pp. 153–170.

- [Her+09] Daniel Hernández Ruipérez, Ana Cristina López Martín, and Fernando Sancho de Salas. “Relative integral functors for singular fibrations and singular partners”. *J. Eur. Math. Soc. (JEMS)* 11.3 (2009), pp. 597–625.
- [HH89] Melvin Hochster and Craig Huneke. “Tight closure and strong F -regularity”. 38. Colloque en l’honneur de Pierre Samuel (Orsay, 1987). 1989, pp. 119–133.
- [HH92] Melvin Hochster and Craig Huneke. “Infinite integral extensions and big Cohen-Macaulay algebras”. *Ann. of Math. (2)* 135.1 (1992), pp. 53–89.
- [Huy06] Daniel Huybrechts. *Fourier-Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006. viii+307.
- [Kat70] Nicholas M. Katz. “Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin”. *Inst. Hautes Études Sci. Publ. Math.* 39 (1970), pp. 175–232.
- [KT25] Tatsuro Kawakami and Burt Totaro. “Endomorphisms of varieties and Bott vanishing”. *J. Algebraic Geom.* 34 (2025), pp. 381–405.
- [Kaw02] Yujiro Kawamata. “ D -equivalence and K -equivalence”. *J. Differential Geom.* 61 (2002), pp. 147–171.
- [Kee99] Seán Keel. “Basepoint freeness for nef and big line bundles in positive characteristic”. *Ann. of Math. (2)* 149.1 (1999), pp. 253–286.
- [Kir92] Frances Kirwan. *Complex algebraic curves*. Vol. 23. London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1992. viii+264.
- [Kol96] János Kollár. *Rational curves on algebraic varieties*. Vol. 32. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. Springer-Verlag, Berlin, 1996. viii+320.
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*. Vol. 134. Cambridge Tracts in Mathematics. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. Cambridge University Press, Cambridge, 1998. viii+254.
- [Lip69] Joseph Lipman. “Rational singularities, with applications to algebraic surfaces and unique factorization”. *Inst. Hautes Études Sci. Publ. Math.* 36 (1969), pp. 195–279.
- [LT81] Joseph Lipman and Bernard Teissier. “Pseudorational local rings and a theorem of Briançon-Skoda about integral closures of ideals”. *Michigan Math. J.* 28.1 (1981), pp. 97–116.
- [LO10] Valery A. Lunts and Dmitri O. Orlov. “Uniqueness of enhancement for triangulated categories”. *J. Amer. Math. Soc.* 23.3 (2010), pp. 853–908.
- [Mum70] David Mumford. *Abelian varieties*. Vol. 5. Tata Institute of Fundamental Research Studies in Mathematics. Oxford University Press, London, 1970. viii+242.
- [Nor82] Madhav V. Nori. “The fundamental group-scheme”. *Proc. Indian Acad. Sci. Math. Sci.* 91.2 (1982), pp. 73–122.
- [Orl02] Dmitri O. Orlov. “Derived categories of coherent sheaves on abelian varieties and equivalences between them”. *Izv. Ross. Akad. Nauk Ser. Mat.* 66.3 (2002), pp. 131–158.
- [Par71] Bodo Pareigis. “When Hopf algebras are Frobenius algebras”. *J. Algebra* 18 (1971), pp. 588–596.
- [PST17] Zsolt Patakfalvi, Karl Schwede, and Kevin Tucker. “Positive characteristic algebraic geometry”. *Surveys on recent developments in algebraic geometry*. Vol. 95. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2017, pp. 33–80.
- [PZ20] Zsolt Patakfalvi and Maciej Zdanowicz. *On the Beauville–Bogomolov decomposition in characteristic $p \geq 0$* . 2020. arXiv: [1912.12742v2](https://arxiv.org/abs/1912.12742v2).
- [SS10] Karl Schwede and Karen E. Smith. “Globally F -regular and log Fano varieties”. *Adv. Math.* 224.3 (2010), pp. 863–894.
- [ST14] Karl Schwede and Kevin Tucker. “On the behavior of test ideals under finite morphisms”. *J. Algebraic Geom.* 23.3 (2014), pp. 399–443.
- [Sin99] Anurag K. Singh. “ \mathbf{Q} -Gorenstein splinter rings of characteristic p are F -regular”. *Math. Proc. Cambridge Philos. Soc.* 127.2 (1999), pp. 201–205.
- [Smi00] Karen E. Smith. “Globally F -regular varieties: applications to vanishing theorems for quotients of Fano varieties”. Vol. 48. Dedicated to William Fulton on the occasion of his 60th birthday. 2000, pp. 553–572.
- [Spe20] David E. Speyer. “Frobenius split subvarieties pull back in almost all characteristics”. *J. Commut. Algebra* 12.4 (2020), pp. 573–579.
- [Sta24] The Stacks project authors. *The Stacks project*. 2024. URL: <https://stacks.math.columbia.edu>.
- [Tot09] Burt Totaro. “Moving codimension-one subvarieties over finite fields”. *Amer. J. Math.* 131.6 (2009), pp. 1815–1833.
- [Ueh04] Hokuto Uehara. “An example of Fourier-Mukai partners of minimal elliptic surfaces”. *Math. Res. Lett.* 11.2-3 (2004), pp. 371–375.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, D-33501 BIELEFELD, GERMANY
 Email address: jkrah@math.uni-bielefeld.de, vial@math.uni-bielefeld.de