# Principles and Application of Gravimetric Analysis

## Introduction

- **Gravimetric analysis** is one of the oldest and most fundamental techniques in quantitative analytical chemistry, relying on the measurement of mass.
- The method involves isolating the analyte by converting it into a highly pure, stable, weighable compound (a **precipitate**).
- Accurate results depend entirely on the high **purity** of the precipitate and the successful conversion of all the analytes into the weighable form.
- A deep understanding of **solubility product (**$K_{sp}$**)** and the factors that influence solubility are essential for controlling the precipitation process to achieve quantitative recovery.
-

## Learning Objectives

By the end of this module, you will be able to:

- Apply the concepts of **solubility products (**$K_{sp}$**)** and factors affecting solubility (e.g., common ion effect, $\mathrm{pH}$, temperature) to control the formation and purity of precipitates in gravimetric analysis.
- Solve **problems** applying the gravimetric analysis principles, including calculating the mass of the analyte, percent composition, and required precipitating agent.

# Key Concepts and Definitions

| Term | Definition |
| --- | --- |
| **Gravimetric Analysis** | A quantitative method where the amount of analyte is determined by accurately measuring the mass of a pure, solid compound containing the analyte. |
| **Precipitate** | The insoluble solid compound formed during the analysis that contains the analyte. |
| **Solubility** | The equilibrium constant for the dissolution of a sparingly soluble ionic compound; determines maximum ion concentration. |
| **Solubility Product ($K_{sp}$)** | The reduction in solubility of a salt when a soluble compound containing one of the same ions is added to the solution. |
| **Common Ion Effect** | The process of heating the precipitate in the solution from which it was formed to yield a purer, more filterable product. |
| **Gravimetric Factor (GF)** | The ratio of the molar mass of the analyte to the molar mass of the weighable precipitate, used for stoichiometric conversion. |

# Detailed Discussion

## Principles of Gravimetric Analysis

Gravimetric analysis follows four main steps: precipitation, filtration, washing/digestion, and weighing.

- **Precipitation:** The analyte is converted into a sparingly soluble compound by adding a precipitating agent. This step is governed by the $K_{sp}$ equilibrium.
- **Digestion:** The precipitate is allowed to stand (often while heated) in the presence of the mother liquor. This process dissolves and re-precipitates smaller, less perfect crystals, leading to the formation of larger, purer crystals that are easier to filter (**Ostwald ripening**).
- **Filtration and Washing:** The solid precipitate is separated from the liquid solution (**mother liquor**) and washed to remove impurities.
- **Drying/Ignition and Weighing:** The precipitate is heated to remove solvent and volatile impurities, converting it into a stable, weighable form. The final mass is measured using an analytical balance.

## Factors Affecting Solubility and Purity

Controlling solubility and purity is the heart of successful gravimetric analysis.

- **Solubility Product ($K_{sp}$):** The smaller the $K_{sp}$ value, the lower the solubility and the more quantitative the precipitation will be.

$$K_{sp} = [Cation]^x [Anion]^y$$

- **$\mathrm{pH}$ Control:** For precipitates whose anion is a weak base (e.g., hydroxide or oxalate), solubility is highly dependent on $\mathrm{pH}$. Decreasing the $\mathrm{pH}$ (adding acid) increases the solubility by removing the anion via protonation.
- **Impurities (Coprecipitation):** The main challenge is contamination of the precipitate by soluble compounds. This error, known as **coprecipitation**, occurs when soluble materials are physically trapped (**occlusion** or **inclusion**) or chemically adsorbed (**surface adsorption**) onto the growing crystal lattice. Methods like digestion and reprecipitation are used to minimize coprecipitation errors.

## Solving Gravimetric Problems

Gravimetric calculations rely on converting the measured mass of the final product back to the initial mass of the analyte using the **Gravimetric Factor (GF)**.

- **Gravimetric Factor (GF):** The stoichiometric ratio of the analyte's molar mass to the precipitate's molar mass.

$$GF = \frac{Molar\ Mass\ of\ Analyte}{Molar\ Mass\ of\ Precipitate} \times \frac{Moles\ Mass\ of\ Analyte}{Moles\ Mass\ of\ Precipitate}$$

- **Calculate Mass of Analyte:** Multiply the measured mass of the precipitate by the GF.

$$Mass\ of\ Analyte = Mass\ of\ Precipitate \times GF$$

- **Determine Percent Composition:** Use the calculated mass of the analyte and the initial mass of the sample.

$$\%\ Analyte = \frac{Mass\ of\ Analyte}{Mass\ of\ Sample} \times 100$$

**Example:** To analyze Fe in an ore, it is precipitated as $Fe(OH)_3$ and ignited to $Fe_2O_3$ (the weighable form). The GF is calculated to convert the mass of $Fe_2O_3$ back to the mass of Fe.

$$GF = \frac{2 \times Molar\ Mass\ Fe}{Molar\ Mass\ Fe_2 0_3}$$

## References:


- Harvey, D. T. (2019). *Analytical Chemistry 2.1.* Retrieved December 6, 2025, from https://chem.libretexts.org/Bookshelves/Analytical_Chemistry/Analytical_Chemistry_2.1_(Harvey
- LibreTexts. (n.d.). *The Quantitative Applications of Gravimetry.* Retrieved December 6, 2025, from https://chem.libretexts.org/Bookshelves/Analytical_Chemistry/Analytical_Chemistry_2.1_(Harvey)/09%3A_Gravimetric_and_Volumetric_Methods/9.02%3A_The_Quantitative_Applications_of_Gravimetry
- Flowers, P., Theopold, K., & Langley, R. (2019). *Chemistry 2e.* Retrieved December 6, 2025, from https://openstax.org/details/books/chemistry-2e