

Adult Health Care Issues in the Angelman Syndrome

July 26, 2007
Charles A. Williams, MD
Raymond C. Phillips Unit
Division of Genetics, University of Florida, Gainesville



Harry Angelman

Syndrome description in 1965



The Angelman ("Happy Puppet") Syndrome

Charles A. Williams and Jaime L. Fries
Department of Pediatrics, Division of Genetics, University of Florida College of Medicine, Gainesville



Literature Reports of Adults with AS

Author	Year	Ages
Williams and Fries	1984	5: 17-33 yrs
Bjerre et al.,	1984	1:75 yrs
Buntinx et al.,	1995	18: >18yrs (5 >35yrs; 1: 47 yrs)
Reish and King	1995	1:50 yrs
Laan et al.,	1996	28: 20-53 yrs
Van Buggenhout	2000	3:32, 49, 52 years
Clayton-Smith	2001	28: 16-40 yrs
Harbord	2001	2: 22, 43 yrs
Stecker and Myers	2003	1:32 yrs
Philippart	2005	1: 74 yrs
Thomson	2006	34: mean 21.6 yrs (5.3-39 yrs)
Total cases		122

Laan et al., Am J Med Genet, 1996



TABLE I. Clinical Findings in 28 Adult Angelman Patients Compared to Those in Children as Reported in the Literature

	Adult AS patients	%	AS Children*	%
Face				
Macrotomia	25/28	100		
Mandibular prognathism	26/27	96	95	
Good visual acuity (without glasses)	20/25	80	89	
Strabismus	16/28	57	42	
Keratoconus	2/28	7		
Neurological findings				
Limb hypertonicity	26/28	93		
Hyperreflexia	25/28	89		
Tremor	11/12	86		
Epileptic seizures	23/28	82	86	
Scoliosis	20/28	71	11	
Truncal hypotonia	13/26	50		
Behavior				
Hypersensitivity	14/28	50		
Fixation on food	6/18	33		
Obesity	4/27	15		

*Percentages reported by Clayton-Smith (1998) (82 patients, almost all children).

Laan et al., Am J Med Genet, 1996



TABLE II. Daily Life Activities in Adult AS Patients

	%
Love to watch TV	24/24
Handle spoon or fork	23/27
Ability to express their will to some extent	23/27
Understanding of simple commands	22/28
Attraction to water	21/27
Use of simple gestures (e.g., pointing)	21/27
Cooperative in (undressing)	19/28
Ability to walk	17/28
Daytime continence for urine	16/28
Undress themselves	14/28
Dress themselves	3/28
Wash themselves	0/28

Surgical Procedures in AS



- Many AS individuals undergo surgery with general anesthesia
 - Fundoplication, G-tube insertion, scoliosis repair, arthrodeses, cleft lip repair, dental procedures, tracheostomy, etc.
 - Seizure management always a concern
 - Post operative course otherwise appears relatively uncomplicated

Reported Surgical Experience in AS


- Bujok and Knapik (Ped Anes, 2004)
 - 12 yr old boy for dental procedure
 - 2 min episode of bradycardia to 40/min
 - Response to atropine was delayed
 - Advise to avoid GABAR interacting agents such as benzodiazepine and limit inhalational halogenated ethers
- Yoshitaka et al, JJDSA, 2005
 - 18 yr male for dental treatment
 - No anesthesia problem
 - Midazolam, thiopental, isoflurane (all affect GABARs)
- Masato K, et al., JJDSA, 2003
 - 2 yr old for adenoidectomy
 - anesthesia with 5% sevoflurane
 - Mild hyperthermia (temp to 37.9)

Anticonvulsant Medication Use

- About 50% use single medications
- Combinations include:
 - Valproate and clonazepam (Klonopin)
 - Valproate and topiramate (Topamax)
 - Valproate and lamotrigine (Lamictal)
 - Others
- Carbamazepine (Tegretol) used infrequently
- Avoid vigabatrin (Sabril), tiagabine (Gabitril)

Carbamazepine*	Carbitrol, Tegretol
Clonazepam	Klonopin
Ethosuximide	Zarontin
Hydantoin	Dilantin
Lamotrigine	Lamictal
Levetiracetam	Keppra
Oxcarbazepine	Trileptal
Tiagabine*	Gabitril
Topiramate	Topamax
valproic acid	Depakene
zonisamide	Zonegran

Tremors in AS



- Purkinje Cells
 - Main output from cerebellum
 - Extremely large
 - Involved in ataxia and tremor disorders
 - Abnormal function in the UBE3A Angelman mouse (Cheron et al., Neurosci, 2005)



Tremors and rigidity in AS

- Present in almost all with AS
- Varies with emotional state, time of day, fatigue, etc.
- May improve or worsen with anticonvulsants and other drugs
 - Harbord, J Clin Neurosci, 2001: levodopa responsive "Parkinsonism" in a 23 and 43 year old
 - Stecker and Meyers, Clin Neurol Neurosurg, 2003: reserpine responsive myoclonus in 32 year old
- May increase with advancing age
- No obvious relationship to seizures on EEG

**ANGELMAN SYNDROME IN THREE ADULT PATIENTS
WITH ATYPICAL PRESENTATION
AND SEVERE NEUROLOGICAL COMPLICATIONS**

BY G.J.C.M. VAN BUCGENHOUT¹*, M.J. DECHERMAKIAN², P. THOM¹,
J.C.M. TRIMBURZEN¹, B.G.A. HANDE² AND J.P. PATTY³



- PH- 52 yrs
 - Institutionalized since age 7
 - Slow and rigid boy with slow tremors
 - MRI cerebellar atrophy
 - 27 clumsy walking, sever spasticity, clonus
 - Carbamazepin, valproate, baclofen
 - Non-ambulatory at 52



- KS- 32 yrs
 - Seizures at 2 years, never ambulatory
 - Institutionalized since age 6?
 - Severe quadriplegia with contractures, scoliosis age 13 yrs
 - Age 29: change in sleep patterns, fell asleep during the day
 - Medication history not noted



- DVO- 49 yrs
 - institutionalized age 28?
 - Semi-ambulatory at 49
 - Lower limb rigidity and spastic walking
 - Day/night sleep patterns disturbed, fell asleep in am
 - History of seizures as child, recurrence at age 29
 - Medication history not noted

Life Span in AS

- Several case reports of relatively long life
- No reports of deaths related to:
 - Diabetes
 - Cancer
 - Autoimmune disorder
 - Neurodegenerative disease
- Life span expected to be decreased more when non-ambulatory and tube fed
- Seizures and cognitive impairment alone do not appear to be obvious risk factors for decreased life span