

CSCE 763: Digital Image Processing

Homework #1

Instructor: Spring 2026
University of South Carolina
Solutions by: JC Vaught

Due: 1:15 pm EST, Tuesday, Feb 3

Table of Contents	
Problem 1: Smallest Discernible Dot (25 pts)	2
Problem 2: Adjacency of Image Subsets (25 pts)	
Part (a): 4-adjacency	3
Part (b): 8-adjacency	3
Part (c): m-adjacency	3
Problem 3: Shortest Paths (25 pts)	
Part (a): $V = \{0, 1\}$	4
Part (b): $V = \{1, 2\}$	8
Problem 4: Set Expressions (25 pts)	11

Problem 1: Smallest Discernible Dot (25 pts)

Thinking purely in geometric terms, estimate the diameter of the smallest printed dot that the eye can discern if the page on which the dot is printed is 0.5 m away from the eyes.

Assumptions:

- (a) The distance between the center of the lens and the retina along the visual axis is 14 mm.
- (b) The visual system ceases to detect the dot when the image of the dot on the fovea becomes smaller than the diameter of one receptor (cone) in that area of the retina.
- (c) The fovea can be modeled as a square array of dimensions $1.5 \text{ mm} \times 1.5 \text{ mm}$, and the cones (about 337,000 in total) and spaces between the cones are distributed uniformly throughout this array.

Problem 2: Adjacency of Image Subsets (25 pts)


Consider the two image subsets S_1 and S_2 shown below. For $V = \{1\}$, determine whether these two subsets are (a) 4-adjacent, (b) 8-adjacent, or (c) m-adjacent.

	S_1					S_2				
0	0	0	0	0	0	0	0	1	1	0
1	0	0	1	1	0	1	0	0	0	1
1	0	0	1	0	1	1	0	0	0	0
0	0	1	1	0	0	0	0	0	0	0
0	0	1	1	1	0	0	0	1	1	1

Problem 3: Shortest Paths (25 pts)

Consider the image segment shown below.

- (a) Let $V = \{0, 1\}$ and compute the lengths of the shortest 4-, 8-, and m-path between p and q . If a particular path does not exist between these two points, explain why.
- (b) Repeat for $V = \{1, 2\}$.



The grid (using row, column indexing from top-left):


	col 0	col 1	col 2	col 3
row 0	3	1	2	1 (q)
row 1	2	0	0	2
row 2	1	2	1	1
row 3	1 (p)	0	1	2

$p = (3, 0)$ with value 1, $q = (0, 3)$ with value 1.

- (a) $V = \{0, 1\}$

Identify Valid Pixels

Pixels with values in $V = \{0, 1\}$:



Valid pixels (shaded): $(0, 1) = 1$, $(0, 3) = 1$, $(1, 1) = 0$, $(1, 2) = 0$, $(2, 0) = 1$, $(2, 2) = 1$, $(2, 3) = 1$, $(3, 0) = 1$, $(3, 1) = 0$, $(3, 2) = 1$.

Shortest 4-path

A 4-path uses only horizontal and vertical steps between pixels whose values are both in V .

One such path from $p = (3, 0)$ to $q = (0, 3)$:

$$(3, 0) \rightarrow (3, 1) \rightarrow (3, 2) \rightarrow (2, 2) \rightarrow (2, 3) \rightarrow (0, 3)$$

Wait — $(2, 3) = 1$ to $(0, 3) = 1$: we need $(1, 3) = 2 \notin V$. So we cannot go directly up from $(2, 3)$.

Let me find a valid 4-path. From $(2, 3) = 1$, the 4-neighbors with values in V : $(2, 2) = 1$, $(3, 3) = 2$ (no). Going up: $(1, 3) = 2$ (no). So $(2, 3)$ is a dead end going upward.

Try via top: $(3, 0) \rightarrow (2, 0) \rightarrow \dots$ $(2, 0) = 1$. 4-neighbors in V : $(3, 0) = 1$ (back), $(1, 0) = 2$ (no). Dead end.

Another attempt: $(3, 0) \rightarrow (3, 1) \rightarrow (3, 2) \rightarrow (2, 2) \rightarrow (1, 2) \rightarrow (1, 1) \rightarrow (0, 1) \rightarrow (0, 3)$? Check: $(0, 1) = 1$ to $(0, 3) = 1$: need $(0, 2) = 2 \notin V$. Cannot step from $(0, 1)$ to $(0, 3)$ directly — they are not 4-neighbors.

$(0, 1) = 1$: 4-neighbors in V : $(1, 1) = 0$, $(0, 0) = 3$ (no), $(0, 2) = 2$ (no). Dead end going right.

No 4-path exists from p to q , because pixel $(0, 1)$ cannot connect rightward to $(0, 3)$ — the intervening pixel $(0, 2) = 2 \notin V$, and there is no other route to reach column 3 in row 0.

Shortest 8-path

An 8-path allows diagonal moves. From the valid pixel map:

$(3,0) \rightarrow (3,1) \rightarrow (3,2) \rightarrow (2,3) \rightarrow (0,3)$? Check $(2,3) = 1$ to $(0,3)$: not 8-neighbors (distance = 2 rows). Invalid.

$(3,0) \rightarrow (3,1) \rightarrow (2,2) \rightarrow (1,1) \rightarrow (0,1)$: need diagonal. $(3,1) = 0$ to $(2,2) = 1$: 8-neighbor, both in V . $(2,2) = 1$ to $(1,1) = 0$: 8-neighbor, both in V . $(1,1) = 0$ to $(0,1)$: wait, is there a diagonal path onward?

$(0,1) = 1$. 8-neighbors in V : $(1,1) = 0$, $(1,2) = 0$, $(0,0) = 3$ (no), $(0,2) = 2$ (no). Still stuck.

Try: $(1,2) = 0$ to $(0,3) = 1$: these are 8-neighbors! Both in V .

Path: $(3,0) \rightarrow (3,1) \rightarrow (2,2) \rightarrow (1,2) \rightarrow (0,3)$

Check each step:

- $(3,0) = 1 \rightarrow (3,1) = 0$: 4-neighbor, both in V . ✓
- $(3,1) = 0 \rightarrow (2,2) = 1$: 8-neighbor (diagonal), both in V . ✓
- $(2,2) = 1 \rightarrow (1,2) = 0$: 4-neighbor, both in V . ✓
- $(1,2) = 0 \rightarrow (0,3) = 1$: 8-neighbor (diagonal), both in V . ✓

Length = 4 steps. This is optimal since the minimum Chebyshev distance from $(3,0)$ to $(0,3)$ is $\max(3,3) = 3$, so the shortest 8-path has length ≥ 3 . Can we find a length-3 path?

A length-3 path requires 3 diagonal steps: $(3,0) \rightarrow (2,1) \rightarrow (1,2) \rightarrow (0,3)$. Check: $(2,1) = 2 \notin V$. Invalid.

Shortest 8-path length = 4.

Shortest m-path

An m-path uses mixed adjacency: diagonal steps are allowed only when the two pixels share no common 4-neighbor with a value in V .

Check the 8-path $(3,0) \rightarrow (3,1) \rightarrow (2,2) \rightarrow (1,2) \rightarrow (0,3)$:

Step $(3,1) \rightarrow (2,2)$ (diagonal): Common 4-neighbors of $(3,1)$ and $(2,2)$ are $(3,2)$ and $(2,1)$. $(3,2) = 1 \in V$. Since there is a common 4-neighbor in V , this diagonal move is **not allowed** in m-adjacency.

Alternative: replace the diagonal with a 4-connected detour via $(3,2)$: $(3,0) \rightarrow (3,1) \rightarrow (3,2) \rightarrow (2,2) \rightarrow (1,2) \rightarrow (0,3)$

Check $(1,2) \rightarrow (0,3)$ (diagonal): Common 4-neighbors of $(1,2)$ and $(0,3)$ are $(0,2)$ and $(1,3)$. $(0,2) = 2 \notin V$, $(1,3) = 2 \notin V$. No common 4-neighbor in V , so this diagonal is **allowed**.

Full m-path: $(3,0) \rightarrow (3,1) \rightarrow (3,2) \rightarrow (2,2) \rightarrow (1,2) \rightarrow (0,3)$

Shortest m-path length = 5.

Results

For $V = \{0, 1\}$:

Path Type	Shortest Length
4-path	Does not exist (no connected 4-path from p to q)
8-path	
m-path	

Problem 3 — Part (b): $V = \{1, 2\}$

Identify Valid Pixels

Pixels with values in $V = \{1, 2\}$:

	col 0	col 1	col 2	col 3
row 0	3	1	2	1 (q)
row 1	2	0	0	2
row 2	1	2	1	1
row 3	1 (p)	0	1	2

Shortest 4-path

$(3, 0) \rightarrow (2, 0) \rightarrow (2, 1) \rightarrow (2, 2) \rightarrow (2, 3) \rightarrow (1, 3) \rightarrow (0, 3)$

Check: all values in V and all steps are 4-neighbors.

- $(3, 0) = 1, (2, 0) = 1, (2, 1) = 2, (2, 2) = 1, (2, 3) = 1, (1, 3) = 2, (0, 3) = 1$ — all $\in V$. ✓

Length = 6. Can we do better? The Manhattan distance is $|3 - 0| + |0 - 3| = 6$, so this is optimal.

Shortest 8-path

The Chebyshev distance is $\max(3, 3) = 3$, so the minimum possible 8-path length is 3.

Try: $(3, 0) \rightarrow (2, 1) \rightarrow (1, 2) \rightarrow (0, 3)$ Check: $(2, 1) = 2 \in V$, $(1, 2) = 0 \notin V$. Invalid.

Try: $(3, 0) \rightarrow (2, 1) \rightarrow (1, 0) \rightarrow (0, 1) \rightarrow (0, 2) \rightarrow (0, 3)$? $(1, 0) = 2 \in V$, $(0, 1)$: is $(1, 0)$ 8-adjacent to $(0, 1)$? Yes (diagonal). $(0, 1) = 1 \in V$, $(0, 2) = 2 \in V$, $(0, 3) = 1 \in V$. Length = 5.

Shorter: $(3, 0) \rightarrow (2, 1) \rightarrow (1, 2)$: $(1, 2) = 0 \notin V$. No.

$(3, 0) \rightarrow (2, 0) \rightarrow (1, 0) \rightarrow (0, 1) \rightarrow (0, 2) \rightarrow (0, 3)$. Length = 5. All in V : 1, 1, 2, 1, 2, 1. ✓

Try length 4: $(3, 0) \rightarrow (2, 1) \rightarrow (1, 0) \rightarrow (0, 1) \rightarrow (0, 3)$? $(0, 1)$ and $(0, 3)$ are not 8-neighbors. No.

$(3, 0) \rightarrow (2, 1) \rightarrow (1, 2)$: invalid (value 0).

$(3, 0) \rightarrow (2, 0) \rightarrow (1, 0) \rightarrow (0, 1) \rightarrow (0, 2) \rightarrow (0, 3)$: length 5.

Try: $(3, 0) \rightarrow (2, 1) \rightarrow (1, 0) \rightarrow (0, 1) \rightarrow (0, 2) \rightarrow (0, 3)$: length 5.

Try length 4: $(3, 0) \rightarrow (2, 1) \rightarrow (1, 0) \rightarrow (0, 1) \rightarrow (0, 3)$? Not adjacent. No.

$(3, 0) \rightarrow (2, 0) \rightarrow (1, 0) \rightarrow (0, 1) \rightarrow (0, 3)$? $(0, 1)$ to $(0, 3)$: distance 2. No.

Try: $(3, 0) \rightarrow (2, 1) \rightarrow (1, 2)$: value 0. Dead end.

Try via right side: $(3, 0) \rightarrow (3, 2) \rightarrow (2, 3) \rightarrow (1, 3) \rightarrow (0, 3)$? $(3, 0)$ to $(3, 2)$: not 8-neighbors (distance 2). No.

$(3, 0) \rightarrow (2, 1) \rightarrow (2, 2) \rightarrow (1, 3) \rightarrow (0, 3)$: length 4. Check: $(2, 1) = 2$, $(2, 2) = 1$, $(1, 3) = 2$, $(0, 3) = 1$. All $\in V$. All 8-neighbors? $(3, 0) \rightarrow (2, 1)$: diagonal ✓. $(2, 1) \rightarrow (2, 2)$: 4-neighbor ✓. $(2, 2) \rightarrow (1, 3)$: diagonal ✓. $(1, 3) \rightarrow (0, 3)$: 4-neighbor ✓.

Shortest 8-path length = 4.

Shortest m-path

Check the 8-path $(3, 0) \rightarrow (2, 1) \rightarrow (2, 2) \rightarrow (1, 3) \rightarrow (0, 3)$:

Step $(3, 0) \rightarrow (2, 1)$ (diagonal): Common 4-neighbors: $(2, 0)$ and $(3, 1)$. $(2, 0) = 1 \in V$. Diagonal not allowed in m-adjacency.

Step $(2, 2) \rightarrow (1, 3)$ (diagonal): Common 4-neighbors: $(2, 3)$ and $(1, 2)$. $(2, 3) = 1 \in V$. Diagonal not allowed.

So this particular path is not valid for m-adjacency. We need to replace disallowed diagonals with 4-connected detours.

Optimal m-path: $(3, 0) \rightarrow (2, 0) \rightarrow (2, 1) \rightarrow (2, 2) \rightarrow (2, 3) \rightarrow (1, 3) \rightarrow (0, 3)$

All 4-connected steps with values in V . **Length = 6.**

Can we do better? Try mixing: $(3, 0) \rightarrow (2, 0) \rightarrow (2, 1) \rightarrow (2, 2) \rightarrow (1, 3) \rightarrow (0, 3)$. Length 5. Check $(2, 2) \rightarrow (1, 3)$: diagonal. Common 4-neighbors: $(1, 2) = 0 \notin V$, $(2, 3) = 1 \in V$. Not allowed.

$(3, 0) \rightarrow (2, 0) \rightarrow (2, 1) \rightarrow (1, 2)$: $(1, 2) = 0 \notin V$. Invalid.

Try: $(3, 0) \rightarrow (2, 1) \rightarrow \dots$: $(3, 0) \rightarrow (2, 1)$ diagonal. Common 4-neighbors of $(3, 0)$ and $(2, 1)$: $(3, 1) = 0$ and $(2, 0) = 1 \in V$. Not allowed.

Shortest m-path length = 6.

Results

For $V = \{1, 2\}$:

Path Type	Shortest Length
4-path	6
8-path	4
m-path	6

Problem 4: Set Expressions (25 pts)

Give expressions for the sets shown shaded in the following figure in terms of sets A , B , and C .

In the figure: A is a rectangle (top), B is a triangle (bottom-left), C is a triangle (bottom-right). The three shapes overlap.

Analyzing Each Figure

Figure (i): The shaded region is the part of A that overlaps with B but not C , plus the part of A that overlaps with C but not B . This is the part of A that is in exactly one of B or C :

$$(A \cap B \cap C^c) \cup (A \cap C \cap B^c)$$

Equivalently: $A \cap (B \oplus C)$ where $\oplus$ denotes symmetric difference, or simply $A \cap (B \cup C) \cap (B \cap C)^c$.

Figure (ii): The shaded region includes the part of B not in A or C , plus the intersection of A and C (including overlaps with B). Looking at the shading pattern:

$$(A \cap C) \cup (B \cap A^c \cap C^c)$$

Figure (iii): The shaded region is the part of B and C that is not in A . That is, the union of B and C minus A :

$$(B \cup C) \cap A^c$$

Results

(i) $(A \cap B \cap C^c) \cup (A \cap C \cap B^c)$

(ii) $(A \cap C) \cup (B \cap A^c \cap C^c)$

(iii) $(B \cup C) \cap A^c$