

4. Linear Regression and Randomized Experiments

ISS5096 || ECI

Jaewon (“Jay-one”) Yoo


National Tsing Hua University

Roadmap

1. Regression with no covariates
2. Linear regression with covariates
3. Cluster randomized experiments

Where are we? Where are we going?

- So far: analysis of experiments with Fisher's and Neyman's approaches.
 - Neyman: Unbiased estimators, (conservative) variances.
 - Fisher: exact test of the sharp null.
- Today: how does the workhorse estimator, OLS, fit into this story?
- Why would we consider using regression?
 - **Simplicity:** known tool that is already very common.
 - **Increased precision:** we may want to add covariates for more precise effect estimates.



Source: *Chapter 13 of The Effect (Textbook 2)* by Nick Huntington-Klein

1/ Regression with no covariates

Analysing Experiments with Regression?

- Q: Under complete randomization, can we use OLS to estimate ATEs
 - Literally, just $\text{lm}(y \sim d)$?
- Recall that the OLS estimator solves the least squares problem:

$$(\hat{\tau}_{\text{ols}}, \hat{\alpha}_{\text{ols}}) = \arg \min_{\tau, \alpha} \sum_{i=1}^n (Y_i - \alpha - \tau D_i)^2 \quad (1)$$

- The coefficient on a binary r.v. is mechanically the diff. in means:

$$\hat{\tau}_{\text{ols}} = \bar{Y}_1 - \bar{Y}_0 = \hat{\tau}_{\text{diff}} \quad (2)$$

- Standard Neyman analysis for unbiasedness, sampling variance.
- Generalized to discrete treatments with > 2 levels.

Justifying the Linear Model

- Mechanically the same, but can we justify the linear model itself?
 - Key assumptions: **linearity** and **mean independence of errors**.
- Some simple manipulation of the consistency assumption:

$$\begin{aligned}Y_i &= D_i Y_i(1) + (1 - D_i) Y_i(0) \\&= Y_i(0) + D_i \{Y_i(1) - Y_i(0)\} \\&= Y_i(0) + D_i \tau_i \\&= \mathbb{E}[Y_i(0)] + D_i \tau + \{Y_i(0) - \mathbb{E}[Y_i(0)]\} + D_i(\tau_i - \tau) \\&= \alpha + D_i \tau + \varepsilon_i\end{aligned}$$

- “Linear” functional form fully justified by consistency alone with:
 - Intercept $\alpha = \mathbb{E}[Y_i(0)]$ is the average control outcome.
 - Slope $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$ is the PATE.
 - Error is deviation for control PO + treatment effect heterogeneity.

Mean Independent Errors

$$\varepsilon_i = (Y_i(0) - \mathbb{E}[Y_i(0)]) + D_i \cdot (\tau_i - \tau)$$

- What about mean independence: $\mathbb{E}[\varepsilon_i | D_i] = 0$?
- Using randomization, we know D_i is independent of $Y_i(1)$, $Y_i(0)$, so:

$$\begin{aligned}\mathbb{E}[\varepsilon_i | D_i] &= \mathbb{E}[\{Y_i(0) - \mathbb{E}[Y_i(0)]\} + D_i \cdot (\tau_i - \tau) | D_i] \\&= \mathbb{E}[Y_i(0) | D_i] - \mathbb{E}[Y_i(0)] + D_i (\mathbb{E}[\tau_i | D_i] - \tau) \\&= \mathbb{E}[Y_i(0)] - \mathbb{E}[Y_i(0)] + D_i \underbrace{(\mathbb{E}[\tau_i] - \tau)}_{\tau = \mathbb{E}[\tau_i]} \\&= 0\end{aligned}$$

- Randomization + consistency $\rightsquigarrow$ linear model.
 - Does not imply homoskedasticity or normal errors, though!

Homoskedasticity

- Software default assumption: **Homoskedasticity**

$$\mathbb{V}[\varepsilon_i \mid \mathbf{D}] = \sigma^2, \quad \forall i$$

- But in general, based on previous error definition:

$$\mathbb{V}[\varepsilon_i \mid \mathbf{D}] = \mathbb{V}[\varepsilon_i \mid D_i] = D_i \sigma_1^2 + (1 - D_i) \sigma_0^2$$

- $\rightsquigarrow$ homoskedasticity true when $\sigma_1^2 = \mathbb{V}[Y_i(1)] = \mathbb{V}[Y_i(0)] = \sigma_0^2$
 - True under constant treatment effects!
- Under homoskedasticity, variance of the OLS estimator is:

$$\mathbb{V}[\hat{\tau}_{\text{ols}} \mid \mathbf{D}] = \frac{\sigma^2}{\sum_{i=1}^n (D_i - \bar{D})^2}$$

Variance Estimation

- “Standard” variance estimator under homoskedasticity:

$$\widehat{V}_{\text{const}} = \frac{\frac{1}{n-2} \sum_{i=1}^n \widehat{\varepsilon}_i^2}{\sum_{i=1}^n (D_i - \bar{D})^2} = \frac{\frac{1}{n-2} \sum_{i=1}^n (Y_i - \widehat{\alpha}_{\text{ols}} - \widehat{\tau}_{\text{ols}} D_i)^2}{\sum_{i=1}^n (D_i - \bar{D})^2}$$

- We can rewrite this as a function of the **pooled** variance $\widehat{\sigma}_{Y|D}^2$:

$$\widehat{V}_{\text{const}} = \widehat{\sigma}_{Y|D}^2 \left(\frac{1}{n_0} + \frac{1}{n_1} \right)$$

$$\widehat{\sigma}_{Y|D}^2 = \frac{1}{n-2} \left(\sum_{i:D_i=0} (Y_i - \bar{Y}_0)^2 + \sum_{i:D_i=1} (Y_i - \bar{Y}_1)^2 \right)$$

- **Inconsistent:** $\widehat{V}_{\text{const}} - \mathbb{V}[\widehat{\tau}] \xrightarrow{p} c \neq 0$ unless
 - Homoskedasticity holds: $\sigma_1^2 = \sigma_0^2$
 - Design is balanced: $n_1 = n_0$

Variance Estimation

- Can we use “standard” variance estimator: $\mathbb{V}[\varepsilon_i | \mathbf{D}] = \sigma^2$

- Inconsistent: $\widehat{\mathbb{V}}_{\text{const}} - \mathbb{V}[\widehat{\tau}] \xrightarrow{P} c \neq 0$

- Bias:

$$\begin{aligned} & \mathbb{E}(\widehat{\mathbb{V}}_{\text{const}}) - \mathbb{V}[\widehat{\tau}] \\ &= \underbrace{\mathbb{E}\left(\frac{\frac{1}{n-2} \sum_{i=1}^n \hat{\varepsilon}_i^2}{\sum_{i=1}^n (D_i - \overline{D})^2}\right)}_{\text{under const. effect assumption}} - \underbrace{\left(\frac{\sigma_1^2}{n_1} + \frac{\sigma_0^2}{n_0}\right)}_{\text{true variance}} \\ &= \frac{(n_1 - n_0)(n - 1)}{n_1 n_0 (n - 2)} (\sigma_1^2 - \sigma_0^2) \neq 0 \end{aligned}$$

- Unless:
 - Homoskedasticity holds: $\sigma_1^2 = \sigma_0^2$
 - Constant effect: $Y_i(1) - Y_i(0) = \text{const.}$
 - $\mathbb{V}[Y_i(1)] = \mathbb{V}[Y_i(0) + \text{const.}] = \mathbb{V}[Y_i(0)]$
 - Design is balanced: $n_1 = n_0$

Robust SEs

- Use robust variance estimator!
- Eicker-Huber-White (EHW) estimator: consistent for $\mathbb{V}(\hat{\tau}_{\text{diff}})$

$$\hat{\mathbb{V}}_{\text{EHW}} = \frac{\tilde{\sigma}_1^2}{n_1} + \frac{\tilde{\sigma}_0^2}{n_0}, \quad \text{where} \quad \tilde{\sigma}_d^2 = \frac{1}{n_d} \sum_{i:D_i=d} (Y_i - \bar{Y}_d)^2$$

- HC2 estimator:

$$\hat{\mathbb{V}}_{\text{HC2}} = \frac{\hat{\sigma}_0^2}{n_0} + \frac{\hat{\sigma}_1^2}{n_1}, \quad \text{where} \quad \hat{\sigma}_d^2 = \frac{1}{n_d - 1} \sum_{i:D_i=d} (Y_i - \bar{Y}_d)^2$$

- Samii & Aronow (2012): HC2 is exactly the Neyman variance estimator.
- $\rightsquigarrow$ **Simple OLS + HC2 = unbiased point and variance estimator.**

In R

```
your_fitted_model <- lm(your_formula, data = your_data)
sandwich::vcovHC(your_fitted_model, type = 'HC2')
# Or
estimatr::lm_robust(your_formula, your_data, se_type = 'HC2')
```

Application in R Using (Sim) AI Exp. Data

```
1 > AI_data <- as_tibble(read.csv(url("https://bit.ly/3FHsusw"))); AI_data
2 # A tibble: 500 x 8
3   treat_ind test_outcome_pre test_outcome_post student_age student_gender tutor_age
4   <int>      <int>          <int>          <int>      <int>      <int>
5 1         1          0            1           11         0         27
6 2         1          0            1            6         1         45
7 3         0          1            1           15         1         54
8 4         0          1            1            6         1         50
9 5         0          1            0            8         1         64
10 6         1          1            1           15         1         43
11 7         1          0            1            7         1         47
12 8         1          1            1           16         0         49
13 9         0          0            1           12         1         42
14 10        1          1            1           11         1         28
15 # ... 490 more rows
16 # ... 2 more variables: years_of_experience <int>, education_level <chr>
17 # Use `print(n = ...)` to see more rows
18
19 > lm1 <- lm(test_outcome_post ~ treat_ind, data = AI_data)
20 > vcovM <- sandwich::vcovHC(lm1, type = 'HC2')
21 > sqrt(vcovM[1,1])
22 [1] 0.02955412
23 > sqrt(vcovM[2,2]) # sqrt(diag(vcovM))
24 [1] 0.04209018
25
26 > # Or
27 > estimatr::lm_robust(test_outcome_post ~ treat_ind, AI_data, se_type = 'HC2')
28       Estimate Std. Error  t value    Pr(>|t|)    CI Lower CI Upper  DF
29 (Intercept) 0.65134100 0.02955412 22.038927 1.233662e-75 0.59327487 0.7094071 498
30 treat_ind    0.03903557 0.04209018  0.927427 3.541541e-01 -0.04366065 0.1217318 498
```

2/ Linear regression with covariates

Adding Covariates

- What if we add covariates to our regression model?

$$(\hat{\tau}_{\text{adj}}, \hat{\alpha}_{\text{adj}}, \hat{\beta}_{\text{adj}}) = \arg \min_{\tau, \alpha, \beta} \sum_{i=1}^n (Y_i - \alpha - \tau D_i - \tilde{\mathbf{X}}_i' \beta)^2$$

- $\tilde{\mathbf{X}}_i = \mathbf{X}_i - \bar{\mathbf{X}}$ are **centered** covariates for notational ease.
- Why might we do this? To increase **precision** of our estimates.
 - We hope $\mathbb{V}[\hat{\tau}_{\text{adj}}] < \mathbb{V}[\hat{\tau}_{\text{diff}}]$ so we have smaller CIs, more powerful tests.
 - Intuition: less residual variation in Y_i after accounting for $\mathbf{X}_i$
- Questions:
 - Is $\hat{\tau}$ still unbiased? Consistent?
 - Should we expect an increase in precision?
 - Controversial! Freedman (2008) “Randomization does not justify the regression model”

OLS is biased, but consistent (Freedman, 2008. Adv. in Appl. Math)

- Agnostic approach: don't assume correctness of the linear model.
 - $\mathbf{X}'_i\beta$ could be badly misspecified.
 - No assumptions about homoskedasticity.
- Under minimal assumptions, OLS is consistent for the linear projection

$$(\hat{\tau}_{\text{adj}}, \hat{\alpha}_{\text{adj}}, \hat{\beta}_{\text{adj}}) \xrightarrow{p} (\tau_0, \alpha_0, \beta_0) = \arg \min_{(\tau, \alpha, \beta)} \mathbb{E} [(Y_i - \alpha - \tau D_i - \tilde{\mathbf{X}}'_i \beta)^2]$$

- $\hat{\tau}_{\text{adj}}$ now **biased** for τ though bias should be small.
- But $\hat{\tau}_{\text{adj}}$ is **consistent**.
 - Intuition: Since $D_i \perp\!\!\!\perp \mathbf{X}_i$, including $\tilde{\mathbf{X}}_i$ won't (asymptotically) affect coefficient on D_i .
- Freedman (2008) also shows bias vanishes as n grows under finite-sample (randomization) inference.

Variance of Adjustment Estimator

- Complete randomization + single, mean-zero covariate X_i
 - Generalizes easily to more covariates.
 - Let $\sigma_{0x} = \text{cov}(Y_i(0), X_i)$ and $\sigma_{1x} = \text{cov}(Y_i(1), X_i)$.
 - Probability of treatment $p = n_1/n$
- Freedman (2008) derived gains from adjusting for X_i using OLS:

$$\mathbb{V}[\widehat{\tau}_{\text{diff}}] - \mathbb{V}[\widehat{\tau}_{\text{adj}}] = \frac{\sigma_{0x}\{\sigma_{0x} + 2(1 - 2p)\sigma_{1x}\}}{np(1 - p)}$$

- Will adjustment decrease the sampling variance?
 - If design is balanced, $p = 1/2$, then adjustment always helps.
 - Design imbalance could lead to adjustment hurting.
- Estimation: EHW (and HC2) robust variance estimators are consistent or asymptotically conservative for $\mathbb{V}[\widehat{\tau}_{\text{adj}}]$

Regression with Full Interactions

- OLS estimator from fully interacted model, $\widehat{\tau}_{\text{inter}}$:

$$Y_i = \alpha + \tau D_i + \widetilde{\mathbf{X}}_i' \beta + D_i \widetilde{\mathbf{X}}_i' \gamma + \varepsilon_i$$

- Equivalent to running separate Y_i on $\widetilde{\mathbf{X}}_i$ in each D_i
- As with non-interacted model, $\widehat{\tau}_{\text{inter}}$ is consistent for τ and asymptotically normal.
- Lin (2013): **fully interacted model will never hurt precision asymptotically.**
 - Freedman critique was right, but Lin shows an easy way to resolve.
- EHW (and HC2) robust variance estimators are consistent or asymptotically conservative.

Linear Regression with Covariates

In R

```
# Step 1: Compute centered covariates
your_data$Xtilde <- NULL

# Step 2: Write down your formula
your_formula <- NULL

# Step 3: Fit the model using lm() or estimatr::lm_robust()
your_fitted_model <- lm(your_formula, data = your_data)

# Step 4: Compute robust standard errors (skip if you used lm_robust)
your_vcov <- sandwich::vcovHC(your_fitted_model, type = 'HC2')

# Step 5: Check the point and se estimate of your coefficients
# (look for tau hat!)
est <- cbind("coef" = your_fitted_model$coef,
             "se" = sqrt(diag(your_vcov)))
```

Example Code

```
1 > AI_data <- AI_data |>
2   mutate(Xtilde = student_age - mean(student_age)) |>
3   select(treat_ind, test_outcome_post, Xtilde); head(AI_data,3)
4 # A tibble: 3 x 3
5   treat_ind test_outcome_post Xtilde
6     <int>         <int>    <dbl>
7 1         1             1 -0.566
8 2         1             1 -5.57
9 3         0             1  3.43
10
11 > estimatr::lm_robust(test_outcome_post ~ treat_ind * Xtilde, data = AI_data)
12               Estimate Std. Error   t value    Pr(>|t|)    CI Lower    CI Upper   DF
13 (Intercept)    0.649860304 0.029645673 21.9209157 5.557487e-75 0.591613722 0.708106886 496
14 treat_ind      0.038364943 0.042245496  0.9081428 3.642438e-01 -0.044637245 0.121367131 496
15 Xtilde         0.008351571 0.008581795  0.9731730 3.309417e-01 -0.008509581 0.025212723 496
16 treat_ind:Xtilde -0.019462905 0.012003266 -1.6214674 1.055529e-01 -0.043046422 0.004120612 496
17
18 > your_fitted_model <- lm(test_outcome_post ~ treat_ind * Xtilde, data = AI_data)
19 > vcovM_adj <- sandwich::vcovHC(your_fitted_model, type = 'HC2'); sqrt(diag(vcovM_adj))
20               (Intercept)      treat_ind      Xtilde treat_ind:Xtilde
21               0.029645673      0.042245496      0.008581795      0.012003266
22
23 > est <- cbind("coef" = your_fitted_model$coef,
24              "se" = sqrt(diag(your_vcov))); est
25               coef      se
26 (Intercept)    0.649860304 0.029645673
27 treat_ind      0.038364943 0.042245496
28 Xtilde         0.008351571 0.008581795
29 treat_ind:Xtilde -0.019462905 0.012003266
```

Summarizing Regression

- Regression with no covariates = standard Neyman analysis.
- Regression with (uninteracted) covariates:
 - Consistent for SATE/PATE.
 - Usually will help with precision, but can hurt.
- Regression with interacted covariates:
 - Consistent for SATE/PATE.
 - Asymptotically will never hurt precision.
- Always use robust/HC2 variance estimators unless you have good reasons.

3/ Cluster randomized experiments


Cf. Angrist & Pischke (2008), *Mostly Harmless Econometrics*, Ch. 8

Clustering Treatments

- Treatment often allocated at a higher level than the data.
 - Counties are treated, but we have individual-level data.
 - Classrooms are treated, but we have student data.
- Has considerable benefits:
 - Often cheaper/easier to implement than individual assignment.
 - Allows for interference within clusters without bias.
- But lots of confusion about how to analyze.
 - More valuable to add more individuals or clusters?
 - What to do with individual-level covariates?


/// : treated

□ : block



/// : treated

□ : cluster



Cluster Randomized Trials

- Setup:
 - Clusters: $k \in \{1, \dots, K\}$
 - Randomly choose K_1 treatment clusters, K_0 control.
 - Each cluster has units $i \in \{1, \dots, m_k\}$ with $\sum_{k=1}^K m_k = n$
 - Treatment assignment at cluster level: $D_{ik} = D_k$
 - Potential outcomes $Y_{ik}(d)$
- Random assignment at the cluster level: $\{Y_{ik}(1), Y_{ik}(0)\} \perp\!\!\!\perp D_j$.
- Quantity of interest still at individual level:

$$\text{SATE} = \frac{1}{n} \sum_{k=1}^K \sum_{i=1}^{m_k} \{Y_{ik}(1) - Y_{ik}(0)\}$$

Analysis of Clustered Experiments

- Simple setting: all clusters have the same size $m_k = m$ for all k .
- Simple difference in means is unbiased:

$$\begin{aligned}\widehat{\tau}_{\text{cl}} &= \frac{1}{mK_1} \sum_{k=1}^K \sum_{i=1}^{m_k} D_k Y_{ik} - \frac{1}{mK_0} \sum_{k=1}^K \sum_{i=1}^{m_k} (1 - D_k) Y_{ik} \\ &= \frac{1}{K_1} \sum_{k=1}^K D_k \bar{Y}_k - \frac{1}{K_0} \sum_{k=1}^K (1 - D_k) \bar{Y}_k\end{aligned}$$

- $\bar{Y}_k$ is the cluster average: $\frac{1}{m} \sum_{i=1}^m Y_{ik}$
 - Unbiasedness follows from Neyman-style analysis at cluster level.
 - Estimator is biased, but consistent (in K) if cluster size varies.
- Neyman-style conservative variance:

$$\mathbb{V}[\widehat{\tau}_{\text{cl}} \mid \mathbf{D}] \leq \frac{\mathbb{V}[\bar{Y}_k(1)]}{K_1} + \frac{\mathbb{V}[\bar{Y}_k(0)]}{K_0} \quad \text{where } \bar{Y}_k(d) = \frac{1}{m} \sum_{i=1}^m Y_{ik}(d)$$

Cluster Robust Standard Errors

- What if we want to use OLS at individual level?
 - Adding individual and unit controls.
- Use **cluster-robust variance estimator** (CRVE) for OLS.
 - Sandwich-type estimator that allows residuals to correlate within clusters.
 - Consistent as the number of clusters grows.
- **Cluster at the treatment assignment level** (no higher or lower).
- Vanilla CRVE is biased, Bell & McCaffrey proposed CR2 adjustment similar to HC2 (usually preferable).

Cluster Randomized Trials: Analysis

In R

```
your_formula <- as.formula("outcome ~ treat + x_tilde1 + x_tilde2")

your_data <- data.frame(outcome, treat,
                        x_tilde1, x_tilde2,
                        cluster)

your_fitted_model <- estimatr::lm_robust(your_formula, data = your_data,
                                       clusters = cluster,
                                       se_type = "CR2")

??estimatr::lm_robust # Check more options for se_type

# Or
your_model <- lm(your_formula, data = your_data)
your_vcov <- clubSandwich::vcovCR(your_model, cluster = your_data$cluster,
                                type = "CR2")
```

Onto the presentations & discussions!

Contact Information:

jaewon.yoo@iss.nthu.edu.tw

<https://j1yoo.github.io/>



Appendix

EHW/Sandwich Variance Estimator

- Eicker-Huber-White (EHW) robust/sandwich variance estimator:

$$\underbrace{\widehat{\mathbf{V}}_{\text{EHW}}}_{\text{sandwich}} = \underbrace{\left(\sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i' \right)^{-1}}_{\text{bread}} \underbrace{\left(\sum_{i=1}^n \hat{\varepsilon}_i^2 \mathbf{x}_i \mathbf{x}_i' \right)}_{\text{meat}} \underbrace{\left(\sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i' \right)^{-1}}_{\text{bread}}$$
$$= (\mathbb{X}'\mathbb{X})^{-1} \left(\sum_{i=1}^n \hat{\varepsilon}_i^2 \mathbf{x}_i \mathbf{x}_i' \right) (\mathbb{X}'\mathbb{X})^{-1} \quad \text{where } \mathbb{X} = [\mathbf{1} \quad \mathbf{D}]$$

HC2 Variance Estimator

- HC2 normalizes residuals by the leverage, h_{ii} :

$$\widehat{\mathbb{V}}_{\text{HC2}} = (\mathbb{X}'\mathbb{X})^{-1} \left(\sum_{i=1}^n \frac{\hat{\varepsilon}_i^2}{1 - h_{ii}} \mathbf{x}_i \mathbf{x}_i' \right) (\mathbb{X}'\mathbb{X})^{-1}$$

- Leverage: $h_{ii} = \mathbf{x}_i (\mathbb{X}'\mathbb{X})^{-1} \mathbf{x}_i'$
- In this setting: $h_{ii} = n_1^{-1}$ if $D_i = 1$ and $h_{ii} = n_0^{-1}$ if $D_i = 0$

Linear Regression and Causality

- Regression: **conditional expectation function** of Y given $\mathbf{X}$

$$\mathbb{E}(Y|\mathbf{X}) = f(\mathbf{X}) = \beta^T \mathbf{X}$$

- Q: When can we interpret coefficients as causal effects?
- Causal model as structural equation model:

$$Y_i(d) = \alpha + \tau d + \varepsilon_i \quad \text{for } d = 0, 1, \text{ where } \mathbb{E}(\varepsilon_i) = 0$$

1. No interference between units
 2. $\mathbb{E}(Y_i(0)) = \alpha$
 3. $Y_i(1) - Y_i(0) = \tau$ for all $i \rightsquigarrow$ **constant unit-level causal effect**
- Heterogeneous treatment effect model:

$$Y_i(d) = \alpha + \tau_i d + \varepsilon_i = \alpha + \tau d + \underbrace{(\tau_i - \tau)d}_{=\varepsilon_i(d)} + \varepsilon_i$$

where $\mathbb{E}(\varepsilon_i) = 0$ and $\tau = \mathbb{E}(\tau_i) = \mathbb{E}(Y_i(1) - Y_i(0))$

- $\mathbb{E}(\varepsilon_i(d)) = 0$ for $d = 0, 1$
- $\alpha = \mathbb{E}(Y_i(0))$

Cost of Clustering

- Standard variance under **individual assignment**:

$$\mathbb{V}[\hat{\tau}_{\text{diff}}] = \frac{\mathbb{V}[Y_{ik}(1)]}{mK_1} + \frac{\mathbb{V}[Y_{ik}(0)]}{mK_0}$$

- Under clustering, the variance of the cluster average inflates:

$$\frac{\mathbb{V}[\bar{Y}_k(1)]}{K_1} = \frac{\mathbb{V}[Y_{ik}(1)]}{mK_1} (1 + (m-1)\rho_1)$$

- Intraclass correlation coefficient** (ICC):

$$\rho_1 = \frac{1}{m(m-1)} \sum_{i=1}^m \sum_{j \neq i} \mathbb{E} \left\{ (Y_{ik}(1) - \bar{Y}(1))(Y_{jk}(1) - \bar{Y}(1)) \right\}$$

- ρ measures how similar units are within clusters.
 - Usually cluster is less efficient because $\rho > 0$.
 - More similarity $\rightsquigarrow$ each unit provides redundant information $\rightsquigarrow$ less efficiency under clustering.