

Building a Spectroscopy Apparatus Around the FP4209 InstaTune

James Usher

DEPARTMENT OF PHYSICS AND ASTRONOMY, BATES COLLEGE, LEWISTON, ME 04240

Building a Spectroscopy Apparatus Around the FP4209 InstaTune

A Senior Thesis

Presented to the Department of Physics and Astronomy

Bates College

in partial fulfillment of the requirements for the

Degree of Bachelor of Arts

by

James Usher

Lewiston, Maine

April 1, 2026

Contents

Acknowledgments

Mom, Dad, your roommate, maybe even your thesis advisor – give credit where credit’s due! For my part, I’d like to thank the Math department for making such a nice L^AT_EX template that we could use as a base for our Physics template.

Introduction

Methane and Climate Sensing

Climate Sensing

Need for model mixing ratios and Lab spectroscopy examples of uses of lab spectroscopy in history and importance in climate policy/action

In addressing climate change, we must frequently take measurements of greenhouse gasses in Earth's atmosphere. To measure the amount of methane in the atmosphere, we must first have a model of the intensity of the absorption spectrum of methane at different temperatures and pressures. Uncertainties in these models, particularly in lineshape functions, propagate to uncertainties in greenhouse gas measurements.

Problems in Modeling Methane's Behavior

Spectroscopic models of greenhouse gasses used in measurement of the atmosphere typically neglect accounting for collisional effects that lead to broadening and narrowing effects of the line shape of the spectra produced by greenhouse gasses.

Ro-Vibrational Transitions and HITRAN Data

Contribution Made by this Work

We take our measurements with a novel digitally controlled IR laser spectrometer created using the FP4209 InstaTune laser module with the intent to characterize the effects of pressure on the line shape function used in modeling the predicted absorption spectrum of methane. The FP4209 scans over a IR wavelength range of 1627.5000 – 1672.4955 nm, which corresponds very well with the ro-vibrational absorption spectrum of methane. Interpolating DAC values and scanning through the range of wavelengths produced by this laser allows us to construct a quasi-continuous profile of the spectrum of the intensity of light absorbed by the methane gas.

This work allows us to take new measurements of the properties of methane gas in a controlled environment, to better understand the spectra we observe in the atmosphere, and to more precisely describe the behavior and amount of methane in our atmosphere.

CHAPTER 1

Theory

Laser Spectroscopy

You can write text in L^AT_EX as you would in any other word processing or typesetting software. You can incorporate *different varieties of typeface*. What's really exciting about writing text in L^AT_EX is that it's very easy to incorporate weird symbols, like α or $\otimes$ or ∞ into your text. Note that the weird symbols appear in between \$ signs – this means that they occur in *math mode*.

You create a new paragraph just by putting in a blank line. Leaving extra space or extra lines in between paragraphs won't change their spacing at all. L^AT_EX will automatically indent your paragraphs, so you don't have to worry about indenting. You can, however prevent it from indenting your paragraphs if you'd like. But that's getting a little ahead of ourselves.

Laser Mechanics

It is beautifully easy to write equations, whether they occur inline with the text $\Psi(x, t) = Ae^{i(kx - \omega t)}$ or as a separate line

$$(1.1) \quad \Psi(x, t) = \sum_{n=1}^{\infty} c_n \psi_n(x) e^{-iE_n t/\hbar}.$$

To continue a paragraph after an equation, make sure that there's no blank line between the equation and your next line of text. Equations that are in a separate line will be numbered, and it's easy to refer to Eq. ?? by labelling your equations

$$(1.2) \quad i\hbar \frac{\partial}{\partial t} \Psi(x, t) = \hat{H} \Psi(x, t).$$

Are you getting tired of writing `\begin{eqnarray}` and `\end{eqnarray}` yet?¹ You can simplify your life by defining abbreviated symbols for commands:

$$(1.3) \quad \langle x | \Psi(t) \rangle = \Psi(x, t).$$

For simplicity, I'd suggest putting all of your new commands in a separate include file.

When writing a derivation with multiple steps, it is often useful to use an equation array

$$\begin{aligned} \int_0^{2\pi} \sin^2 x dx &= \int_0^{2\pi} \frac{1}{2} (1 - \cos 2x) dx \\ &= \frac{1}{2} \left(x - \frac{\sin 2x}{2} \right) \Big|_0^{2\pi} \\ &= \pi. \end{aligned}$$

¹Notice how I made a } by putting a backslash in front of the symbol. The curly brace has special meaning in L^AT_EX, so if you want to make one, you have to let the program know that you want the symbol.

Note that you can suppress equation numbering, which is often desirable in derivations!

The syntax for equation arrays is similar to the syntax for arrays within equations. For example,

(1.4)

$$\hat{H} = \begin{pmatrix} \Delta & \Omega & 0 \\ \Omega^* & 0 & \kappa \\ 0 & \kappa^* & -\Delta \end{pmatrix}$$

There are many more complicated things you can do with equations, but this should be enough to get started!

Distributed Bragg Reflector (DBR) Lasers

If you want to make a table, you'll need to specify the justification for each of the columns; `TeX` will automatically calculate the column widths for you.

Name	Age	Town	Time
Joe Schmoe	23	Rumford	1:04
Mary Q. Contrary	44	Yarmouth	1:12
Baxter Boulevard	32	Portland	1:42
Lass Plaiice	60	Lewiston	1:57

TABLE 1. Not too many people showed up for the first, and probably last, annual Androscoggin swimming race.

- You might also, for example, want to
- (1) make lists

(2) number your lists

• use bullet points

• find better items to put in your bullet points




FIGURE 1.1. This is a very bad figure.

test citation [11]

test citation [9]


test citation [10]

test citation [4]
test citation [7]
test citation [1]
test citation [8]
test citation [5]
test citation [6]
test citation [3]
citation [2]

CHAPTER 2

Method

Laser Functionality Program Sequencing



Communication Protocol
Data Collection

W00000

CHAPTER 3

Analysis

DAC LUT Discontinuities

Laser Precision

Spectroscopic Measurements

WOOOOO

Appendix A

Long derivations, marginally relevant details, and the like can be relegated to your appendix, which unlike the mammalian counterpart will never require removal.

Bibliography

- [1] Rami Arieli. *Rami Arieli: "The Laser Adventure,"*. URL: <https://perg.phys.ksu.edu/vqm/laserweb/ch-5/f5s1p5.htm>.
- [2] T. Delahaye et al. "Measurement and Modeling of Air-Broadened Methane Absorption in the MERLIN Spectral Region at Low Temperatures". In: *Journal of Geophysical Research: Atmospheres* 124.6 (2019), pp. 3556–3564. DOI: <https://doi.org/10.1029/2018JD028917>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2018JD028917>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD028917>.
- [3] T. Delahaye et al. "Precise methane absorption measurements in the 1.64μm spectral region for the MERLIN mission". In: *Journal of Geophysical Research: Atmospheres* 121.12 (2016), pp. 7360–7370. DOI: <https://doi.org/10.1002/2016JD025024>. eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2016JD025024>. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016JD025024>.
- [4] S E Gustafsson, M N Khan, and J Van Eijk. "Wavelength scanning and mirror alignment of a new multipass Fabry-Perot interferometer". In: *Journal of Physics E: Scientific Instruments* 12.11 (Nov. 1979), p. 1100. DOI: 10.1088/0022-3735/12/11/020. URL: <https://doi.org/10.1088/0022-3735/12/11/020>.
- [5] Ronald K. Hanson, R. Mitchell Spearrin, and Christopher S. Goldenstein. *Spectroscopy and Optical Diagnostics for Gases*. en. Cham: Springer International Publishing, 2016. ISBN: 9783319232515. DOI: 10.1007/978-3-319-23252-2. URL: <https://link.springer.com/10.1007/978-3-319-23252-2>.
- [6] Vilas Jangale. "Real-time characterization of natural gas and opportunity fuels by NIR absorption spectroscopy". PhD thesis. Nov. 2012. DOI: 10.13140/RG.2.2.11453.67045.
- [7] Štěpán Kunc. "Study of the Magnetically Induced QED Birefringence of the Vacuum in experiment OSQAR". PhD thesis. Apr. 2018.
- [8] D.A. Long et al. "The effects of advanced spectral line shapes on atmospheric carbon dioxide retrievals". In: *Journal of Quantitative Spectroscopy and Radiative Transfer* 291 (2022), p. 108324. ISSN: 0022-4073. DOI: <https://doi.org/10.1016/j.jqsrt.2022.108324>. URL: <https://www.sciencedirect.com/science/article/pii/S002240732200259X>.
- [9] Javad Shemshad, Saïed Aminossadati, and M. Kizil. "A review of developments in near infrared methane detection based on tunable diode laser". In: *Sensors and Actuators B: Chemical* 171–172 (Aug. 2012), pp. 77–92. DOI: 10.1016/j.snb.2012.06.018.
- [10] "The Fabry Perot Resonator". In: *Laser Resonators and Beam Propagation: Fundamentals, Advanced Concepts and Applications*. New York, NY: Springer New York, 2005, pp. 189–215. ISBN: 978-0-387-25110-3. DOI: 10.1007/0-387-25110-3_5. URL: https://doi.org/10.1007/0-387-25110-3_5.
- [11] Magdalena Zatorska, Grzegorz Gomółka, and Michał Nikodem. "Near-infrared quartz-enhanced photoacoustic spectroscopy system for ppb-level methane detection". In: *Optics Continuum* 2 (Jan. 2023), pp. 266–273. DOI: 10.1364/OPTCON.477782.