

Variational Bayes Applied to Multimodal Distributions in B Physics

S. Jahn¹ F. Beaujean² C. Bobeth³


¹Exzellenzcluster Universe
Technische Universität München

²Exzellenzcluster Universe
Ludwig-Maximilians-Universität München

³Institute for Advanced Study
Technische Universität München

March 12
DPG Wuppertal 2015

effective Lagrangian for $b \rightarrow s\ell^+\ell^-$ (SM):


$$\mathcal{L}_{int} = \frac{4G_F}{\sqrt{2}} \frac{\alpha_e}{4\pi} V_{tb} V_{ts}^* \sum_i C_i \mathcal{O}_i + \dots + \text{h.c.}$$

$$\mathcal{O}_9 = [\bar{s}\gamma_\mu P_L \quad b] [\bar{\ell}\gamma^\mu \ell] \qquad \mathcal{O}_{10} = [\bar{s}\gamma_\mu P_L \quad b] [\bar{\ell}\gamma^\mu \gamma_5 \ell]$$

effective Lagrangian for $b \rightarrow s \ell^+ \ell^-$ (**beyond**-SM):

$$\mathcal{L}_{int} = \frac{4G_F}{\sqrt{2}} \frac{\alpha_e}{4\pi} V_{tb} V_{ts}^* \sum_i \mathcal{C}_i \mathcal{O}_i + \dots + \text{h.c.}$$

$$\mathcal{O}_9^{(\prime)} = [\bar{s} \gamma_\mu P_{L(R)} b] [\bar{\ell} \gamma^\mu \ell]$$

$$\mathcal{O}_{10}^{(\prime)} = [\bar{s} \gamma_\mu P_{L(R)} b] [\bar{\ell} \gamma^\mu \gamma_5 \ell]$$



$$\mathcal{O}_S^{(\prime)} = [\bar{s} P_{R(L)} b] [\bar{\ell} \ell]$$

$$\mathcal{O}_P^{(\prime)} = [\bar{s} P_{R(L)} b] [\bar{\ell} \gamma_5 \ell]$$

$$\mathcal{O}_T = [\bar{s} \sigma_{\mu\nu} b] [\bar{\ell} \sigma^{\mu\nu} \ell]$$

$$\mathcal{O}_{T5} = [\bar{s} \sigma_{\mu\nu} b] [\bar{\ell} \sigma^{\mu\nu} \gamma_5 \ell]$$

- $B \rightarrow K\mu^+\mu^-$ angular distribution

- $B_s \rightarrow \mu^+\mu^-$ branching fraction

- $B \rightarrow K\mu^+\mu^-$ angular distribution

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta} = \frac{3}{4}(1 - F_H) \sin^2\theta + \frac{1}{2}F_H + A_{FB} \cos\theta$$

- $B_s \rightarrow \mu^+\mu^-$ branching fraction

- $B \rightarrow K \mu^+ \mu^-$ angular distribution

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta} = \frac{3}{4}(1 - F_H) \sin^2\theta + \frac{1}{2}F_H + A_{FB} \cos\theta$$

$$A_{FB} \propto \text{Re}[(C_P + C'_P)C_{T5}^* + (C_S + C'_S)C_T^* + \mathcal{O}(m_\ell/\sqrt{q^2})] / \Gamma$$

- $B_s \rightarrow \mu^+ \mu^-$ branching fraction

- $B \rightarrow K \mu^+ \mu^-$ angular distribution

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta} = \frac{3}{4}(1 - F_H) \sin^2\theta + \frac{1}{2}F_H + A_{FB} \cos\theta$$

$$A_{FB} \propto \text{Re}[(C_P + C'_P) C_{T5}^* + (C_S + C'_S) C_T^* + \mathcal{O}(m_\ell/\sqrt{q^2})] / \Gamma$$
$$F_H \propto \left[\dots (|C_T|^2 + |C_{T5}|^2) + \dots (|C_S + C'_S|^2 + |C_P + C'_P|^2) + \mathcal{O}(m_\ell/\sqrt{q^2}) \right] / \Gamma$$

- $B_s \rightarrow \mu^+ \mu^-$ branching fraction

- $B \rightarrow K \mu^+ \mu^-$ angular distribution

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta} = \frac{3}{4}(1 - F_H) \sin^2\theta + \frac{1}{2}F_H + A_{FB} \cos\theta$$


$$A_{FB} \propto \text{Re}[(C_P + C'_P) C_{T5}^* + (C_S + C'_S) C_T^* + \mathcal{O}(m_\ell/\sqrt{q^2})] / \Gamma$$


$$F_H \propto \left[\dots (|C_T|^2 + |C_{T5}|^2) + \dots (|C_S + C'_S|^2 + |C_P + C'_P|^2) + \mathcal{O}(m_\ell/\sqrt{q^2}) \right] / \Gamma$$

$$F_H^{\text{SM}} = \mathcal{O}(m_\ell^2/q^2) \quad A_{FB}^{\text{SM}} = 0$$

- $B_s \rightarrow \mu^+ \mu^-$ branching fraction

- $B \rightarrow K \mu^+ \mu^-$ angular distribution

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta} = \frac{3}{4}(1 - F_H) \sin^2\theta + \frac{1}{2}F_H + A_{FB} \cos\theta$$




$$A_{FB} \propto \text{Re}[(C_P + C'_P)C_{T5}^* + (C_S + C'_S)C_T^* + \mathcal{O}(m_\ell/\sqrt{q^2})] / \Gamma$$

$$F_H \propto \left[\dots (|C_T|^2 + |C_{T5}|^2) + \dots (|C_S + C'_S|^2 + |C_P + C'_P|^2) + \mathcal{O}(m_\ell/\sqrt{q^2}) \right] / \Gamma$$

$$F_H^{\text{SM}} = \mathcal{O}(m_\ell^2/q^2) \quad A_{FB}^{\text{SM}} = 0$$

- $B_s \rightarrow \mu^+ \mu^-$ branching fraction

$$\mathcal{B}(B_s \rightarrow \mu^+ \mu^-) \propto |C_S - C'_S|^2 + |(C_P - C'_P) + \frac{2m_\ell}{M_{B_s}}(C_{10} - C'_{10})|^2$$

we want:

$$P(\boldsymbol{\theta}|\mathcal{D}, M) \stackrel{\text{Bayes}}{\propto} P(\mathcal{D}|\boldsymbol{\theta}, M)P(\boldsymbol{\theta}|M)$$

we want:



$$P(\boldsymbol{\theta}|\mathcal{D}, M) \stackrel{\text{Bayes}}{\propto} P(\mathcal{D}|\boldsymbol{\theta}, M)P(\boldsymbol{\theta}|M)$$

split theory and experiment - *observables* **O**:

$$P(\mathcal{D}|\boldsymbol{\theta}, M) = P(\mathcal{D}|\mathbf{O}(\boldsymbol{\theta}, M))$$

we want:

$$P(\boldsymbol{\theta}|\mathcal{D}, \mathbf{M}) \stackrel{\text{Bayes}}{\propto} P(\mathcal{D}|\boldsymbol{\theta}, \mathbf{M})P(\boldsymbol{\theta}|\mathbf{M})$$

split theory and experiment - *observables* \mathbf{O} :

$$P(\mathcal{D}|\boldsymbol{\theta}, \mathbf{M}) = P(\mathcal{D}|\mathbf{O}(\boldsymbol{\theta}, \mathbf{M}))$$

theory

experiment

we want:

$$P(\boldsymbol{\theta}|\mathcal{D}, M) \stackrel{\text{Bayes}}{\propto} P(\mathcal{D}|\boldsymbol{\theta}, M)P(\boldsymbol{\theta}|M)$$

split theory and experiment - *observables* \mathbf{O} :

$$P(\mathcal{D}|\boldsymbol{\theta}, M) = P(\mathcal{D}|\mathbf{O}(\boldsymbol{\theta}, M))$$

theory

calculate observables
 $\mathbf{O}(\boldsymbol{\theta}, M)$

experiment



we want:

$$P(\theta|\mathcal{D}, M) \stackrel{\text{Bayes}}{\propto} P(\mathcal{D}|\theta, M)P(\theta|M)$$

split theory and experiment - *observables* \mathbf{O} :

$$P(\mathcal{D}|\theta, M) = P(\mathcal{D}|\mathbf{O}(\theta, M))$$

theory

calculate observables
 $\mathbf{O}(\theta, M)$



experiment

measure observables
 $P(\mathcal{D}|\mathbf{O})$



we want:

$$P(\theta|\mathcal{D}, M) \stackrel{\text{Bayes}}{\propto} P(\mathcal{D}|\theta, M)P(\theta|M)$$

split theory and experiment - *observables* \mathbf{O} :

$$P(\mathcal{D}|\theta, M) = P(\mathcal{D}|\mathbf{O}(\theta, M), \text{ ~~\theta, M~~ })$$

assumption

theory

calculate observables
 $\mathbf{O}(\theta, M)$



experiment

measure observables
 $P(\mathcal{D}|\mathbf{O}, \text{ ~~\theta, M~~ })$



- $B \rightarrow K\mu^+\mu^-$: \mathcal{B} , A_{FB} , F_H
 - LHCb 2014 (arXiv:1403.8044 , arXiv:1403.8045)
 - CDF 2012
(http://www-cdf.fnal.gov/physics/new/bottom/120628.blessed-b2smumu_96)
- $B_s \rightarrow \mu^+\mu^-$: \mathcal{B}
 - LHCb+CMS 2014 (arXiv:1411.4413)
- $B \rightarrow K^*\mu^+\mu^-$: \mathcal{B}
 - LHCb 2013 (arXiv:1304.6325)
 - CMS 2013 (arXiv:1308.3409)
 - CDF 2012
(http://www-cdf.fnal.gov/physics/new/bottom/120628.blessed-b2smumu_96)

scan parameters

- Wilson coefficients $\mathcal{C}_{10}^{(\prime)}, \mathcal{C}_S^{(\prime)}, \mathcal{C}_P^{(\prime)}, \mathcal{C}_T$, and \mathcal{C}_{T5}

scan parameters

- Wilson coefficients $C_{10}^{(\prime)}, C_S^{(\prime)}, C_P^{(\prime)}, C_T$, and C_{T5}

nuisance parameters

- CKM matrix (4 parameters)
- charm and bottom quark mass (2 parameters)
- form factors
 - $B \rightarrow K$ (5 parameters)
 - $B \rightarrow K^*$ (6 parameters)
- B_s decay constant f_{B_s}
- subleading corrections (11 parameters)

scan parameters

- Wilson coefficients $\mathcal{C}_{10}^{(\prime)}, \mathcal{C}_S^{(\prime)}, \mathcal{C}_P^{(\prime)}, \mathcal{C}_T$, and \mathcal{C}_{T5}

nuisance parameters

- CKM matrix (4 parameters)
- charm and bottom quark mass (2 parameters)
- form factors
 - $B \rightarrow K$ (5 parameters)
 - $B \rightarrow K^*$ (6 parameters)
- B_s decay constant f_{B_s}
- subleading corrections (11 parameters)







theory calculation $\mathcal{O}(\theta, \mathbf{M})$:

open-source implementation: EOS-package

<http://project.het.physik.tu-dortmund.de/eos/>





Joint fit of $\mathcal{C}_{10}^{(j)}$, $\mathcal{C}_S^{(j)}$, $\mathcal{C}_P^{(j)}$, \mathcal{C}_T , and \mathcal{C}_{T5}







Joint fit of $\mathcal{C}_{10}^{(I)}$, $\mathcal{C}_S^{(I)}$, $\mathcal{C}_P^{(I)}$, \mathcal{C}_T , and \mathcal{C}_{T5}



- first *simultaneous* fit







Joint fit of $\mathcal{C}_{10}^{(\prime)}$, $\mathcal{C}_S^{(\prime)}$, $\mathcal{C}_P^{(\prime)}$, \mathcal{C}_T , and \mathcal{C}_{T5}



- first *simultaneous* fit
- interference $\mathcal{C}_{10}^{(\prime)} \leftrightarrow \mathcal{C}_{S,P}^{(\prime)}$ in $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$

Joint fit of $\mathcal{C}_{10}^{(\prime)}, \mathcal{C}_S^{(\prime)}, \mathcal{C}_P^{(\prime)}, \mathcal{C}_T$, and \mathcal{C}_{T5}



- first *simultaneous* fit

- interference $\mathcal{C}_{10}^{(\prime)} \leftrightarrow \mathcal{C}_{S,P}^{(\prime)}$ in $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$

⇒ larger uncertainty

than obtained for fixed $\mathcal{C}_{10}^{(\prime)} = \mathcal{C}_{10}^{(\prime)\text{SM}}$


arXiv:1205.5811,

arXiv:1206.0273,


arXiv:1407.7044




Nuisance parameters



Nuisance parameters




Difficulties




Difficulties

- curse of dimensionality




Difficulties

- curse of dimensionality
- multimodality



Difficulties


- curse of dimensionality
- multimodality
- degeneracies




Difficulties

- curse of dimensionality
- multimodality
- degeneracies

no standard
algorithm so
far



Sampling algorithm



<https://pypi.python.org/pypi/pypmc>

- model-independent search for new physics

- model-independent search for new physics
- **simultaneous** fit of $\mathcal{C}_S^{(\prime)}$, $\mathcal{C}_P^{(\prime)}$, \mathcal{C}_T , \mathcal{C}_{T5} , and $\mathcal{C}_{10}^{(\prime)}$
 - ⇒ updated, data-driven constraints

- model-independent search for new physics
- **simultaneous** fit of $\mathcal{C}_S^{(\prime)}, \mathcal{C}_P^{(\prime)}, \mathcal{C}_T, \mathcal{C}_{T5}$, and $\mathcal{C}_{10}^{(\prime)}$
 - ⇒ updated, data-driven constraints
- no significant deviation from the SM

- model-independent search for new physics
- **simultaneous** fit of $\mathcal{C}_S^{(\prime)}$, $\mathcal{C}_P^{(\prime)}$, \mathcal{C}_T , \mathcal{C}_{T5} , and $\mathcal{C}_{10}^{(\prime)}$
 - ⇒ updated, data-driven constraints
- no significant deviation from the SM
- need better theoretical control (form factors, subleading QCDF)

- model-independent search for new physics
- **simultaneous** fit of $\mathcal{C}_S^{(\prime)}$, $\mathcal{C}_P^{(\prime)}$, \mathcal{C}_T , \mathcal{C}_{T5} , and $\mathcal{C}_{10}^{(\prime)}$
 - ⇒ updated, data-driven constraints
- no significant deviation from the SM
- need better theoretical control (form factors, subleading QCDF)
- sampling algorithm to handle multimodality in $\text{dim} = \mathcal{O}(40)$

$B \rightarrow K^* \mu^+ \mu^-$ angular analysis

$B \rightarrow K^* \mu^+ \mu^-$ angular analysis

- $J_{6c}, (J_{1s} - 3J_{2s}), (J_{1c} - J_{2c})$
SM-contributions suppressed $\mathcal{O}\left(\frac{m_\ell}{\sqrt{q^2}}\right)$

$B \rightarrow K^* \mu^+ \mu^-$ angular analysis

- $J_{6c}, (J_{1s} - 3J_{2s}), (J_{1c} - J_{2c})$
SM-contributions suppressed $\mathcal{O}\left(\frac{m_\ell}{\sqrt{q^2}}\right)$

- LHCb currently assumes $\mathcal{C}_{S,P,T,T5}^{(\prime)} = 0 \Rightarrow P(\mathcal{D}|\mathbf{O}, \overbrace{\theta, M}^{\text{invalid}})$

<http://www.physi.uni-heidelberg.de/Forschung/he/LHCb/documents/WorkshopNeckarzMar14/NeckarzimmernKstmumuExp.pdf>

$B \rightarrow K^* \mu^+ \mu^-$ angular analysis

- $J_{6c}, (J_{1s} - 3J_{2s}), (J_{1c} - J_{2c})$
 SM-contributions suppressed $\mathcal{O}\left(\frac{m_\ell}{\sqrt{q^2}}\right)$
- LHCb currently assumes $\mathcal{C}_{S,P,T,T5}^{(\prime)} = 0 \Rightarrow P(\mathcal{D}|\mathbf{O}, \overbrace{\theta, \mathbf{M}}^{\text{invalid}})$
<http://www.physi.uni-heidelberg.de/Forschung/he/LHCb/documents/WorkshopNeckarzMar14/NeckarzimmerKstmumuExp.pdf>
- paper work in progress