

MATH20212 Cheat Sheet

1 Rings

A **ring** is a set R and two binary operations, written $+$ and \times , on R which satisfies the following conditions:

- (R1) $\langle R, + \rangle$ is an abelian group with identity 0
- (R2) \times is associative
- (R3) \times is distributive over $+$
- (R4) there exists an element $1 \in R$, different from 0, that is an identity for \times

Let R be a ring and $S \subseteq R$. Then S is a **subring** of R if it is a ring in its own right with respect to the same addition and multiplication as in R and S contains 1_R .

Subring Test: Let R be a ring and $S \subseteq R$, then S is a subring of R , iff:

- (i) $1 \in S$
- (ii) $r + s, r \times s \in S$, for all $r, s \in S$
- (iii) $-r \in S$ for all $r \in S$

Let R be a ring. The **ring of polynomials** $R[X]$ in the indeterminate X is defined as follows:

Elements: formal linear combinations of the form $\sum_{i \geq 0} a_i X^i$ with $a_i \in R$ for $i = 0, 1, \dots$

Equality: $\sum_{i \geq 0} a_i X^i = \sum_{i \geq 0} b_i X^i \iff a_i = b_i$ for all $i \geq 0$

Addition: $\sum_{i \geq 0} a_i X^i + \sum_{i \geq 0} b_i X^i = \sum_{i \geq 0} (a_i + b_i) X^i$

Multiplication: $(\sum_{i \geq 0} a_i X^i)(\sum_{i \geq 0} b_i X^i) = \sum_{k \geq 0} (\sum_{i+j=k} a_i b_j) X^k$

Zero element is $\sum_{i \geq 0} 0 X^i = 0$ and the **one** is $1X^0 + \sum_{i \geq 1} 0 X^i = 1$

For a polynomial $f = \sum_{i \geq 0} a_i X^i$, we define the **degree** of f , denoted $\deg(f)$, to be the largest i such that $a_i \neq 0$ and we let $\deg(f) = -\infty$ if $f = 0$.

2 Integral Domains and Fields

The **characteristic**, $\text{char}(R)$, of a ring R is the least positive integer n such that $n \cdot 1 = 0$. If there is no such n , then the characteristic of R is defined to be 0.

A non-zero element $r \in R$ is a **zero-divisor** if there is a non-zero element $s \in R$ with $rs = 0$ or $sr = 0$.

The ring R is a **domain** if, for all $r, s \in R$, $rs = 0 \implies r = 0$ or $s = 0$, so a domain is a ring with **no** zero-divisors. A commutative domain is called an **integral domain**.

A **division ring** is a ring in which every non-zero element has a right inverse and a left inverse. In this case, these inverses are the same. We write r^{-1} for this **inverse** of r and say that r is **invertible** or that r is a **unit**. A **field** is a commutative division ring.

An element r of a ring R is **nilpotent** if there is some integer $n \geq 1$ with $r^n = 0$ and the least such n is the **index of nilpotence** of r . An element $r \in R$ is **idempotent** if $r^2 = r$ - and 0 and 1 are idempotent in any ring.