

# MATH20122 Cheat Sheet

## 1 Definitions and Examples

**metric space:** a **metric space**  $(X, d)$  consists of a non-empty set  $X$  and a non-negative real valued **metric** (*distance function*)  $d : X \times X \rightarrow \mathbb{R}^{\geq}$  which satisfies the following axioms:

- (i)  $d(x, y) = 0 \iff x = y$  for all  $x, y \in X$
- (ii)  $d(x, y) = d(y, x)$  for all  $x, y \in X$
- (iii)  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in X$  (the *triangle inequality*)

**subspace:** given any subset  $W \subseteq X$ , the restriction of  $d$  to  $W$  determines the subspace  $(W, d := d|_W)$  of  $(X, d)$

**open ball:** for any metric space  $(X, d)$ , the open ball of radius  $r > 0$  around any  $x \in X$  is  $B_r(x) := \{y : d(y, x) < r\}$

**closed ball:** for any metric space  $(X, d)$ , the open ball of radius  $r > 0$  around any  $x \in X$  is  $\bar{B}_r(x) := \{y : d(y, x) \leq r\}$

---

**euclidean n-space:**  $(\mathbb{R}^n, d_2)$  consists of all real  $n$ -dimensional vectors  $x = (x_1, \dots, x_n)$ , equipped with the **euclidean metric**  $d_2(x, y) = ((x_1 - y_1)^2 + \dots + (x_n - y_n)^2)^{1/2}$  where the positive square root is understood

**taxicab metric:**  $d_1$  is given on  $\mathbb{R}^n$  is given by  $d_1(x, y) = |x_1 - y_1| + \dots + |x_n - y_n|$

$$\text{discrete metric : } d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{otherwise} \end{cases}$$

**isometry:** for any two metric spaces  $(X, d_X)$  and  $(Y, d_Y)$ , a bijection  $f : X \rightarrow Y$  is an isometry whenever  $d_X(x, y) = d_Y(f(x), f(y))$  for all  $x, y \in X$

**standard metric:**  $d_{\mathbb{C}}$  on the complex numbers  $\mathbb{C}$  is given by  $d_{\mathbb{C}}(z, z') = |z - z'|$

---

**graph:**  $\Gamma := (V, E)$  consists of a set  $V$  of **vertices** and a set  $E$  of **edges**

**path:** a **path** in  $\Gamma$  from  $u$  to  $w$  is a finite sequence of edges  $\pi(u, w) = (uv_1, v_1v_2, \dots, v_{n-2}v_{n-1}, v_{n-1}w)$  with length  $n$

**path connected:** a graph is path connected whenever there is a path joining *any* pair of vertices

**edge metric:**  $e$  on the vertex set  $V$  of a path connected graph is defined by  $e(u, w) = \min_{\pi(u, w)} l(\pi(u, w))$

**alphabet:** a finite set  $A$  of **letters** and a finite sequence of letters is a **word** in  $A$ . the vertex set  $W$  of the associated **word graph**  $\Gamma(A)$  consists of all possible words in  $A$ . word  $w_1$  and  $w_2$  are joined by an edge iff they differ by one of (i) inserting or deleting a letter (ii) swapping two adjacent letters (iii) replacing one letter with another

**word metric:**  $d_w$  on  $W$  is the edge metric on the associated word graph

---

**binary sequences:**  $X = \{0, 1\}^{\infty}$  is the set of all infinite binary sequences  $x = x_0x_1\dots$  where  $x_n = 0$  or  $1$  for all  $n \geq 0$

$$d_{\min}(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1/2^n & \text{if } n = \min\{m : x_m \neq y_m\} \end{cases}$$
$$d^*(x, y) = \sum_{j=0}^{\infty} \frac{|x_j - y_j|}{2^j}$$

---

**bounded:** a real valued function  $f$  on a closed interval  $[a, b] \subset \mathbb{R}$  is bounded whenever  $\exists K, |f(x)| \leq K, \forall x \in [a, b]$

let  $X$  denote the set of all bounded  $f : [a, b] \rightarrow \mathbb{R}$  then

$$d_{\sup}(f, g) = \sup_{x \in [a, b]} |f(x) - g(x)| \text{ with } (X, d_{\sup}) \text{ denoted by } \mathcal{B}[a, b]$$

let  $Y$  denote the set of all continuous  $f : [a, b] \rightarrow \mathbb{R}$  then

$$d_1(f, g) = \int_a^b |f(t) - g(t)| dt \text{ with } (Y, d_1) \text{ denoted by } \mathcal{L}_1[0, 1]$$

let  $X$  denote the set of all closed intervals  $[a, b]$  in the euclidean line

**interval metric:**  $d_H$  on  $X$  is given by  $d_H([a, b], [r, s]) = \max\{|r - a|, |s - b|\}$

$$d_\infty((x_1, x_2), (y_1, y_2)) = \max\{|x_1 - y_1|, |x_2 - y_2|\}$$

let  $X$  be the set of infinite sequences  $(a_i : i \geq 0)$  of reals, such that  $\sum_i a_i$  is absolutely convergent

$$d_1((a_i), (b_i)) = \sum_{i \geq 0} |a_i - b_i|$$

**cartesian product:** of two metric spaces  $(X, d)$  and  $(X', d')$  is the set  $X \times X'$  with one of the metrics

1.  $d_a((x, x'), (y, y')) = d(x, y) + d'(x', y')$
2.  $d_b((x, x'), (y, y')) = (d(x, y)^2 + d'(x', y')^2)^{1/2}$
3.  $d_c((x, x'), (y, y')) = \max\{d(x, y), d'(x', y')\}$

**lipschitz equivalent:** two metrics  $d$  and  $e$  on a given set  $X$  are lipschitz equivalent whenever there exists positive constants  $h, k \in \mathbb{R}$  such that  $he(x, y) \leq d(x, y) \leq ke(x, y)$  for every  $x, y \in X$

**theorem:** the metrics  $d_a, d_b, d_c$  on  $X \times X'$  are lipschitz equivalent

## 2 Open and Closed Sets

in this section, let  $(X, d)$  be a any metric space and  $U \subseteq X$

**interior point:**  $u \in U$  such that  $\exists \varepsilon > 0, B_\varepsilon(u) \subseteq U$ , the **interior** of  $U$  is the subset  $U^\circ \subseteq U$  of all interior points and if  $U^\circ = U$ , then  $U$  is **open** in  $X$

**proposition:** every open ball  $B_r(x)$  is open in  $X$

**theorem:** given any two subsets  $U, V \subseteq X$ , the following hold:

- $U \subseteq V \implies U^\circ \subseteq V^\circ$
- $U^\circ$  is open in  $X$
- $(U^\circ)^\circ = U^\circ$
- $U^\circ$  is the largest subset of  $U$ , open in  $X$

**theorem:** the sets  $X$  and  $\emptyset$  are open in  $X$ , so are an arbitrary union  $U = \bigcup_{i \in I} U_i$  of open sets  $U_i$ , and a finite intersection  $U' = U'_1 \cap \dots \cap U'_m$  of open sets  $U'_j$

**closure point:**  $x \in X$  is a closure point of  $U \subseteq X$  if  $B_\varepsilon(x) \cap U$  is non-empty for every  $\varepsilon > 0$ , the **closure** of  $U$  is the superset  $\overline{U} \supseteq U$  of all closure points and if  $\overline{U} = U$ , then  $U$  is **closed** in  $X$

**proposition:** a set  $V$  is closed in  $X$  iff its complement  $U := X \setminus V$  is open

**corollary:** every closed ball  $\overline{B}_r(x)$  is closed in  $X$

**partially open ball:** in  $X$  is a set  $P_r(x) := B_r(x) \cup P$ , where  $P$  is a proper subset of  $\{p : d(x, p) = r\}$

**theorem:** given any two subsets  $U, V \subseteq X$ , the following hold:

- $U \subseteq V \implies \overline{U} \subseteq \overline{V}$
- $\overline{V}$  is closed in  $X$
- $\overline{\overline{V}} = \overline{V}$
- $\overline{V}$  is the smallest set containing  $V$ , closed in  $X$

**theorem:** the sets  $X$  and  $\emptyset$  are closed in  $X$ , so are an arbitrary intersection  $V = \bigcap_{i \in I} V_i$  of closed sets  $V_i$  and a finite union  $V' = V'_1 \cup \dots \cup V'_m$  of closed sets  $V'_j$

---

**sequence:** a sequence in  $X$  is a function  $s : \mathbb{N} \rightarrow X$  and it is standard practise to write  $s(n)$  as  $x_n$  and display the sequence as  $(x_n : n \geq 1)$

**converges:** a sequence  $(x_n)$  converges to the point  $x \in X$  whenever  $\forall \varepsilon > 0, \exists N \in \mathbb{N}$ , such that  $n \geq N \implies d(x, x_n) < \varepsilon$  and in this situation, we say that  $x$  is the **limit** of  $(x_n)$

**theorem:** in any metric space  $(X, d)$ , the limit of a convergent sequence is unique

**theorem:** suppose that  $Y \subseteq X$  and  $y \in X$ , then  $y$  lies in  $\overline{Y}$  iff there exists a sequence  $(y_n)$  in  $Y$  such that  $y_n \rightarrow y$  as  $n \rightarrow \infty$

**cauchy sequence:** in any metric space  $(X, d)$ , a cauchy sequence  $(x_n)$  satisfies  $\forall \varepsilon > 0, \exists N \in \mathbb{N}$ , such that  $m, n \geq N \implies d(x_n, x_m) < \varepsilon$

**dense:** a subset  $Y$  is dense in  $(X, d)$  whenever  $\overline{Y} = X$

---

**bounded:** a subset  $A$  of the metric space  $(X, d)$  is bounded whenever there exists  $x_0 \in X$  and  $M \in \mathbb{R}$  such that  $d(x, x_0) \leq M$  for every  $x \in A$ . a function  $f : S \rightarrow X$  is bounded whenever its image  $f(S) \subset X$  is a bounded, for any set

**diameter:** the diameter,  $\text{diam}(A)$  of a bounded non-empty set  $A \subseteq X$  is the real number  $\sup\{d(x, y) : x, y \in A\}$

**eucildean (n - 1)-sphere of radius r:**  $\{x : |x| = r\}$

**boundary point:**  $x \in X$  of  $A$  is one for which every open ball  $B_\varepsilon(x)$  meets both  $A$  and  $X \setminus A$ , the **boundary**  $\delta A$  of  $A$  is the set of all such boundary points

**theorem:** any subset  $A$  of  $(X, d)$  satisfies:

- $A \setminus \delta A = \overline{A} \setminus \delta A = A^\circ$
- $\delta A = \delta(X \setminus A)$
- $\delta A$  is closed in  $X$